

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

CANDIDATE NAME					
CENTRE NUMBER			ANDIDATE UMBER		



CHEMISTRY 9701/21

Paper 2 Structured Questions AS Core

May/June 2012

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use			
1			
2			
3			
4			
5			
Total			

This document consists of 11 printed pages and 1 blank page.



Answer **all** the questions in the spaces provided.

For Examiner's Use

1	Oxides are compounds which usually contain oxygen combined with one other element.						
	Oxides are classified as follows.						
		acidic	alkaline	amp	hoteric	basic	
		ng these term third period of t				ne oxides of th	e elements of
	Na ₂ O	MgO	Al_2O_3	SiO ₂	P ₄ O ₁₀	SO ₂	Cl ₂ O ₇
							acidic
							[4]
	(b) Give the names of two elements from sodium to chlorine which form more than one oxide.						nore than one
	(c) Soc	lium reacts with	n water.				
	(i)	Describe, as for with water.	ılly as you can	ı, what you wo	uld see when	a piece of sod	lium is reacted
	(ii)	Write an equa	tion for the rea	action of sodiu	ım with water		
							[4]

(d)	Sul	fur dioxide is present in small, but significant, amounts in the Earth's atmosphere.
	(i)	State one way by which sulfur dioxide enters the atmosphere.
	(ii)	Give the formula of another sulfur compound which is formed in the atmosphere from sulfur dioxide.
	(iii)	What are the environmental consequences of the compound you have identified in $\mbox{(ii)}$?
		[3]
(e)		fur dioxide is used as a food preservative. at property of sulfur dioxide enables it to act in this way?
		[1]
(f)	OC	other sulfur compound which is present in the Earth's atmosphere is carbonyl sulfide, S. The sequence of atoms in the molecule is oxygen-carbon-sulfur and the molecule ot cyclic.
	(i)	Draw a 'dot-and-cross' diagram of the OCS molecule. Show outer electrons only.
	(ii)	Suggest a value for the O-C-S bond angle.
		[2]
		[Total: 15]

Use

For Examiner's 2 Ammonium sulfate, $(NH_4)_2SO_4$, is widely used as a fertiliser.

For Examiner's Use

In order to determine its percentage purity, a sample of ammonium sulfate fertiliser was analysed by reacting a known amount with an excess of NaOH(aq) and then titrating the unreacted NaOH with dilute HC *l.*

(a) Ammonium sulfate reacts with NaOH in a 1:2 ratio. Complete and balance the equation for this reaction.

$$(NH_4)_2SO_4 + 2NaOH \rightarrow \dots NH_3 + \dots + \dots$$

[2]

(b) A 5.00 g sample of a fertiliser containing $(NH_4)_2SO_4$ was warmed with 50.0 cm³ (an excess) of 2.00 mol dm⁻³ NaOH.

When all of the ammonia had been driven off, the solution was cooled.

The remaining NaOH was then titrated with 1.00 mol dm⁻³ HC *l* and 31.2 cm³ were required for neutralisation.

(i) Write a balanced equation for the reaction between NaOH and HC1.

.....

(ii) Calculate the amount, in moles, of HCl in 31.2 cm³ of 1.00 mol dm⁻³ HCl.

(iii) Calculate the amount, in moles, of NaOH in 50.0 cm³ of 2.00 mol dm⁻³ NaOH.

(iv) Use your answers to (i), (ii) and (iii) to calculate the amount, in moles, of NaOH used up in the reaction with $(NH_4)_2SO_4$.

(v)	Use your answer to (iv) and the equation in (a) to calculate the amount, in moles, of $(NH_4)_2SO_4$ that reacted with NaOH.	For Examiner's Use
(vi)	Use your answer to (v) to calculate the mass of (NH $_4$) $_2$ SO $_4$ that reacted with NaOH.	
(vii)	Hence, calculate the percentage purity of the ammonium sulfate fertiliser.	
	[7]	
	[Total: 9]	

[Turn over © UCLES 2012 9701/21/M/J/12

3 Methanol, CH₃OH, is considered to be a possible alternative to fossil fuels, particularly for use in vehicles.

For Examiner's Use

Methanol can be produced from fossil fuels and from agricultural waste. It can also be synthesised from carbon dioxide and hydrogen.

(a)	Define, with the aid of an equation which includes state symbols, the standard enthalpy
	change of formation of carbon dioxide.

equation	 	 	 	
definition	 	 	 	
	 	 	 	 [3]

(b) Relevant ΔH_f^{\bullet} values for the reaction that synthesises methanol are given in the table.

compound	$\Delta H_{\rm f}^{\rm e}/{\rm kJmol^{-1}}$
CO ₂ (g)	-394
CH₃OH(g)	-201
H ₂ O(g)	-242

(i) Use these values to calculate $\Delta H_{\text{reaction}}^{\bullet}$ for this synthesis of methanol.

Include a sign in your answer.

$$CO_2(g) + 3H_2(g) \rightleftharpoons CH_3OH(g) + H_2O(g)$$

$\Delta H_{\text{reaction}}^{\bullet} =$:KJ mol ⁻ '

(ii) Suggest **one** possible environmental advantage of this reaction. Explain your answer.

.....

[5]

[Total: 14]

For Examiner's Use

(c) The synthesis of methanol is carried out at about 500 K with a pressure of between 40 and 100 atmospheres (between 4×10^6 Pa and 10×10^7 Pa) and using a catalyst. The use of such conditions will affect both the rate of reaction and the equilibrium yield.


In the spaces below, explain the effects of higher temperature, higher pressure, and the use of a catalyst on the **equilibrium yield** of methanol.

gner temperature	
fect	
planation	
gher pressure	
fect	
planation	
se of catalyst	
fect	
planation	
	 6]
	']

© UCLES 2012 9701/21/M/J/12 **[Turn over**

4 One method of preparing ethene in a school or college laboratory is from ethanol by using the apparatus shown below.

For Examiner's Use



(a) (i)	Write a balanced equation for this reaction.
(ii)	What type of reaction is this?
(iii)	Give the chemical name of a reagent other than pumice that could be used to carry out this reaction. It is not necessary to use the same apparatus.
	[3]

(b) Ethene is bubbled into two separate test-tubes, one containing aqueous hydrogen bromide and the other containing cold, dilute acidified potassium manganate(VII).

In **each** case, describe any colour changes you would see and give the structural formula of the organic product.

	aqueous hydrogen bromide	cold, dilute acidified potassium manganate(VII)
colour at start		
colour after reaction		
structural formula of organic product		

[4]

	9	
(c) C	yclohexene has the following structural formula.	
(i) What is the molecular formula of cyclohexene?	
(ii	Draw the structural formula of the compound formed when cyclohexene is reacted with bromine.	ed
(iii) State as fully as you can what <i>type of reaction</i> this is.	
(iv) Draw the structural formula of the compound formed when cyclohexene is reacted with hot concentrated acidified potassium manganate(VII).	ed
		[5]
	[Total: 1	12]

For Examiner's Use

Organic compounds which contain oxygen may contain alcohol, aldehyde, carboxylic acid, ester or ketone functional groups. The functional groups may be identified by their reactions 5 with specific reagents.

For Examiner's Use

Compound	X has the	emnirical	formula	CH O and	$dM \cap fQ \cap$

	'	3				
С	ompoi	and X has the empirical formula CH_2O and M_r of 90.				
(a) The	ere is no reaction when X is treated with NaHCO ₃ .				
	Wh	at functional group does this test show to be not present in X ?				
		[1]				
(b	(b) When 0.600 g of X is reacted with an excess of Na, 160 cm³ of H ₂ , measured at room temperature and pressure, is produced.					
	(i)	What functional group does this reaction show to be present in X ?				
	(ii)	Use the data to calculate the amount, in moles, of hydrogen $atoms$ produced from $0.600g$ of ${\bf X}$.				
	(iii)	Hence, show that each molecule of X contains two of the functional groups you have given in (i).				

[4]

For

Examiner's

Use

www.maxpapers.com 11 (c) When **X** is warmed with Fehling's reagent, a brick red precipitate is formed. Treatment of **X** with 2,4-dinitrophenylhydrazine reagent produces an orange solid. (i) What functional group do these reactions show to be present in X? Draw the displayed formula of this functional group. (ii) Use your answers to (b)(i), (b)(ii) and (c)(i) to deduce the structural formula of X. (iii) What is the structural formula of the organic product of the reaction of X with Fehling's reagent? [3] (d) Compound X can be both oxidised and reduced. (i) Give the structural formula of the compound formed when **X** is reacted with NaBH_A under suitable conditions. (ii) Give the structural formula of the compound formed when **X** is heated under reflux with acidified $K_2Cr_2O_7$.

[2]

[Total: 10]

12

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.