UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education – Advanced Subsidiary Level and

Advanced Level

CHEMISTRY

Paper 2 Structured Questions AS Core



9701/02

October/November 2004

1 hour 15 minutes

Candidates answer on the Question Paper.
Additional Materials:
Data Booklet

Candidate Name		
Centre Number	Candidate Number	

READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number in the spaces provided at the top of this page. Write in dark blue or black pen in the spaces provided on the Question Paper.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all questions.

The number of marks is given in brackets [] at the end of each question or part question.

You may lose marks if you do not show your working or if you do not use appropriate units.

DO NOT WRITE IN THE BARCODE.

DO NOT WRITE IN THE GREY AREAS BETWEEN THE PAGES.

If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given on this page.

Stick your personal label here, if provided.

For Exam	iner's Use
1	
2	
3	
4	
5	
TOTAL	

This document consists of 11 printed pages and 1 blank page.

SP (AT) S83011/2 © UCLES 2004



[Turn over

For Examiner's Use

Нус	iroge	en lodide dissociates into its elements according to the equation below.
		$2HI(g) \rightleftharpoons H_2(g) + I_2(g)$
(a)	Wri	te the expression for the equilibrium constant, K_c .
		[1]
(b)		20 °C the equilibrium mixture contains 1.47 mol dm $^{-3}$ of HI(g), 0.274 mol dm $^{-3}$ each $I_2(g)$ and $I_2(g)$.
	Cal	culate the value of K_c for the equilibrium at 120 °C.
		[1]
		[1]
(c)		[1] gest and explain why it would be more difficult to determine ${\it K_c}$ for this equilibrium at m temperature.
(c)		gest and explain why it would be more difficult to determine \mathcal{K}_c for this equilibrium at
(c)		gest and explain why it would be more difficult to determine \mathcal{K}_c for this equilibrium at
(c)		gest and explain why it would be more difficult to determine \mathcal{K}_c for this equilibrium at
(c)		gest and explain why it would be more difficult to determine \mathcal{K}_c for this equilibrium at
(c)		gest and explain why it would be more difficult to determine \mathcal{K}_c for this equilibrium at
(c)		igest and explain why it would be more difficult to determine K_c for this equilibrium at m temperature.
		gest and explain why it would be more difficult to determine \mathcal{K}_c for this equilibrium at m temperature.
		igest and explain why it would be more difficult to determine K_c for this equilibrium at m temperature.
		igest and explain why it would be more difficult to determine K_c for this equilibrium at m temperature.
		igest and explain why it would be more difficult to determine K_c for this equilibrium at m temperature.
		igest and explain why it would be more difficult to determine K_c for this equilibrium at m temperature.
		igest and explain why it would be more difficult to determine K_c for this equilibrium at m temperature.

(ii)	Use bond	energies	from	the	Data	Booklet	to	calculate	ΔH	for	the	following
	dissociation	٦.										

For Examiner's Use

$$2H\mathrm{I}(g) \to H_2(g) + \mathrm{I}_2(g)$$

[3]

(e) HI dissolved in water behaves as a strong acid.

(i) Explain what is meant by a *strong* acid.

.....

(ii) Complete the equation.

 $\mathsf{HI} + \mathsf{H_2O} \rightarrow \dots \dots + \dots \dots$

(iii) Identify the conjugate base of HI in this equation.

.....[3]

[Total : 10]

The table below gives data on some oxides of elements in Period 3 of the Periodic Table. 2

For
Examiner's
Use

oxide	Na ₂ O	MgO	Al_2O_3	SiO ₂	P ₄ O ₁₀	SO ₃
melting point/K	1193	3125	2345	1883	853	290
boiling point/K	1548	3873	3253	2503	_	318

(a)	Write an equation for the reaction of aluminium with oxygen to form aluminium oxide.
	[1]
(b)	Drawing diagrams where appropriate, suggest in terms of structure and bonding explanations for the following.

(i) the high melting point and boiling point of ${\rm A}\it{l}_{2}{\rm O}_{3}$

(ii) the low boiling point of ${\rm SO}_3$

(iii) the melting point of ${\rm SiO_2}$ is much higher than that of ${\rm P_4O_{10}}$

(c)	Wat	er was added to each of the oxides in the table.	For
	Cho	osing a suitable oxide in each case, write an equation for the formation of	Examiner Use
	(i)	an alkaline solution,	
	(ii)	an acidic solution[2]	
		[Total : 10]	

3	(a)	(i)	Describe, with the aid of a fully labelled diagram, the industrial electrolysis of brine (aqueous NaC1). State what the electrodes are made of and show clearly the inlet and the outlets.	For Examiner's Use
		(ii)	Write equations for the reactions at each electrode, giving state symbols.	
		(11)	anode	
			cathode	
	((iii)	Explain in terms of changes in oxidation number why redox processes take place at the electrodes.	
			anode	
			cathode	
	((iv)	Name the chemical which is produced in solution by this electrolysis.	
		(v)	Suggest two large scale uses of this chemical.	
			[10]	

(b)	-	drochloric acid is manufactured by burning the hydrogen formed in this electrolysis in brine and dissolving the product in water.	For Examiner's
	(i)	Construct an equation for the burning of hydrogen in chlorine.	Use
	(ii)	When the product of (i) dissolves in water there is a change in bonding. Explain with the aid of an equation what change in bonding has occurred.	
		[2]	
(c)	Des	scribe, with the aid of equations including state symbols, what happens when	
	(i)	hydrochloric acid is added to aqueous silver nitrate,	
	(ii)	an excess of aqueous ammonia is added to the resulting mixture.	
		[E]	
		[5]	
		[Total : 17]	

4 Some perfumes and scents of flowers and fruit contain compounds which are structural isomers. Two such examples are citronellol and geraniol.

For Examiner's Use

citronellol

geraniol

- (a) Confirm that citronellol and geraniol are isomers by calculating their molecular formula and their relative molecular mass, M_r .
 - (i) Molecular formula
- (b) Name two functional groups present in **both** molecules.
 - (i)
 - (ii)[3]

Citronellol and geraniol also show stereo isomerism.

(c) On the diagram of the structure of citronellol above, draw a circle around a chiral carbon atom.

For Examiner's Use

d)	(i)	Draw the other cis -trans isomer of geraniol. [In parts (d) and (f) use R - to represent a part of the molecule.]
	(ii)	Explain why geraniol has no optical isomers.
		[2]
e)	Stat	te what you would expect to see if citronellol was reacted with aqueous bromine.
		[1]
f)		w structures of the organic products when geraniol reacts with each of the following gents.
	(i)	an excess of H ⁺ /Cr ₂ O ₇ ²⁻ under reflux
	(ii)	ethanoic acid in the presence of an acidic catalyst

[4]

[Total : 13]
[Turn over

2-Hydro synthes	xypropanoic acid (lactic acid), $\mathrm{CH_3CH(OH)CO_2H}$, can be prepared in a two-stage is from ethanal, $\mathrm{CH_3CHO}$.	For Examiner's Use			
	In the first stage, ethanal reacts with hydrogen cyanide, HCN, in the presence of ar NaCN catalyst to produce a cyanohydrin.				
(i)	Write an equation for the reaction of ethanal and HCN, giving the displayed formula of the product.				
(ii)	State what type of reaction this is.				
(iii)	Describe the mechanism of this reaction.				
	[5]				
(b) In t	he second stage, the product from (a) is converted into lactic acid.				
(i)	Write the equation for this stage.				
(ii)	State what type of reaction this is.				
	[2]				

5

(c) In this synthesis 4.40 g of ethanal were used and at the end 5.40 g of lactic acid were obtained.

Examiner's Use

Calculate the percentage yield of lactic acid.

[3]

[Total : 10]

BLANK PAGE

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES) which is itself a department of the University of Cambridge.

