Centre Number

Candidate Number

Name

## UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

CHEMISTRY 9701/04

Paper 4 Structured Questions A2 Core

October/November 2004

1 hour 15 minutes

Candidates answer on the Question Paper. Additional Materials: Data Booklet

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen in the spaces provided on the Question Paper.

You may use a pencil for any diagrams, graphs or rough working.

You may use a calculator.

Do not use staples, paper clips, highlighters, glue or correction fluid.

## Answer all questions.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

You may lose marks if you do not show your working or if you do not use appropriate units.

If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given at the top of this page.

Stick your personal label here, if provided.

For Exam	iner's Use
1	
2	
3	
4	
5	
6	
7	
Total	

This document consists of 12 printed pages.





[Turn over

For Examiner's Use

1 Sulphuric acid is a strong dibasic acid, which ionises in solution as follows.


$$H_2SO_4(aq) \Longrightarrow 2H^+(aq) + SO_4^{2-}(aq)$$

(a) The organic base guanidine contains carbon, nitrogen and hydrogen. Its reaction with acids can be represented as follows.

$$B(aq) + H^+(aq) \Longrightarrow BH^+(aq)$$

where B represents the molecule of guanidine.

When a 25.0 cm<sup>3</sup> sample of dilute sulphuric acid was titrated against a solution of guanidine, the following titration curve was obtained.



Use this curve to answer the following questions.

(i)	Is guanidine a strong or a weak base? Explain your answer.
(ii)	The pH at the start of the titration was 0.70. Calculate the [H <sup>+</sup> ], and hence the concentration of sulphuric acid, at the start of the titration.




	(iii)	Calculate the concentration of guanidine in the solution in mol dm <sup>-3</sup> .
	(iv)	The guanidine solution contained 8.68 g of the base per dm $^3$ . Use your answer to (iii) calculate the $M_{\rm r}$ of guanidine.
(b)	fluo	e of the major industrial uses of sulphuric acid is to convert phosphate rock (calcium rophosphate(V)) into 'superphosphate' for use as a fertiliser. The process can be resented by the following partially balanced equation.
	2 Ca	$_{5}(PO_{4})_{3}F$ + $7H_{2}SO_{4}$ $\longrightarrow$ $CaSO_{4}$ + $Ca(H_{2}PO_{4})_{2}$ + HF 'superphosphate'
	(i)	Balance the above equation.
	(ii)	Use your balanced equation to calculate the mass of $\rm H_2SO_4$ required to manufacture 1.0 kg of superphosphate fertiliser.
		[4]
(c)	Solu	utions of hydrogenphosphates make useful buffers for biochemical experiments.
		$H_2PO_4^- \rightleftharpoons HPO_4^{2-} + H^+$
	(i)	Explain what is meant by the term buffer solution.
	(ii)	Calculate the pH of a buffer solution that contains $0.20\mathrm{moldm^{-3}}\mathrm{NaH_2PO_4}$ and $0.10\mathrm{moldm^{-3}}\mathrm{Na_2HPO_4}$ . $[K_\mathrm{a}(\mathrm{H_2PO_4^-})=6.3\mathrm{x}10^{-8}\mathrm{moldm^{-3}}]$
		[3]
		[Total: 13]



For Examiner's Use

2 The diagram shows a laboratory illustration of a simple hydrogen-oxygen fuel cell.



when the cell operates.	(a)
[1]	
) State the polarity (+ or -) of the left hand (oxygen) electrode[1]	(b)
Use the Data Booklet to calculate the voltage produced by this cell.	(c)
[1]	
Only a very small current can be drawn from this laboratory cell. Suggest <b>one</b> way in which it could be modified to enable a larger current to be drawn from it.	(d)
[1]	
A fuel cell in an orbiting satellite is required to produce a current of 0.010 A for 400 days. Calculate the mass of hydrogen that will be needed.	(e)
[3]	

Use

	(1)	compared to hydrocarbon fuels such as petrol.
		advantage:
		disadvantage:
		[2]
		[Total: 9]
3	me <sup>s</sup>	cium sulphate is a major by-product of flue gas desulphurisation, which is an important thod of decreasing the emission of acid-rain gases from power stations. It is used ensively in plaster and cement. Both magnesium sulphate and barium sulphate find uses nedicine.
	Des	scribe and explain the variation in the solubilities of the Group II sulphates in water.
		[4]
		[Total: 4]

(a)	Exp	plain what is meant by the term transition element.
		[1]
(b)	(i)	How do the atomic radii of the transition elements vary from chromium to copper?
	(ii)	Predict, with a reason, the variation in the densities of the transition elements from chromium to copper.
		[3]
(c)	Cor	nplete the following electronic configuration of the Cu <sup>2+</sup> ion.
		1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup>
(d)	Cop	oper ions in aqueous solution are pale blue, due to the formation of a complex ion.
	(i)	Explain what is meant by the term complex ion.
	(ii)	Draw the structure of the complex ion formed in a solution of Cu <sup>2+</sup> (aq).

[2]




4

(e)	When dilute aqueous ammonia is added to a solution of Cu <sup>2+</sup> (aq), the colour changes as a new complex ion is formed.
	(i) State the colour of the new complex
	(ii) Write an equation showing the formation of the new complex.
	[2]
(f)	When concentrated hydrochloric acid is added to a solution of $Cu^{2+}(aq)$ , the colour changes to yellow-green. On adding water, the colour returns to pale blue.
	Suggest an explanation for these changes.
	[3]
	[Total: 12]

studyguide.pk

- 5 This question is concerned with organochlorine compounds.
  - (a) State the conditions needed to produce the two compounds A and B.



(i) conditions for reaction I

.....

(ii) conditions for reaction II

.....[2]

**(b)** State the reagent needed to carry out the following reaction.

C

reagent for reaction III: ......[1]


- (c) The three chloro-compounds A, B and C vary in their ease of hydrolysis.
  - (i) Place a tick in the box corresponding to the correct relative rates of hydrolysis. [the symbol '>' means 'faster than']

	place <b>one</b> tick only in this column
A > B > C	
A > C > B	
B > A > C	
B > C > A	
C > B > A	
C > A > B	

(ii)	Suggest an explanation for these differences in reactivity.
	[3

For Examiner's Use

(d) Draw the structural formulae of the organic products of the following reactions of compound  ${\bf C}$ .



[3]

[Total: 9]

6 Compounds **D** and **E** are both ketones.

(ii)

products

$\sim$ 1.1 $\sim$ 1.	$\Gamma \cap \cap \cap$	$\cdots$
.H (.F	1 (.()(.	H <sub>2</sub> CH <sub>2</sub>
$\mathcal{L}$	いっししし	1 1201 12

D

CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>COCH<sub>3</sub>

Ε

(a) State which one of these compound reacts with alkaline aqueous iodine, and draw the structural formulae of the products formed during this reactions.(i) compound (D or E) ......

(b)	The reduction of <b>D</b> with NaBH <sub>4</sub> produces just one alcohol, but a similar reduction of <b>E</b> produces two isomers in equal amounts. Explain these observations, drawing structures where appropriate.

[3]


[Total: 6]



7 Both phenol and phenylamine react similarly with aqueous bromine.

(a)	State two observations	you would	make when	these reaction	ons take place.
-----	------------------------	-----------	-----------	----------------	-----------------


- **(b)** Describe a simple test-tube reaction you could use to distinguish between phenol and phenylamine.
- (c) The compound 3-aminobenzoic acid can be prepared by the following series of reactions.



Suggest suitable reagents and conditions for reaction IV, reaction V,

reaction VI. ......[4]

[Total: 7]

Every reasonable effort has been made to trace all copyright holders where the publishers (i.e. UCLES) are aware that third-party material has been reproduced. The publishers would be pleased to hear from anyone whose rights they have unwittingly infringed.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

