

CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Level

MARK SCHEME for the October/November 2013 series

9701 CHEMISTRY

9701/51

Paper 5 (Planning, Analysis, Evaluation),
maximum raw mark 30



This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	51

Question	Expected Answer	Mark
1 (a) (i)	(The temperature would) decrease The lattice enthalpy is more negative/exothermic than the (sum of the) enthalpies/energies of hydration.	1 1
(ii)	 <p>OR</p> <p>OR ecf from 1(a)(i)</p> 	2

Page 3	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	51

(b) (i)	concentration/concentration change	
(ii)	temperature change/decrease in temperature (allow ecf on (a)(i))	1
(c)	Diagram shows a container labelled with its capacity (between 25 cm ³ and 250 cm ³) and with the thermometer in a solution.	1
	The apparatus is insulated and has a lid.	1
	Thermometer range must include 25 °C and with a precision of between 0.1 °C and 0.5 °C.	1
(d)	A minimum of 5 workable experiments using masses or concentrations.	1
	Measures initial and final temperatures.	1
	Measures a volume of water AND the volume of water will fit into container labelled in (c) .	1
	States a mass which is the maximum for a volume of water stated.	1
(e)	Ammonium nitrate may cause a fire/explosion so must not be ground up OR dilute to less than 0.5 mol dm ⁻³ before disposal.	1
(f)	Columns must include units: Mass of ammonium nitrate used / any mass unit Volume / mass of water used / any volume or mass units Initial temperature / °C Final temperature / °C Temperature fall / change in temperature / °C Concentration of ammonium nitrate / any concentration units	
	Four columns correct	1
	Five or six columns correct	1
		[Total: 15]

Page 4	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	51

2 (a)	<table><tr><th>F</th><th>G</th><th>H solubility</th></tr><tr><th>D–C / g</th><th>C–B / g</th><th>[(F × 100)] / G / g/ 100 g</th></tr><tr><td>1.25</td><td>25.00</td><td>5.00</td></tr><tr><td>1.25</td><td>20.00</td><td>6.25</td></tr><tr><td>5.00</td><td>25.00</td><td>20.00</td></tr><tr><td>7.76</td><td>19.40</td><td>40.00</td></tr><tr><td>11.11</td><td>23.00</td><td>48.30</td></tr><tr><td>11.75</td><td>25.00</td><td>47.00</td></tr><tr><td>9.62</td><td>21.00</td><td>45.81</td></tr><tr><td>9.10</td><td>20.00</td><td>45.50</td></tr><tr><td>11.25</td><td>25.00</td><td>45.00</td></tr><tr><td>13.35</td><td>30.00</td><td>44.50</td></tr></table>	F	G	H solubility	D–C / g	C–B / g	[(F × 100)] / G / g/ 100 g	1.25	25.00	5.00	1.25	20.00	6.25	5.00	25.00	20.00	7.76	19.40	40.00	11.11	23.00	48.30	11.75	25.00	47.00	9.62	21.00	45.81	9.10	20.00	45.50	11.25	25.00	45.00	13.35	30.00	44.50	
	F	G	H solubility																																			
	D–C / g	C–B / g	[(F × 100)] / G / g/ 100 g																																			
	1.25	25.00	5.00																																			
	1.25	20.00	6.25																																			
	5.00	25.00	20.00																																			
	7.76	19.40	40.00																																			
	11.11	23.00	48.30																																			
	11.75	25.00	47.00																																			
	9.62	21.00	45.81																																			
	9.10	20.00	45.50																																			
	11.25	25.00	45.00																																			
	13.35	30.00	44.50																																			
Heading for final column calculating the solubility is given correctly with units.	1																																					
All data is to 2 decimal places. Allow 1 error.	1																																					
Data in final column is correct. Allow 1 error in computation.	1																																					
(b)	The x-axis must start at zero and be labelled 'temperature / °C' OR T / °C and y-axis as 'solubility (of sodium sulfate) g/100g'. Plotted points must cover at least half the grid in both directions.	1																																				
All 10 points plotted correctly.	1																																					
First (left-hand) curve is smooth passing through (or extremely close to) all the points and does not deviate to accommodate a mis-plot or incorrect point. Curve intersects with a second curve at or above the candidate's solubility for experiment 5.	1																																					
Second curve (right hand) is smooth passing through (or extremely close to) all the points and does not deviate to accommodate a mis-plot or incorrect point.	1																																					
(c)	the temperature is read correctly the solubility is read correctly	1 1																																				
(d) (i)	Cross is on the 40 g / 100 g line and to the right of the point plotted at 30 °C.	1																																				
(ii)	Transition temperature would be higher as intersection of curves would be at a higher temperature.	1																																				
(e)	Solubility is 47.6 (g / 100g) 1.2% OR 1.21% OR 1.28% OR 1.3%	1 1																																				

Page 5	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	51

(f)	$\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ endothermic because solubility increases with increasing temperature (or reverse argument) Na_2SO_4 exothermic because solubility decreases with increasing temperature (or reverse argument)	
	For endothermic and exothermic correctly assigned For providing the correct reasons	1 1
		[Total: 15]

CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Level

MARK SCHEME for the October/November 2013 series

9701 CHEMISTRY

9701/52

Paper 5 (Planning, Analysis, Evaluation),
maximum raw mark 30



This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	52

Question	Expected Answer	Mark
1 (a) (i)	(The temperature would) decrease The lattice enthalpy is more negative/exothermic than the (sum of the) enthalpies/energies of hydration.	1 1
(ii)	 <p>OR</p> <p>OR ecf from 1(a)(i)</p> 	2

Page 3	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	52

(b) (i)	concentration/concentration change	
(ii)	temperature change/decrease in temperature (allow ecf on (a)(i))	1
(c)	Diagram shows a container labelled with its capacity (between 25 cm ³ and 250 cm ³) and with the thermometer in a solution.	1
	The apparatus is insulated and has a lid.	1
	Thermometer range must include 25 °C and with a precision of between 0.1 °C and 0.5 °C.	1
(d)	A minimum of 5 workable experiments using masses or concentrations.	1
	Measures initial and final temperatures.	1
	Measures a volume of water AND the volume of water will fit into container labelled in (c) .	1
	States a mass which is the maximum for a volume of water stated.	1
(e)	Ammonium nitrate may cause a fire/explosion so must not be ground up OR dilute to less than 0.5 mol dm ⁻³ before disposal.	1
(f)	Columns must include units: Mass of ammonium nitrate used / any mass unit Volume / mass of water used / any volume or mass units Initial temperature / °C Final temperature / °C Temperature fall / change in temperature / °C Concentration of ammonium nitrate / any concentration units	
	Four columns correct	1
	Five or six columns correct	1
		[Total: 15]

Page 4	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	52

2	(a)	<table><tr><th>F</th><th>G</th><th>H solubility</th></tr><tr><th>D–C / g</th><th>C–B / g</th><th>[(F × 100)] / G / g/ 100 g</th></tr><tr><td>1.25</td><td>25.00</td><td>5.00</td></tr><tr><td>1.25</td><td>20.00</td><td>6.25</td></tr><tr><td>5.00</td><td>25.00</td><td>20.00</td></tr><tr><td>7.76</td><td>19.40</td><td>40.00</td></tr><tr><td>11.11</td><td>23.00</td><td>48.30</td></tr><tr><td>11.75</td><td>25.00</td><td>47.00</td></tr><tr><td>9.62</td><td>21.00</td><td>45.81</td></tr><tr><td>9.10</td><td>20.00</td><td>45.50</td></tr><tr><td>11.25</td><td>25.00</td><td>45.00</td></tr><tr><td>13.35</td><td>30.00</td><td>44.50</td></tr></table>	F	G	H solubility	D–C / g	C–B / g	[(F × 100)] / G / g/ 100 g	1.25	25.00	5.00	1.25	20.00	6.25	5.00	25.00	20.00	7.76	19.40	40.00	11.11	23.00	48.30	11.75	25.00	47.00	9.62	21.00	45.81	9.10	20.00	45.50	11.25	25.00	45.00	13.35	30.00	44.50	
		F	G	H solubility																																			
		D–C / g	C–B / g	[(F × 100)] / G / g/ 100 g																																			
		1.25	25.00	5.00																																			
		1.25	20.00	6.25																																			
		5.00	25.00	20.00																																			
		7.76	19.40	40.00																																			
		11.11	23.00	48.30																																			
		11.75	25.00	47.00																																			
		9.62	21.00	45.81																																			
		9.10	20.00	45.50																																			
		11.25	25.00	45.00																																			
		13.35	30.00	44.50																																			
Heading for final column calculating the solubility is given correctly with units.	1																																						
All data is to 2 decimal places. Allow 1 error.	1																																						
Data in final column is correct. Allow 1 error in computation.	1																																						
(b)	The x-axis must start at zero and be labelled ‘temperature / °C’ OR T / °C and y-axis as ‘solubility (of sodium sulfate) g/100g’. Plotted points must cover at least half the grid in both directions.	1																																					
	All 10 points plotted correctly.	1																																					
	First (left-hand) curve is smooth passing through (or extremely close to) all the points and does not deviate to accommodate a mis-plot or incorrect point. Curve intersects with a second curve at or above the candidate’s solubility for experiment 5.	1																																					
	Second curve (right hand) is smooth passing through (or extremely close to) all the points and does not deviate to accommodate a mis-plot or incorrect point.	1																																					
	(c)	the temperature is read correctly the solubility is read correctly	1 1																																				
(d) (i)	Cross is on the 40 g / 100 g line and to the right of the point plotted at 30 °C.	1																																					
(ii)	Transition temperature would be higher as intersection of curves would be at a higher temperature.	1																																					
(e)	Solubility is 47.6 (g / 100g) 1.2% OR 1.21% OR 1.28% OR 1.3%	1 1																																					

Page 5	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	52

(f)	$\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ endothermic because solubility increases with increasing temperature (or reverse argument) Na_2SO_4 exothermic because solubility decreases with increasing temperature (or reverse argument)	
	For endothermic and exothermic correctly assigned For providing the correct reasons	1 1
		[Total: 15]

CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Level

MARK SCHEME for the October/November 2013 series

9701 CHEMISTRY

9701/53

Paper 5 (Planning, Analysis and Evaluation),
maximum raw mark 30

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	53

Question	Expected Answer	Mark
1 (a) (i)	(The solubility of air) decreases (as the temperature is increased).	1
	Dissolving is exothermic so an increase in temperature will promote the reverse reaction (or backwards or right to left).	1
(ii)	Axes are labelled AND graph is a curve/straight line showing a decrease in solubility with temperature.	1
	Graph is scaled and starts at 0 °C on the solubility axis AND goes through point (25 °C, 19 cm ³ dm ⁻³), AND ends at point (100 °C, 0 cm ³ dm ⁻³) AND provided there is no maximum or minimum in the curve. Units required for this mark.	1
(b)	(i) temperature	1
	(ii) solubility (of air).	
	Total 5	
2 (a)	Diagram shows a container with both chemicals named and attached to a syringe connected without leaks.	1
	Container shows the catalyst and hydrogen peroxide separated ready to mix.	1
	100 cm ³ of oxygen is 100/24000 = 0.00417 (mol) (0.004166666)	1
	Mol of H ₂ O ₂ is 2 × mole answer above = 0.00834 (0.00833333) (ecf on alternative volume of oxygen used)	1
	Volume of hydrogen peroxide is therefore (answer above × 1000)/0.30 (27.8 cm ³) (27.78 cm ³) (27.7777773 cm ³) Allow 28 cm ³ (units are required) (reverse calculation also accepted)	1
	The reaction is timed from the moment of mixing to the collection of a stated volume of oxygen.	1
(b)	surface area of catalyst.	1
(c)	Axes are labelled (vol and time or min or s etc.) AND graph is a curve starting at the origin, shows the steepest slope at the start and slowing down as reaction proceeds	1
	x-axis has numerical scale from the origin to at least 3 min and graph shows no change in volume of oxygen after 2 minutes. The time axis must be scaled and have a unit of min or s.	1

Page 3	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9701	53

Question	Expected Answer	Mark
(d)	lead(IV) oxide causes (skin) irritation AND wear (safety) gloves OR Either catalyst or both catalysts (or names or powders or solids) are poisonous AND wear a face mask OR do in a fume cupboard.	1
	Total 10	
3 (a)	$M_r(\text{CCl}_4) = 154$ AND $M_r(\text{C}_2\text{H}_2\text{Cl}_4) = 168$	1
(b) (i)	Mol of $\text{CCl}_4 = 0.0469$ AND mol of $\text{C}_2\text{H}_2\text{Cl}_4 = 0.0531$ AND mole fraction = $0.0469 / (0.0469 + 0.0531) = 0.469$	1
(ii)	Mol of $\text{CCl}_4 = 0.0918$ AND mol of $\text{C}_2\text{H}_2\text{Cl}_4 = 0.00821$ AND mole fraction = $0.0918 / (0.0918 + 0.00821) = 0.918$	1
(c)	The x-axis must start at zero and be labelled as 'mole fraction' with no units and y-axis as temperature or $T/^\circ\text{C}$. Plotted points must cover at least half the grid in both directions. All 16 points plotted correctly . The vapour line is a continuous curve of best fit that does not deviate to accommodate an off curve point. The liquid line is a continuous curve of best fit that does not deviate to accommodate an off curve point.	1 1 1
(d)	Anomalous point is circled at $T = 99.3^\circ\text{C}$ (for the vapour curve). Analysis was made at a temperature that was too low.	1 1
(e)	Horizontal line drawn from 0.500 on the liquid mol fraction curve to meet vapour curve. Correctly reads the value from the vapour curve.	1 1
(f)	For $\text{C}_2\text{H}_2\text{Cl}_4$ $(0.5/120) \times 100 = 0.417\%$ OR $(1.0/120) \times 100 = 0.833\%$ For CCl_4 $(0.5/76) \times 100 = 0.658\%$ OR $(1.0/76) \times 100 = 1.316\%$	1 1
(g) (i)	CCl_4	1
(ii)	Vapour produced when a mixture is heated has a greater proportion of CCl_4 than the mixture/liquid.	1
	Total 15	