UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

General Certificate of Education

Advanced Subsidiary Level and Advanced Level

CHEMISTRY

Paper 2 Structured Questions AS Core



9701/02

May/June 2005

1 hour 15 minutes

Candidates answer on the Question Paper. Additional Materials: Data Booklet

Candidate Name						
Centre Number			Candidate Number			

READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number in the spaces at the top of this page.

Write in dark blue or black pen in the spaces provided on the Question Paper.

You may use a pencil for any diagrams, graphs, or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all questions.

The number of marks is given in brackets [] at the end of each question or part question.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

You may use a calculator.

DO NOT WRITE IN THE BARCODE.

DO NOT WRITE IN THE GREY AREAS BETWEEN THE PAGES.

For Exam	iner's Use
1	
2	
3	
4	
5	
TOTAL	

This document consists of 10 printed pages and 2 blank pages.

SP (MML 8078 3/04) S92060/2.1
© UCLES 2005
STUCYCUCE-DK

Answer all the questions in the space provided.

For Examiner's Use

				nas one.								
(a)	Ехр	lain the mea	aning c	of the term	isotop	e.						
											[2	21
(b)	The		mon isc								tope of coba	-
		the <i>Data E</i> of ⁵⁹ Co.	Booklet	to comple	ete the	table	below to	sho	w the a	atomic stru	ucture of ⁵⁶ F	-e
						n	umber of					
		isotop	е	proto	ns	r	neutrons		ele	ectrons	-	
		⁵⁶ Fe										
		⁵⁹ Co										
												וכי
(c)	A sa	ample of iro	n has t	he followir	ng isoto	pic c	ompositic	on by	y mass	S.	Į.	3]
(c)	A sa	ample of iro		he followir	ng isoto	pic c	ompositic		y mass 57	5.	Į.	o]
(c)	A sa	ample of iro	isotop							3.	Į.	ی]
(c)	A sa	ample of iro	isotop	pe mass y mass	54 5.84	ļ	56		57	3.	Į.	S]
(c)			isotop	pe mass y mass	54 5.84	ļ	56		57	5.		
(c)			isotop	pe mass y mass	54 5.84	ļ	56		57	S.	Į.	
(c)			isotop	pe mass y mass	54 5.84	ļ	56		57	S.		
(c)		Define the	isotop	oe mass y mass elative ato	54 5.84 mic ma	PSS.	56 91.68	2	57		iron to thre	

[5]

[Total: 10]

BLANK PAGE

2 Sulphur and its compounds are found in volcanoes, in organic matter and in minerals.

For Examiner's Use

Sulphuric acid, an important industrial chemical, is manufactured from sulphur by the Contact process. There are three consecutive reactions in the Contact process which are essential.

Coo	Jiiliai	•
(a)		e a balanced equation (using \rightleftharpoons where appropriate) for each of these reactions ne correct sequence .
	1	
	2	
	3	[4]
(b)	Wha	at catalyst is used?
		[1]
		n sulphide, H ₂ S, is a foul-smelling compound found in the gases from volcanoes. n sulphide is covalent, melting at –85 °C and boiling at –60 °C.
(c)	(i)	Draw a 'dot-and-cross' diagram to show the structure of the H ₂ S molecule.
	(ii)	Predict the shape of the H ₂ S molecule.
	(iii)	Oxygen and sulphur are both in Group VI of the Periodic Table.
		Suggest why the melting and boiling points of water, $\rm H_2O$, are much higher than those of $\rm H_2S$.
		[4]

[Total: 17]

For Examiner's Use

•	Hydrogen sulphide burns with a blue flame in an excess of oxygen to form sulphur dioxide and water.						
(d)	(i)	Write a balanced equation for the complete combustion of H ₂ S.					
	(ii)	What is the change in the oxidation number of sulphur in this reaction?					
		from to					
	(iii)	What volume of oxygen, measured at room temperature and pressure, is required for the complete combustion of 8.65 g of $\rm H_2S$? Give your answer to two decimal places.					
		[5]					
Hyd few	roge S ²⁻ i	n sulphide is a weak diprotic (dibasic) acid. Its solution in water contains HS ⁻ and a ons.					
(e)	(i)	What is meant by the term weak acid?					
	(ii)	Write an equation, with state symbols, for the ${\it first}$ ionisation of ${\rm H_2S}$ when it dissolves in water.					
		[3]					


studyguide.pk Turn over

For Examiner's Use

3 Magnesium is the eighth most common element in the Earth's crust.

The metal is widely used in alloys which are light and strong.

Some reactions of magnesium and its compounds are shown in the reaction scheme below.



(a) Identify, by name or formula, compounds A to F.

Α	٠.		 	٠.		 	 	•	 •		 		•	 •	 •		 •	 ٠.	-	
В		 •	 			 •					 			 •				 	•	
С			 								 			 •				 	•	
D		 •	 				 				 		-	 •				 	•	
Ε		 •	 				 				 		-	 •				 	•	
F	 	 	 				 											 		

[6]



(b)	(i)	Construct balanced equations for the following reactions. magnesium to compound A				
		compound C to compound D				
		compound F to compound D				
	(ii)	Suggest a balanced equation for the effect of heat on compound E .				
		[4] [Total: 10]				

4 Compound **G**, in which R- represents the rest of the molecule, was made for use as a tear gas in World War 2.

For Examiner's Use

compound **G**

Compound **G** was made by the following sequence of reactions.

stage I		stage II		stage III	
R-CH ₃ stage I →	R-CH ₂ Cl	→	R-CH ₂ CN	→	R-CHBrCN

(a) (i) For stage I and for stage II, state the reagent(s) and condition(s) used to carry out each change.

stage I	reagent(s)
	condition(s)
stage II	reagent(s)
	condition(s)

(ii) Suggest the reagent(s) and condition(s) necessary to carry out stage III.

reagent(s)	
condition(s)	
	[6]

studyguide.pk

Compound **G** was not actually used in World War 2 and stocks of it had to be destroyed safely. The following sequence of reactions was used in this process.

For Examiner's Use

R-CHBrCN	stage IV ➤ R-CHBrCO ₂ H	stage V	R-CH(OH)CO ₂ H
			stage VI
		ا	√ R-CH₂CO₂H

stages I to VI which you consider to be chiral.

			R-CH ₂ CO ₂ H
(b)		_	IV and for stage V state the reagent(s) and condition(s) necessary to bring ch reaction.
	stage	IV	reagent(s)
			condition(s)
	stage	V	reagent(s)
			condition(s) [4]
(c)	The force		sequence of stages I to VI involves some compounds which contain chiral
	(i) E	xpla	ain what is meant by the term <i>chiral centre</i> .
	(ii) D	raw	displayed formulae for the isomers of one compound in the full sequence of

[3]

[Total: 13]

[2]

5 A student obtained the following results when analysing an organic compound, H.

For Examiner's Use

test		observation
test 1	relative molecular mass	72
test 2	% composition by mass	C, 66.7%; H, 11.1%; O, 22.2%
test 3	reactions with Br ₂ (aq)	Br ₂ decolourised
test 4	reaction with Na(s)	H ₂ (g) evolved
test 5	reaction with warm Cr ₂ O ₇ ²⁻ /H ⁺	green colour observed

The student allowed test 5 to go to completion and then investigated the **product** of test 5 with the following result.

test 6	reaction with 2,4-dinitrophenylhydrazine	no reaction
--------	--	-------------

(a) Calculate the molecular formula of H.

(b)	Wha	at can be deduced about the nature of H by the following tests?	
	(i)	test 3	
	(ii)	test 4	
			[2]
(c)	(i)	What functional group would have given a positive result in test 6?	
	(ii)	What functional group is shown to be present in H by tests 5 and 6?	

d)	On testing a sample of H , the student found that it was not chiral.				
	H did, however, show <i>cis-trans</i> isomerism.	Examiner's Use			
	How does <i>cis-trans</i> isomerism arise in an organic molecule?				
	[2]				
(e)	Use all of the information above to draw labelled, displayed formulae of the stereoisomers of compound H .				

[2]

[Total: 10]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

