Centre Number

Candidate Number

Name

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

CHEMISTRY 9701/04

Paper 4 Structured Questions A2 Core

October/November 2005

1 hour 15 minutes

Candidates answer on the Question Paper. Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number in the spaces at the top of this page. Write in dark blue or black pen in the spaces provided on the Question Paper. You may use a pencil for any diagrams, graphs, or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all questions.

The number of marks is given in brackets [] at the end of each question or part question. You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

You may use a calculator.

If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given at the top of this page.

Stick your personal label here, if provided.

For Exam	iner's Use
1	
2	
3	
4	
5	
TOTAL	

This document consists of 10 printed pages and 2 blank pages.

SPA (MML 8078 3/04 © UCLES 2005




[Turn over

⊢or
Examiner's
Use

- Silver bromide, AgBr, is widely used in photography. In a photographic film, AgBr crystals are precipitated into a gelatine base as 'grains' of diameter about 1×10^{-6} m.
 - (a) Calculate the approximate number of silver ions contained in a grain of AgBr of mass $2.5\times10^{-12}\,g$.

(b) AgBr is only sparingly soluble in water. The [Ag⁺] in a saturated solution of AgBr can be estimated by measuring the $E_{\rm cell}$ of the following cell.



(i) In the spaces below, identify what the four letters ${\bf A}-{\bf D}$ in the above diagram represent.

A C

(ii) Predict how the potential of the right hand electrode might vary as [Ag+] is decreased.

In its saturated solution, [AgBr(aq)] = 7.1×10^{-7} mol dm⁻³.

(iii) Write an expression for the solubility product of AgBr, and calculate its value, including units.

.....

[7]

(c)	(i)	(i) Write a chemical equation representing the lattice energy of AgBr.		
	(ii) Use the following data to calculate a value for the lattice energy of AgBr(s).			
		first ionisation energy of silver $= +731 \text{ kJ mol}^{-1}$ electron affinity of bromine $= -325 \text{ kJ mol}^{-1}$ enthalpy change of atomisation of silver $= +285 \text{ kJ mol}^{-1}$ enthalpy change of atomisation of bromine $= +112 \text{ kJ mol}^{-1}$ enthalpy change of formation of AgBr(s) $= -100 \text{ kJ mol}^{-1}$		
	(iii)	How might the lattice energy of AgCl compare to that of AgBr? Explain your answer.		
		[4]		
-		graphy a bromide ion absorbs a photon and releases an electron which reduces a to a silver atom.		
		$Br^- \longrightarrow Br + e^-$		
		$Ag^+ + e^- \longrightarrow Ag$		
(d)		dict whether it would require more energy or less energy to initiate this process in gCl emulsion, compared to a AgBr emulsion. Explain your answer.		



[Total: 14] | [Turn over

For
Examiner's
Use

(a)	What do you understand by the term standard electrode potential?		
	[2]		
(b)	By reference to relevant E^{\ominus} data in the <i>Data Booklet</i> , explain how the halogen/halide electrode potentials relate to the relative reactivity of the halogens as oxidising agents.		
	ro.		
	[2]		
(c)	Use data from the <i>Data Booklet</i> to construct redox equations, and calculate the standard cell potentials, for the reactions between		
	(i) Acidified H ₂ O ₂ (aq) and KI(aq),		
	(ii) $Cl_2(aq) + SO_2(aq)$.		
	[4]		
(d)	Use data from the <i>Data Booklet</i> to predict the likely product of the reaction between		
(α)	I_2 (aq) and tin metal, writing a balanced equation for the reaction.		
	[2]		
	[Total: 10]		


2

BLANK PAGE



For Examiner's Use

3 (a) (i) Use the following sets of axes to sketch graphs of the variations in the melting points and the electrical conductivities of the Group IV elements.



For
Examiner's
Use

(c)		ne one ceramic based on silicon(IV) oxide, and explain what properties of the le make it suitable for this use.
		[1]
(d)	Tin(II) oxide reacts with both acids and alkalis.
	(i)	What name is given to this property of an oxide?
	(ii)	Write suitable equations to show these two reactions of tin(II) oxide.
		[3]
		[Total: 13]

For Examiner's Use

4 Rodinol is used as a photographic developer. In alkaline solution it is a mild reducing agent, providing electrons according to the following half equation.

OH
$$+ 2 OH^ + H_2O + NH_3 + 2e^-$$


$$E^{\Theta} = +0.65 V$$
rodinol

Rodinol 'develops' a latent photographic image by reducing activated silver bromide grains to silver metal and bromide ions.

(a) Construct a balanced equation for the reaction between rodinol and AgBr.

	[1]
(b)	Suggest, with a reason, how the basicity of rodinol might compare to that of ammonia.
	[2]

(c) Suggest structural formulae for the compounds E, F and G in the following chart of the reactions of rodinol.




[3]



For Examiner's Use

(d) Rodinol can be synthesised from phenol by the following route.



(i)	Suggest reagents and conditions for step I.

(ii)	What type of reaction is step II?

.....

(iii) Place a tick in the box by the most suitable reagent for step II.

(place a tick in one box only)

$$Cr_2O_7^{2-} + H^+ + heat$$

[3]

(e) Rodinol is also an important intermediate in the commercial production of the analgesic drug *paracetamol*.

(i) Name two functional groups in paracetamol.

 	 •••••	•••••	

(ii) Suggest a reagent to convert rodinol into paracetamol.

[3] [Total: 12]

Examiner's Use

5 Hydrophilic polymers find important uses in the manufacture of contact lenses and wound dressings. Their chemical structures allow them to bond with water molecules, which keeps them soft and flexible. Sections of two hydrophilic polymers are shown below.

(a)	VVII	at type of polymensation has produced	
	(i)	polymer H?	
	(ii)	polymer J ?[2]	
(b)	Wh	at type of attractions might occur between these polymers and molecules of water?	
		[1]	
(c)	Chains of polymer H can be 'cross-linked', i.e. joined together, by reaction with a sma bifunctional molecule.		
	(i)	Which one of the following molecules would be most suitable for such cross-linking?	
		(place a tick in one box only)	
		HOCH ₂ CH ₂ OH	
		H ₂ NCH ₂ CH ₂ NH ₂	

HOCH₂CH₂CO₂H HO₂CCH₂CH₂CO₂H H2NCH2CH2CO2H

(ii) What type of bond would be formed during the cross-linking?

[2]



For
Examiner's
Use

(d)	(i)	Suggest the reagents and conditions needed to hydrolyse polymer monomers.	J into its	
	(ii)	Draw the structural formulae of the two products of this hydrolysis reaction	on.	
			[3]	
(e) The last compound in the list in (c)(i) above is 3-aminopropanoic acid. This can be made from potassium chloroethanoate by the following 3-stage route.				
C <i>I</i> CH₂C0	D ₂ - K	(+ KCN stage II → H ₂ NCH ₂ Cl	H ₂ CO ₂ -K+	
			stage III	

- (i) In the box above write the structure of the intermediate in this route.
- (ii) Suggest reagents and conditions for

stage II

[3]

[Total: 11]

H₂NCH₂CH₂CO₂H



BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

