

CANDIDATE  
NAME

|  |
|--|
|  |
|--|

CENTRE  
NUMBER

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

CANDIDATE  
NUMBER

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

**ADDITIONAL MATHEMATICS**

Paper 2

**0606/22**

**May/June 2014**

**2 hours**

Candidates answer on the Question Paper.

Additional Materials:      Electronic calculator

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES.**

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 80.

This document consists of **14** printed pages and **2** blank pages.

**Mathematical Formulae****1. ALGEBRA***Quadratic Equation*

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

*Binomial Theorem*

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

**2. TRIGONOMETRY***Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

*Formulae for  $\triangle ABC$* 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

- 1 Without using a calculator, express  $\frac{(2 + \sqrt{5})^2}{\sqrt{5} - 1}$  in the form  $a + b\sqrt{5}$ , where  $a$  and  $b$  are constants to be found. [4]

- 
- 2 Find the values of  $k$  for which the line  $y + kx - 2 = 0$  is a tangent to the curve  $y = 2x^2 - 9x + 4$ . [5]

3 (i) Given that  $x + 1$  is a factor of  $3x^3 - 14x^2 - 7x + d$ , show that  $d = 10$ . [1]

(ii) Show that  $3x^3 - 14x^2 - 7x + 10$  can be written in the form  $(x + 1)(ax^2 + bx + c)$ , where  $a$ ,  $b$  and  $c$  are constants to be found. [2]

(iii) Hence solve the equation  $3x^3 - 14x^2 - 7x + 10 = 0$ . [2]

- 4 (i) Express  $12x^2 - 6x + 5$  in the form  $p(x - q)^2 + r$ , where  $p$ ,  $q$  and  $r$  are constants to be found. [3]

- (ii) Hence find the greatest value of  $\frac{1}{12x^2 - 6x + 5}$  and state the value of  $x$  at which this occurs. [2]
-

- 5 (i) Find and simplify the first three terms of the expansion, in ascending powers of  $x$ , of  $(1 - 4x)^5$ . [2]

- (ii) The first three terms in the expansion of  $(1 - 4x)^5(1 + ax + bx^2)$  are  $1 - 23x + 222x^2$ . Find the value of each of the constants  $a$  and  $b$ . [4]

6 (a) (i) State the value of  $u$  for which  $\lg u = 0$ . [1]

(ii) Hence solve  $\lg|2x + 3| = 0$ . [2]

(b) Express  $2 \log_3 15 - (\log_a 5)(\log_3 a)$ , where  $a > 1$ , as a single logarithm to base 3. [4]

7 Differentiate with respect to  $x$

(i)  $x^4 e^{3x}$ , [2]

(ii)  $\ln(2 + \cos x)$ , [2]

(iii)  $\frac{\sin x}{1 + \sqrt{x}}$ . [3]



- 8 The line  $y = x - 5$  meets the curve  $x^2 + y^2 + 2x - 35 = 0$  at the points  $A$  and  $B$ . Find the exact length of  $AB$ . [6]

9 A curve is such that  $\frac{dy}{dx} = (2x + 1)^{\frac{1}{2}}$ . The curve passes through the point (4, 10).


(i) Find the equation of the curve. [4]

(ii) Find  $\int y dx$  and hence evaluate  $\int_0^{1.5} y dx$ . [5]

10 Two variables  $x$  and  $y$  are connected by the relationship  $y = Ab^x$ , where  $A$  and  $b$  are constants.

- (i) Transform the relationship  $y = Ab^x$  into a straight line form. [2]

An experiment was carried out measuring values of  $y$  for certain values of  $x$ . The values of  $\ln y$  and  $x$  were plotted and a line of best fit was drawn. The graph is shown on the grid below.



- (ii) Use the graph to determine the value of  $A$  and the value of  $b$ , giving each to 1 significant figure. [4]

- (iii) Find  $x$  when  $y = 220$ . [2]

**11** The functions  $f$  and  $g$  are defined, for real values of  $x$  greater than 2, by

$$f(x) = 2^x - 1,$$

$$g(x) = x(x + 1).$$

**(i)** State the range of  $f$ .

[1]

**(ii)** Find an expression for  $f^{-1}(x)$ , stating its domain and range.

[4]

- (iii) Find an expression for  $gf(x)$  and explain why the equation  $gf(x) = 0$  has no solutions. [4]
-

**12** A curve has equation  $y = x^3 - 9x^2 + 24x$ .

- (i) Find the set of values of  $x$  for which  $\frac{dy}{dx} \geq 0$ . [4]

The normal to the curve at the point on the curve where  $x = 3$  cuts the  $y$ -axis at the point  $P$ .

- (ii) Find the equation of the normal and the coordinates of  $P$ . [5]



**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.