

NUMBER



ADDITIONAL MATHEMATICS

0606/12

Paper 1

NAME

CENTRE

NUMBER

October/November 2011

2 hours

Candidates answer on the Question Paper.

Additional Materials:

Electronic calculator

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 80.

For Exam	iner's Use
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
Total	

This document consists of 16 printed pages.



Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \cdot$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where *n* is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$.

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\csc^2 A = 1 + \cot^2 A$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^{2} = b^{2} + c^{2} - 2bc \cos A$$
$$\Delta = \frac{1}{2} bc \sin A$$

1 Show that $\frac{1}{\tan \theta + \cot \theta} = \sin \theta \cos \theta$.


[3] For Examiner's Use

2 Find the coordinates of the points where the line 2y = x - 1 meets the curve $x^2 + y^2 = 29$. [5]


For Examiner's Use

	[1]	For Examiner's Use
Using the result of part (i), and the substitution $u = \log_2 x$, find the values of x which sat the equation $\log_2 x = 3 - 2 \log 2$.		
1	Using the result of part (i), and the substitution $u = \log_2 x$, find the values of x which satithe equation $\log_2 x = 3 - 2 \log_x 2$.	Using the result of part (i), and the substitution $u = \log_2 x$, find the values of x which satisfy the equation $\log_2 x = 3 - 2 \log_x 2$. [4]

	U	
4	A curve has equation $y = (3x^2 + 15)^{\frac{2}{3}}$. Find the equation of the normal to the curve at the point where $x = 2$.	For Examiner
		Use
		1



For Examiner's Use



(i) Express
$$y^2$$
 in terms of 2^x .

(ii) Find the value of
$$x$$
 when $y = 6$.

6	(i)	Given that	$(3+x)^5 + (3-x)^5 = A + Bx^2 + Cx^4$, find the value of A, of B and of C.	[4]	For
					Examiner's
					IIco

(ii) Hence, using the substitution $y = x^2$, solve, for x, the equation

$$(3+x)^5 + (3-x)^5 = 1086.$$
 [4]

7 (i) Show that $\frac{\left(4-\sqrt{x}\right)^2}{\sqrt{x}}$ can be written in the form $px^{-\frac{1}{2}}+q+rx^{\frac{1}{2}}$, where p, q and r are integers to be found. [3]

For Examiner's Use

(ii) A curve is such that $\frac{dy}{dx} = \frac{\left(4 - \sqrt{x}\right)^2}{\sqrt{x}}$ for x > 0. Given that the curve passes through the point (9, 30), find the equation of the curve. [5]

© UCLES 2011 0606/12/O/N/11 **[Turn over**

8 The line CD is the perpendicular bisector of the line joining the point A(-1, -5) and the point B(5,3).

For Examiner's Use

(i) Find the equation of the line *CD*.

[4]

(ii)			
	find the coordinates of \hat{D} .	[3]	Examiner's
			Use
(***)		50 3	
(iii)	Find the area of the triangle <i>CAD</i> .	[2]	

9 (i) Given that $y = x \sin 4x$, find $\frac{dy}{dx}$.

[3] For Examiner's Use

(ii) Hence find $\int x \cos 4x \, dx$ and evaluate $\int_0^{\frac{\pi}{8}} x \cos 4x \, dx$.

[6]

10 (i) Solve $2 \sec^2 x = 5 \tan x + 5$, for $0^\circ < x < 360^\circ$.

For Examiner's Use

[5]

(ii) Solve
$$\sqrt{2} \sin \left(\frac{y}{2} + \frac{\pi}{3} \right) = 1$$
, for $0 < y < 4\pi$ radians.

[5]

11 Answer only **one** of the following two alternatives.

For
Examiner's
$II_{S\rho}$

EITHER

A curve has equation $y = e^{-x} (A\cos 2x + B\sin 2x)$. At the point (0, 4) on the curve, the gradient of the tangent is 6.

- (i) Find the value of A. [1]
- (ii) Show that B = 5. [5]
- (iii) Find the value of x, where $0 < x < \frac{\pi}{2}$ radians, for which y has a stationary value. [5]

OR

A curve has equation $y = \frac{\ln(x^2 - 1)}{x^2 - 1}$, for x > 1.

- (i) Show that $\frac{dy}{dx} = \frac{kx(1 \ln(x^2 1))}{(x^2 1)^2}$, where k is a constant to be found. [4]
- (ii) Hence find the approximate change in y when x increases from $\sqrt{5}$ to $\sqrt{5} + p$, where p is small. [2]
- (iii) Find, in terms of e, the coordinates of the stationary point on the curve. [5]

Start your answer to Question 11 here.				
Indicate which question you are answering.	EITHER			
indicate which question you are answering.	OR			

Continue your answer to Question 11 here.	For Examiner's Use

Continue your answer here if necessary.	For
	Examiner's Use

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.