


AS Final Exam : Revision 6 Vectors

P1 June 08

- 10 Relative to an origin O , the position vectors of points A and B are $2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$ and $3\mathbf{i} - 2\mathbf{j} + p\mathbf{k}$ respectively.
- (i) Find the value of p for which OA and OB are perpendicular. [2]
- (ii) In the case where $p = 6$, use a scalar product to find angle AOB , correct to the nearest degree. [3]
- (iii) Express the vector \overrightarrow{AB} in terms of p and hence find the values of p for which the length of AB is 3.5 units. [4]

P1 Nov 08

10



The diagram shows a cube $OABCDEFG$ in which the length of each side is 4 units. The unit vectors \mathbf{i} , \mathbf{j} and \mathbf{k} are parallel to \overrightarrow{OA} , \overrightarrow{OC} and \overrightarrow{OD} respectively. The mid-points of OA and DG are P and Q respectively and R is the centre of the square face $ABFE$.

- (i) Express each of the vectors \overrightarrow{PR} and \overrightarrow{PQ} in terms of \mathbf{i} , \mathbf{j} and \mathbf{k} . [3]
- (ii) Use a scalar product to find angle QPR . [4]
- (iii) Find the perimeter of triangle PQR , giving your answer correct to 1 decimal place. [3]

P1 June 07

9 Relative to an origin O , the position vectors of the points A and B are given by

$$\overrightarrow{OA} = \begin{pmatrix} 4 \\ 1 \\ -2 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OB} = \begin{pmatrix} 3 \\ 2 \\ -4 \end{pmatrix}.$$


(i) Given that C is the point such that $\overrightarrow{AC} = 2\overrightarrow{AB}$, find the unit vector in the direction of \overrightarrow{OC} . [4]

The position vector of the point D is given by $\overrightarrow{OD} = \begin{pmatrix} 1 \\ 4 \\ k \end{pmatrix}$, where k is a constant, and it is given that $\overrightarrow{OD} = m\overrightarrow{OA} + n\overrightarrow{OB}$, where m and n are constants.

(ii) Find the values of m , n and k . [4]


P1 Nov 07

10



The diagram shows a cube $OABCDEFG$ in which the length of each side is 4 units. The unit vectors \mathbf{i} , \mathbf{j} and \mathbf{k} are parallel to \overrightarrow{OA} , \overrightarrow{OG} and \overrightarrow{OD} respectively. The mid-points of OA and DG are P and Q respectively and R is the centre of the square face $ABFE$.

- (i) Express each of the vectors \overrightarrow{PR} and \overrightarrow{PQ} in terms of \mathbf{i} , \mathbf{j} and \mathbf{k} . [3]
- (ii) Use a scalar product to find angle QPR . [4]
- (iii) Find the perimeter of triangle PQR , giving your answer correct to 1 decimal place. [3]



The diagram shows the roof of a house. The base of the roof, $OABC$, is rectangular and horizontal with $OA = CB = 14$ m and $OC = AB = 8$ m. The top of the roof DE is 5 m above the base and $DE = 6$ m. The sloping edges OD , CD , AE and BE are all equal in length.

Unit vectors \mathbf{i} and \mathbf{j} are parallel to OA and OC respectively and the unit vector \mathbf{k} is vertically upwards.

(i) Express the vector \overrightarrow{OD} in terms of \mathbf{i} , \mathbf{j} and \mathbf{k} , and find its magnitude. [4]

(ii) Use a scalar product to find angle DOB . [4]

P1 Nov 06

4 The position vectors of points A and B are $\begin{pmatrix} -3 \\ 6 \\ 3 \end{pmatrix}$ and $\begin{pmatrix} -1 \\ 2 \\ 4 \end{pmatrix}$ respectively, relative to an origin O .

(i) Calculate angle AOB . [3]

(ii) The point C is such that $\overrightarrow{AC} = 3\overrightarrow{AB}$. Find the unit vector in the direction of \overrightarrow{OC} . [4]

P1 June 05

11 Relative to an origin O , the position vectors of the points A and B are given by

$$\overrightarrow{OA} = 2\mathbf{i} + 3\mathbf{j} - \mathbf{k} \quad \text{and} \quad \overrightarrow{OB} = 4\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}.$$

(i) Use a scalar product to find angle AOB , correct to the nearest degree. [4]

(ii) Find the unit vector in the direction of \overrightarrow{AB} . [3]

(iii) The point C is such that $\overrightarrow{OC} = 6\mathbf{j} + p\mathbf{k}$, where p is a constant. Given that the lengths of \overrightarrow{AB} and \overrightarrow{AC} are equal, find the possible values of p . [4]

P1 Nov 05

- 4 Relative to an origin O , the position vectors of points P and Q are given by

$$\overrightarrow{OP} = \begin{pmatrix} -2 \\ 3 \\ 1 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OQ} = \begin{pmatrix} 2 \\ 1 \\ q \end{pmatrix},$$

where q is a constant.

- (i) In the case where $q = 3$, use a scalar product to show that $\cos POQ = \frac{1}{7}$. [3]
- (ii) Find the values of q for which the length of \overrightarrow{PQ} is 6 units. [4]

P1 June 04

- 9 Relative to an origin O , the position vectors of the points A , B , C and D are given by

$$\overrightarrow{OA} = \begin{pmatrix} 1 \\ 3 \\ -1 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 3 \\ -1 \\ 3 \end{pmatrix}, \quad \overrightarrow{OC} = \begin{pmatrix} 4 \\ 2 \\ p \end{pmatrix} \quad \text{and} \quad \overrightarrow{OD} = \begin{pmatrix} -1 \\ 0 \\ q \end{pmatrix},$$

where p and q are constants. Find


- (i) the unit vector in the direction of \overrightarrow{AB} , [3]
- (ii) the value of p for which angle $AOC = 90^\circ$, [3]
- (iii) the values of q for which the length of \overrightarrow{AD} is 7 units. [4]

P1 Nov 04

- 8 The points A and B have position vectors $\mathbf{i} + 7\mathbf{j} + 2\mathbf{k}$ and $-5\mathbf{i} + 5\mathbf{j} + 6\mathbf{k}$ respectively, relative to an origin O .
- (i) Use a scalar product to calculate angle AOB , giving your answer in radians correct to 3 significant figures. [4]
- (ii) The point C is such that $\overrightarrow{AB} = 2\overrightarrow{BC}$. Find the unit vector in the direction of \overrightarrow{OC} . [4]

P1 June 03

- 8 The points A , B , C and D have position vectors $3\mathbf{i} + 2\mathbf{k}$, $2\mathbf{i} - 2\mathbf{j} + 5\mathbf{k}$, $2\mathbf{j} + 7\mathbf{k}$ and $-2\mathbf{i} + 10\mathbf{j} + 7\mathbf{k}$ respectively.
- (i) Use a scalar product to show that BA and BC are perpendicular. [4]
- (ii) Show that BC and AD are parallel and find the ratio of the length of BC to the length of AD . [4]




The diagram shows a triangular prism with a horizontal rectangular base $ADFC$, where $CF = 12$ units and $DF = 6$ units. The vertical ends ABC and DEF are isosceles triangles with $AB = BC = 5$ units. The mid-points of BE and DF are M and N respectively. The origin O is at the mid-point of AC .

Unit vectors \mathbf{i} , \mathbf{j} and \mathbf{k} are parallel to OC , ON and OB respectively.

- (i) Find the length of OB . [1]
- (ii) Express each of the vectors \overrightarrow{MC} and \overrightarrow{MN} in terms of \mathbf{i} , \mathbf{j} and \mathbf{k} . [3]
- (iii) Evaluate $\overrightarrow{MC} \cdot \overrightarrow{MN}$ and hence find angle CMN , giving your answer correct to the nearest degree. [4]

P1 June 02



The diagram shows a solid cylinder standing on a horizontal circular base, centre O and radius 4 units. The line BA is a diameter and the radius OC is at 90° to OA . Points O' , A' , B' and C' lie on the upper surface of the cylinder such that OO' , AA' , BB' and CC' are all vertical and of length 12 units. The mid-point of BB' is M .

Unit vectors \mathbf{i} , \mathbf{j} and \mathbf{k} are parallel to OA , OC and OO' respectively.

(i) Express each of the vectors \overrightarrow{MO} and $\overrightarrow{MC'}$ in terms of \mathbf{i} , \mathbf{j} and \mathbf{k} . [3]

(ii) Hence find the angle OMC' . [4]

P1 Nov 02

7 Given that $\mathbf{a} = \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} 2 \\ 6 \\ 3 \end{pmatrix}$ and $\mathbf{c} = \begin{pmatrix} p \\ p \\ p+1 \end{pmatrix}$, find

(i) the angle between the directions of \mathbf{a} and \mathbf{b} , [4]

(ii) the value of p for which \mathbf{b} and \mathbf{c} are perpendicular. [3]