

CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level

MARK SCHEME for the May/June 2013 series

9709 MATHEMATICS

9709/22

Paper 2, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – May/June 2013	9709	22

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol ∇ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – May/June 2013	9709	22

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only - often written by a “fortuitous” answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through ✓” marks. MR is not applied when the candidate misreads his own figures - this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – May/June 2013	9709	22

- 1 Integrate and obtain term of the form $k \ln(7 - 2x)$ M1
 State $y = -2 \ln(7 - 2x) (+c)$ A1
 Evaluate c DM1
 Obtain answer $y = -2 \ln(7 - 2x) + 2$ A1✓ [4]
- 2 Either State or imply non-modular inequality $(x - 8)^2 > (2x - 4)^2$, or M1
 corresponding equation or pair of linear equations M1
 Make reasonable solution attempt at a quadratic, or solve two linear equations A1
 Obtain critical values 4 and -4 A1
 State correct answer $-4 < x < 4$
- Or Obtain one critical value, e.g. $x = 4$, by solving a linear equation (or inequality) or B1
 from a graphical method or by inspection B2
 Obtain the other critical value similarly B1 [4]
 State correct answer $-4 < x < 4$
- 3 (i) Substitute $x = -1$ and equate to zero M1
 Obtain answer $a = 7$ A1 [2]
- (ii) Substitute $x = -3$ and evaluate expression M1
 Obtain answer 18 A1 [2]
- 4 (i) State or imply $(y + 1) \log 5 = 3x \log 2$ M1
 State that this is of the form $ay = bx + c$ and thus a straight line, or equivalent A1 [2]
- (ii) State gradient is $\frac{3 \ln 2}{\ln 5}$, or equivalent, e.g. $3 \log_5 2$ B1
 State $(0, -1)$ B1 [2]
- 5 (i) State $3 \frac{dy}{dx}$ as derivative of $3y$, or equivalent B1
 State $4xy + 2x^2 \frac{dy}{dx}$ as a derivative of $2x^2y$, or equivalent B1
 Equate derivative of LHS to zero and solve for $\frac{dy}{dx}$ M1
 Obtain given answer correctly A1 [4]
- (ii) Substitute $x = 2$ into given equation and solve for y M1
 Obtain gradient $= \frac{12}{5}$ correctly A1
 Form equation of the normal at their point, using negative recip of their $\frac{dy}{dx}$ M1
 State correct equation of normal $5x + 12y + 2 = 0$ or equivalent A1 [4]
- 6 (i) Make a recognisable sketch of a relevant graph, e.g. $y = 3e^x$ or $y = 8 - 2x$ B1
 Sketch a second relevant graph and justify the given statement B1 [2]

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – May/June 2013	9709	22

- (ii) Consider sign of $3e^x - 8 + 2x$ at $x = 0.7$ and $x = 0.8$, or equivalent M1
 Complete the argument correctly with appropriate calculations A1 [2]
 ($f(0.7) = -0.559$, $f(0.8) = 0.277$ or equivalent)

- (iii) Show that given equation is equivalent to $x = \ln\left(\frac{8-2x}{3}\right)$, or vice versa B1 [1]

- (iv) Use the iterative formula correctly at least once M1
 Obtain final answer 0.768 A1
 Show sufficient iterations to justify its accuracy to 3 d.p.

$x_0 = 0.7$	$x_0 = 0.75$	$x_0 = 0.8$
0.78846	0.77319	0.75769
0.76129	0.76603	0.77082
0.76971	0.76825	0.76676
0.76711	0.76756	0.76802
0.76791		0.76763
0.76766		

or show there is a sign change in the interval (0.7675, 0.7685) B1 [3]

- 7 (a) Obtain one term of form ke^{2x-1} with any non-zero k M1
 Obtain correct integral $x + \frac{1}{2}e^{2x-1}$ A1
 Substitute limits, giving exact values M1
 Correct answer $\frac{1}{2}e^3 + 1$ A1 [4]

- (b) Use product or quotient rule M1*
 Obtain correct derivative in any form A1
 Equate derivative to zero and solve for x M1*
 Obtain $\tan 2x = 1$ dep
 Obtain $x = \frac{\pi}{8}$ A1 [5]

- 8 (i) Use correct $\sin(A - B)$ and $\cos(A - B)$ formula M1
 Substitute exact values for $\cos 30^\circ$ etc. M1
 Obtain given answer correctly A1 [3]

- (ii) State $2\operatorname{cosec} x = 3\cot^2 x - 2$ B1
 Use $\cot^2 x = \operatorname{cosec}^2 x - 1$ M1
 Attempt solution of quadratic equation in $\operatorname{cosec} x$ or $\sin x$ M1
 ($3\operatorname{cosec}^2 x - 2\operatorname{cosec} x - 5 = 0$ or $5\sin^2 x = 2\sin x - 3 = 0$)
 Obtain $\sin x = \frac{3}{5}$ or -1 A1✓
 Obtain one correct answer for $\sin^{-1}\left(\frac{3}{5}\right)$ A1
 Obtain remaining 2 answers from 36.9° , 143.1° , 270° and no others in the range
 [Ignore answers outside the given range] A1 [6]
 SC If only answer given is 270° B1