

**UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS**

GCE Advanced/Advanced Subsidiary Level

**MARK SCHEME for the May/June 2006 question paper**

**9709 MATHEMATICS**

**9709/07**

**Paper 7**

**Maximum raw mark 50**

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were initially instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the *Report on the Examination*.

The minimum marks in these components needed for various grades were previously published with these mark schemes, but are now instead included in the Report on the Examination for this session.

- CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2006 question papers for most IGCSE and GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

## Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
  - A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
  - B Mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
  - The symbol  $\surd$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
  - Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking  $g$  equal to 9.8 or 9.81 instead of 10.

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only - often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

### **Penalties**

MR -1	A penalty of MR -1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures - this is regarded as an error in accuracy. An MR-2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA -1	This is deducted from A or B marks in the case of premature approximation. The PA -1 penalty is usually discussed at the meeting.

Page 1	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – May/June 2006	9709	07

<b>1</b> $99.2 \pm 2.576 \times \frac{2.3}{\sqrt{200}}$ $= (98.8, 99.6)$	M1 M1 A1 <b>3</b>	One of $99.2 + zs/\sqrt{n}$ or $99.2 - zs/\sqrt{n}$ seen For $z = 2.576$ or $z = 2.326$ rounding to correct answer or equivalent
<b>2</b> mean $= 0.75 \times 53.2 + 0.25 \times 78$ $= 59.4$ var $= 0.75^2 \times 9.3^2 + 0.25^2 \times 5.1^2$ $= 50.27$ sd $= 7.09$	B1 M1 M1 A1 <b>4</b>	Correct answer $0.75^2$ and $0.25^2$ multiplied by var or sd $0.75$ or $0.75^2$ mult by $9.3^2$ and $0.25$ or $0.25^2$ mult by $5.1^2$ correct answer
<b>3 (i)</b> Normal mean 6 variance $3.6/120$ (0.03)	B1 B1 B1 <b>3</b>	Correct mean Correct variance
<b>(ii)</b> $P(\bar{X} > 6.1) = 1 - \Phi\left(\frac{6.1 - 6}{\sqrt{3.6/120}}\right)$ $= 1 - \Phi(0.5773)$ $= 1 - 0.7181$ $= 0.282$	M1 M1 A1 <b>3</b>	Standardising must have $\sqrt{120}$ Correct area ie $< 0.5$ Correct answer
<b>4</b> $D - W \sim N(4.2, 6.3^2 + 4.8^2)$ $P(D - W > 3) = 1 - \Phi\left(\frac{3 - 4.2}{\sqrt{62.73}}\right)$ $= \Phi(0.152)$ $= 0.560$  $P(D - W < -3) = \Phi\left(\frac{-3 - 4.2}{\sqrt{62.73}}\right)$ $= 1 - \Phi(0.909)$ $= 1 - 0.8182$ $= 0.182$ total prob $= 0.742$	B1 M1* M1 dep* M1 A1 A1 <b>6</b>	$6.3^2 + 4.8^2$ or 62.7 seen considering $P(D - W > 3)$ or considering $P(W - D > 3)$ , or $P(D - W < -3)$ and standardising correct prob area for their prob considered above ie $> 0.5$ or $< 0.5$ depending Considering the other version not considered above 0.560 or 0.182 seen correct answer

Page 2	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – May/June 2006	9709	07

<b>5 (i)</b> $\int_0^1 4x^k dx = 1$ $\left[ \frac{4x^{k+1}}{k+1} \right]_0^1 = 1$ $k = 3$ <b>AG</b>	M1	Equating to 1 and attempting to integrate
	A1 <b>2</b>	Correct answer legitimately obtained
<b>(ii)</b> $E(X) = \int_0^1 4x^4 dx$ $= \left[ \frac{4x^5}{5} \right]_0^1 = 0.8$ <b>AG</b> $\text{Var}(X) = \int_0^1 4x^5 dx - 0.8^2$ $= \left[ \frac{4x^6}{6} \right]_0^1 - 0.8^2 = 0.0267$	M1	Attempting to evaluate $\int_0^1 4x^4 dx$ with or without limits
	A1	Correct answer legitimately obtained
	M1	Attempt at integral of $x^2 f(x) - 0.8^2$
	A1 <b>4</b>	Correct answer (accept 0.027)
<b>(iii)</b> $\int_0^{q_3} 4x^k dx = 0.75$ $q_3 = \sqrt[4]{0.75} \quad (= 0.931)$ $= 0.931$	M1	Finding UQ by solving integral = 0.75
	A1ft <b>2</b>	Correct UQ
<b>(iv)</b> $q_1 = \sqrt[4]{0.25} \quad (= 0.707)$ <b>IQ range</b> = 0.223 ( 0.22349)	B1	Correct LQ
	B1ft <b>2</b>	Accept 0.223 or 0.224



Page 3	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – May/June 2006	9709	07

<b>6 (i)</b> $1 - e^{-0.6}(1 + 0.6)$ $= 1 - 0.878$ $= 0.122$	M1 B1 A1	3	Poisson calc $1 - P(0, 1)$ Correct numerical expression for $P(0, 1)$ Correct answer
<b>(ii)</b> $(e^{-0.6})(0.97)$ $= 0.532$	M1 B1	2	Multiplying $P(0)$ for skirt by 0.97 Correct answer
<b>(iii)</b> $P(F, F) = (1 - e^{-0.6})(0.03)$ $= 0.0135 (0.01354)$	M1 A1	2	Finding $P(F, F)$ Correct answer
<b>(iv)</b> $X$ approx $P(300 \times 0.01354) \sim P(4.062)$ $P(X < 3) = e^{-4.062} \left( 1 + 4.062 + \frac{4.062^2}{2} \right)$ $= 0.229$	M1 A1ft A1	3	Using poisson approx, $\lambda = 300 \times$ their $P(F, F)$ Correct numerical expression for $P(X < 3)$ with their $\lambda$ Correct answer
<b>7 (i)</b> mean = 6.53 (6.525) variance = $\frac{1}{39} \left( 1815 - \frac{261^2}{40} \right)$ $= 2.87$	B1 M1 A1	3	Substituting in formula from tables Correct answer
<b>(ii)</b> $H_1: \mu < 7.2$ test statistic $z = \frac{6.525 - 7.2}{\sqrt{2.871/40}}$ $= -2.52$ critical value $z = \pm 1.645$ or $\pm 1.64$ or $\pm 1.65$ reduction in cars exceeding speed limit	B1 M1 A1 B1 B1ft	5	Correct $H_1$ Standardising attempt must have $\sqrt{40}$ Correct test statistic Correct CV or finding area on LHS of their $z$ Correct conclusion, reject $H_0$ not enough, must compare + with + or – with –
<b>(iii)</b> saying there is a reduction in the number of speeding cars, when there isn't normal curve with mean 7.2 shown and small shaded area on LHS shaded area on LHS labelled as 5% and Type I error.	B1 B1 B1	3	