

MARK SCHEME for the October/November 2013 series

9709 MATHEMATICS

9709/61

Paper 6, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	61

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol ∇ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	61

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only – often written by a ‘fortuitous’ answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through ✓” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	61

5 (i) $z = -1.406$ $\frac{c - 14.2}{3.6} = -1.406$ $c = 9.14$	B1 M1 A1 3	Rounding to ± 1.41 seen Standardising allow sq rt no cc Correct answer
(ii) $P\left(\frac{15 - 14.2}{3.6}\right) < z < \left(\frac{16 - 14.2}{3.6}\right)$ $= \Phi(0.5) - \Phi(0.222)$ $= 0.6915 - 0.5879$ $= 0.1036$ $P(\text{at least } 2) = 1 - P(0, 1)$ $= 1 - (0.8964)^7 - (0.8964)^6(0.1036)$ $= 1 - 0.8413$ $= 0.159$	M1 M1 A1 M1 M1 A1 6	2 attempts at standardising no cc no sq rt Subt two Φ s (indep mark) Needn't be entirely accurate, rounding to 0.10 Binomial term with ${}^7C_r p^r (1-p)^{7-r}$ seen $r \neq 0$ any $p < 1$ $1 - P(0)$, $1 - P(1)$, $1 - P(0, 1)$ seen their p Correct answer accept 3sf rounding to 0.16
6 (i) M R O 3 1 2 = ${}^7C_3 \times {}^5C_1 \times {}^8C_2 = 4900$ 3 2 1 = ${}^7C_3 \times {}^5C_2 \times {}^8C_1 = 2800$ 2 2 2 = ${}^7C_2 \times {}^5C_2 \times {}^8C_2 = 5880$ Total = 13580	M1 M1 A1 A1 4	Summing more than one 3term option involving combs (can be added) Mult 3 combs only (indep) 1 option correct unsimplified Correct answer
(ii) 4 groups in 4! ways 3 mountain in 3! ways 2 ordinary in 2! ways $4! \times 3! \times 2 = 288$	M1 M1 A1 3	4! seen mult by something Mult by 3! for racing or 2! for ordinary Correct answer
(iii) e.g. s O x x x x O s s s Ordinary in 2! Rest of bikes in 4! Bikes and spaces 5 groups in 5 ways $2! \times 4! \times 5 = 240$	M1 M1 A1 3	2! or 4! seen mult Mult by 5 (ssssb) Correct answer

Page 6	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	61

7	(i) if throw H then smallest score is 2 $P(T, 1) = 1/2 \times 1/4 = 1/8$ AG	B1 B1	2	Or equivalent																		
	(ii) $P(3)$ from two dice = 2/16 seen $P(H, 3) = 1/2 \times 2/16 = 2/32$ $P(T, 3) = 1/2 \times 1/4 = 1/8$ So $P(3) = 6/32 = 3/16$ AG	B1 M1 A1 A1	 4	From (1, 2) and (2, 1) Summing $P(H, 3)$ and $P(T, 3)$ One correct Correct answer must see clear reasoning																		
	(iii) <table border="1"><tr><td>X</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td><td>8</td></tr><tr><td>Prob</td><td></td><td>5/32</td><td></td><td>7/32</td><td></td><td>3/32</td><td></td><td></td></tr></table>	X	1	2	3	4	5	6	7	8	Prob		5/32		7/32		3/32			B1 B1 B1	 3	One correct prob A second correct prob A third correct prob
	X	1	2	3	4	5	6	7	8													
Prob		5/32		7/32		3/32																
(iv) $P(Q \cap R) = 0$ or ‘if you throw a tail you can’t get a 7’ Yes they are exclusive	M1 A1dep	 2	Stating $P(Q \cap R) = 0$ or implying by words Dep on previous M																			

MARK SCHEME for the October/November 2013 series

9709 MATHEMATICS

9709/62

Paper 6, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	62

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol ∇ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously ‘correct’ answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	62

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no 'follow through' from a previous error is allowed)
CWO	Correct Working Only – often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become 'follow through' marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	62

1	$P(x < -2.4) = P\left(z < \frac{-2.4 - 1.5}{3.2}\right)$	M1		Standardising no cc can have sq
	$= P(z < -1.219)$	M1		Correct area, i.e. < 0.5
	$= 1 - 0.8886$ $= 0.111$	A1	[3]	Correct answer rounding to 0.111
2	(i) $P(C \cap < 50) = 0.35 \times 0.2 = 0.07$	B1	[1]	
	(ii) $P(C \mid < 50) = \frac{P(C \cap < 50)}{P(< 50)}$	M1	[4]	Summing three 2-factor products seen anywhere (can omit the 1)
	$= \frac{0.35 \times 0.2}{0.25 \times 0.3 + 0.35 \times 0.2 + 0.4(\times 1)}$	A1		0.545 (unsimplified) seen as num or denom of a fraction
	$= \frac{0.07}{0.545}$	M1		Attempt at $P(C \cap < 50)$ as 2-factor prod only seen as num or denom of a fraction
	$= 0.128 \text{ (14/109)}$	A1		Correct answer
3	(i) $z = 0.878$ $\frac{190 - 160}{\sigma} = 0.878$ $\sigma = 34.2$	B1		$\pm 0.878, 0.88$, rounding to 0.88 seen $(190 - 160)/\sigma = \text{something}$
		M1		
		A1	[3]	Correct answer
	(ii) $P(\text{at least } 1) = 1 - P(0)$	M1		Using $1 - P(0)$, $1 - P(0, 1)$, $P(1, 2 \dots 12)$ or $P(2, \dots 12)$ with $p = 0.19$ or 0.81 , terms must be evaluated to get the M1
	$= 1 - (0.81)^{12} = 0.920$	A1	[2]	Correct answer accept 0.92
4	(i) number = $1.5 \times 50 = 75$ (AG)	B1	[1]	Must see 1.5×50
	(ii) freqs are 10, 25, 50, 75, 30 (15, 15)	M1		Attempt at freqs not fd
		A1		Correct freqs
	Mean = $(10 \times 125 + 25 \times 162.5 + 50 \times 187.5 + 75 \times 225 + 30 \times 300)/190$ $= 40562.5/190 = 213 \text{ (213.48 ...)}$	M1		attempt at mid points not cw or ucb or lcb
		A1		correct mean
	$sd^2 = 10 \times 125^2 + 25 \times 162.5^2 + 50 \times 187.5^2 + 75 \times 225^2 + 30 \times 300^2)/190 - (213.48 \dots)^2$	M1		subst their Σfx^2 in correct variance formula
	$sd = 46.5 \text{ or } 46.6$	A1	[6]	
	(iii) have used the mid-point of each interval and not the raw data	B1	[1]	

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	62

5	(i) $P(4, 5, 6) = (0.22)^4(0.78)^4 {}^8C_4 + (0.22)^5(0.78)^3 {}^8C_5 + (0.22)^6(0.78)^2 {}^8C_6$ $= 0.0763$	M1		[3]	Bin term with ${}_8C_r p^r (1-p)^{8-r}$ seen $r \neq 0$ any $p < 1$ Summing 2 or 3 bin probs $p = 0.22$, $n = 8$ Correct answer
		M1			
		A1			
		B1			
		B1ft			
		M1			
	(ii) prob = 0.13 mean = $300 \times 0.13 = 39$ var = $300 \times 0.13 \times 0.87 = 33.93$ $P(30 < x < 50) = P\left(\frac{30.5 - 39}{\sqrt{33.93}} < z < \frac{49.5 - 39}{\sqrt{33.93}}\right)$ $= P(-1.4592 < z < 1.8026)$ $= \Phi(1.8026) + \Phi(1.4592) - 1$ $= 0.9643 + 0.9278 - 1 = 0.892$	M1		[6]	Correct prob can be implied Correct unsimplified np and npq ft wrong 0.13 Standardising a value need sq rt Cont correction 30.5 / 31.5 or 48.5/49.5 only Correct area $\Phi_1 + \Phi_2 - 1$ oe Rounding to correct answer SC $P(31, \dots, 49) = {}^{300}C_{31}(0.13)^{31}(0.87)^{269} + \dots + {}^{300}C_{49}$ etc.) B1B1
		M1			
		M1			
		M1			
		A1			
6	(i) 1663200 (ii) M xxxxxxxxx M Number of ways = $\frac{9!}{3!2!} = 30240$ (iii) 4 vowels together = $8! \times 4/2!2! = 40320$ $1663200 - 40320 = 1622880$ (iv) Exactly 2 Es $4C2 = 6$ Exactly 3 Es $4C1 = 4$ Total = 10 ways OR $5C2 = 10$	B1		[1]	9! or 9P9 seen
		M1			
		A1			
		M1			
		M1			
		B1			
		M1			
		B1			
		A1			
		M2		[3]	Summing 2 options One option correct Correct answer M1 for $k5C2$ Correct ans
		A1			

Page 6	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	62

7	(i) options (3, 4, 4,) or (4, 3, 4) or (4, 4, 3) Probs $(4/10 \times 6/9 \times 5/8) \times 3C1$ $= 360/720$ $= \frac{1}{2}$ AG	M1			Summing three 3-factor options oe
		M1			$10 \times 9 \times 8$ seen in denom
		A1	[3]		Correct answer
	OR $\frac{{}_6C_2 \times {}_4C_1}{{}_{10}C_3} = \frac{1}{2}$ AG	M1			One of $6C2$ or $4C1$ seen in num
		M1			$10C3$ in denom
		A1			Correct answer
	(ii)	B1	[4]		9, 10, 11, 12 only seen
		B1			One correct prob other than P(11), with or without replacement
		B1			Another correct prob
		B1			Σ all 4 probs = 1
		B1			
	(iii) $P(R) = 0.5$ $P(S) = 0.4$ $P(R \cap S) = 120/720$ $P(R \cap S) = 120/720 \neq P(R) \times P(S)$ Not indep	B1	[3]		$P(R \cap S) = 120/720$ (1/6)
		M1			Numerical attempt to compare $P(R \text{ and } S)$ with $P(R) \times P(S)$ provided $P(R \cap S) \neq 1/5$
	(iv) $P(R \cap S) \neq 0$ or there is an overlap between R and S (34,4) Not exclusive $\Sigma xf/\Sigma f$	A1ft			Correct conclusion ft wrong $P(R \cap S) \neq 1/5$, $P(S)$ correct
		B1ft	[1]		Correct answer following correct reasoning ft wrong non zero $P(R \cap S)$

MARK SCHEME for the October/November 2013 series

9709 MATHEMATICS

9709/63

Paper 6, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	63

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol ∇ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously ‘correct’ answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	63

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no 'follow through' from a previous error is allowed)
CWO	Correct Working Only – often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become 'follow through' marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	63

1 bars are not touching oe Area not rep by frequency, not used fd, not labelled fd	B1	Sensible reason involving not touching, no gaps, class boundaries, group data not continuous (may be the negative)
	B1 2	Must be frequency density oe. Wrong height not sufficient. (Best 2 reasons awarded)
2 $P(13.6 < X < 14.8) = P\left(\frac{13.6 - 14}{0.52} < z < \frac{14.8 - 14}{0.52}\right)$ $= P(-0.7692 < z < 1.538)$ $= \Phi(1.538) - [1 - \Phi(0.7692)]$ $= 0.9380 - [1 - 0.7791]$ $= 0.7171$ $P(8) = (0.7171)^8 (0.2829)^2 {}_{10}C_8$ $= 0.252$	M1 M1 A1 M1 A1 5	Standardising 1 expression, no cc, no sq rt, no sq, \pm , mean on num. $\Phi 1 + \Phi 2 - 1$ (indep) oe ($\Phi 2 - \Phi 1$ if cc used) Correct probability rounding to 0.72 here Binomial expression $10C_8 p^8 q^2$, $\Sigma p + q = 1$, any p Correct answer (rounding to 0.252)
3 (i) $(p =)0.85$ $P(< 12) = 1 - P(12, 13, 14)$ $= 1 - [(0.85)^{12} (0.15)^2 {}_{14}C_{12} + (0.85)^{13} (0.15) {}_{14}C_{13} + (0.85)^{14}]$ $= 1 - 0.6479$ $= 0.352$	B1 M1 A1 3	$(p =)0.85$ oe seen anywhere Summing 2 or 3 consistent bin probs, any $p < 1$, $n = 14$ (or summing 12 or 13 consistent bin probs) Correct answer
(ii) $(0.85)^n \geq 0.1$ $n \leq 14.2$ $n = 14$	M1 M1 A1 3	Eqn or inequality in 0.85(or 0.15), n , 0.1, n as a power Attempt to solve (can be implied) if n a power Correct answer – must be equals, not approx. MR allowed for 0.01, M1M1A0 max.
4 (i) $(220 \times 20 + 118 \times 25)/45$ $= 163$	M1 A1 2	Mult by 20 and 25 and dividing their sum by 45 Correct answer, 163.3 or $490/3$ oe acceptable
(ii) $\Sigma x_o^2 / 20 - 220^2 = 32^2$ $\Sigma x_o^2 = 988480$ $\Sigma x_l^2 / 25 - 118^2 = 12^2$ $\Sigma x_l^2 = 351700$ $\Sigma x_o^2 + \Sigma x_l^2 = 1340180$ New var $= 1340180/45 - (7350/45)^2$ $= 3100 - 3120$	M1 A1 A1 M1 A1 5	Subst in correct variance formula Correct Σx_o^2 correct Σx_l^2 Subst their combined results in correct var formula Correct answer

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	63

<p>5 (a) $P(X < q + 82) = 0.72$ $z = 0.583$ $\frac{\pm q}{7.4} \text{ or } \frac{\pm 2q}{7.4} = z \text{ or probability (o.e.)}$ $q = 4.31$</p>	<p>M1 M1 A1 3</p>	<p>Rounding to ± 0.58 or ± 0.15 seen Standardising, no cc, no sq, no sq rt correct answer</p>
<p>(b) $\frac{0.5\mu - \mu}{\sigma} = \frac{\pm 0.5\mu}{\sigma}$ $\frac{0.2\sigma^2}{\sigma} = -0.2\sigma = -0.580$ $\sigma = 2.90$ $\mu = 3.36$</p>	<p>M1 B1 M1 A1 4</p>	<p>Standardising attempt some μ/σ allow cc, sq rt, sq Can be implied ± 0.580 seen (accept ± 0.58) substituting to eliminate μ or σ, arriving at numerical solution, any z value or probability – not dependent both answers correct, accept 2.9</p>
<p>6 (i) $\frac{8!}{3!2!2!}$ $= 1680$</p>	<p>M1 A1 2</p>	<p>8! Divided by at least one of 3!2!2! oe Correct answer</p>
<p>(ii) $5!$ $= 120$</p>	<p>M1 A1 2</p>	<p>5! Seen (not added, may be divided/multiplied) Correct answer</p>
<p>(iii) $\frac{5!4!}{3!2!2!}$ $= 120$</p>	<p>B1 M1 A1 3</p>	<p>5! Or 4! Seen in sum or product in numerator (denominator may be 1) $\frac{k5!4!}{3!2!2!}$ in a numerical expression Correct final answer</p>
<p>(iv) GG with AA, AE, EE, RA, RE, RT, TA, TE, $= 8$ ways GGG with A, E, R, T = 4 ways Total = 12 ways</p>	<p>M1 A1 A1 3</p>	<p>Summing 2 options (could be lists) 1 correct option Correct answer</p>

Page 6	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9709	63

7	<p>(i) $P(\text{same}) = P(1, 1) + P(3, 3) + P(5, 5)$</p> $= \frac{2}{9} \times \frac{1}{8} + \frac{4}{9} \times \frac{3}{8} + \frac{3}{9} \times \frac{2}{8}$ <p>$= 5/18 \text{ (0.278)}$</p> <p>Alt. method:</p> $\frac{2C_2 + 4C_2 + 3C_2}{9C_2}$ <p>or $\frac{2 \times 1 + 3 \times 4 + 2 \times 3}{9C_2 \times 2} \text{ oe}$</p>	M1	Summing 3 two-factor options								
		M1	Multiplying terms by one less in the numerator or denominator								
		A1 3	Correct answer								
			M1 for numerator, M1 for denominator, A1 correct answer								
	<p>(ii) $P(5, \bar{5}) + P(\bar{5}, 5)$</p> $= \frac{3}{9} \times \frac{6}{8} + \frac{6}{9} \times \frac{3}{8} = \frac{36}{72} = \frac{1}{2} \text{ or } 0.5$ <p>Alt. method:</p> $\frac{6C_1 \times 3C_1 (\times 2)}{9C_2 (\times 2)} \text{ oe}$	M1	Mult 2 probs whose numerators sum to 9 o.e.								
		M1	Summing 2 options or mult by 2 (may be 4 options)								
		A1 3	Correct answer								
			M1 for numerator, M1 for denominator, A1 correct answer								
	<p>(iii) $P(5 \cap \bar{5}) = \frac{3}{9} \times \frac{6}{8} = \frac{1}{4}$</p> $P(\bar{5}) = \frac{1}{4} + \frac{6}{9} \times \frac{5}{8} = 48/72 = 0.6666$ $P(5_1 \bar{5}_2) = \frac{1/4}{48/72} = 3/8$ $= 0.375$	M1	Attempt at P(5 and not 5) seen as numerator or denominator of a fraction								
		M1	Attempt at P(not 5) sum of 2 two-factor terms seen anywhere								
		A1	Correct P($\bar{5}$) as numerator or denominator in fraction								
		A1 4	Correct answer								
	<p>(iv)</p> <table border="1"><tr><td>x</td><td>0</td><td>1</td><td>2</td></tr><tr><td>$P(X = x)$</td><td>5/12</td><td>1/2</td><td>1/12</td></tr></table> <p>$P(0) = P(\bar{5}, \bar{5}) = \frac{6}{9} \times \frac{5}{8} = 30/72 \text{ (5/12)}$ (0.4166)</p> <p>$P(1) = 0.5$ from part (ii)</p> <p>$P(2) = 6/72 \text{ (1/12)}$ (0.0833) from part (i)</p>	x	0	1	2	$P(X = x)$	5/12	1/2	1/12	B1	Values 0, 1, 2 seen in table with at least 1 prob
x	0	1	2								
$P(X = x)$	5/12	1/2	1/12								
		B1	Correct P(0) unsimplified								
		B1ft 3	If $x=0,1,2(,3)$ ft $\Sigma p = 1$, no –ve values, all probabilities <1								