



CANDIDATE NAME				
CENTRE NUMBER		CANDIDATE NUMBER		

PHYSICS 9702/34

Advanced Practical Skills 2

May/June 2011

2 hours

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer both questions.

You will be allowed to work with the apparatus for a maximum of one hour for each question.

You are expected to record all your observations as soon as these observations are made, and to plan the presentation of the records so that it is not necessary to make a fair copy of them.

You may lose marks if you do not show your working or if you do not use appropriate units.

Additional answer paper and graph paper should be submitted only if it becomes necessary to do so.

You are reminded of the need for good English and clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use					
1					
2					
Total					

This document consists of 10 printed pages and 2 blank pages.



You may not need to use all of the materials provided.

For Examiner's Use

- 1 In this experiment, you will investigate the angle through which a loaded beaker rolls as a turning force is applied.
 - (a) The apparatus has been assembled as shown in Fig. 1.1.




Fig. 1.1

- (b) Make sure that the beaker is positioned so that the masses do not touch the rails.
- (c) Using the set square, measure and record the angle x, as shown in Fig. 1.2.




Fig. 1.2

 $x = \dots [1]$

(d)	(i)	Hook the mass	hanger	on	the	string	loop.	Record	the	mass	m that	is	suspend	led
		from the loop.												

For Examiner's Use

m =	

- (ii) Wait for the beaker to stop moving, making sure that the beaker is positioned so that the masses do not touch the rails.
- (iii) Using the set square, measure and record the angle y, as shown in Fig. 1.3.




Fig. 1.3

<i>y</i> =	 [1]]
,		

(iv) Calculate θ , where $\theta = y - x$.

$\theta =$	
<i>U</i> –	

(e)	Rep and	ange m by adding masses to the hanger and repeat (d)(ii), (d)(iii) and (d)(iv). Deat this procedure until you have six sets of values for m (the total suspended matangle y . Under the values for θ (using your answer from (c)) and θ .	ass)	For Examiner's Use
			[9]	
(f)	(i)	Plot a graph of $\sin \theta$ on the <i>y</i> -axis against <i>m</i> on the <i>x</i> -axis.	[3]	
	(ii)	Draw the straight line of best fit.	[1]	
	(iii)	Determine the gradient and <i>y</i> -intercept of this line.		
		gradient =		
		<i>y</i> -intercept =	[2]	

5

For Examiner's Use

(g)	(i)	Unhook the masses from the string loop and remove the beaker from the rails.	For
	(ii)	Take measurements to determine the radius r of the beaker.	Examiner's Use
		r =[1]	
(h)	It is	suggested that the relationship between θ and m is	
		$\sin\theta = \frac{r m}{a} + b$	
	whe	ere a and b are constants.	
	Usi unit	ng your answers from (f)(iii) and (g)(ii) , determine the value of <i>a</i> . Give an appropriate .	
		a =[2]	
		a =[2]	

You may not need to use all of the materials provided.

For Examiner's Use

- 2 In this experiment you will investigate the motion of a mass suspended from a rubber band.
 - (a) (i) Suspend the rubber band from the retort stand and hang the 100g mass hanger from the rubber band, as shown in Fig. 2.1




Fig. 2.1

(ii)	Determine	and	record	the	radius	R	of the	suspended	mass	hanger	at	its	widest
	point.												

R =[2]

(iii) Estimate the percentage uncertainty in *R*.

percentage uncertainty =[1]

8 (b) (i) Twist the mass hanger about half a turn and release it so that it turns between For positions A and B, as shown in Fig. 2.2. Examiner's Use Fig. 2.2 (ii) Take measurements to determine the time T for the mass hanger to rotate from A to B and back to A. (This may be determined accurately by using the time for several turns.) $T = \dots [2]$ (c) For a mass hanger of mass m and radius R, it is suggested that T is related to a quantity C, where $C = mR^2$. Calculate the value of C for this mass hanger. Give an appropriate unit. *C* =[1]

(d)	(i) Remove the 100 g mass hanger and suspend the 50 g mass hanger from the rubb band.					
	(ii)	Repeat (a)(ii), (b) and (c) for this new suspended mass hanger.	Use			
		R =				
		K =				
		<i>T</i> =				
		<i>i</i> –				
		$C = \dots $ [4]				
(e)	(i)	It is suggested that the relationship between <i>T</i> and <i>C</i> is				
(-)	(-)	$T^2 = k C$				
		where <i>k</i> is a constant.				
		Using your data, calculate two values of <i>k</i> .				
		first value of <i>k</i> =				
		second value of $k = \dots$				
		Second value of k =[1]				

	(ii)	Explain whether your results support the suggested relationship in (e)(i) .	For Examiner's Use
		[1]	
(f)	(i)	Describe four sources of uncertainty or limitations of the procedure for this experiment.	
		1	
		2	
		3	
		4	
		[4]	
	(ii)	Describe four improvements that could be made to this experiment. You may suggest the use of other apparatus or different procedures.	
		1	
		2	
		3	
		4	

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.