

Concentration of Measure

Sudeep Kamath



CIRM workshop, 25 Jan 2016

What is concentration?

“A random variable that depends in a smooth way on many independent random variables (but not too much on any of them) is essentially constant.”

- M. Talagrand, 1996.

What is concentration?

“A random variable that depends in a smooth way on many independent random variables (but not too much on any of them) is essentially constant.”

- M. Talagrand, 1996.

If Z is a function of many independent variables X_1, X_2, \dots, X_n ,
how large are typical deviations of Z ?

What is concentration?

“A random variable that depends in a smooth way on many independent random variables (but not too much on any of them) is essentially constant.”

- M. Talagrand, 1996.

If Z is a function of many independent variables X_1, X_2, \dots, X_n ,
how large are typical deviations of Z ?

Goal: Quantify by bounding for $t > 0$,

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t]$$

Applications

Concentration of measure has far-reaching consequences in

- Pure and applied probability,
- High-dimensional statistics,
- Functional analysis,
- Computer science,
- Machine learning,
- Statistical physics,
- Complex graphs and networks,
- Information theory, communication and coding theory.


Approaches for Proving Concentration


- *The martingale approach*: Hoeffding (1963), Azuma (1967), Milman and Schechtman (1986), Shamir and Spencer (1987) and McDiarmid (1989, 1998), Sipser and Spielman (1996), Richardson and Urbanke (2001)
- *Talagrand's inequalities for product measures*: Talagrand (1996).
- *Entropy method and log-Sobolev inequalities*: Ledoux (1996), Massart (1998), Lugosi et al. (1999, 2001)
- *Transportation method*: Ahlswede, Gács and Körner (1976), Marton (1986, 1996, 1997), Dembo (1997), Villani (2003, 2008)
- *Stein's method of exchangeable pairs*: Chatterjee (2007), Chatterjee and Dey (2010), Goldstein et al. (2011, 2014)

Approaches for Proving Concentration


- *The martingale approach*: Hoeffding (1963), Azuma (1967), Milman and Schechtman (1986), Shamir and Spencer (1987) and McDiarmid (1989, 1998), Sipser and Spielman (1996), Richardson and Urbanke (2001)
- *Talagrand's inequalities for product measures*: Talagrand (1996).
- *Entropy method and log-Sobolev inequalities*: Ledoux (1996), Massart (1998), Lugosi et al. (1999, 2001)
- *Transportation method*: Ahlswede, Gács and Körner (1976), Marton (1986, 1996, 1997), Dembo (1997), Villani (2003, 2008)
- *Stein's method of exchangeable pairs*: Chatterjee (2007), Chatterjee and Dey (2010), Goldstein et al. (2011, 2014)

We will focus on the entropy method and transportation method where information theoretic methods shine.







Undergraduate ("informal")	




Undergraduate ("informal")	Graduate (formal probability)




Undergraduate ("informal")	Graduate (formal probability)
Probability density,	Radon-Nikodym derivative,




Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral




Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability




Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability
$\mathbb{E}[X Y]$ conditioning on random variables	$\mathbb{E}[X \sigma(Y)]$ conditioning on σ -fields



Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability
$\mathbb{E}[X Y]$ conditioning on random variables	$\mathbb{E}[X \sigma(Y)]$ conditioning on σ -fields
Convergence in distribution	Also: almost sure, in L^1 , in probability



Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability
$\mathbb{E}[X Y]$ conditioning on random variables	$\mathbb{E}[X \sigma(Y)]$ conditioning on σ -fields
Convergence in distribution	Also: almost sure, in L^1 , in probability
None	Monotone and dominated convergence theorem



Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability
$\mathbb{E}[X Y]$ conditioning on random variables	$\mathbb{E}[X \sigma(Y)]$ conditioning on σ -fields
Convergence in distribution	Also: almost sure, in L^1 , in probability
None	Monotone and dominated convergence theorem
None	Non-measurable subsets of \mathbb{R}

Why this tutorial?

Why this tutorial?

- Many results easy to appreciate from the undergraduate view

Why this tutorial?

- Many results easy to appreciate from the undergraduate view
- Non-asymptotic results: easy to use

Why this tutorial?

- Many results easy to appreciate from the undergraduate view
- Non-asymptotic results: easy to use
- Use basic information-theoretic ideas

Why this tutorial?

- Many results easy to appreciate from the undergraduate view
- Non-asymptotic results: easy to use
- Use basic information-theoretic ideas

Role of information theory

“The emphasis put on information theoretic methods is one main feature of the exposition and there is considerable benefit in this approach for a number of fundamental results [...]”
- M. Ledoux, foreword to ‘Concentration Inequalities’ by Boucheron, Lugosi, Massart.

Roadmap

Roadmap

- Monday: Variance bounds

Roadmap

- Monday: Variance bounds
- Tuesday: Information inequalities

Roadmap

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities

Roadmap

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities
- Friday: Transportation method

Roadmap

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities
- Friday: Transportation method

Thanks to Ramon van Handel, Igal Sason, Max Raginsky

Roadmap

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities
- Friday: Transportation method

Thanks to Ramon van Handel, Igal Sason, Max Raginsky

Reference: 'Concentration Inequalities' by Boucheron, Lugosi, Massart

Roadmap

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities
- Friday: Transportation method

Thanks to Ramon van Handel, Igal Sason, Max Raginsky

Reference: 'Concentration Inequalities' by Boucheron, Lugosi, Massart

Slides available on my homepage:

<http://www.princeton.edu/~sukamath/concentration.pdf>

Variance bounds

Variance bounds

Say Z is a function of independent random variables X_1, X_2, \dots, X_n . An upper bound on $\text{Var}(Z)$ gives tail bounds as:

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t] \leq \frac{\text{Var}(Z)}{t^2}$$

Variance bounds

Say Z is a function of independent random variables X_1, X_2, \dots, X_n . An upper bound on $\text{Var}(Z)$ gives tail bounds as:

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t] \leq \frac{\text{Var}(Z)}{t^2}$$

Probability of Z being within 10 standard deviations, i.e.
 $t = 10\sqrt{\text{Var}(Z)}$ of $\mathbb{E}Z$ is at least 99%

Variance bounds

Say Z is a function of independent random variables X_1, X_2, \dots, X_n . An upper bound on $\text{Var}(Z)$ gives tail bounds as:

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t] \leq \frac{\text{Var}(Z)}{t^2}$$

Probability of Z being within 10 standard deviations, i.e.
 $t = 10\sqrt{\text{Var}(Z)}$ of $\mathbb{E}Z$ is at least 99%

Trivial example

Let $Z = X_1 + X_2 + \dots + X_n$ where $\{X_i\}_{i=1}^n$ are independent and identically distributed (i.i.d.) with finite variance. Then,

$$\mathbb{E}Z = n\mathbb{E}X_1 \quad \text{Var}(Z) = n \text{Var}(X_1)$$

Variance bounds

Say Z is a function of independent random variables X_1, X_2, \dots, X_n . An upper bound on $\text{Var}(Z)$ gives tail bounds as:

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t] \leq \frac{\text{Var}(Z)}{t^2}$$

Probability of Z being within 10 standard deviations, i.e.
 $t = 10\sqrt{\text{Var}(Z)}$ of $\mathbb{E}Z$ is at least 99%

Trivial example

Let $Z = X_1 + X_2 + \dots + X_n$ where $\{X_i\}_{i=1}^n$ are independent and identically distributed (i.i.d.) with finite variance. Then,

$$\mathbb{E}Z = n\mathbb{E}X_1 \quad \text{Var}(Z) = n \text{Var}(X_1)$$

Mean = $\Theta(n)$, Standard Deviation = $O(\sqrt{n})$.

Variance bounds: sharper truths

Variance bounds: sharper truths

Spectral norm of a random matrix

Populate an $m \times n$ matrix A by independent entries, each taking values in $[0, 1]$. The random variable $Z = \|A\|$ satisfies

Variance bounds: sharper truths

Spectral norm of a random matrix

Populate an $m \times n$ matrix A by independent entries, each taking values in $[0, 1]$. The random variable $Z = \|A\|$ satisfies

$$\text{Var}(Z) \leq 1$$

Variance bounds: sharper truths

Spectral norm of a random matrix

Populate an $m \times n$ matrix A by independent entries, each taking values in $[0, 1]$. The random variable $Z = \|A\|$ satisfies

$$\text{Var}(Z) \leq 1$$

Plug-in entropy estimation

Let Z be the estimate of entropy of an unknown distribution defined by the entropy of the empirical distribution from drawing n independent samples.

Variance bounds: sharper truths

Spectral norm of a random matrix

Populate an $m \times n$ matrix A by independent entries, each taking values in $[0, 1]$. The random variable $Z = \|A\|$ satisfies

$$\text{Var}(Z) \leq 1$$

Plug-in entropy estimation

Let Z be the estimate of entropy of an unknown distribution defined by the entropy of the empirical distribution from drawing n independent samples.

$$\text{Var}(Z) \leq \frac{\log^2 n}{n}$$

High-level idea

High-level idea

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable

High-level idea

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable
- Not obvious this is possible

High-level idea

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable
- Not obvious this is possible
- When it is, we say the quantity *tensorizes*

High-level idea

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable
- Not obvious this is possible
- When it is, we say the quantity *tensorizes*
- Quantities that tensorize behave well in high dimension

High-level idea

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable
- Not obvious this is possible
- When it is, we say the quantity *tensorizes*
- Quantities that tensorize behave well in high dimension
- **Variance is such a quantity!**

Tensorization of variance

Tensorization of variance

Let $Z = f(X_1, X_2, \dots, X_n)$ where X_1, X_2, \dots, X_n are independent random variables.

Tensorization of variance

Let $Z = f(X_1, X_2, \dots, X_n)$ where X_1, X_2, \dots, X_n are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n)$$

Tensorization of variance

Let $Z = f(X_1, X_2, \dots, X_n)$ where X_1, X_2, \dots, X_n are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

Tensorization of variance

Let $Z = f(X_1, X_2, \dots, X_n)$ where X_1, X_2, \dots, X_n are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

Tensorization of variance

Let $Z = f(X_1, X_2, \dots, X_n)$ where X_1, X_2, \dots, X_n are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

$$g_i(x^{(i)}) = \text{Var}(f(x_1, \dots, X_i, \dots, x_n))$$

Tensorization of variance

Let $Z = f(X_1, X_2, \dots, X_n)$ where X_1, X_2, \dots, X_n are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

$$g_i(x^{(i)}) = \text{Var}(f(x_1, \dots, X_i, \dots, x_n)) \implies \text{Var}^{(i)}(Z) = g_i(X^{(i)})$$

Tensorization of variance

Let $Z = f(X_1, X_2, \dots, X_n)$ where X_1, X_2, \dots, X_n are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

$$g_i(x^{(i)}) = \text{Var}(f(x_1, \dots, X_i, \dots, x_n)) \implies \text{Var}^{(i)}(Z) = g_i(X^{(i)})$$

Tensorization of variance (Efron-Stein-Steele inequality)

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)]$$

Tensorization of variance

Let $Z = f(X_1, X_2, \dots, X_n)$ where X_1, X_2, \dots, X_n are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

$$g_i(x^{(i)}) = \text{Var}(f(x_1, \dots, X_i, \dots, x_n)) \implies \text{Var}^{(i)}(Z) = g_i(X^{(i)})$$

Tensorization of variance (Efron-Stein-Steele inequality)

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)] = \sum_{i=1}^n \mathbb{E}[(Z - \mathbb{E}^{(i)} Z)^2]$$

Recall: if $Y \in [a, b]$, then $\text{Var}(Y) \leq$

Recall: if $Y \in [a, b]$, then $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2]$

Recall: if $Y \in [a, b]$, then $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2] \leq \frac{(b - a)^2}{4}$

from $u = \frac{1}{2}(a + b)$.

Recall: if $Y \in [a, b]$, then $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2] \leq \frac{(b - a)^2}{4}$

from $u = \frac{1}{2}(a + b)$.

Simplest application: Bounded differences inequality

Suppose

$$|f(x_1, \dots, x_i, \dots, x_n) - f(x_1, \dots, x'_i, \dots, x_n)| \leq c_i.$$

Recall: if $Y \in [a, b]$, then $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2] \leq \frac{(b - a)^2}{4}$

from $u = \frac{1}{2}(a + b)$.

Simplest application: Bounded differences inequality

Suppose

$$|f(x_1, \dots, x_i, \dots, x_n) - f(x_1, \dots, x'_i, \dots, x_n)| \leq c_i.$$

$$\text{Then, } \text{Var}(f(X)) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)] \leq \frac{1}{4} \sum_{i=1}^n c_i^2$$

Recall: if $Y \in [a, b]$, then $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2] \leq \frac{(b - a)^2}{4}$

from $u = \frac{1}{2}(a + b)$.

Simplest application: Bounded differences inequality

Suppose

$$|f(x_1, \dots, x_i, \dots, x_n) - f(x_1, \dots, x'_i, \dots, x_n)| \leq c_i.$$

$$\text{Then, } \text{Var}(f(X)) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)] \leq \frac{1}{4} \sum_{i=1}^n c_i^2$$

Tight if $f(X) = \sum_{i=1}^n X_i$ with X_i equiprobable on $\{-1, +1\}$

Bin packing problem

Bin packing problem

Let $X_1, X_2, \dots, X_n \in [0, 1]$ be i.i.d.

Bin packing problem

Let $X_1, X_2, \dots, X_n \in [0, 1]$ be i.i.d.

Let Z be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

Bin packing problem

Let $X_1, X_2, \dots, X_n \in [0, 1]$ be i.i.d.

Let Z be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

Changing one X_i changes $Z = f(X)$ by at most $c_i = 1$

Bin packing problem

Let $X_1, X_2, \dots, X_n \in [0, 1]$ be i.i.d.

Let Z be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

Changing one X_i changes $Z = f(X)$ by at most $c_i = 1$

$$\text{Therefore, } \text{Var}(Z) \leq \frac{1}{4} \sum_{i=1}^n c_i^2 = \frac{n}{4} .$$

Bin packing problem

Let $X_1, X_2, \dots, X_n \in [0, 1]$ be i.i.d.

Let Z be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

Changing one X_i changes $Z = f(X)$ by at most $c_i = 1$

$$\text{Therefore, } \text{Var}(Z) \leq \frac{1}{4} \sum_{i=1}^n c_i^2 = \frac{n}{4} .$$

$$\text{However, } \mathbb{E}Z \geq \mathbb{E} \left[\sum_{i=1}^n X_i \right] = n\mathbb{E}X_1 .$$

Bin packing problem

Let $X_1, X_2, \dots, X_n \in [0, 1]$ be i.i.d.

Let Z be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

Changing one X_i changes $Z = f(X)$ by at most $c_i = 1$

$$\text{Therefore, } \text{Var}(Z) \leq \frac{1}{4} \sum_{i=1}^n c_i^2 = \frac{n}{4} .$$

$$\text{However, } \mathbb{E}Z \geq \mathbb{E} \left[\sum_{i=1}^n X_i \right] = n\mathbb{E}X_1.$$

Standard deviation = $O(\sqrt{n})$, Mean = $\Theta(n)$.

Plug-in entropy estimation

Plug-in entropy estimation

Entropy of a distribution $p = (p_1, p_2, \dots, p_k)$ is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Plug-in entropy estimation

Entropy of a distribution $p = (p_1, p_2, \dots, p_k)$ is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let X_1, X_2, \dots, X_n be independent samples from p

Plug-in entropy estimation

Entropy of a distribution $p = (p_1, p_2, \dots, p_k)$ is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let X_1, X_2, \dots, X_n be independent samples from p

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|$$

Plug-in entropy estimation

Entropy of a distribution $p = (p_1, p_2, \dots, p_k)$ is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let X_1, X_2, \dots, X_n be independent samples from p

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

Plug-in entropy estimation

Entropy of a distribution $p = (p_1, p_2, \dots, p_k)$ is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let X_1, X_2, \dots, X_n be independent samples from p

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate X_i affects two of the \hat{p}_r 's.

Plug-in entropy estimation

Entropy of a distribution $p = (p_1, p_2, \dots, p_k)$ is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let X_1, X_2, \dots, X_n be independent samples from p

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate X_i affects two of the \hat{p}_r 's.

$$\left| a \log \frac{1}{a} - b \log \frac{1}{b} \right| \leq \frac{\log n}{n} \text{ if } |a - b| = \frac{1}{n}.$$

Plug-in entropy estimation

Entropy of a distribution $p = (p_1, p_2, \dots, p_k)$ is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let X_1, X_2, \dots, X_n be independent samples from p

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate X_i affects two of the \hat{p}_r 's.

$$\left| a \log \frac{1}{a} - b \log \frac{1}{b} \right| \leq \frac{\log n}{n} \text{ if } |a - b| = \frac{1}{n}.$$

$$\text{Thus, } \text{Var}(Z) \leq \sum_{i=1}^n c_i^2 / 4 = (\log^2 n) / n$$

Plug-in entropy estimation

Entropy of a distribution $p = (p_1, p_2, \dots, p_k)$ is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let X_1, X_2, \dots, X_n be independent samples from p

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate X_i affects two of the \hat{p}_r 's.

$$\left| a \log \frac{1}{a} - b \log \frac{1}{b} \right| \leq \frac{\log n}{n} \text{ if } |a - b| = \frac{1}{n}.$$

$$\text{Thus, } \text{Var}(Z) \leq \sum_{i=1}^n c_i^2 / 4 = (\log^2 n) / n$$

But Z is not really concentrated at $H(p)$ unless $n \gtrsim k$.

Plug-in entropy estimation

Entropy of a distribution $p = (p_1, p_2, \dots, p_k)$ is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let X_1, X_2, \dots, X_n be independent samples from p

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate X_i affects two of the \hat{p}_r 's.

$$\left| a \log \frac{1}{a} - b \log \frac{1}{b} \right| \leq \frac{\log n}{n} \text{ if } |a - b| = \frac{1}{n}.$$

$$\text{Thus, } \text{Var}(Z) \leq \sum_{i=1}^n c_i^2 / 4 = (\log^2 n) / n$$

But Z is not really concentrated at $H(p)$ unless $n \gtrsim k$.
For $n \ll k$, Z is concentrated but somewhere else!

Warning

Warning

- We have shown bounds on deviation of Z from $\mathbb{E}Z$

Warning

- We have shown bounds on deviation of Z from $\mathbb{E}Z$
- But say nothing about $\mathbb{E}Z$ itself!!

Warning

- We have shown bounds on deviation of Z from $\mathbb{E}Z$
- But say nothing about $\mathbb{E}Z$ itself!!
- Estimating magnitude and fluctuations are two quite distinct problems

Warning

- We have shown bounds on deviation of Z from $\mathbb{E}Z$
- But say nothing about $\mathbb{E}Z$ itself!!
- Estimating magnitude and fluctuations are two quite distinct problems
- We have a general theorem for bounding fluctuations and elementary ideas can often bound sensitivity to coordinates, even if the function itself is complicated

Warning

- We have shown bounds on deviation of Z from $\mathbb{E}Z$
- But say nothing about $\mathbb{E}Z$ itself!!
- Estimating magnitude and fluctuations are two quite distinct problems
- We have a general theorem for bounding fluctuations and elementary ideas can often bound sensitivity to coordinates, even if the function itself is complicated
- No such general principle for estimating $\mathbb{E}Z$

Warning

- We have shown bounds on deviation of Z from $\mathbb{E}Z$
- But say nothing about $\mathbb{E}Z$ itself!!
- Estimating magnitude and fluctuations are two quite distinct problems
- We have a general theorem for bounding fluctuations and elementary ideas can often bound sensitivity to coordinates, even if the function itself is complicated
- No such general principle for estimating $\mathbb{E}Z$
- Can estimate $\mathbb{E}Z$ from Monte Carlo methods if Z is concentrated

Recall that $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$

Recall that $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$

So, $\text{Var}^{(i)}(Z) = \inf_{f_i(x^{(i)})} \mathbb{E}^{(i)}[(Z - f_i(X^{(i)}))^2]$

Recall that $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$

So, $\text{Var}^{(i)}(Z) = \inf_{f_i(x^{(i)})} \mathbb{E}^{(i)}[(Z - f_i(X^{(i)}))^2]$

Let $Z_i = f_i(X^{(i)})$ for any function f_i .

Recall that $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$

$$\text{So, } \text{Var}^{(i)}(Z) = \inf_{f_i(x^{(i)})} \mathbb{E}^{(i)}[(Z - f_i(X^{(i)}))^2]$$

Let $Z_i = f_i(X^{(i)})$ for any function f_i .

$$\text{Then, } \text{Var}^{(i)}(Z) \leq \mathbb{E}^{(i)}[(Z - Z_i)^2]$$

Recall that $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$

$$\text{So, } \text{Var}^{(i)}(Z) = \inf_{f_i(x^{(i)})} \mathbb{E}^{(i)}[(Z - f_i(X^{(i)}))^2]$$

Let $Z_i = f_i(X^{(i)})$ for any function f_i .

$$\text{Then, } \text{Var}^{(i)}(Z) \leq \mathbb{E}^{(i)}[(Z - Z_i)^2]$$

Variant : “guess functions”

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)] \leq \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2]$$

Convex Lipschitz functions

Suppose $f : [a, b]^n \mapsto \mathbb{R}$ is convex, differentiable and L -Lipschitz,
i.e. $|f(x) - f(y)| \leq L\|x - y\|_2 \ \forall \ x, y$.

Convex Lipschitz functions

Suppose $f : [a, b]^n \mapsto \mathbb{R}$ is convex, differentiable and L -Lipschitz,
i.e. $|f(x) - f(y)| \leq L\|x - y\|_2 \ \forall \ x, y$.

Choose $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$ with inf attained at X'_i

Convex Lipschitz functions

Suppose $f : [a, b]^n \mapsto \mathbb{R}$ is convex, differentiable and L -Lipschitz,
i.e. $|f(x) - f(y)| \leq L\|x - y\|_2 \ \forall \ x, y$.

Choose $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$ with inf attained at X'_i

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$

Convex Lipschitz functions

Suppose $f : [a, b]^n \mapsto \mathbb{R}$ is convex, differentiable and L -Lipschitz,
i.e. $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$.

Choose $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$ with inf attained at X'_i

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i) \qquad 0 \leq Z - Z_i \leq -\frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$

Convex Lipschitz functions

Suppose $f : [a, b]^n \mapsto \mathbb{R}$ is convex, differentiable and L -Lipschitz,
i.e. $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$.

Choose $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$ with inf attained at X'_i

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i) \qquad 0 \leq Z - Z_i \leq -\frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$

$$0 \leq (Z - Z_i)^2 \leq \left| \frac{\partial f}{\partial x_i}(X) \right|^2 (b - a)^2$$

Convex Lipschitz functions

Suppose $f : [a, b]^n \mapsto \mathbb{R}$ is convex, differentiable and L -Lipschitz,
i.e. $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$.

Choose $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$ with inf attained at X'_i

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i) \qquad 0 \leq Z - Z_i \leq -\frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$

$$0 \leq (Z - Z_i)^2 \leq \left| \frac{\partial f}{\partial x_i}(X) \right|^2 (b - a)^2$$

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2] \leq L^2(b - a)^2$$

Convex Lipschitz functions

Suppose $f : [a, b]^n \mapsto \mathbb{R}$ is convex, differentiable and L -Lipschitz,
i.e. $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$.

Choose $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$ with inf attained at X'_i

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i) \qquad 0 \leq Z - Z_i \leq -\frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$

$$0 \leq (Z - Z_i)^2 \leq \left| \frac{\partial f}{\partial x_i}(X) \right|^2 (b - a)^2$$

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2] \leq L^2(b - a)^2$$

Differentiability assumption unnecessary:
convolve f with a smooth kernel.

Spectral norm of a random matrix

Populate an $m \times n$ matrix A by independent entries, each taking values in $[0, 1]$.

Spectral norm of a random matrix

Populate an $m \times n$ matrix A by independent entries, each taking values in $[0, 1]$.

The function $f : [0, 1]^{m \times n} \mapsto \mathbb{R}$, given by

$$Z = f(A) = \|A\| = \sup_{\|v\|_2=1} \|Av\|_2$$

Spectral norm of a random matrix

Populate an $m \times n$ matrix A by independent entries, each taking values in $[0, 1]$.

The function $f : [0, 1]^{m \times n} \mapsto \mathbb{R}$, given by

$$Z = f(A) = \|A\| = \sup_{\|v\|_2=1} \|Av\|_2$$

is convex and 1-Lipschitz

Spectral norm of a random matrix

Populate an $m \times n$ matrix A by independent entries, each taking values in $[0, 1]$.

The function $f : [0, 1]^{m \times n} \mapsto \mathbb{R}$, given by

$$Z = f(A) = \|A\| = \sup_{\|v\|_2=1} \|Av\|_2$$

is convex and 1-Lipschitz

(hint: spectral norm \leq Frobenius norm)

Spectral norm of a random matrix

Populate an $m \times n$ matrix A by independent entries, each taking values in $[0, 1]$.

The function $f : [0, 1]^{m \times n} \mapsto \mathbb{R}$, given by

$$Z = f(A) = \|A\| = \sup_{\|v\|_2=1} \|Av\|_2$$

is convex and 1-Lipschitz

(hint: spectral norm \leq Frobenius norm)

With $a = 0, b = 1, L = 1$ in previous result, we get

$$\text{Var}(Z) \leq 1.$$