

# *Concentration of Measure*

Sudeep Kamath



CIRM workshop, 25 Jan 2016

## What is concentration?

“A random variable that depends in a smooth way on many independent random variables (but not too much on any of them) is essentially constant.”

- M. Talagrand, 1996.

## What is concentration?

“A random variable that depends in a smooth way on many independent random variables (but not too much on any of them) is essentially constant.”

- M. Talagrand, 1996.

If  $Z$  is a function of many independent variables  $X_1, X_2, \dots, X_n$ ,  
how large are typical deviations of  $Z$ ?

## What is concentration?

“A random variable that depends in a smooth way on many independent random variables (but not too much on any of them) is essentially constant.”

- M. Talagrand, 1996.

If  $Z$  is a function of many independent variables  $X_1, X_2, \dots, X_n$ ,  
how large are typical deviations of  $Z$ ?

Goal: Quantify by bounding for  $t > 0$ ,

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t]$$

## *Applications*

Concentration of measure has far-reaching consequences in

- Pure and applied probability,
- High-dimensional statistics,
- Functional analysis,
- Computer science,
- Machine learning,
- Statistical physics,
- Complex graphs and networks,
- Information theory, communication and coding theory.


## *Approaches for Proving Concentration*

- *The martingale approach*: Hoeffding (1963), Azuma (1967), Milman and Schechtman (1986), Shamir and Spencer (1987) and McDiarmid (1989, 1998), Sipser and Spielman (1996), Richardson and Urbanke (2001)
- *Talagrand's inequalities for product measures*: Talagrand (1996).
- *Entropy method and log-Sobolev inequalities*: Ledoux (1996), Massart (1998), Lugosi et al. (1999, 2001)
- *Transportation method*: Ahlswede, Gács and Körner (1976), Marton (1986, 1996, 1997), Dembo (1997), Villani (2003, 2008)
- *Stein's method of exchangeable pairs*: Chatterjee (2007), Chatterjee and Dey (2010), Goldstein et al. (2011, 2014)


## *Approaches for Proving Concentration*

- *The martingale approach*: Hoeffding (1963), Azuma (1967), Milman and Schechtman (1986), Shamir and Spencer (1987) and McDiarmid (1989, 1998), Sipser and Spielman (1996), Richardson and Urbanke (2001)
- *Talagrand's inequalities for product measures*: Talagrand (1996).
- *Entropy method and log-Sobolev inequalities*: Ledoux (1996), Massart (1998), Lugosi et al. (1999, 2001)
- *Transportation method*: Ahlswede, Gács and Körner (1976), Marton (1986, 1996, 1997), Dembo (1997), Villani (2003, 2008)
- *Stein's method of exchangeable pairs*: Chatterjee (2007), Chatterjee and Dey (2010), Goldstein et al. (2011, 2014)


We will focus on the entropy method and transportation method where information theoretic methods shine.









Undergraduate ("informal")	




Undergraduate ("informal")	Graduate (formal probability)




Undergraduate ("informal")	Graduate (formal probability)
Probability density,	Radon-Nikodym derivative,




Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral




Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability



Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability
$\mathbb{E}[X Y]$ conditioning on random variables	$\mathbb{E}[X \sigma(Y)]$ conditioning on $\sigma$ -fields




Undergraduate ("informal")	Graduate (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability
$\mathbb{E}[X Y]$ conditioning on random variables	$\mathbb{E}[X \sigma(Y)]$ conditioning on $\sigma$ -fields
Convergence in distribution	Also: almost sure, in $L^1$ , in probability



<b>Undergraduate</b> ("informal")	<b>Graduate</b> (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability
$\mathbb{E}[X Y]$ conditioning on random variables	$\mathbb{E}[X \sigma(Y)]$ conditioning on $\sigma$ -fields
Convergence in distribution	Also: almost sure, in $L^1$ , in probability
None	Monotone and dominated convergence theorem





<b>Undergraduate</b> ("informal")	<b>Graduate</b> (formal probability)
Probability density, Riemann integral	Radon-Nikodym derivative, Lebesgue integral
Conditional probability	Regular conditional probability
$\mathbb{E}[X Y]$ conditioning on random variables	$\mathbb{E}[X \sigma(Y)]$ conditioning on $\sigma$ -fields
Convergence in distribution	Also: almost sure, in $L^1$ , in probability
None	Monotone and dominated convergence theorem
None	Non-measurable subsets of $\mathbb{R}$

*Why this tutorial?*

## *Why this tutorial?*

- Many results easy to appreciate from the undergraduate view

## *Why this tutorial?*

- Many results easy to appreciate from the undergraduate view
- Non-asymptotic results: easy to use

## *Why this tutorial?*

- Many results easy to appreciate from the undergraduate view
- Non-asymptotic results: easy to use
- Use basic information-theoretic ideas

## *Why this tutorial?*

- Many results easy to appreciate from the undergraduate view
- Non-asymptotic results: easy to use
- Use basic information-theoretic ideas

### Role of information theory

“The emphasis put on information theoretic methods is one main feature of the exposition and there is considerable benefit in this approach for a number of fundamental results [...]”  
- M. Ledoux, foreword to ‘Concentration Inequalities’ by Boucheron, Lugosi, Massart.

# *Roadmap*

## *Roadmap*

- Monday: Variance bounds



## *Roadmap*

- Monday: Variance bounds
- Tuesday: Information inequalities

## *Roadmap*

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities

## *Roadmap*

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities
- Friday: Transportation method

## *Roadmap*

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities
- Friday: Transportation method

Thanks to Ramon van Handel, Igal Sason, Max Raginsky

## *Roadmap*

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities
- Friday: Transportation method

Thanks to Ramon van Handel, Igal Sason, Max Raginsky

Reference: 'Concentration Inequalities' by Boucheron, Lugosi, Massart

## *Roadmap*

- Monday: Variance bounds
- Tuesday: Information inequalities
- Thursday: Entropy method and log-Sobolev inequalities
- Friday: Transportation method

Thanks to Ramon van Handel, Igal Sason, Max Raginsky

Reference: 'Concentration Inequalities' by Boucheron, Lugosi, Massart

Slides available on my homepage:

<http://www.princeton.edu/~sukamath/concentration.pdf>

## *Variance bounds*

## *Variance bounds*

Say  $Z$  is a function of independent random variables  $X_1, X_2, \dots, X_n$ . An upper bound on  $\text{Var}(Z)$  gives tail bounds as:

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t] \leq \frac{\text{Var}(Z)}{t^2}$$



## *Variance bounds*

Say  $Z$  is a function of independent random variables  $X_1, X_2, \dots, X_n$ . An upper bound on  $\text{Var}(Z)$  gives tail bounds as:

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t] \leq \frac{\text{Var}(Z)}{t^2}$$

Probability of  $Z$  being within 10 standard deviations, i.e.

$t = 10\sqrt{\text{Var}(Z)}$  of  $\mathbb{E}Z$  is at least 99%

## Variance bounds

Say  $Z$  is a function of independent random variables  $X_1, X_2, \dots, X_n$ . An upper bound on  $\text{Var}(Z)$  gives tail bounds as:

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t] \leq \frac{\text{Var}(Z)}{t^2}$$

Probability of  $Z$  being within 10 standard deviations, i.e.  
 $t = 10\sqrt{\text{Var}(Z)}$  of  $\mathbb{E}Z$  is at least 99%

### Trivial example

Let  $Z = X_1 + X_2 + \dots + X_n$  where  $\{X_i\}_{i=1}^n$  are independent and identically distributed (i.i.d.) with finite variance. Then,

$$\mathbb{E}Z = n\mathbb{E}X_1 \quad \text{Var}(Z) = n \text{Var}(X_1)$$

## Variance bounds

Say  $Z$  is a function of independent random variables  $X_1, X_2, \dots, X_n$ . An upper bound on  $\text{Var}(Z)$  gives tail bounds as:

$$\mathbb{P}[|Z - \mathbb{E}Z| \geq t] \leq \frac{\text{Var}(Z)}{t^2}$$

Probability of  $Z$  being within 10 standard deviations, i.e.  
 $t = 10\sqrt{\text{Var}(Z)}$  of  $\mathbb{E}Z$  is at least 99%

### Trivial example

Let  $Z = X_1 + X_2 + \dots + X_n$  where  $\{X_i\}_{i=1}^n$  are independent and identically distributed (i.i.d.) with finite variance. Then,

$$\mathbb{E}Z = n\mathbb{E}X_1 \quad \text{Var}(Z) = n \text{Var}(X_1)$$

Mean =  $\Theta(n)$ , Standard Deviation =  $O(\sqrt{n})$ .

*Variance bounds: sharper truths*

## *Variance bounds: sharper truths*

### Spectral norm of a random matrix

Populate an  $m \times n$  matrix  $A$  by independent entries, each taking values in  $[0, 1]$ . The random variable  $Z = \|A\|$  satisfies

## *Variance bounds: sharper truths*

### Spectral norm of a random matrix

Populate an  $m \times n$  matrix  $A$  by independent entries, each taking values in  $[0, 1]$ . The random variable  $Z = \|A\|$  satisfies

$$\text{Var}(Z) \leq 1$$

## *Variance bounds: sharper truths*

### Spectral norm of a random matrix

Populate an  $m \times n$  matrix  $A$  by independent entries, each taking values in  $[0, 1]$ . The random variable  $Z = \|A\|$  satisfies

$$\text{Var}(Z) \leq 1$$

### Plug-in entropy estimation

Let  $Z$  be the estimate of entropy of an unknown distribution defined by the entropy of the empirical distribution from drawing  $n$  independent samples.

## *Variance bounds: sharper truths*

### Spectral norm of a random matrix

Populate an  $m \times n$  matrix  $A$  by independent entries, each taking values in  $[0, 1]$ . The random variable  $Z = \|A\|$  satisfies

$$\text{Var}(Z) \leq 1$$

### Plug-in entropy estimation

Let  $Z$  be the estimate of entropy of an unknown distribution defined by the entropy of the empirical distribution from drawing  $n$  independent samples.

$$\text{Var}(Z) \leq \frac{\log^2 n}{n}$$



## *High-level idea*

## *High-level idea*

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable

## *High-level idea*

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable
- Not obvious this is possible

## *High-level idea*

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable
- Not obvious this is possible
- When it is, we say the quantity *tensorizes*

## *High-level idea*

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable
- Not obvious this is possible
- When it is, we say the quantity *tensorizes*
- Quantities that tensorize behave well in high dimension

## *High-level idea*

- Obtain a bound for a function of many random variables by bounds for functions of each individual random variable
- Not obvious this is possible
- When it is, we say the quantity *tensorizes*
- Quantities that tensorize behave well in high dimension
- Variance is such a quantity!

# *Tensorization of variance*

## *Tensorization of variance*

Let  $Z = f(X_1, X_2, \dots, X_n)$  where  $X_1, X_2, \dots, X_n$  are independent random variables.



## *Tensorization of variance*

Let  $Z = f(X_1, X_2, \dots, X_n)$  where  $X_1, X_2, \dots, X_n$  are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n)$$

## *Tensorization of variance*

Let  $Z = f(X_1, X_2, \dots, X_n)$  where  $X_1, X_2, \dots, X_n$  are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

## *Tensorization of variance*

Let  $Z = f(X_1, X_2, \dots, X_n)$  where  $X_1, X_2, \dots, X_n$  are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

## *Tensorization of variance*

Let  $Z = f(X_1, X_2, \dots, X_n)$  where  $X_1, X_2, \dots, X_n$  are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

$$g_i(x^{(i)}) = \text{Var}(f(x_1, \dots, X_i, \dots, x_n))$$

## *Tensorization of variance*

Let  $Z = f(X_1, X_2, \dots, X_n)$  where  $X_1, X_2, \dots, X_n$  are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

$$g_i(x^{(i)}) = \text{Var}(f(x_1, \dots, X_i, \dots, x_n)) \implies \text{Var}^{(i)}(Z) = g_i(X^{(i)})$$

## *Tensorization of variance*

Let  $Z = f(X_1, X_2, \dots, X_n)$  where  $X_1, X_2, \dots, X_n$  are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

$$g_i(x^{(i)}) = \text{Var}(f(x_1, \dots, X_i, \dots, x_n)) \implies \text{Var}^{(i)}(Z) = g_i(X^{(i)})$$

Tensorization of variance (Efron-Stein-Steele inequality)

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)]$$

## Tensorization of variance

Let  $Z = f(X_1, X_2, \dots, X_n)$  where  $X_1, X_2, \dots, X_n$  are independent random variables.

$$X^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n), \quad \mathbb{E}^{(i)}[\cdot] := \mathbb{E}[\cdot | X^{(i)}]$$

$$\text{Var}^{(i)}(Z) := \text{Var}(Z | X^{(i)})$$

$$g_i(x^{(i)}) = \text{Var}(f(x_1, \dots, X_i, \dots, x_n)) \implies \text{Var}^{(i)}(Z) = g_i(X^{(i)})$$

Tensorization of variance (Efron-Stein-Steele inequality)

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)] = \sum_{i=1}^n \mathbb{E}[(Z - \mathbb{E}^{(i)} Z)^2]$$





Recall: if  $Y \in [a, b]$ , then  $\text{Var}(Y) \leq$

Recall: if  $Y \in [a, b]$ , then  $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2]$

Recall: if  $Y \in [a, b]$ , then  $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2] \leq \frac{(b - a)^2}{4}$

from  $u = \frac{1}{2}(a + b)$ .

Recall: if  $Y \in [a, b]$ , then  $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2] \leq \frac{(b - a)^2}{4}$

from  $u = \frac{1}{2}(a + b)$ .

Simplest application: Bounded differences inequality

Suppose

$$|f(x_1, \dots, x_i, \dots, x_n) - f(x_1, \dots, x'_i, \dots, x_n)| \leq c_i.$$

Recall: if  $Y \in [a, b]$ , then  $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2] \leq \frac{(b - a)^2}{4}$

from  $u = \frac{1}{2}(a + b)$ .

### Simplest application: Bounded differences inequality

Suppose

$$|f(x_1, \dots, x_i, \dots, x_n) - f(x_1, \dots, x'_i, \dots, x_n)| \leq c_i.$$

$$\text{Then, } \text{Var}(f(X)) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)] \leq \frac{1}{4} \sum_{i=1}^n c_i^2$$

Recall: if  $Y \in [a, b]$ , then  $\text{Var}(Y) \leq \inf_u \mathbb{E}[(Y - u)^2] \leq \frac{(b - a)^2}{4}$

from  $u = \frac{1}{2}(a + b)$ .

### Simplest application: Bounded differences inequality

Suppose

$$|f(x_1, \dots, x_i, \dots, x_n) - f(x_1, \dots, x'_i, \dots, x_n)| \leq c_i.$$

$$\text{Then, } \text{Var}(f(X)) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)] \leq \frac{1}{4} \sum_{i=1}^n c_i^2$$

Tight if  $f(X) = \sum_{i=1}^n X_i$  with  $X_i$  equiprobable on  $\{-1, +1\}$

## Bin packing problem

## Bin packing problem

Let  $X_1, X_2, \dots, X_n \in [0, 1]$  be i.i.d.



## Bin packing problem

Let  $X_1, X_2, \dots, X_n \in [0, 1]$  be i.i.d.

Let  $Z$  be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

## Bin packing problem

Let  $X_1, X_2, \dots, X_n \in [0, 1]$  be i.i.d.

Let  $Z$  be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

Changing one  $X_i$  changes  $Z = f(X)$  by at most  $c_i = 1$

## Bin packing problem

Let  $X_1, X_2, \dots, X_n \in [0, 1]$  be i.i.d.

Let  $Z$  be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

Changing one  $X_i$  changes  $Z = f(X)$  by at most  $c_i = 1$

$$\text{Therefore, } \text{Var}(Z) \leq \frac{1}{4} \sum_{i=1}^n c_i^2 = \frac{n}{4} .$$

## Bin packing problem

Let  $X_1, X_2, \dots, X_n \in [0, 1]$  be i.i.d.

Let  $Z$  be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

Changing one  $X_i$  changes  $Z = f(X)$  by at most  $c_i = 1$

$$\text{Therefore, } \text{Var}(Z) \leq \frac{1}{4} \sum_{i=1}^n c_i^2 = \frac{n}{4} .$$

$$\text{However, } \mathbb{E}Z \geq \mathbb{E} \left[ \sum_{i=1}^n X_i \right] = n\mathbb{E}X_1 .$$

## Bin packing problem

Let  $X_1, X_2, \dots, X_n \in [0, 1]$  be i.i.d.

Let  $Z$  be the minimum number of bins in to which they can be packed so that each bin adds to at most 1.

Changing one  $X_i$  changes  $Z = f(X)$  by at most  $c_i = 1$

$$\text{Therefore, } \text{Var}(Z) \leq \frac{1}{4} \sum_{i=1}^n c_i^2 = \frac{n}{4} .$$

$$\text{However, } \mathbb{E}Z \geq \mathbb{E} \left[ \sum_{i=1}^n X_i \right] = n\mathbb{E}X_1.$$

Standard deviation =  $O(\sqrt{n})$ , Mean =  $\Theta(n)$ .

## Plug-in entropy estimation

## Plug-in entropy estimation

Entropy of a distribution  $p = (p_1, p_2, \dots, p_k)$  is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

## Plug-in entropy estimation

Entropy of a distribution  $p = (p_1, p_2, \dots, p_k)$  is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let  $X_1, X_2, \dots, X_n$  be independent samples from  $p$



## Plug-in entropy estimation

Entropy of a distribution  $p = (p_1, p_2, \dots, p_k)$  is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let  $X_1, X_2, \dots, X_n$  be independent samples from  $p$

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|$$

## Plug-in entropy estimation

Entropy of a distribution  $p = (p_1, p_2, \dots, p_k)$  is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let  $X_1, X_2, \dots, X_n$  be independent samples from  $p$

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

## Plug-in entropy estimation

Entropy of a distribution  $p = (p_1, p_2, \dots, p_k)$  is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let  $X_1, X_2, \dots, X_n$  be independent samples from  $p$

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate  $X_i$  affects two of the  $\hat{p}_r$ 's.

## Plug-in entropy estimation

Entropy of a distribution  $p = (p_1, p_2, \dots, p_k)$  is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let  $X_1, X_2, \dots, X_n$  be independent samples from  $p$

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate  $X_i$  affects two of the  $\hat{p}_r$ 's.

$$\left| a \log \frac{1}{a} - b \log \frac{1}{b} \right| \leq \frac{\log n}{n} \text{ if } |a - b| = \frac{1}{n}.$$

## Plug-in entropy estimation

Entropy of a distribution  $p = (p_1, p_2, \dots, p_k)$  is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let  $X_1, X_2, \dots, X_n$  be independent samples from  $p$

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate  $X_i$  affects two of the  $\hat{p}_r$ 's.

$$\left| a \log \frac{1}{a} - b \log \frac{1}{b} \right| \leq \frac{\log n}{n} \text{ if } |a - b| = \frac{1}{n}.$$

$$\text{Thus, } \text{Var}(Z) \leq \sum_{i=1}^n c_i^2 / 4 = (\log^2 n) / n$$

## Plug-in entropy estimation

Entropy of a distribution  $p = (p_1, p_2, \dots, p_k)$  is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let  $X_1, X_2, \dots, X_n$  be independent samples from  $p$

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate  $X_i$  affects two of the  $\hat{p}_r$ 's.

$$\left| a \log \frac{1}{a} - b \log \frac{1}{b} \right| \leq \frac{\log n}{n} \text{ if } |a - b| = \frac{1}{n}.$$

$$\text{Thus, } \text{Var}(Z) \leq \sum_{i=1}^n c_i^2 / 4 = (\log^2 n) / n$$

But  $Z$  is not really concentrated at  $H(p)$  unless  $n \gtrsim k$ .

## Plug-in entropy estimation

Entropy of a distribution  $p = (p_1, p_2, \dots, p_k)$  is defined as

$$H(p) = \sum_{r=1}^k p_r \log \frac{1}{p_r}$$

Let  $X_1, X_2, \dots, X_n$  be independent samples from  $p$

$$\text{Let } \hat{p}_r = \frac{1}{n} |\{i : X_i = r\}|, \quad Z = \sum_{r=1}^k \hat{p}_r \log \frac{1}{\hat{p}_r}$$

A change in any one co-ordinate  $X_i$  affects two of the  $\hat{p}_r$ 's.

$$\left| a \log \frac{1}{a} - b \log \frac{1}{b} \right| \leq \frac{\log n}{n} \text{ if } |a - b| = \frac{1}{n}.$$

$$\text{Thus, } \text{Var}(Z) \leq \sum_{i=1}^n c_i^2 / 4 = (\log^2 n) / n$$

But  $Z$  is not really concentrated at  $H(p)$  unless  $n \gtrsim k$ .  
For  $n \ll k$ ,  $Z$  is concentrated but somewhere else!

*Warning*



## *Warning*

- We have shown bounds on deviation of  $Z$  from  $\mathbb{E}Z$

## *Warning*

- We have shown bounds on deviation of  $Z$  from  $\mathbb{E}Z$
- But say nothing about  $\mathbb{E}Z$  itself!!

## *Warning*

- We have shown bounds on deviation of  $Z$  from  $\mathbb{E}Z$
- But say nothing about  $\mathbb{E}Z$  itself!!
- Estimating magnitude and fluctuations are two quite distinct problems

## Warning

- We have shown bounds on deviation of  $Z$  from  $\mathbb{E}Z$
- But say nothing about  $\mathbb{E}Z$  itself!!
- Estimating magnitude and fluctuations are two quite distinct problems
- We have a general theorem for bounding fluctuations and elementary ideas can often bound sensitivity to coordinates, even if the function itself is complicated

## Warning

- We have shown bounds on deviation of  $Z$  from  $\mathbb{E}Z$
- But say nothing about  $\mathbb{E}Z$  itself!!
- Estimating magnitude and fluctuations are two quite distinct problems
- We have a general theorem for bounding fluctuations and elementary ideas can often bound sensitivity to coordinates, even if the function itself is complicated
- No such general principle for estimating  $\mathbb{E}Z$

## Warning

- We have shown bounds on deviation of  $Z$  from  $\mathbb{E}Z$
- But say nothing about  $\mathbb{E}Z$  itself!!
- Estimating magnitude and fluctuations are two quite distinct problems
- We have a general theorem for bounding fluctuations and elementary ideas can often bound sensitivity to coordinates, even if the function itself is complicated
- No such general principle for estimating  $\mathbb{E}Z$
- Can estimate  $\mathbb{E}Z$  from Monte Carlo methods if  $Z$  is concentrated



Recall that  $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$



Recall that  $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$

So,  $\text{Var}^{(i)}(Z) = \inf_{f_i(x^{(i)})} \mathbb{E}^{(i)}[(Z - f_i(X^{(i)}))^2]$

Recall that  $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$

So,  $\text{Var}^{(i)}(Z) = \inf_{f_i(x^{(i)})} \mathbb{E}^{(i)}[(Z - f_i(X^{(i)}))^2]$

Let  $Z_i = f_i(X^{(i)})$  for any function  $f_i$ .

Recall that  $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$

$$\text{So, } \text{Var}^{(i)}(Z) = \inf_{f_i(x^{(i)})} \mathbb{E}^{(i)}[(Z - f_i(X^{(i)}))^2]$$

Let  $Z_i = f_i(X^{(i)})$  for any function  $f_i$ .

$$\text{Then, } \text{Var}^{(i)}(Z) \leq \mathbb{E}^{(i)}[(Z - Z_i)^2]$$

Recall that  $\text{Var}(Z) = \inf_u \mathbb{E}[(Z - u)^2]$

$$\text{So, } \text{Var}^{(i)}(Z) = \inf_{f_i(x^{(i)})} \mathbb{E}^{(i)}[(Z - f_i(X^{(i)}))^2]$$

Let  $Z_i = f_i(X^{(i)})$  for any function  $f_i$ .

$$\text{Then, } \text{Var}^{(i)}(Z) \leq \mathbb{E}^{(i)}[(Z - Z_i)^2]$$

Variant : “guess functions”

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[\text{Var}^{(i)}(Z)] \leq \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2]$$

## *Convex Lipschitz functions*

Suppose  $f : [a, b]^n \mapsto \mathbb{R}$  is convex, differentiable and  $L$ -Lipschitz,  
i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \ \forall \ x, y$ .

## *Convex Lipschitz functions*

Suppose  $f : [a, b]^n \mapsto \mathbb{R}$  is convex, differentiable and  $L$ -Lipschitz,  
i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \ \forall \ x, y$ .

Choose  $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$  with inf attained at  $X'_i$

## *Convex Lipschitz functions*

Suppose  $f : [a, b]^n \mapsto \mathbb{R}$  is convex, differentiable and  $L$ -Lipschitz,  
i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \ \forall \ x, y$ .

Choose  $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$  with inf attained at  $X'_i$

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$

## Convex Lipschitz functions

Suppose  $f : [a, b]^n \mapsto \mathbb{R}$  is convex, differentiable and  $L$ -Lipschitz,  
i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$ .

Choose  $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$  with inf attained at  $X'_i$

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i) \qquad 0 \leq Z - Z_i \leq -\frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$



## Convex Lipschitz functions

Suppose  $f : [a, b]^n \mapsto \mathbb{R}$  is convex, differentiable and  $L$ -Lipschitz,  
i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$ .

Choose  $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$  with inf attained at  $X'_i$

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i) \qquad 0 \leq Z - Z_i \leq -\frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$

$$0 \leq (Z - Z_i)^2 \leq \left| \frac{\partial f}{\partial x_i}(X) \right|^2 (b - a)^2$$

## Convex Lipschitz functions

Suppose  $f : [a, b]^n \mapsto \mathbb{R}$  is convex, differentiable and  $L$ -Lipschitz,  
i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$ .

Choose  $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$  with inf attained at  $X'_i$

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i) \qquad 0 \leq Z - Z_i \leq -\frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$

$$0 \leq (Z - Z_i)^2 \leq \left| \frac{\partial f}{\partial x_i}(X) \right|^2 (b - a)^2$$

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2] \leq L^2(b - a)^2$$

## Convex Lipschitz functions

Suppose  $f : [a, b]^n \mapsto \mathbb{R}$  is convex, differentiable and  $L$ -Lipschitz,  
i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$ .

Choose  $Z_i = \inf_{x_i} f(X_1, X_2, \dots, x_i, \dots, X_n)$  with inf attained at  $X'_i$

$$Z_i \geq Z + \frac{\partial f}{\partial x_i}(X)(X'_i - X_i) \qquad 0 \leq Z - Z_i \leq -\frac{\partial f}{\partial x_i}(X)(X'_i - X_i)$$

$$0 \leq (Z - Z_i)^2 \leq \left| \frac{\partial f}{\partial x_i}(X) \right|^2 (b - a)^2$$

$$\text{Var}(Z) \leq \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2] \leq L^2(b - a)^2$$

Differentiability assumption unnecessary:  
convolve  $f$  with a smooth kernel.



## Spectral norm of a random matrix

Populate an  $m \times n$  matrix  $A$  by independent entries, each taking values in  $[0, 1]$ .

## Spectral norm of a random matrix

Populate an  $m \times n$  matrix  $A$  by independent entries, each taking values in  $[0, 1]$ .

The function  $f : [0, 1]^{m \times n} \mapsto \mathbb{R}$ , given by

$$Z = f(A) = \|A\| = \sup_{\|v\|_2=1} \|Av\|_2$$

## Spectral norm of a random matrix

Populate an  $m \times n$  matrix  $A$  by independent entries, each taking values in  $[0, 1]$ .

The function  $f : [0, 1]^{m \times n} \mapsto \mathbb{R}$ , given by

$$Z = f(A) = \|A\| = \sup_{\|v\|_2=1} \|Av\|_2$$

is convex and 1-Lipschitz

## Spectral norm of a random matrix

Populate an  $m \times n$  matrix  $A$  by independent entries, each taking values in  $[0, 1]$ .

The function  $f : [0, 1]^{m \times n} \mapsto \mathbb{R}$ , given by

$$Z = f(A) = \|A\| = \sup_{\|v\|_2=1} \|Av\|_2$$

is convex and 1-Lipschitz

(hint: spectral norm  $\leq$  Frobenius norm)



## Spectral norm of a random matrix

Populate an  $m \times n$  matrix  $A$  by independent entries, each taking values in  $[0, 1]$ .

The function  $f : [0, 1]^{m \times n} \mapsto \mathbb{R}$ , given by

$$Z = f(A) = \|A\| = \sup_{\|v\|_2=1} \|Av\|_2$$

is convex and 1-Lipschitz

(hint: spectral norm  $\leq$  Frobenius norm)

With  $a = 0, b = 1, L = 1$  in previous result, we get

$$\text{Var}(Z) \leq 1.$$



Recall if  $Z, Z'$  are i.i.d., then

Recall if  $Z, Z'$  are i.i.d., then

$$\text{Var}(Z) = \frac{1}{2}\mathbb{E}[(Z - Z')^2]$$

Recall if  $Z, Z'$  are i.i.d., then

$$\text{Var}(Z) = \frac{1}{2} \mathbb{E}[(Z - Z')^2]$$

If  $Z = f(X_1, \dots, X_i, \dots, X_n)$ , and  $Z_i = f(X_1, \dots, X'_i, \dots, X_n)$   
where  $X'_i$  is an independent copy of  $X_i$ , then

Recall if  $Z, Z'$  are i.i.d., then

$$\text{Var}(Z) = \frac{1}{2} \mathbb{E}[(Z - Z')^2]$$

If  $Z = f(X_1, \dots, X_i, \dots, X_n)$ , and  $Z_i = f(X_1, \dots, X'_i, \dots, X_n)$   
where  $X'_i$  is an independent copy of  $X_i$ , then

$$\text{Var}^{(i)}(Z) = \frac{1}{2} \mathbb{E}^{(i)}[(Z - Z_i)^2]$$

Recall if  $Z, Z'$  are i.i.d., then

$$\text{Var}(Z) = \frac{1}{2} \mathbb{E}[(Z - Z')^2]$$

If  $Z = f(X_1, \dots, X_i, \dots, X_n)$ , and  $Z_i = f(X_1, \dots, X'_i, \dots, X_n)$   
where  $X'_i$  is an independent copy of  $X_i$ , then

$$\text{Var}^{(i)}(Z) = \frac{1}{2} \mathbb{E}^{(i)}[(Z - Z_i)^2]$$

Variant: “resampling coordinates”

Recall if  $Z, Z'$  are i.i.d., then

$$\text{Var}(Z) = \frac{1}{2} \mathbb{E}[(Z - Z')^2]$$

If  $Z = f(X_1, \dots, X_i, \dots, X_n)$ , and  $Z_i = f(X_1, \dots, X'_i, \dots, X_n)$   
where  $X'_i$  is an independent copy of  $X_i$ , then

$$\text{Var}^{(i)}(Z) = \frac{1}{2} \mathbb{E}^{(i)}[(Z - Z_i)^2]$$

Variant: “resampling coordinates”

$$\text{Var}(Z) \leq \frac{1}{2} \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2]$$



Recall if  $Z, Z'$  are i.i.d., then

$$\text{Var}(Z) = \frac{1}{2}\mathbb{E}[(Z - Z')^2] = \mathbb{E}[(Z - Z')_+^2], \quad (a)_+ = \max\{a, 0\}$$

If  $Z = f(X_1, \dots, X_i, \dots, X_n)$ , and  $Z_i = f(X_1, \dots, X'_i, \dots, X_n)$   
where  $X'_i$  is an independent copy of  $X_i$ , then

$$\text{Var}^{(i)}(Z) = \frac{1}{2}\mathbb{E}^{(i)}[(Z - Z_i)^2]$$

Variant: “resampling coordinates”

$$\text{Var}(Z) \leq \frac{1}{2} \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2]$$

Recall if  $Z, Z'$  are i.i.d., then

$$\text{Var}(Z) = \frac{1}{2}\mathbb{E}[(Z - Z')^2] = \mathbb{E}[(Z - Z')_+^2], \quad (a)_+ = \max\{a, 0\}$$

If  $Z = f(X_1, \dots, X_i, \dots, X_n)$ , and  $Z_i = f(X_1, \dots, X'_i, \dots, X_n)$   
where  $X'_i$  is an independent copy of  $X_i$ , then

$$\text{Var}^{(i)}(Z) = \frac{1}{2}\mathbb{E}^{(i)}[(Z - Z_i)^2] = \mathbb{E}^{(i)}[(Z - Z_i)_+^2]$$

Variant: “resampling coordinates”

$$\text{Var}(Z) \leq \frac{1}{2} \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2]$$

Recall if  $Z, Z'$  are i.i.d., then

$$\text{Var}(Z) = \frac{1}{2}\mathbb{E}[(Z - Z')^2] = \mathbb{E}[(Z - Z')_+^2], \quad (a)_+ = \max\{a, 0\}$$

If  $Z = f(X_1, \dots, X_i, \dots, X_n)$ , and  $Z_i = f(X_1, \dots, X'_i, \dots, X_n)$   
where  $X'_i$  is an independent copy of  $X_i$ , then

$$\text{Var}^{(i)}(Z) = \frac{1}{2}\mathbb{E}^{(i)}[(Z - Z_i)^2] = \mathbb{E}^{(i)}[(Z - Z_i)_+^2]$$

Variant: “resampling coordinates”

$$\text{Var}(Z) \leq \frac{1}{2} \sum_{i=1}^n \mathbb{E}[(Z - Z_i)^2] = \sum_{i=1}^n \mathbb{E}[(Z - Z_i)_+^2]$$

## Largest eigenvalue of a random matrix

Let  $A$  be an  $n \times n$  symmetric matrix with independent entries  $X_{ij}$ ,  $1 \leq i \leq j \leq n$  independent,  $-1 \leq X_{ij} \leq 1$ .

## Largest eigenvalue of a random matrix

Let  $A$  be an  $n \times n$  symmetric matrix with independent entries  $X_{ij}$ ,  $1 \leq i \leq j \leq n$  independent,  $-1 \leq X_{ij} \leq 1$ .

Let  $Z = \lambda_{\max}(A) = \max_{\|w\|=1} w^T A w = u^T A u$   
for some  $u$  that depends on the  $X_{ij}$ 's.

## Largest eigenvalue of a random matrix

Let  $A$  be an  $n \times n$  symmetric matrix with independent entries  $X_{ij}$ ,  $1 \leq i \leq j \leq n$  independent,  $-1 \leq X_{ij} \leq 1$ .

$$\text{Let } Z = \lambda_{\max}(A) = \max_{\|w\|=1} w^T A w = u^T A u$$

for some  $u$  that depends on the  $X_{ij}$ 's.

$$Z \in [-1, n] \text{ (exercise)}$$

## Largest eigenvalue of a random matrix

Let  $A$  be an  $n \times n$  symmetric matrix with independent entries  $X_{ij}$ ,  $1 \leq i \leq j \leq n$  independent,  $-1 \leq X_{ij} \leq 1$ .

Let  $Z = \lambda_{\max}(A) = \max_{\|w\|=1} w^T A w = u^T A u$   
for some  $u$  that depends on the  $X_{ij}$ 's.

$Z \in [-1, n]$  (exercise)

Let  $Z_{ij}$  denote  $\lambda_{\max}$  for the matrix  $\bar{A}^{ij}$  which is same as the matrix  $A$  except  $X_{ij} = X_{ji}$  gets replaced by an independent copy  $X'_{ij} = X'_{ji}$ .

## Largest eigenvalue of a random matrix

$$Z - Z_{ij} = u^T A u - \max_{\|w\|=1} w^T \bar{A}^{ij} w$$



## Largest eigenvalue of a random matrix

$$\begin{aligned} Z - Z_{ij} &= u^T A u - \max_{\|w\|=1} w^T \bar{A}^{ij} w \\ &\leq u^T A u - u^T \bar{A}^{ij} u \end{aligned}$$

## Largest eigenvalue of a random matrix

$$\begin{aligned} Z - Z_{ij} &= u^T A u - \max_{\|w\|=1} w^T \bar{A}^{ij} w \\ &\leq u^T A u - u^T \bar{A}^{ij} u \\ &\leq 2 \cdot |u_i| \cdot |u_j| \cdot |X_{ij} - X'_{ij}| \end{aligned}$$

## Largest eigenvalue of a random matrix

$$\begin{aligned} Z - Z_{ij} &= u^T A u - \max_{\|w\|=1} w^T \bar{A}^{ij} w \\ &\leq u^T A u - u^T \bar{A}^{ij} u \\ &\leq 2 \cdot |u_i| \cdot |u_j| \cdot |X_{ij} - X'_{ij}| \\ &\leq 4 \cdot |u_i| \cdot |u_j| \end{aligned}$$

## Largest eigenvalue of a random matrix

$$\begin{aligned} Z - Z_{ij} &= u^T A u - \max_{\|w\|=1} w^T \bar{A}^{ij} w \\ &\leq u^T A u - u^T \bar{A}^{ij} u \\ &\leq 2 \cdot |u_i| \cdot |u_j| \cdot |X_{ij} - X'_{ij}| \\ &\leq 4 \cdot |u_i| \cdot |u_j| \end{aligned}$$

$$\sum_{ij} (Z - Z_{ij})_+^2 \leq 16 \sum_{ij} u_i^2 u_j^2 = 16 \cdot \|u\|^2 \cdot \|u\|^2 = 16.$$

## Largest eigenvalue of a random matrix

$$\begin{aligned} Z - Z_{ij} &= u^T A u - \max_{\|w\|=1} w^T \bar{A}^{ij} w \\ &\leq u^T A u - u^T \bar{A}^{ij} u \\ &\leq 2 \cdot |u_i| \cdot |u_j| \cdot |X_{ij} - X'_{ij}| \\ &\leq 4 \cdot |u_i| \cdot |u_j| \end{aligned}$$

$$\sum_{ij} (Z - Z_{ij})_+^2 \leq 16 \sum_{ij} u_i^2 u_j^2 = 16 \cdot \|u\|^2 \cdot \|u\|^2 = 16.$$

Thus,  $\text{Var}(Z) \leq 16$ .

## *Suboptimality warning*

## *Suboptimality warning*

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

## *Suboptimality warning*

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

$\text{Var}(\lambda_{\max}(A)) \sim n^{-1/3}$ , i.e. *superconcentration*.



## *Suboptimality warning*

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

$$\text{Var}(\lambda_{\max}(A)) \sim n^{-1/3}, \text{ i.e. } \textit{superconcentration}.$$

- We don't get the optimal bound by a general theorem

## *Suboptimality warning*

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

$$\text{Var}(\lambda_{\max}(A)) \sim n^{-1/3}, \text{ i.e. } \textit{superconcentration}.$$

- We don't get the optimal bound by a general theorem
- But ...

## *Suboptimality warning*

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

$$\text{Var}(\lambda_{\max}(A)) \sim n^{-1/3}, \text{ i.e. } \textit{superconcentration}.$$

- We don't get the optimal bound by a general theorem
- But ...
  - we didn't employ any random matrix theory

## *Suboptimality warning*

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

$$\text{Var}(\lambda_{\max}(A)) \sim n^{-1/3}, \text{ i.e. } \textit{superconcentration}.$$

- We don't get the optimal bound by a general theorem
- But ...
  - we didn't employ any random matrix theory
  - we didn't carry out any detailed analysis

## *Suboptimality warning*

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

$$\text{Var}(\lambda_{\max}(A)) \sim n^{-1/3}, \text{ i.e. } \textit{superconcentration}.$$

- We don't get the optimal bound by a general theorem
- But ...
  - we didn't employ any random matrix theory
  - we didn't carry out any detailed analysis
- Still, we obtained a genuinely non-trivial result

## *Suboptimality warning*

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

$$\text{Var}(\lambda_{\max}(A)) \sim n^{-1/3}, \text{ i.e. } \textit{superconcentration}.$$

- We don't get the optimal bound by a general theorem
- But ...
  - we didn't employ any random matrix theory
  - we didn't carry out any detailed analysis
- Still, we obtained a genuinely non-trivial result
- In many cases, these are sufficient since they provide bounds of optimal order

## *Suboptimality warning*

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

$$\text{Var}(\lambda_{\max}(A)) \sim n^{-1/3}, \text{ i.e. } \textit{superconcentration}.$$

- We don't get the optimal bound by a general theorem
- But ...
  - we didn't employ any random matrix theory
  - we didn't carry out any detailed analysis
- Still, we obtained a genuinely non-trivial result
- In many cases, these are sufficient since they provide bounds of optimal order
- E.g. here,  $\text{Var}(\lambda_{\max}(A))$  can be  $1/4$  if entries are not identically distributed

## Suboptimality warning

In fact, if  $X_{ij}$ 's are i.i.d. equiprobable on  $\{-1, +1\}$ , then

$$\text{Var}(\lambda_{\max}(A)) \sim n^{-1/3}, \text{ i.e. } \textit{superconcentration}.$$

- We don't get the optimal bound by a general theorem
- But ...
  - we didn't employ any random matrix theory
  - we didn't carry out any detailed analysis
- Still, we obtained a genuinely non-trivial result
- In many cases, these are sufficient since they provide bounds of optimal order
- E.g. here,  $\text{Var}(\lambda_{\max}(A))$  can be  $1/4$  if entries are not **identically distributed**
- Can a general principle capture superconcentration? Active research area





A Poincaré inequality says “ $\text{variance}(f) \lesssim c\mathbb{E}[\|\text{gradient}(f)\|^2]$ ” for a suitable notion of gradient.

A Poincaré inequality says “ $\text{variance}(f) \lesssim c\mathbb{E}[\|\text{gradient}(f)\|^2]$ ” for a suitable notion of gradient. It is closely associated with mixing in Markov processes

A Poincaré inequality says “ $\text{variance}(f) \lesssim c\mathbb{E}[\|\text{gradient}(f)\|^2]$ ” for a suitable notion of gradient. It is closely associated with mixing in Markov processes

### Gaussian Poincaré inequality

If  $X \sim \mathcal{N}(0, I_n)$ , and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable, then

A Poincaré inequality says “ $\text{variance}(f) \lesssim c\mathbb{E}[\|\text{gradient}(f)\|^2]$ ” for a suitable notion of gradient. It is closely associated with mixing in Markov processes

### Gaussian Poincaré inequality

If  $X \sim \mathcal{N}(0, I_n)$ , and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable, then  $\text{Var}(f(X)) \leq \mathbb{E} [\|\nabla f(X)\|^2]$

A Poincaré inequality says “ $\text{variance}(f) \lesssim c\mathbb{E}[\|\text{gradient}(f)\|^2]$ ” for a suitable notion of gradient. It is closely associated with mixing in Markov processes

### Gaussian Poincaré inequality

If  $X \sim \mathcal{N}(0, I_n)$ , and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable, then  $\text{Var}(f(X)) \leq \mathbb{E} [\|\nabla f(X)\|^2]$

### Corollary of Gaussian Poincaré inequality

A Poincaré inequality says “ $\text{variance}(f) \lesssim c\mathbb{E}[\|\text{gradient}(f)\|^2]$ ” for a suitable notion of gradient. It is closely associated with mixing in Markov processes

### Gaussian Poincaré inequality

If  $X \sim \mathcal{N}(0, I_n)$ , and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable, then  $\text{Var}(f(X)) \leq \mathbb{E} [\|\nabla f(X)\|^2]$

### Corollary of Gaussian Poincaré inequality

If  $X \sim \mathcal{N}(0, I_n)$  and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is  $L$ -Lipschitz, i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$ , then

A Poincaré inequality says “ $\text{variance}(f) \lesssim c\mathbb{E}[\|\text{gradient}(f)\|^2]$ ” for a suitable notion of gradient. It is closely associated with mixing in Markov processes

### Gaussian Poincaré inequality

If  $X \sim \mathcal{N}(0, I_n)$ , and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable, then  $\text{Var}(f(X)) \leq \mathbb{E} [\|\nabla f(X)\|^2]$

### Corollary of Gaussian Poincaré inequality

If  $X \sim \mathcal{N}(0, I_n)$  and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is  $L$ -Lipschitz, i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$ , then

$$\text{Var}(f(X)) \leq L^2$$



A Poincaré inequality says “ $\text{variance}(f) \lesssim c\mathbb{E}[\|\text{gradient}(f)\|^2]$ ” for a suitable notion of gradient. It is closely associated with mixing in Markov processes

### Gaussian Poincaré inequality

If  $X \sim \mathcal{N}(0, I_n)$ , and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable, then  $\text{Var}(f(X)) \leq \mathbb{E} [\|\nabla f(X)\|^2]$

### Corollary of Gaussian Poincaré inequality

If  $X \sim \mathcal{N}(0, I_n)$  and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is  $L$ -Lipschitz, i.e.  $|f(x) - f(y)| \leq L\|x - y\|_2 \forall x, y$ , then

$$\text{Var}(f(X)) \leq L^2$$

Note: The bounds are tight if  $f$  is linear!

## *Gaussian Poincaré: proof*

## *Gaussian Poincaré: proof*

First, consider a 1-dimensional Gaussian  $X \sim \mathcal{N}(0, 1)$ .

## *Gaussian Poincaré: proof*

First, consider a 1-dimensional Gaussian  $X \sim \mathcal{N}(0, 1)$ .

Let  $f : \mathbb{R} \mapsto \mathbb{R}$  be twice continuously differentiable with compact support. Let  $\sup_x |f''(x)| = K$ .

## *Gaussian Poincaré: proof*

First, consider a 1-dimensional Gaussian  $X \sim \mathcal{N}(0, 1)$ .

Let  $f : \mathbb{R} \mapsto \mathbb{R}$  be twice continuously differentiable with compact support. Let  $\sup_x |f''(x)| = K$ .

Let  $Y_1, Y_2, \dots, Y_m$  be independent and equiprobable on  $\{-1, +1\}$ .

## *Gaussian Poincaré: proof*

First, consider a 1-dimensional Gaussian  $X \sim \mathcal{N}(0, 1)$ .

Let  $f : \mathbb{R} \mapsto \mathbb{R}$  be twice continuously differentiable with compact support. Let  $\sup_x |f''(x)| = K$ .

Let  $Y_1, Y_2, \dots, Y_m$  be independent and equiprobable on  $\{-1, +1\}$ .

$$\text{Let } S_m = \frac{1}{\sqrt{m}} (Y_1 + Y_2 + \dots + Y_m).$$

## *Gaussian Poincaré: proof*

First, consider a 1-dimensional Gaussian  $X \sim \mathcal{N}(0, 1)$ .

Let  $f : \mathbb{R} \mapsto \mathbb{R}$  be twice continuously differentiable with compact support. Let  $\sup_x |f''(x)| = K$ .

Let  $Y_1, Y_2, \dots, Y_m$  be independent and equiprobable on  $\{-1, +1\}$ .

$$\text{Let } S_m = \frac{1}{\sqrt{m}} (Y_1 + Y_2 + \dots + Y_m).$$

$$\text{Var}^{(i)}(f(S_m))$$

## *Gaussian Poincaré: proof*

First, consider a 1-dimensional Gaussian  $X \sim \mathcal{N}(0, 1)$ .

Let  $f : \mathbb{R} \mapsto \mathbb{R}$  be twice continuously differentiable with compact support. Let  $\sup_x |f''(x)| = K$ .

Let  $Y_1, Y_2, \dots, Y_m$  be independent and equiprobable on  $\{-1, +1\}$ .

Let  $S_m = \frac{1}{\sqrt{m}} (Y_1 + Y_2 + \dots + Y_m)$ .

$$\text{Var}^{(i)}(f(S_m)) = \frac{1}{4} \left( f \left( S_m - \frac{Y_i}{\sqrt{m}} + \frac{1}{\sqrt{m}} \right) - f \left( S_m - \frac{Y_i}{\sqrt{m}} - \frac{1}{\sqrt{m}} \right) \right)^2$$



## *Gaussian Poincaré: proof*

First, consider a 1-dimensional Gaussian  $X \sim \mathcal{N}(0, 1)$ .

Let  $f : \mathbb{R} \mapsto \mathbb{R}$  be twice continuously differentiable with compact support. Let  $\sup_x |f''(x)| = K$ .

Let  $Y_1, Y_2, \dots, Y_m$  be independent and equiprobable on  $\{-1, +1\}$ .

$$\text{Let } S_m = \frac{1}{\sqrt{m}} (Y_1 + Y_2 + \dots + Y_m).$$

$$\begin{aligned} \text{Var}^{(i)}(f(S_m)) &= \frac{1}{4} \left( f \left( S_m - \frac{Y_i}{\sqrt{m}} + \frac{1}{\sqrt{m}} \right) - f \left( S_m - \frac{Y_i}{\sqrt{m}} - \frac{1}{\sqrt{m}} \right) \right)^2 \\ &\leq \frac{1}{4} \left( \frac{2}{\sqrt{m}} |f'(S_m)| + \frac{2K}{m} \right)^2 \end{aligned}$$

## Gaussian Poincaré: proof

First, consider a 1-dimensional Gaussian  $X \sim \mathcal{N}(0, 1)$ .

Let  $f : \mathbb{R} \mapsto \mathbb{R}$  be twice continuously differentiable with compact support. Let  $\sup_x |f''(x)| = K$ .

Let  $Y_1, Y_2, \dots, Y_m$  be independent and equiprobable on  $\{-1, +1\}$ .

Let  $S_m = \frac{1}{\sqrt{m}} (Y_1 + Y_2 + \dots + Y_m)$ .

$$\begin{aligned} \text{Var}^{(i)}(f(S_m)) &= \frac{1}{4} \left( f \left( S_m - \frac{Y_i}{\sqrt{m}} + \frac{1}{\sqrt{m}} \right) - f \left( S_m - \frac{Y_i}{\sqrt{m}} - \frac{1}{\sqrt{m}} \right) \right)^2 \\ &\leq \frac{1}{4} \left( \frac{2}{\sqrt{m}} |f'(S_m)| + \frac{2K}{m} \right)^2 = \frac{1}{m} \left( |f'(S_m)| + \frac{K}{\sqrt{m}} \right)^2 \end{aligned}$$

## *Gaussian Poincaré: proof*

## *Gaussian Poincaré: proof*

$$\mathrm{Var}(f(S_m)) \leq \sum_{i=1}^m \mathbb{E} \mathrm{Var}^{(i)}(f(S_m))$$

## *Gaussian Poincaré: proof*

$$\mathrm{Var}(f(S_m)) \leq \sum_{i=1}^m \mathbb{E} \mathrm{Var}^{(i)}(f(S_m)) \leq \mathbb{E} \left[ \left( |f'(S_m)| + \frac{K}{\sqrt{m}} \right)^2 \right]$$

## *Gaussian Poincaré: proof*

$$\mathrm{Var}(f(S_m)) \leq \sum_{i=1}^m \mathbb{E} \mathrm{Var}^{(i)}(f(S_m)) \leq \mathbb{E} \left[ \left( |f'(S_m)| + \frac{K}{\sqrt{m}} \right)^2 \right]$$

As  $m \rightarrow \infty$ , we have  $S_m \rightarrow X \sim \mathcal{N}(0, 1)$  in distribution by the Central Limit Theorem.

## *Gaussian Poincaré: proof*

$$\mathrm{Var}(f(S_m)) \leq \sum_{i=1}^m \mathbb{E} \mathrm{Var}^{(i)}(f(S_m)) \leq \mathbb{E} \left[ \left( |f'(S_m)| + \frac{K}{\sqrt{m}} \right)^2 \right]$$

As  $m \rightarrow \infty$ , we have  $S_m \rightarrow X \sim \mathcal{N}(0, 1)$  in distribution by the Central Limit Theorem.

Since  $f$  and  $f'$  are continuous and bounded, we get

$$\mathrm{Var}(f(X)) \leq \mathbb{E} [f'(X)^2]$$

## *Gaussian Poincaré: proof*

$$\mathrm{Var}(f(S_m)) \leq \sum_{i=1}^m \mathbb{E} \mathrm{Var}^{(i)}(f(S_m)) \leq \mathbb{E} \left[ \left( |f'(S_m)| + \frac{K}{\sqrt{m}} \right)^2 \right]$$

As  $m \rightarrow \infty$ , we have  $S_m \rightarrow X \sim \mathcal{N}(0, 1)$  in distribution by the Central Limit Theorem.

Since  $f$  and  $f'$  are continuous and bounded, we get

$$\mathrm{Var}(f(X)) \leq \mathbb{E} [f'(X)^2]$$

Extend to all continuously differentiable functions by



## Gaussian Poincaré: proof

$$\mathrm{Var}(f(S_m)) \leq \sum_{i=1}^m \mathbb{E} \mathrm{Var}^{(i)}(f(S_m)) \leq \mathbb{E} \left[ \left( |f'(S_m)| + \frac{K}{\sqrt{m}} \right)^2 \right]$$

As  $m \rightarrow \infty$ , we have  $S_m \rightarrow X \sim \mathcal{N}(0, 1)$  in distribution by the Central Limit Theorem.

Since  $f$  and  $f'$  are continuous and bounded, we get

$$\mathrm{Var}(f(X)) \leq \mathbb{E} [f'(X)^2]$$

Extend to all continuously differentiable functions by

- Truncation of  $f$  to  $[-M, M]$  and apply dominated convergence theorem as  $M \rightarrow \infty$

## Gaussian Poincaré: proof

$$\mathrm{Var}(f(S_m)) \leq \sum_{i=1}^m \mathbb{E} \mathrm{Var}^{(i)}(f(S_m)) \leq \mathbb{E} \left[ \left( |f'(S_m)| + \frac{K}{\sqrt{m}} \right)^2 \right]$$

As  $m \rightarrow \infty$ , we have  $S_m \rightarrow X \sim \mathcal{N}(0, 1)$  in distribution by the Central Limit Theorem.

Since  $f$  and  $f'$  are continuous and bounded, we get

$$\mathrm{Var}(f(X)) \leq \mathbb{E} [f'(X)^2]$$

Extend to all continuously differentiable functions by

- Truncation of  $f$  to  $[-M, M]$  and apply dominated convergence theorem as  $M \rightarrow \infty$
- Smoothen truncated  $f$  by convolution with a sharply concentrated twice differentiable kernel with compact support

## *Gaussian Poincaré: proof*

## *Gaussian Poincaré: proof*

Now, if  $X \sim \mathcal{N}(0, I_n)$  is an  $n$ -dimensional Gaussian vector

## *Gaussian Poincaré: proof*

Now, if  $X \sim \mathcal{N}(0, I_n)$  is an  $n$ -dimensional Gaussian vector  
and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable,

## *Gaussian Poincaré: proof*

Now, if  $X \sim \mathcal{N}(0, I_n)$  is an  $n$ -dimensional Gaussian vector  
and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable,  
use tensorization of variance again.

## *Gaussian Poincaré: proof*

Now, if  $X \sim \mathcal{N}(0, I_n)$  is an  $n$ -dimensional Gaussian vector  
and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable,  
use tensorization of variance again.

$$\mathrm{Var}(f(X)) \leq \sum_{i=1}^n \mathbb{E} \left[ \mathrm{Var}^{(i)}(f(X)) \right]$$

## *Gaussian Poincaré: proof*

Now, if  $X \sim \mathcal{N}(0, I_n)$  is an  $n$ -dimensional Gaussian vector  
and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable,  
use tensorization of variance again.

$$\begin{aligned}\mathrm{Var}(f(X)) &\leq \sum_{i=1}^n \mathbb{E} \left[ \mathrm{Var}^{(i)}(f(X)) \right] \\ &\leq \sum_{i=1}^n \mathbb{E} \left[ \mathbb{E}^{(i)} \left| \frac{\partial f}{\partial x_i}(X) \right|^2 \right]\end{aligned}$$



## *Gaussian Poincaré: proof*

Now, if  $X \sim \mathcal{N}(0, I_n)$  is an  $n$ -dimensional Gaussian vector  
and  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is continuously differentiable,  
use tensorization of variance again.

$$\begin{aligned}\mathrm{Var}(f(X)) &\leq \sum_{i=1}^n \mathbb{E} \left[ \mathrm{Var}^{(i)}(f(X)) \right] \\ &\leq \sum_{i=1}^n \mathbb{E} \left[ \mathbb{E}^{(i)} \left| \frac{\partial f}{\partial x_i}(X) \right|^2 \right] \\ &= \mathbb{E} [\|\nabla f(X)\|^2]\end{aligned}$$