

Newman-Penrose Formalism Calculations

Stephani et al., Equation (15.12_k0)

Metric

$$ds^2 = x_1^2 x_3^2 dx_1 \otimes dx_1 + x_3^2 dx_2 \otimes dx_2 + \left(\frac{9 x_3^4}{\Lambda^2 x_3^8 + 12 \Lambda m x_3^5 + 6 \Lambda x_3^4 e^2 + 36 m^2 x_3^2 + 36 m x_3 e^2 + 9 e^4} \right) dx_3 \otimes dx_3 + \left(-\frac{\Lambda^2 x_3^8 + 12 \Lambda m x_3^5 + 6 \Lambda x_3^4 e^2 + 36 m^2 x_3^2 + 36 m x_3 e^2 + 9 e^4}{9 x_3^4} \right) dx_4 \otimes dx_4$$

Coordinates

$$(x^\mu) = (x_1, x_2, x_3, x_4)$$

Metric Functions

$$\Sigma = x_1$$

Null Tetrad

Covariant components

$$l = \left(-\frac{3\sqrt{2}}{2\left(\Lambda x_3^2 + \frac{6m}{x_3} + \frac{3e^2}{x_3^2}\right)} \right) dx_3 + \left(-\frac{1}{6}\sqrt{2}\Lambda x_3^2 - \frac{\sqrt{2}m}{x_3} - \frac{\sqrt{2}e^2}{2x_3^2} \right) dx_4$$

$$n = \left(\frac{3\sqrt{2}}{2\left(\Lambda x_3^2 + \frac{6m}{x_3} + \frac{3e^2}{x_3^2}\right)} \right) dx_3 + \left(-\frac{1}{6}\sqrt{2}\Lambda x_3^2 - \frac{\sqrt{2}m}{x_3} - \frac{\sqrt{2}e^2}{2x_3^2} \right) dx_4$$

$$m = \left(\frac{1}{2}\sqrt{2}x_1x_3 \right) dx_1 + \left(\frac{1}{2}i\sqrt{2}x_3 \right) dx_2$$

$$\bar{m} = \left(\frac{1}{2} \sqrt{2} x_1 x_3 \right) dx_1 + \left(-\frac{1}{2} i \sqrt{2} x_3 \right) dx_2$$

Contravariant components

$$l = \left(-\frac{1}{6} \sqrt{2} \Lambda x_3^2 - \frac{\sqrt{2} m}{x_3} - \frac{\sqrt{2} e^2}{2 x_3^2} \right) dx_3 + \left(\frac{3 x_3^2}{\sqrt{2} \Lambda x_3^4 + 6 \sqrt{2} m x_3 + 3 \sqrt{2} e^2} \right) dx_4$$

$$n = \left(\frac{1}{6} \sqrt{2} \Lambda x_3^2 + \frac{\sqrt{2} m}{x_3} + \frac{\sqrt{2} e^2}{2 x_3^2} \right) dx_3 + \left(\frac{3 x_3^2}{\sqrt{2} \Lambda x_3^4 + 6 \sqrt{2} m x_3 + 3 \sqrt{2} e^2} \right) dx_4$$

$$m = \left(\frac{\sqrt{2}}{2 x_1 x_3} \right) dx_1 + \left(\frac{i \sqrt{2}}{2 x_3} \right) dx_2$$

$$\bar{m} = \left(\frac{\sqrt{2}}{2 x_1 x_3} \right) dx_1 + \left(-\frac{i \sqrt{2}}{2 x_3} \right) dx_2$$

Directional Derivatives

$$D = -\frac{1}{6} \sqrt{2} \Lambda x_3^2 - \frac{\sqrt{2} m}{x_3} - \frac{\sqrt{2} e^2}{2 x_3^2} \partial_{x_3} + \frac{3 x_3^2}{\sqrt{2} \Lambda x_3^4 + 6 \sqrt{2} m x_3 + 3 \sqrt{2} e^2} \partial_{x_4}$$

$$\Delta = \frac{1}{6} \sqrt{2} \Lambda x_3^2 + \frac{\sqrt{2} m}{x_3} + \frac{\sqrt{2} e^2}{2 x_3^2} \partial_{x_3} + \frac{3 x_3^2}{\sqrt{2} \Lambda x_3^4 + 6 \sqrt{2} m x_3 + 3 \sqrt{2} e^2} \partial_{x_4}$$

$$\delta = \frac{\sqrt{2}}{2 x_1 x_3} \partial_{x_1} + \frac{i \sqrt{2}}{2 x_3} \partial_{x_2}$$

$$\bar{\delta} = \frac{\sqrt{2}}{2 x_1 x_3} \partial_{x_1} - \frac{i \sqrt{2}}{2 x_3} \partial_{x_2}$$

Spin Coefficients

$$\rho = \frac{1}{6} \sqrt{2} \Lambda x_3 + \frac{\sqrt{2} m}{x_3^2} + \frac{\sqrt{2} e^2}{2 x_3^3}$$

$$\sigma = 0$$

$$\kappa = 0$$

$$\mu = \frac{1}{6} \sqrt{2} \Lambda x_3 + \frac{\sqrt{2} m}{x_3^2} + \frac{\sqrt{2} e^2}{2 x_3^3}$$

$$\lambda = 0$$

$$\nu = 0$$

$$\alpha = 0$$

$$\beta = 0$$

$$\tau = 0$$

$$\pi = 0$$

$$\epsilon = -\frac{1}{6} \sqrt{2} \Lambda x_3 + \frac{\sqrt{2} m}{2 x_3^2} + \frac{\sqrt{2} e^2}{2 x_3^3}$$

$$\gamma = -\frac{1}{6} \sqrt{2} \Lambda x_3 + \frac{\sqrt{2} m}{2 x_3^2} + \frac{\sqrt{2} e^2}{2 x_3^3}$$

Ricci Tensor Components

$$\Phi_{00} = 0$$

$$\Phi_{01} = 0$$

$$\Phi_{02} = 0$$

$$\Phi_{10} = 0$$

$$\Phi_{11} = \frac{5}{36} \Lambda^2 x_3^2 - \frac{\Lambda m}{3 x_3} - \frac{\Lambda e^2}{6 x_3^2} + \frac{2 m^2}{x_3^4} + \frac{5 m e^2}{x_3^5} + \frac{9 e^4}{4 x_3^6}$$

$$\Phi_{12} = 0$$

$$\Phi_{20} = 0$$

$$\Phi_{21} = 0$$

$$\Phi_{22} = 0$$

$$\Lambda = -\frac{5}{36} \Lambda^2 x_3^2 - \frac{\Lambda m}{3 x_3} - \frac{\Lambda e^2}{18 x_3^2} - \frac{m e^2}{3 x_3^5} - \frac{e^4}{4 x_3^6}$$

Weyl Tensor Components

$$\Psi_0 = 0$$

$$\Psi_1 = 0$$

$$\Psi_2 = \frac{1}{18} \Lambda^2 x_3^2 + \frac{\Lambda e^2}{9 x_3^2} + \frac{4 m^2}{x_3^4} + \frac{20 m e^2}{3 x_3^5} + \frac{5 e^4}{2 x_3^6}$$

$$\Psi_3 = 0$$

$$\Psi_4 = 0$$

Newman-Penrose Equations

Equations are vanished.

Bianchi Identities

Equations are vanished.

Petrov Type

Type "D"