



ROYAL COLLEGE  
RED CROSS SOCIETY

# SPORTS INJURIES




## Contents

<b>Ankle Sprain .....</b>	3
<b>Knee injury: ACL tear .....</b>	9
<b>Fractures.....</b>	14
<b>Groin pull .....</b>	19
<b>Hamstring Injury.....</b>	24
<b>Achilles tendon rupture.....</b>	30
<b>Dislocated shoulder.....</b>	35
<b>Patellofemoral Syndrome.....</b>	39
<b>Hip Flexor Strain .....</b>	44
<b>Rotator cuff Injury .....</b>	48
<b>Shin splints.....</b>	52
<b>Ankle Strain .....</b>	56
<b>Tennis Elbow .....</b>	60
<b>Dislocated Finger .....</b>	64

## Ankle Sprain

- An ankle sprain is an injury to the tough bonds of tissue (ligaments) that surround and connect the bones of the leg to the foot.
- There are 3 types of ankle sprain,



### ➤ Causes


- Activities that involve explosive side-to-side motion such as tennis, basketball, soccer, rugby football, badminton.
- Carelessness and inadequate attention paid to movements of the ankles which causes unbalanced movements.
- Slow neuron muscular response to an off-balance position.
- Running on uneven surfaces.



- Weak or lax ligaments that join together the bones of the ankle joint.
- Shoes with inadequate heel support.



## ➤ Symptoms



- Swelling of the ankle



- Bruising of the ankle



- Tenderness of the ankle



- Inability to put weight on the affected ankle



- Pain in the ankle

## ➤ Providing first aid

1



Ask what happened at the moment that person was injured.

This may be difficult, especially if he/she is in a great deal of pain. However, their experience at the moment of injury may provide clues.

- Determine the severeness of the sprained ankle.


2



Grade I

Grade II

Grade III



A fracture is a bone injury that is especially common with high-speed ankle injuries. The symptoms are often similar to a grade III sprain.

**A fracture will require X-rays and professional treatment.**



3



Should get plenty of rest and should not put weight on injured ankle.

4



It helps to reduce swelling. You should apply ice every 20 to 30 minutes, 3 to 4 times per day.

5



Use elastic bandages to wrap the ankle, but not too tightly.

6



Elevate the foot with pillows as necessary to reduce swelling above the level of the heart.


7



You may need crutches, a cane, or a walker.

You may want to consider using a bandage or ankle brace to immobilize the ankle.

## ➤ Prevention



Performing strengthening exercises.




Warm up before exercising.



Avoiding high heels and wear sturdy, quality footwear.



Whether you've experienced ankle sprains before or not, bracing your ankles will reduce your chances of spraining them during athletic activities.




Build your diet around fish and other lean proteins, as well as fresh fruits and vegetables to help protect your joints.



Maintain a healthy weight to put less pressure on your joints.  
Carrying extra weight on your body puts added pressure on your joints, especially your ankles.


## Knee injury: ACL tear

- An ACL injury is a tear of the **anterior cruciate ligament (ACL)** which is one of the major ligaments in your knee. ACL injuries most commonly occur during sports such as soccer, basketball, rugby football etc.



### ➤ Causes

- Sports and fitness activities that can put stress on the knee.
- Suddenly slowing down and changing direction of an activity.
- Pivoting with your foot firmly planted.
- Landing awkwardly from a jump.
- Stopping suddenly.
- Receiving a direct blow to the knee or collision, such as a rugby tackle.




## ➤ Symptoms



## ➤ Providing first aid

- Be aware that there are three grades of ACL injury.



2



Protection is meant to prevent further injury.

3



General rest is necessary for healing.

4




Ice the wounded area by using cold packs with a barrier between them and your skin 3 to 4 times per day for 20 minutes at a time.

5



Wrap an elastic badge or compression wrap around your knee but not too tightly.

6



Lie down with your knee propped up on pillow and keep it above the level of the heart.

7



Once you do immediate care of your knee, you should reach out to a doctor.

## ➤ Prevention

- Do exercises that strengthen leg muscles, particularly hamstring exercises to ensure an overall balance in leg muscle strength.



Heel slide



Quad Sets



Passive knee extension



Wall squat with a ball



Balance and reach exercise A



Balance and reach exercise B



Knee stabilization: A



Knee stabilization: B



Knee stabilization: C



Knee stabilization: D



Resisted terminal knee extension




Warm up your muscles before exercising.



Practice proper jumping  
and landing techniques.



Recognize that certain activities,  
cause more ACL injuries.



Using a brace can help to  
protect and support your knee.



Add anti-inflammatory  
foods to your diet.



Keep your weight under  
control




Cool down after working out

## Fractures

- A bone fracture is a medical condition in which there is a partial or complete break in the continuity of the bone. In more severe cases, the bone may be broken into several pieces.

### Types of Bone Fractures



### ➤ Causes

- Practicing incorrect training or sport technique.
- Having too rapid of a training program
- Changing the surface
- Running on a track or road with sloped surface.
- Having a poor diet that has inadequate caloric intake for volume of sport.
- Having a low vitamin D level.



## ➤ Symptoms



Swelling and discoloration at the site of fracture.



Tenderness or “pinpoint pain” when touched on the bone



Pain that begins after starting an activity.

Pain that's present throughout the activity and does not go away after the activity has ended.

Pain which occurs while at rest, during normal activity.

## ➤ Providing first aid

- Move the patient to a safe location and check for other injuries.

1



2



Control any bleeding.

3



Immediately ask for medical help.

- In the meantime,

4



Immobilize the injured area. Don't try to realign the bone or push a bone that's sticking out back in.

5



Wrap an ice pack in a towel and apply for 20 minutes at a time. Never put ice directly on the skin.

6



Avoid weight-bearing activities and elevate the injured area.

## ➤ Prevention



Make sure to properly warm up and cool down before and after activities.



Strengthen your bones by building bone tissue through exercise.



Wear proper running shoes. Running shoes should be replaced each year or when they become worn out.



Start new sports activities slowly and gradually increase the time, speed and distance.



Once you feel pain, stop exercising.

Only return to exercise if you are pain-free.




Be aware of the risk factors that can lead to a fracture



Increase your intake of calcium.

Almost all of the calcium in your body gets stored in your bones and teeth, making them stronger.




Get plenty of vitamin D each day.

Your body needs vitamin D to absorb calcium correctly.

## Groin pull

- A groin pull or groin strain results from putting too much stress on muscles in your groin and thigh. If these muscles are tensed too forcefully or too suddenly, they can get over-stretched or torn.



### ➤ Causes


- Groin strain is usually caused by muscles in the groin being contracted or stretched with too much force.
- In practice, this often happens during sports where the leg is rotated or moved quickly upwards or sideways.
- It may also happen when muscles are being overused or are not warmed up. This is one of the reasons why it often affects athletes.




## ➤ Symptoms



Pain, a deep ache or spasms in the groin area or lower abdomen.



Swelling and bruising in the groin area



Weakness in the leg when trying to walk, climb stairs, or move the leg.


Limping when walking.

Difficulty performing daily activities that require standing and walking.


## ➤ Providing first aid

- Be aware that there are three grades of groin pull.


1



GRADE 1 TEAR



GRADE 2 TEAR



GRADE 3 TEAR

2



3




The severity of your groin injury determines how long you need to refrain from physical activity.

4



5



Compression helps to further reduce swelling and stabilize the injured muscle.

This helps to prevent swelling and to encourage adequate blood flow.


6




Once you do immediate care, seek medical attention.

## ➤ Prevention


- Do exercises that strengthen groin muscles.




Hip adductor stretch




Hamstring stretch in wall




Side-lying leg lift (cross over)




Straight leg raise



Hip flexion



Hip flexion



Hip abduction

Hip adduction




Always warm up your legs and groin muscles before physical activity.



Do exercises regularly to maintain healthy muscles.



Wear shoes with good support that fit well.



Always increase the intensity of your physical activity slowly.




Allow any injuries to heal before doing more physical activity.



Stop exercising if you feel pain or tightness in your groin or the inside of your thigh.

## Hamstring Injury

- A hamstring injury is a strain or tear to the tendons or large muscles at the back of the thigh.



### ➤ Causes

- Not warming up properly before exercising
- Being out of shape or overdoing it. Weak muscles are less able to handle the stress of exercise, making them more likely to get injured.
- An imbalance in the size of your leg muscles.
- Poor techniques when doing activities.
- Returning to activities too quickly after an injury.



## ➤ Symptoms



Sharp pain at the back of the thigh.



Swelling and deformity.



Sometimes bruising and discoloration can be seen along the back of the thigh.



Pain and discomfort when sitting.



Difficulty moving and bearing weight.

Following a hamstring injury, it may be hard or impossible to continue activity.

## ➤ Providing first aid

- Be aware that there are three grades of hamstring strain.

1

### HAMSTRING STRAIN

#### THREE GRADES OF HAMSTRING INJURY

GRADE 1



GRADE 2



GRADE 3



2



Get medical help for severe injuries.

3



Get off your feet as soon as you can after injury, even if you only feel mild twinges.

4



Apply cold packs to your hamstring for up to 20 minutes every 2 to 3 hours.

5



Compress or bandage the thigh to limit any swelling and movement that could cause further damage.

6



To reduce swelling, sit or lie down and prop the leg up on a tall object, so the site of the injury is higher than the heart.

7



Use crutches or a cane to immobilize the injured leg.

8



After basic first aid treatments, seek medical attention.

## ➤ Prevention

- Do hamstring stretches before you do any physical activity.

1 DOWNWARD  
FACING DOG



2 TRIANGLE



3 STANDING WIDE  
LEGGED FOLD



4 NINJA



5 PYRAMID



6 STANDING  
SPLIT



7 RAGDOLL



8 DOWNWARD  
FACING DOG



9 HALF SPLIT



10 SEATED FORWARD  
FOLD




11 RECLINED BIG  
TOE



12 CORPSE




Warm up properly before  
exercise or intense  
physical activity.



Do not overexert your  
hamstrings if you have a history  
of hamstring injuries.

- Do exercises that strengthen hamstring muscles.




If you feel pain in your thigh,  
stop your activity immediately.



Reduce your weight if you are  
overweight or obese.

## Achilles tendon rupture

- Achilles tendon rupture is an injury that affects the back of your lower leg. It mainly occurs in people playing recreational sports, but it can happen to anyone.
- The Achilles tendon is a strong fibrous cord that connects the muscles in the back of your calf to your heel bone. If you overstretch your Achilles tendon, it can tear (rupture) completely or just partially.



### ➤ Causes

- Ruptures often are caused by a sudden increase in the stress on your Achilles tendon.
- Common examples include:
  - Increasing the intensity of sports participation, especially in sports that involve jumping.
  - Falling from a height.
  - Stepping into a hole.



## ➤ Symptoms




Pain and stiffness along the Achilles tendon.



Feel for swelling on the Achilles tendon.



Tenderness when touched on the heel.



A popping or snapping sound from injured area.



Get medical help immediately to diagnose the injury after check for symptoms.

## ➤ Providing first aid

1



You have to ask medical help immediately.

- In the meantime,

2



Rest the injured area and keep weight off it.

3



Ice the injured area to ease swelling.

4



Compress injury by gently wrapping with an elastic bandage.



5



Elevate injured area above the heart level, if possible.


## ➤ Prevention

- Calf-strengthening exercises can also help the muscle and tendon absorb more force and prevent injury.



Warm up before exercising.

Alternate your routine. Alternating between high-impact exercises with lower impact exercise.



Assess what exercises may cause injury.



Use a protective heel pad specially designed to alleviate stress on the Achilles tendon.



Stop your activity if you notice any pain in your Achilles tendons and the surrounding area.



Wear proper shoes.




Maintain a healthy weight.

Extra weight adds stress to the Achilles tendons and may contribute to serious injuries.

## Dislocated shoulder

- A dislocated shoulder is an injury in which your upper arm bone pops out of the cup-shaped socket that's part of your shoulder blade. The shoulder is the body's most mobile joint, which makes it susceptible to dislocation.

### Shoulder Dislocation



Normal  
anatomy

Anterior  
dislocation

Posterior  
dislocation

### ➤ Causes

- Any hard fall onto the shoulder.
- Forceful hitting, lifting, or throwing.
- A hit to an outstretched arm.
- Athletes at risk of shoulder dislocations include:
  - Football players
  - Hockey players
  - Badminton players
  - Tennis players




## ➤ Symptoms



Severe shoulder pain.



Limited motion of the shoulder.



A distortion in the contour of the shoulder.



A hard knob under the skin near the shoulder.



Shoulder bruising or abrasions if an impact has caused your injury.

## ➤ Providing first aid

1



Get medical help right away for a shoulder that appears dislocated.

- While you're waiting for medical attention:

2



Advise the casualty to stay still. Don't try to move the dislocated joint or force it back into place.

3



Ice the dislocated shoulder to reduce pain and swelling.

4



Compress the shoulder by strapping it.

5




Use a sling or shoulder immobilizer to prevent further injury.

## ➤ Prevention



Warm up properly before exercise.



Exercise regularly to maintain flexibility in joints and muscles.



Wear protective gear when you play contact sports.



Be aware of the risk factors that can lead to a dislocated shoulder.

- Once you've dislocated your shoulder joint, you may be more susceptible to future shoulder dislocations. To avoid a recurrence, follow the specific strength and stability exercises that you and your doctor have discussed for your injury.




Resisted shoulder adduction



Resisted shoulder flexion




Resisted shoulder extension



Latissimus dorsi strengthening

## Patellofemoral Syndrome

- Knee injuries or Patellofemoral syndrome, is caused by the kneecap repeatedly moving against the leg bone. This movement damages the kneecap's tissues and causes pain. Basketball, cycling, swimming, rugby football, volleyball and running are the most common sports where these injuries occur.



### ➤ Causes


- Running or jumping sports puts repetitive stress on your knee joint, which can cause irritation under the kneecap.
- Muscle imbalances or weaknesses.
- Trauma to the kneecap, such as a dislocation or fracture, has been linked to patellofemoral pain syndrome.



## ➤ Symptoms



Painful crepitus of the knee.




Crackling or popping sounds  
in the knee.



Swelling.

## ➤ Providing first aid

1



Rest your knee for a few  
days after you begin to notice  
the pain.

2



Ice your knee to ease pain  
and swelling.

3




Wrap your knee.  
Use an elastic bandage,  
patellar straps, or sleeves to  
give it extra support.

4



Elevate your leg on a pillow  
when you sit or lie down.

5



Wear medical shoe inserts.

6



See a doctor after the basic  
first aid treatments.

## ➤ Prevention



Warm up before running or  
other exercises.



Wear a brace to protect  
knee.



Practice shoe smarts. Make sure your shoes fit well and provide good shock absorption.

Lose excess weight. If you're overweight, losing weight relieves stress on your knees.









Be aware of the risk factors.




If you feel pain in your thigh, stop your activity immediately.

- Maintain strength. Strong quadriceps and hip abductor muscles help keep the knee balanced during activity.

**Hamstring stretch****Side-lying leg lift****Quadriceps stretch****Quadriceps isometrics****Straight leg raise****Step-up****Wall squat with ball****A****B****A****B****C****D****Resisted knee extension****Knee stabilization**

## Hip Flexor Strain

- The hip flexor is a group of muscles that assist with the upward movement of your leg or knee. An injury occurs when these muscles are torn or stretched too far.



### ➤ Causes

- A hip flexor strain is typically caused by overuse.
- Other causes of hip flexor tears or strains:
  - Extended periods sitting (muscles in the hip stay contracted for a long period of time).
  - Weaknesses in muscles surrounding the hip flexors.
  - Acute contraction of the muscles.



## ➤ Symptoms



Sudden, sharp pain in the hip or pelvis after trauma to the area.



A cramping or clenching sensation in the muscles of the upper leg area.



Swelling or bruising around the hip or thigh area.



Reduced mobility and discomfort when moving.



Tightness or stiffness around the hip area.



Inability to continue kicking, jumping, or sprinting.

## ➤ Providing first aid

- Be aware that there are three grades of hip flexor strain.

1



HEALTHY HIP FLEXOR

GRADE 1 TEAR

GRADE 2 TEAR


GRADE 3 TEAR

2



Take a break from unnecessary activities.

3




Apply ice or a cold pack to the hip immediately.

4



Wrap your hip with an elastic bandage in order to reduce swelling.

5



Elevate your hip on a pillow when you are lying down.

6



Walk with a cane or crutches to take the pressure off your hip.

7



See a doctor after the immediate treatment.

## ➤ Prevention



Warm up and cool down before and after work outs.



Maintain a proper weight.




Wear proper shoes.



Stop your activity if you feel any pain.

## Rotator cuff Injury

- The rotator cuff is a group of muscles and tendons that surround the shoulder joint, keeping the head of your upper arm bone firmly within the shallow socket of the shoulder. A rotator cuff injury can cause a dull ache in the shoulder, which often worsens with use of the arm away from the body.



### ➤ Causes

- Rotator cuff disease may be the result of either a substantial injury to the shoulder or to progressive degeneration or wear and tear of the tendon tissue.
- Repetitive overhead activity, heavy lifting over a prolonged period of time may irritate damage the tendon.
- Excessive or awkward heavy lifting, pushing or pulling.
- Falling onto an out-stretched arm.
- Overuse injury commonly seen in throwing and racquet sports such as cricket, baseball and tennis.



## ➤ Symptoms



Pain in the shoulder



Progressive weakness of the shoulder.



Pain or weakness when lift or extend the arm.



Listen for grating, clicking, or popping when moving shoulder.



Redness, swelling, or tenderness around the shoulder joint.

## ➤ Providing first aid

1



Rest the shoulder by avoiding painful activities.

2



Apply ice to decrease swelling.

3



A sling may help support the arm rest the rotator cuff muscles.

4



After basic first aid treatments, seek medical attention.

## ➤ Prevention



Warm up and stretch your shoulders before exercising.



Avoid activities with repetitive overhead arm action.




Keep good posture.

Be aware of the risk factors that can lead to a rotator cuff injury.


- Do rotator cuff strengthening exercises to help the muscle and tendon absorb more force and prevent injury.



Tubing exercise for external rotation



Tubing exercise for internal rotation




Scaption



Sidelying external rotation




Horizontal abduction



Push-ups with a plus

## Shin splints

- Shin splints are injuries that happen to the shin bone(tibia), the large bone in the front of the lower leg. This injury is medical known as medial tibial stress syndrome.



### ➤ Causes

- shin splints often occur in athletes who have recently intensified or changed their training routines.
- Anatomical abnormalities (eg: flat foot syndrome)
- Activities that involve sudden movements to the legs such as tennis, football, badminton.
- Inadequate calcium in bones.
- Running on uneven surfaces.
- Tackles to the shin and kicking heavy objects with the shin.



## ➤ Symptoms



Pain along the shin bone.



Swelling in the lower leg.



Tenderness of the inner side  
of the shin bone.



Pain when moving the leg.



Inability to put weight  
on the leg.

## ➤ Providing first aid

1



Rest by taking an adequate amount of time to heal and avoiding physical activity.

2



Use ice packs to reduce swellings and use it for 20 minutes intervals about 4 times per day.

3



Wear elastic compression bandages.

4



See a doctor after the basic first aid treatments.

## ➤ Prevention



Wearing shoes that fit well and offer good support.




Warming up before exercising.



Know when to stop activities.



Replace your shoes once a year or when they are worn out.



Fit your shoes with shock-absorbing insoles to reduce impact.



Increasing exercise intensity gradually.



Work out on soft surfaces.

## Ankle Strain

- An ankle strain refers to damage to muscles and tendons in the ankle as a result of being pulled or stretched too far.



### ➤ Causes

- Repetitive movement of the muscles and tendons over a long period of time.
- Weak muscles or tendons that cross the Ankle joint.
- Activities that involve lifting weights and the sports that require physical contact (contact sports) such as martial arts, rugby football, Soccer etc.
- Improper body mechanics with any activity.



## ➤ Symptoms



Swelling of ankle.



Limited flexibility of ankle.



Muscle spasm of ankle.



Bruising of the ankle.

## ➤ Providing first aid

1



Rest by taking an adequate amount of time to heal and avoiding physical activity.

2



Ice your ankle 4 to 8 times per day for 20 minutes at a time.

3



Wrap an elastic bandage around the ankle to prevent swelling and inflammation, which delay healing.

4



Use pillow or other devices to raise the ankle while you rest.

5



Keeping pressure off your ankle by supporting it with crutches or a cane.

6



Consult a doctor after the basic first aid treatment.

## ➤ Prevention



Warm up before exercising.



Keep your movement low impact.



Wear proper shoes that fit well and offer good support.



Replace your shoes once a year or when they are worn out.




Recognize that certain activities, cause more ankle strain.



Maintain a healthy weight to put less pressure on your joints.


## Tennis Elbow

- Tennis Elbow is inflammation or, in some cases, micro tearing of the tendons that join the forearm muscles on the outside of the elbow. The forearm muscles and tendons become damaged from overuse; repeating the same motions again and again. This leads to pain and tenderness on the outside of the elbow.



### ➤ Causes

- Tennis elbow is mostly caused by overusing your forearm due to a repetitive or strenuous activity. It can also sometimes occur after banging or knocking your elbow. If the muscles in your forearm are strained, tiny tears and inflammation can develop near the bony lump (lateral epicondyle) on the outside of your elbow.



## ➤ Symptoms

- Will feel pain while,



On the outside of the elbow,  
which may travel down the  
forearm when lifting or  
bending your arm.



When gripping small objects.



When twisting your forearm.

## ➤ Providing first aid

1



Avoiding all activity that could potentially aggravate your injury.

2



Apply the ice to your elbow region for approximately 10-20 minutes.

3



An elbow brace may be recommended to help and support elbow tendons

4



Raise and rest your arm so that it is above the level of your heart.

5



Get medical help after providing first aid.

## ➤ Prevention



Do proper warm ups and stretches before engaging in sports.



Use sports equipment with bearable weight and flexibility.



Follow proper techniques when playing sports such as Tennis.




Wear a brace to protect the elbow.



Avoid heavy lifting over a prolonged period of time.

## Dislocated Finger

- Every finger has three joints. The thumb has two joints. These joints allow our fingers to bend and straighten. When any two bones are forced out of place at the joint, such as by a traumatic sports injury or a fall, the finger becomes dislocated.



### ➤ Causes

- Sports injuries such as getting hit on the finger by a ball.
- Fingers getting caught on sports equipment.
- Falling on to the ground and using hands to break the fall.
- Accidents like closing a door on the finger.
- Some people are born with weak ligaments they are also prone to finger dislocations.



## ➤ Symptoms



Numbness of the finger.



Crookedness of the finger.




Swelling of the finger.



Bruising of the finger.



Pain in the finger.



Inability to move the finger.

## ➤ Providing First aid


- Seek medical help as soon as possible.

1



- In the meantime,

2




Do not try to move the bone  
back into place.

3



Control if there is bleeding.

4



Remove any jewellery rings  
from the affected finger.

5




Apply an ice pack to the injured  
finger.

6



Elevate the hand above the level of your heart.

7



Immobilize the affected finger with a splint to prevent any further injury.

## ➤ Prevention



Wear protective gloves when playing sports.



Removing rings or other jewellery when playing sports.



Practice hand exercises prescribed by a physiotherapist to prevent further injuries.

## Acknowledgement

- To our Principal Mr. M. V. S. Gunathilaka, Senior Games Master Mr. M. A. M. Riyaz, Senior Master of Clubs and Societies Mr. T. D. C. P. Amaratunga, Teacher in Charge Mrs. Jasmine Wickramatilake and Assistant Teacher in Charge Mrs. Saranga Rajapakshe for their guidance.
- To Dr. Ajith Tennakoon, President of the Royal College Doctors' Association, Dr. Thushantha Wijesinghe and the Royal College Doctors' Association for their assistance.
- To the Royal College Prefects' Council 20/21, Deputy Head Prefect of Clubs and Societies and Clubs Committee for their guidance and support.