


# Lectures 12: Savitch's Theorem



# Lectures 12: Savitch's Theorem

We will:

# Lectures 12: Savitch's Theorem

We will:

- Prove Savitch's Theorem:  $SPACE(f(n)) \subseteq SPACE((f(n))^2)$ .

# Lectures 12: Savitch's Theorem

We will:

- Prove Savitch's Theorem:  $SPACE(f(n)) \subseteq SPACE((f(n))^2)$ .
- Conclude that  $PSPACE = NPSPACE$ .

# Savitch's Theorem

## Theorem

$$f(n) \geq n \implies \text{NSPACE}(f(n)) \subseteq \text{SPACE}(f^2(n)).$$

## Proof.

We convert a NDTM machine  $N$  into an equivalent DTM  $M$  while only squaring the space.

# Savitch's Theorem

## Theorem

$$f(n) \geq n \implies \text{NSPACE}(f(n)) \subseteq \text{SPACE}(f^2(n)).$$

## Proof.

We convert a NDTM machine  $N$  into an equivalent DTM  $M$  while only squaring the space.

For configurations  $c, c'$  and  $k \in \mathbb{N}$ , we want to know if  $N$  can get from  $c$  to  $c'$  in at most  $k$  steps.

# Savitch's Theorem

## Theorem

$$f(n) \geq n \implies \text{NSPACE}(f(n)) \subseteq \text{SPACE}(f^2(n)).$$

## Proof.

We convert a NDTM machine  $N$  into an equivalent DTM  $M$  while only squaring the space.

For configurations  $c, c'$  and  $k \in \mathbb{N}$ , we want to know if  $N$  can get from  $c$  to  $c'$  in at most  $k$  steps. Let  $c = (w, q, \ell)$  and  $c' = (w', q', \ell')$

|      | Step     | State     | Head  | $T_0$     | $T_1$     | $T_2$     | ... | $T_{n-1}$   | $T_n$     | ...      | $T_{f(n)}$   |
|------|----------|-----------|-------|-----------|-----------|-----------|-----|-------------|-----------|----------|--------------|
| $c$  | 0        | $q$       | $h$   | $w_0$     | $w_1$     | $w_2$     |     | $w_n$       | $w_{n+1}$ |          | $w_{f(n)}$   |
|      | 0        | $q_{i_0}$ | $h_0$ | $w_{0,0}$ | $w_{0,1}$ | $w_{0,2}$ |     | $w_{0,n-1}$ | $w_{0,n}$ |          | $w_{0,f(n)}$ |
|      | 1        | $q_{i_1}$ | $h_1$ | $w_{1,0}$ | $w_{1,1}$ | $w_{1,2}$ |     | $w_{1,n-1}$ | $w_{1,n}$ |          | $w_{1,f(n)}$ |
|      | 2        | $q_{i_2}$ | $h_2$ | $w_{2,0}$ | $w_{2,1}$ | $w_{2,2}$ |     | $w_{2,n-1}$ | $w_{2,n}$ |          | $w_{2,f(n)}$ |
|      | $\vdots$ |           |       |           |           |           |     |             |           | $\ddots$ |              |
|      | $k$      | $q_{i_k}$ | $h_k$ | $w_{k,0}$ | $w_{k,1}$ | $w_{k,2}$ |     | $w_{k,n-1}$ | $w_{k,n}$ |          | $w_{k,f(n)}$ |
| $c'$ | $k$      | $q'$      | $h'$  | $w'_0$    | $w'_1$    | $w'_2$    |     | $w'_{n-1}$  | $w'_n$    |          | $w'_{f(n)}$  |

# Savitch's Theorem

## Theorem

$$f(n) \geq n \implies \text{NSPACE}(f(n)) \subseteq \text{SPACE}(f^2(n)).$$

## Proof.

We convert a NDTM machine  $N$  into an equivalent DTM  $M$  while only squaring the space.

For configurations  $c, c'$  and  $k \in \mathbb{N}$ , we want to know if  $N$  can get from  $c$  to  $c'$  in at most  $k$  steps. Let  $c = (w, q, \ell)$  and  $c' = (w', q', \ell')$

|      | Step     | State     | Head  | $T_0$     | $T_1$     | $T_2$     | ... | $T_{n-1}$   | $T_n$     | ...      | $T_{f(n)}$   |
|------|----------|-----------|-------|-----------|-----------|-----------|-----|-------------|-----------|----------|--------------|
| $c$  | 0        | $q$       | $h$   | $w_0$     | $w_1$     | $w_2$     |     | $w_n$       | $w_{n+1}$ |          | $w_{f(n)}$   |
|      | 0        | $q_{i_0}$ | $h_0$ | $w_{0,0}$ | $w_{0,1}$ | $w_{0,2}$ |     | $w_{0,n-1}$ | $w_{0,n}$ |          | $w_{0,f(n)}$ |
|      | 1        | $q_{i_1}$ | $h_1$ | $w_{1,0}$ | $w_{1,1}$ | $w_{1,2}$ |     | $w_{1,n-1}$ | $w_{1,n}$ |          | $w_{1,f(n)}$ |
|      | 2        | $q_{i_2}$ | $h_2$ | $w_{2,0}$ | $w_{2,1}$ | $w_{2,2}$ |     | $w_{2,n-1}$ | $w_{2,n}$ |          | $w_{2,f(n)}$ |
|      | $\vdots$ |           |       |           |           |           |     |             |           | $\ddots$ |              |
|      | $k$      | $q_{i_k}$ | $h_k$ | $w_{k,0}$ | $w_{k,1}$ | $w_{k,2}$ |     | $w_{k,n-1}$ | $w_{k,n}$ |          | $w_{k,f(n)}$ |
| $c'$ | $k$      | $q'$      | $h'$  | $w'_0$    | $w'_1$    | $w'_2$    |     | $w'_{n-1}$  | $w'_n$    |          | $w'_{f(n)}$  |

Almost looks like a ladder!



# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

Function Reachable( $c, c', k$ )      # Can we reach from  $c$  to  $c'$  in  $k$  steps?

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

Function  $\text{Reachable}(c, c', k)$       # Can we reach from  $c$  to  $c'$  in  $k$  steps?

If ( $k = 1$ )

Return ( $c'$  can be reached from  $c$  in  $M$ )  $\vee$  ( $c = c'$ )

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

Function  $\text{Reachable}(c, c', k)$       # Can we reach from  $c$  to  $c'$  in  $k$  steps?

If ( $k = 1$ )

Return ( $c'$  can be reached from  $c$  in  $M$ )  $\vee$  ( $c = c'$ )

For each possible configuration  $c_{\text{mid}}$

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

Function  $\text{Reachable}(c, c', k)$     # Can we reach from  $c$  to  $c'$  in  $k$  steps?

If ( $k = 1$ )

Return ( $c'$  can be reached from  $c$  in  $M$ )  $\vee$  ( $c = c'$ )

For each possible configuration  $c_{\text{mid}}$

If ( $\text{Reachable}(c, c_{\text{mid}}, \lceil k/2 \rceil) \wedge \text{Reachable}(c_{\text{mid}}, c', \lfloor k/2 \rfloor)$ )

Return T

Return F

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

- We can tweak the machine to get a single accepting  $c_Y$  configuration by reversing the head and erasing the tape before halting.

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

- We can tweak the machine to get a single accepting  $c_Y$  configuration by reversing the head and erasing the tape before halting.
- Since  $N$  uses  $f(n)$  space, there are just  $C = |Q| \times f(n) \times |\Gamma|^{f(n)}$  is the number of possible configurations.

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

- We can tweak the machine to get a single accepting  $c_Y$  configuration by reversing the head and erasing the tape before halting.
- Since  $N$  uses  $f(n)$  space, there are just  $C = |Q| \times f(n) \times |\Gamma|^{f(n)}$  is the number of possible configurations.
- We start by calling  $\text{Reachable}(c_0, c_Y, C)$ , where  $c_0$  encodes the (input-specific!) starting configuration.

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

- We can tweak the machine to get a single accepting  $c_Y$  configuration by reversing the head and erasing the tape before halting.
- Since  $N$  uses  $f(n)$  space, there are just  $C = |Q| \times f(n) \times |\Gamma|^{f(n)}$  is the number of possible configurations.
- We start by calling  $\text{Reachable}(c_0, c_Y, C)$ , where  $c_0$  encodes the (input-specific!) starting configuration. **Why is this enough to determine if  $N$  accepts?**



# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

- We can tweak the machine to get a single accepting  $c_Y$  configuration by reversing the head and erasing the tape before halting.
- Since  $N$  uses  $f(n)$  space, there are just  $C = |Q| \times f(n) \times |\Gamma|^{f(n)}$  is the number of possible configurations.
- We start by calling  $\text{Reachable}(c_0, c_Y, C)$ , where  $c_0$  encodes the (input-specific!) starting configuration. **Why is this enough to determine if  $N$  accepts?**
- The depth of the recursion is  $\log_2(C) = O(f(n))$ .

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

- We can tweak the machine to get a single accepting  $c_Y$  configuration by reversing the head and erasing the tape before halting.
- Since  $N$  uses  $f(n)$  space, there are just  $C = |Q| \times f(n) \times |\Gamma|^{f(n)}$  is the number of possible configurations.
- We start by calling  $\text{Reachable}(c_0, c_Y, C)$ , where  $c_0$  encodes the (input-specific!) starting configuration. **Why is this enough to determine if  $N$  accepts?**
- The depth of the recursion is  $\log_2(C) = O(f(n))$ .
- Each time we store one configuration, requiring  $O(f(n))$  space.

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

- We can tweak the machine to get a single accepting  $c_Y$  configuration by reversing the head and erasing the tape before halting.
- Since  $N$  uses  $f(n)$  space, there are just  $C = |Q| \times f(n) \times |\Gamma|^{f(n)}$  is the number of possible configurations.
- We start by calling  $\text{Reachable}(c_0, c_Y, C)$ , where  $c_0$  encodes the (input-specific!) starting configuration. **Why is this enough to determine if  $N$  accepts?**
- The depth of the recursion is  $\log_2(C) = O(f(n))$ .
- Each time we store one configuration, requiring  $O(f(n))$  space.
- Therefore, the total space is  $O(f^2(n))$ . □

# Savitch's Theorem:

$$f(n) \geq n \implies NSPACE(f(n)) \subseteq SPACE(f^2(n))$$

## Proof (Cont.)

- We can tweak the machine to get a single accepting  $c_Y$  configuration by reversing the head and erasing the tape before halting.
- Since  $N$  uses  $f(n)$  space, there are just  $C = |Q| \times f(n) \times |\Gamma|^{f(n)}$  is the number of possible configurations.
- We start by calling  $\text{Reachable}(c_0, c_Y, C)$ , where  $c_0$  encodes the (input-specific!) starting configuration. **Why is this enough to determine if  $N$  accepts?**
- The depth of the recursion is  $\log_2(C) = O(f(n))$ .
- Each time we store one configuration, requiring  $O(f(n))$  space.
- Therefore, the total space is  $O(f^2(n))$ . □

**Conclusion:**  $PSPACE = NPSPACE$ .

## Definition

A problem  $B$  is PSPACE-Complete if:

- $B \in PSPACE$ .
- For all  $A \in PSPACE : A \leq_p B$ .


# PSPACE-Completeness

## Definition

A problem  $B$  is PSPACE-Complete if:

- $B \in PSPACE$ .
- For all  $A \in PSPACE : A \leq_p B$ .

To ponder: why do we use  $\leq_p$ ?



# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE$  – *Complete*, and assuming that  $TQBF \in NP$ , which of the following are true:

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE$  – *Complete*, and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ .



# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE$  – *Complete*, and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ .
- $NP = PSPACE$ .

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE$  – *Complete*, and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ .
- $NP = PSPACE$ .
- $P = NP$ .

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE$  – *Complete*, and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ .
- $NP = PSPACE$ .
- $P = NP$ .
- $NP = co-NP$ .

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE$  – *Complete*, and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ .
- $NP = PSPACE$ .
- $P = NP$ .
- $NP = co-NP$ .
- $co-NP = PSPACE$ .

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE - Complete$ , and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ .
- $NP = PSPACE$ .
- $P = NP$ .
- $NP = co-NP$ .
- $co-NP = PSPACE$ .
- SAT is PSPACE-Complete.

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE - Complete$ , and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ .
- $NP = PSPACE$ .
- $P = NP$ .
- $NP = co-NP$ .
- $co-NP = PSPACE$ .
- SAT is PSPACE-Complete.
- BEQ is NP-Complete.

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE - Complete$ , and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ .
- $NP = PSPACE$ .
- $P = NP$ .
- $NP = co-NP$ .
- $co-NP = PSPACE$ .
- SAT is PSPACE-Complete.
- BEQ is NP-Complete.

Answer on Mentimeter:



# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE - Complete$ , and assuming that  $TQBF \in NP$ , which of the following are true:



# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE$  – *Complete*, and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ . No, this doesn't imply that  $TQBF \in P$ .

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE - Complete$ , and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ . No, this doesn't imply that  $TQBF \in P$ .
- $NP = PSPACE$ . Yes, all PSPACE problems are reducible to TQBF as its PSPACE-Complete.

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE - Complete$ , and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ . No, this doesn't imply that  $TQBF \in P$ .
- $NP = PSPACE$ . Yes, all PSPACE problems are reducible to TQBF as its PSPACE-Complete.
- $P = NP$ . No.

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE - Complete$ , and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ . No, this doesn't imply that  $TQBF \in P$ .
- $NP = PSPACE$ . Yes, all PSPACE problems are reducible to TQBF as its PSPACE-Complete.
- $P = NP$ . No.
- $NP = co-NP$ . Yes! PSPACE is closed under complementation.

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE - Complete$ , and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ . No, this doesn't imply that  $TQBF \in P$ .
- $NP = PSPACE$ . Yes, all PSPACE problems are reducible to TQBF as its PSPACE-Complete.
- $P = NP$ . No.
- $NP = co-NP$ . Yes! PSPACE is closed under complementation.
- $co-NP = PSPACE$ . Yes.

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE$  – Complete, and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ . No, this doesn't imply that  $TQBF \in P$ .
- $NP = PSPACE$ . Yes, all PSPACE problems are reducible to TQBF as its PSPACE-Complete.
- $P = NP$ . No.
- $NP = co-NP$ . Yes! PSPACE is closed under complementation.
- $co-NP = PSPACE$ . Yes.
- SAT is PSPACE-Complete. Yes.

# PSPACE-Completeness of TQBF

Knowing that  $TQBF \in PSPACE$  – Complete, and assuming that  $TQBF \in NP$ , which of the following are true:

- $P = PSPACE$ . No, this doesn't imply that  $TQBF \in P$ .
- $NP = PSPACE$ . Yes, all PSPACE problems are reducible to TQBF as its PSPACE-Complete.
- $P = NP$ . No.
- $NP = co-NP$ . Yes! PSPACE is closed under complementation.
- $co-NP = PSPACE$ . Yes.
- SAT is PSPACE-Complete. Yes.
- BEQ is NP-Complete. Yes!  $\overline{BEQ}$  is NP-Complete (why?). For any  $L \in NP$ , we have thus  $\overline{L} \leq_p TQBF \leq_p \overline{BEQ}$  which gives  $L \leq_p BEQ$ .

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch.

Recall the proof of the Cook-Levin theorem:



# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch.

Recall the proof of the Cook-Levin theorem:

| Step     | State          | Head       | $T_0$        | $T_1$        | $T_2$        | ... | $T_{n-1}$         | $T_n$        | ...      | $T_N$        |
|----------|----------------|------------|--------------|--------------|--------------|-----|-------------------|--------------|----------|--------------|
| 0        | $q_0$          | 0          | $w_0$        | $w_1$        | $w_2$        |     | $w_{n-1}$         | $\sqcup$     |          | $\sqcup$     |
| 0        | $q_{i_0}$      | $h_0$      | $w_{0,0}$    | $w_{0,1}$    | $w_{0,2}$    |     | $w_{0,n-1}$       | $w_{0,n}$    |          | $w_{0,N}$    |
| 1        | $q_{i_1}$      | $h_1$      | $w_{1,0}$    | $w_{1,1}$    | $w_{1,2}$    |     | $w_{1,n-1}$       | $w_{1,n}$    |          | $w_{1,N}$    |
| 2        | $q_{i_2}$      | $h_2$      | $w_{2,0}$    | $w_{2,1}$    | $w_{2,2}$    |     | $w_{2,n-1}$       | $w_{2,n}$    |          | $w_{2,N}$    |
| $\vdots$ |                |            |              |              |              |     |                   |              | $\ddots$ |              |
| $f(N)$   | $q_{i_{f(N)}}$ | $h_{f(N)}$ | $w_{f(N),0}$ | $w_{f(N),1}$ | $w_{f(N),2}$ |     | $w_{f(N),f(N)-1}$ | $w_{f(N),n}$ |          | $w_{f(N),N}$ |

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch.

Recall the proof of the Cook-Levin theorem:

| Step     | State          | Head       | $T_0$        | $T_1$        | $T_2$        | ... | $T_{n-1}$         | $T_n$        | ...      | $T_N$        |
|----------|----------------|------------|--------------|--------------|--------------|-----|-------------------|--------------|----------|--------------|
| 0        | $q_0$          | 0          | $w_0$        | $w_1$        | $w_2$        |     | $w_{n-1}$         | $\sqcup$     |          | $\sqcup$     |
| 0        | $q_{i_0}$      | $h_0$      | $w_{0,0}$    | $w_{0,1}$    | $w_{0,2}$    |     | $w_{0,n-1}$       | $w_{0,n}$    |          | $w_{0,N}$    |
| 1        | $q_{i_1}$      | $h_1$      | $w_{1,0}$    | $w_{1,1}$    | $w_{1,2}$    |     | $w_{1,n-1}$       | $w_{1,n}$    |          | $w_{1,N}$    |
| 2        | $q_{i_2}$      | $h_2$      | $w_{2,0}$    | $w_{2,1}$    | $w_{2,2}$    |     | $w_{2,n-1}$       | $w_{2,n}$    |          | $w_{2,N}$    |
| $\vdots$ |                |            |              |              |              |     |                   |              | $\ddots$ |              |
| $f(N)$   | $q_{i_{f(N)}}$ | $h_{f(N)}$ | $w_{f(N),0}$ | $w_{f(N),1}$ | $w_{f(N),2}$ |     | $w_{f(N),f(N)-1}$ | $w_{f(N),n}$ |          | $w_{f(N),N}$ |

The challenge is that now we have a PSPACE machine, so while space is bounded by  $N = n^{o(1)}$ , the time could be  $f(N) = 2^{O(N)} = 2^{n^{O(1)}}$ , so the same reduction would not work (the formula will have exponential size).

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

| Step   | State      | Head       | $T_0$        | $T_1$        | $T_2$        | ... | $T_{n-1}$      | $T_n$        | ... | $T_N$        |
|--------|------------|------------|--------------|--------------|--------------|-----|----------------|--------------|-----|--------------|
| 0      | $q_0$      | 0          | $w_0$        | $w_1$        | $w_2$        |     | $w_{n-1}$      | $\square$    |     | $\square$    |
| 0      | $q_0$      | $h_0$      | $w_{0,0}$    | $w_{0,1}$    | $w_{0,2}$    |     | $w_{0,n-1}$    | $w_{0,n}$    |     | $w_{0,N}$    |
| 1      | $q_1$      | $h_1$      | $w_{1,0}$    | $w_{1,1}$    | $w_{1,2}$    |     | $w_{1,n-1}$    | $w_{1,n}$    |     | $w_{1,N}$    |
| 2      | $q_2$      | $h_2$      | $w_{2,0}$    | $w_{2,1}$    | $w_{2,2}$    |     | $w_{2,n-1}$    | $w_{2,n}$    |     | $w_{2,N}$    |
| ...    |            |            |              |              |              |     |                |              | ... |              |
| $f(N)$ | $q_{f(N)}$ | $h_{f(N)}$ | $w_{f(N),0}$ | $w_{f(N),1}$ | $w_{f(N),2}$ |     | $w_{f(N),n-1}$ | $w_{f(N),n}$ |     | $w_{f(N),N}$ |

Instead, we can think about this similarly to a word ladder.

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

| Step   | State      | Head       | $T_0$        | $T_1$        | $T_2$        | ... | $T_{n-1}$      | $T_n$        | ... | $T_N$        |
|--------|------------|------------|--------------|--------------|--------------|-----|----------------|--------------|-----|--------------|
| 0      | $q_0$      | 0          | $w_0$        | $w_1$        | $w_2$        |     | $w_{n-1}$      | $\square$    |     | $\square$    |
| 0      | $q_0$      | $h_0$      | $w_{0,0}$    | $w_{0,1}$    | $w_{0,2}$    |     | $w_{0,n-1}$    | $w_{0,n}$    |     | $w_{0,N}$    |
| 1      | $q_1$      | $h_1$      | $w_{1,0}$    | $w_{1,1}$    | $w_{1,2}$    |     | $w_{1,n-1}$    | $w_{1,n}$    |     | $w_{1,N}$    |
| 2      | $q_2$      | $h_2$      | $w_{2,0}$    | $w_{2,1}$    | $w_{2,2}$    |     | $w_{2,n-1}$    | $w_{2,n}$    |     | $w_{2,N}$    |
| ...    |            |            |              |              |              |     |                |              |     |              |
| $f(N)$ | $q_{f(N)}$ | $h_{f(N)}$ | $w_{f(N),0}$ | $w_{f(N),1}$ | $w_{f(N),2}$ |     | $w_{f(N),n-1}$ | $w_{f(N),n}$ |     | $w_{f(N),N}$ |

Instead, we can think about this similarly to a word ladder.

Namely, given configurations  $c, c'$  and a bound on the number of steps  $b$ , we construct the following QBF  $\phi_{c,c',b}$ , which is true if and only if  $c'$  is reachable from  $c$  within at most  $b$  steps.

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

| Step   | State      | Head       | $T_0$        | $T_1$        | $T_2$        | ... | $T_{n-1}$      | $T_n$        | ... | $T_N$        |
|--------|------------|------------|--------------|--------------|--------------|-----|----------------|--------------|-----|--------------|
| 0      | $q_0$      | 0          | $w_0$        | $w_1$        | $w_2$        |     | $w_{n-1}$      | $\square$    |     | $\square$    |
| 0      | $q_0$      | $h_0$      | $w_{0,0}$    | $w_{0,1}$    | $w_{0,2}$    |     | $w_{0,n-1}$    | $w_{0,n}$    |     | $w_{0,N}$    |
| 1      | $q_1$      | $h_1$      | $w_{1,0}$    | $w_{1,1}$    | $w_{1,2}$    |     | $w_{1,n-1}$    | $w_{1,n}$    |     | $w_{1,N}$    |
| 2      | $q_2$      | $h_2$      | $w_{2,0}$    | $w_{2,1}$    | $w_{2,2}$    |     | $w_{2,n-1}$    | $w_{2,n}$    |     | $w_{2,N}$    |
| ...    |            |            |              |              |              |     |                |              | ... |              |
| $f(N)$ | $q_{f(N)}$ | $h_{f(N)}$ | $w_{f(N),0}$ | $w_{f(N),1}$ | $w_{f(N),2}$ |     | $w_{f(N),n-1}$ | $w_{f(N),n}$ |     | $w_{f(N),N}$ |

Instead, we can think about this similarly to a word ladder.

Namely, given configurations  $c, c'$  and a bound on the number of steps  $b$ , we construct the following QBF  $\phi_{c,c',b}$ , which is true if and only if  $c'$  is reachable from  $c$  within at most  $b$  steps.

$$\phi_{c,c',b} = \exists c_{\text{mid}}$$

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

| Step   | State      | Head       | $T_0$        | $T_1$        | $T_2$        | ... | $T_{n-1}$      | $T_n$        | ... | $T_N$        |
|--------|------------|------------|--------------|--------------|--------------|-----|----------------|--------------|-----|--------------|
| 0      | $q_0$      | 0          | $w_0$        | $w_1$        | $w_2$        |     | $w_{n-1}$      | $\square$    |     | $\square$    |
| 0      | $q_0$      | $h_0$      | $w_{0,0}$    | $w_{0,1}$    | $w_{0,2}$    |     | $w_{0,n-1}$    | $w_{0,n}$    |     | $w_{0,N}$    |
| 1      | $q_1$      | $h_1$      | $w_{1,0}$    | $w_{1,1}$    | $w_{1,2}$    |     | $w_{1,n-1}$    | $w_{1,n}$    |     | $w_{1,N}$    |
| 2      | $q_2$      | $h_2$      | $w_{2,0}$    | $w_{2,1}$    | $w_{2,2}$    |     | $w_{2,n-1}$    | $w_{2,n}$    |     | $w_{2,N}$    |
| ...    |            |            |              |              |              |     |                |              | ... |              |
| $f(N)$ | $q_{f(N)}$ | $h_{f(N)}$ | $w_{f(N),0}$ | $w_{f(N),1}$ | $w_{f(N),2}$ |     | $w_{f(N),n-1}$ | $w_{f(N),n}$ |     | $w_{f(N),N}$ |

Instead, we can think about this similarly to a word ladder.

Namely, given configurations  $c, c'$  and a bound on the number of steps  $b$ , we construct the following QBF  $\phi_{c,c',b}$ , which is true if and only if  $c'$  is reachable from  $c$  within at most  $b$  steps.

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \phi_{c,c_{\text{mid}},b/2} \wedge \phi_{c_{\text{mid}},c',b/2} \right]$$

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

| Step   | State      | Head       | $T_0$        | $T_1$        | $T_2$        | ... | $T_{n-1}$      | $T_n$        | ... | $T_N$        |
|--------|------------|------------|--------------|--------------|--------------|-----|----------------|--------------|-----|--------------|
| 0      | $q_0$      | 0          | $w_0$        | $w_1$        | $w_2$        |     | $w_{n-1}$      | $\square$    |     | $\square$    |
| 0      | $q_0$      | $h_0$      | $w_{0,0}$    | $w_{0,1}$    | $w_{0,2}$    |     | $w_{0,n-1}$    | $w_{0,n}$    |     | $w_{0,N}$    |
| 1      | $q_1$      | $h_1$      | $w_{1,0}$    | $w_{1,1}$    | $w_{1,2}$    |     | $w_{1,n-1}$    | $w_{1,n}$    |     | $w_{1,N}$    |
| 2      | $q_2$      | $h_2$      | $w_{2,0}$    | $w_{2,1}$    | $w_{2,2}$    |     | $w_{2,n-1}$    | $w_{2,n}$    |     | $w_{2,N}$    |
| ...    |            |            |              |              |              |     |                |              | ... |              |
| $f(N)$ | $q_{f(N)}$ | $h_{f(N)}$ | $w_{f(N),0}$ | $w_{f(N),1}$ | $w_{f(N),2}$ |     | $w_{f(N),n-1}$ | $w_{f(N),n}$ |     | $w_{f(N),N}$ |

Instead, we can think about this similarly to a word ladder.

Namely, given configurations  $c, c'$  and a bound on the number of steps  $b$ , we construct the following QBF  $\phi_{c,c',b}$ , which is true if and only if  $c'$  is reachable from  $c$  within at most  $b$  steps.

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \phi_{c,c_{\text{mid}},b/2} \wedge \phi_{c_{\text{mid}},c',b/2} \right] \quad (\text{Assuming } b > 1)$$

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

| Step   | State      | Head       | $T_0$        | $T_1$        | $T_2$        | ... | $T_{n-1}$      | $T_n$        | ... | $T_N$        |
|--------|------------|------------|--------------|--------------|--------------|-----|----------------|--------------|-----|--------------|
| 0      | $q_0$      | 0          | $w_0$        | $w_1$        | $w_2$        |     | $w_{n-1}$      | $\square$    |     | $\square$    |
| 0      | $q_0$      | $h_0$      | $w_{0,0}$    | $w_{0,1}$    | $w_{0,2}$    |     | $w_{0,n-1}$    | $w_{0,n}$    |     | $w_{0,N}$    |
| 1      | $q_1$      | $h_1$      | $w_{1,0}$    | $w_{1,1}$    | $w_{1,2}$    |     | $w_{1,n-1}$    | $w_{1,n}$    |     | $w_{1,N}$    |
| 2      | $q_2$      | $h_2$      | $w_{2,0}$    | $w_{2,1}$    | $w_{2,2}$    |     | $w_{2,n-1}$    | $w_{2,n}$    |     | $w_{2,N}$    |
| ...    |            |            |              |              |              |     |                |              |     |              |
| $f(N)$ | $q_{f(N)}$ | $h_{f(N)}$ | $w_{f(N),0}$ | $w_{f(N),1}$ | $w_{f(N),2}$ |     | $w_{f(N),n-1}$ | $w_{f(N),n}$ |     | $w_{f(N),N}$ |

Instead, we can think about this similarly to a word ladder.

Namely, given configurations  $c, c'$  and a bound on the number of steps  $b$ , we construct the following QBF  $\phi_{c,c',b}$ , which is true if and only if  $c'$  is reachable from  $c$  within at most  $b$  steps.

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \phi_{c,c_{\text{mid}},b/2} \wedge \phi_{c_{\text{mid}},c',b/2} \right] \quad (\text{Assuming } b > 1)$$

Each configuration is encoded with the same  $x_{t,i,\gamma}, y_{t,s}, z_{t,h}$  variables as in the CL theorem.



# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

| Step   | State      | Head       | $T_0$        | $T_1$        | $T_2$        | ... | $T_{n-1}$      | $T_n$        | ... | $T_N$        |
|--------|------------|------------|--------------|--------------|--------------|-----|----------------|--------------|-----|--------------|
| 0      | $q_0$      | 0          | $w_0$        | $w_1$        | $w_2$        |     | $w_{n-1}$      | $\square$    |     | $\square$    |
| 0      | $q_0$      | $h_0$      | $w_{0,0}$    | $w_{0,1}$    | $w_{0,2}$    |     | $w_{0,n-1}$    | $w_{0,n}$    |     | $w_{0,N}$    |
| 1      | $q_1$      | $h_1$      | $w_{1,0}$    | $w_{1,1}$    | $w_{1,2}$    |     | $w_{1,n-1}$    | $w_{1,n}$    |     | $w_{1,N}$    |
| 2      | $q_2$      | $h_2$      | $w_{2,0}$    | $w_{2,1}$    | $w_{2,2}$    |     | $w_{2,n-1}$    | $w_{2,n}$    |     | $w_{2,N}$    |
| ...    |            |            |              |              |              |     |                |              |     |              |
| $f(N)$ | $q_{f(N)}$ | $h_{f(N)}$ | $w_{f(N),0}$ | $w_{f(N),1}$ | $w_{f(N),2}$ |     | $w_{f(N),n-1}$ | $w_{f(N),n}$ |     | $w_{f(N),N}$ |

Instead, we can think about this similarly to a word ladder.

Namely, given configurations  $c, c'$  and a bound on the number of steps  $b$ , we construct the following QBF  $\phi_{c,c',b}$ , which is true if and only if  $c'$  is reachable from  $c$  within at most  $b$  steps.

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \phi_{c,c_{\text{mid}},b/2} \wedge \phi_{c_{\text{mid}},c',b/2} \right] \quad (\text{Assuming } b > 1)$$

Each configuration is encoded with the same  $x_{t,i,\gamma}, y_{t,s}, z_{t,h}$  variables as in the CL theorem.

For  $b = 1$  we explicitly write the formula of the  $t$ 'th step yielding the  $t + 1$ 'th

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \phi_{c,c_{\text{mid}},b/2} \wedge \phi_{c_{\text{mid}},c',b/2} \right] \quad (\text{Assuming } b > 1)$$

The reduction outputs  $\phi_{c_0,c_Y,f(N)}$ .

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?

$$\phi_{c_0, c_Y, f(N)}$$


# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?




# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?




# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?




# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?





## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?

We created an  $\Omega(f(N)) = 2^{\Omega(n)}$  sized formula!

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?

We created an  $\Omega(f(N)) = 2^{\Omega(n)}$  sized formula!

Why is this not surprising?

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \phi_{c,c_{\text{mid}},b/2} \wedge \phi_{c_{\text{mid}},c',b/2} \right]$$

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?

We created an  $\Omega(f(N)) = 2^{\Omega(n)}$  sized formula!

Why is this not surprising?

$$\phi_{c,c',b} = \exists c_{\text{mid}} [ \phi_{c,c_{\text{mid}},b/2} \wedge \phi_{c_{\text{mid}},c',b/2} ]$$

We were using only  $\exists$  quantifiers (essentially, created a CNF!)

# PSPACE-Completeness of TQBF

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

How big is the formula?

We created an  $\Omega(f(N)) = 2^{\Omega(n)}$  sized formula.

Why is this not surprising?

$$\phi_{c,c',b} = \exists c_{\text{mid}} [ \phi_{c,c_{\text{mid}},b/2} \wedge \phi_{c_{\text{mid}},c',b/2} ]$$

We were using only  $\exists$  quantifiers (essentially, created a CNF!)

# PSPACE-Completeness of TQBF (2nd attempt)

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

We can use the “abbreviation trick”:

# PSPACE-Completeness of TQBF (2nd attempt)

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

We can use the “abbreviation trick”:

$$\phi_{c,c',b} = \exists c_{\text{mid}}$$

# PSPACE-Completeness of TQBF (2nd attempt)

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

We can use the “abbreviation trick”:

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \forall (c_i, c_j) \in \{(c, c_{\text{mid}}), (c_{\text{mid}}, c')\} : \right.$$

# PSPACE-Completeness of TQBF (2nd attempt)

## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

We can use the “abbreviation trick”:

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \forall (c_i, c_j) \in \{(c, c_{\text{mid}}), (c_{\text{mid}}, c')\} : \phi_{c_i, c_j, b/2} \right].$$



# PSPACE-Completeness of TQBF (2nd attempt)


## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

We can use the “abbreviation trick”:

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \forall (c_i, c_j) \in \{(c, c_{\text{mid}}), (c_{\text{mid}}, c')\} : \phi_{c_i, c_j, b/2} \right].$$



# PSPACE-Completeness of TQBF (2nd attempt)


## Theorem

*TQBF is PSPACE-Complete.*

## Proof Sketch. (Cont.)

We can use the “abbreviation trick”:

$$\phi_{c,c',b} = \exists c_{\text{mid}} \left[ \forall (c_i, c_j) \in \{(c, c_{\text{mid}}), (c_{\text{mid}}, c')\} : \phi_{c_i, c_j, b/2} \right].$$



We have  $\log f(N) = n^{O(1)}$  levels, each has a polynomial size QBF.

# Lectures 12: Recap

Today we saw:

# Lectures 12: Recap

Today we saw:

- Savitch's theorem, that shows that  $NSPACE(S(N)) \subseteq SPACE(S^2(N))$ .

# Lectures 12: Recap

Today we saw:

- Savitch's theorem, that shows that  $NSPACE(S(N)) \subseteq SPACE(S^2(N))$ .
- This implies that  $NPSPACE \subseteq PSPACE$ .

# Lectures 12: Recap

Today we saw:

- Savitch's theorem, that shows that  $NSPACE(S(N)) \subseteq SPACE(S^2(N))$ .
- This implies that  $NPSPACE \subseteq PSPACE$ .
- That TQBF is PSPACE-Complete.

# Lectures 12: Recap

Today we saw:

- Savitch's theorem, that shows that  $NSPACE(S(N)) \subseteq SPACE(S^2(N))$ .
- This implies that  $NPSPACE \subseteq PSPACE$ .
- That TQBF is PSPACE-Complete.
- To ponder: would the proof work for an NDTM machine  $M$ ?

# Lectures 12: Recap

Today we saw:

- Savitch's theorem, that shows that  $NSPACE(S(N)) \subseteq SPACE(S^2(N))$ .
- This implies that  $NPSPACE \subseteq PSPACE$ .
- That TQBF is PSPACE-Complete.
- To ponder: would the proof work for an NDTM machine  $M$ ?

Next lecture: the Log-space class.