

Physics 106a — Classical Mechanics

Michael Cross

California Institute of Technology

Fall Term, 2013

Lecture 6 Equilibria and Small Oscillations

Outline

- Lagrangian Approach
 - Equilibria
 - Dynamics near an equilibrium
 - Simple harmonic motion
 - Undriven
 - Driven
- Add Dissipation


Solution strategy

Look for simple solutions

- Time independent solution in some coordinate system (equilibrium, fixed point)
- Behavior near equilibrium (linearize equations about equilibrium point)
 - Unstable equilibrium: deviation from equilibrium grows exponentially
 - Stable equilibrium: deviation from equilibrium remains small (may decay)
 - Study small oscillations near stable equilibrium (simple harmonic motion)
- Study change in behavior of equilibria as parameters change (bifurcation theory)

Resonance

- Tacoma Narrows bridge [Movie, Article: Am. J. Phys. 59, 118 (1991)]
- Rings of Saturn



Damped simple harmonic oscillator

Mechanical systems usually have some dissipation or damping. A common form (*not* valid for solid on solid friction) is a force proportional to the negative of the velocity. This leads to the equation of motion

$$\ddot{q} + \frac{1}{Q}\dot{q} + q = 0$$

with Q^{-1} proportional to the dissipation coefficient.

Q is the *quality factor* of the oscillator

- large Q — high quality, small dissipation
- small Q — low quality, high dissipation

Damped simple harmonic oscillator

$$\ddot{q} + \frac{1}{Q}\dot{q} + q = 0$$

The general solution is

$$q(t) = A_+ e^{\lambda_+ t} + A_- e^{\lambda_- t}$$

with λ_{\pm} the roots of the polynomial $\lambda^2 + Q^{-1}\lambda + 1 = 0$

$$\lambda_{\pm} = -\frac{1}{2Q} \pm \sqrt{\frac{1}{4Q^2} - 1}.$$

The behavior depends on the sign inside the $\sqrt{\quad}$

$$Q > \frac{1}{2} \quad \text{Underdamped} \quad q(t) = A e^{-\frac{t}{2Q}} \cos(\omega' t + \phi) \text{ with } \omega' = \sqrt{1 - \frac{1}{4Q^2}}$$

$$Q < \frac{1}{2} \quad \text{Overdamped} \quad q(t) = A_+ e^{\lambda_+ t} + A_- e^{\lambda_- t} \text{ with } \lambda_+, \lambda_- < 0$$

$$Q = \frac{1}{2} \quad \text{Critically damped} \quad q(t) = (A + Bt)e^{-t} \text{ (special case)}$$

Damped simple harmonic oscillator

Energy $E = \frac{1}{2}(q^2 + \dot{q}^2)$ decays as

$$\dot{E} = \dot{q}(\ddot{q} + q) = -\frac{1}{Q}\dot{q}^2 < 0$$

For large Q oscillatory motion, the energy decays little in one cycle, and averaging over one cycle gives

$$\langle V \rangle = \langle \frac{1}{2}q^2 \rangle \simeq \langle T \rangle = \langle \frac{1}{2}\dot{q}^2 \rangle \simeq \frac{1}{2}\langle E \rangle$$

and so the average energy decays as

$$\langle \dot{E} \rangle \simeq -\frac{\langle E \rangle}{Q}$$

- exponential decay with a $1/e$ time of Q (in units of ω_0^{-1})
- $\simeq Q/2\pi$ oscillations in this decay time
- fractional energy loss per period (cycle) is $\simeq 2\pi/Q$.

Damped simple harmonic oscillator

Sometimes, this type of dissipation force proportional to velocity is shoe-horned into a Lagrangian type approach by introducing a dissipation function $\mathcal{D}(\{\dot{q}_k\})$ such that the equations of motion are

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_k} \right) - \frac{\partial L}{\partial q_k} + \frac{\partial \mathcal{D}}{\partial \dot{q}_k} = 0$$

For the damped simple harmonic oscillator the dissipation function (using the scaled time) would be

$$\mathcal{D} = \frac{1}{2} Q^{-1} \dot{q}^2$$

The rate of energy dissipation is $2\mathcal{D}$

Driven, damped simple harmonic oscillator

Oscillatory forcing

$$\ddot{q} + \frac{1}{Q}\dot{q} + q = F(t) \quad \text{with} \quad F(t) = \cos \omega t = \operatorname{Re}[e^{i\omega t}]$$

- Calculate the response $q_c(t)$ to the forcing $F_c(t) = e^{i\omega t}$, and take the real part at the end.
- For $F_c(t)$ the steady state solution is

$$q_c(t) = A_c e^{i\omega t} = A e^{i(\omega t + \phi)}$$

with

$$A_c = \frac{1}{(-\omega^2 + 1) + \frac{i\omega}{Q}}$$

- This gives the solution for $F(t) = \cos \omega t$ as $q(t) = A \cos(\omega t + \phi)$.

Driven, damped simple harmonic oscillator

The amplitude is given by

$$A^2 = \frac{1}{(\omega^2 - 1)^2 + \omega^2/Q^2} \simeq \frac{1}{4} \frac{1}{(\omega - 1)^2 + 1/(4Q^2)}$$

For large Q the response is a sharp *resonance peak* centered at $\omega_r \simeq 1$, with full width at half intensity (half height in A^2) $1/Q$ and amplitude at resonance Q larger than the static response


The phase is

$$\phi = -\tan^{-1} \left(\frac{\omega/Q}{1 - \omega^2} \right)$$

and varies from slightly negative for low frequencies, passing rapidly through $-\pi/2$ on resonance, and decreasing to $-\pi$ for large frequencies.


Single molecule mass sensor

Roukes group, Caltech, Nature Nanotechnology 4, 445, (2009)



Single molecule mass sensor

Roukes group, Caltech, Nature Nanotechnology 4, 445, (2009)



Single molecule mass sensor

Roukes group, Caltech, Nature Nanotechnology 4, 445, (2009)

