

Hence

$$M_2 < AF(T+1)^{1-A(\sigma-\frac{1}{2}-\delta)} \leq AF(T+1)(\log^{\frac{1}{2}}T)^{-A(\sigma-\frac{1}{2}-\delta)}.$$

This gives the required result if $\sigma \leq \frac{1}{2}$, and for $\frac{1}{2} \leq \sigma \leq 2$ it is trivial, if the A is small enough.

LEMMA γ . For $\sigma > \frac{1}{2}$, $0 < \xi < \frac{1}{2}t$,

$$\log \zeta(s) = i \int_{t-\xi}^{t+\xi} \frac{S(y)}{s-\frac{1}{2}-iy} dy + O\left(\frac{\phi(2t)}{\xi}\right) + O(1). \quad (14.13.3)$$

We have

$$\begin{aligned} \int_{t+\xi}^{2t} \frac{S(y)}{s-\frac{1}{2}-iy} dy &= \left[\frac{S_1(y)}{s-\frac{1}{2}-iy} \right]_{t+\xi}^{2t} - i \int_{t+\xi}^{2t} \frac{S_1(y)}{(s-\frac{1}{2}-iy)^2} dy \\ &= O\left(\frac{\phi(2t)}{\xi}\right) + O\left(\phi(2t) \int_{t+\xi}^{2t} \frac{dy}{(\sigma-\frac{1}{2})^2 + (y-t)^2}\right) = O\left(\frac{\phi(2t)}{\xi}\right), \end{aligned}$$

and similarly for the integral over $(\frac{1}{2}t, t-\xi)$. The result therefore follows from (14.12.4).

Proof of Theorem 14.13. By Lemmas α and γ ,

$$\begin{aligned} \log \zeta(s) &= O\{\phi(4t)\log t\}^{\frac{1}{2}} \int_{t-\xi}^{t+\xi} \frac{dy}{\{(\sigma-\frac{1}{2})^2 + (y-t)^2\}^{\frac{1}{2}}} + O\left(\frac{\phi(2t)}{\xi}\right) + O(1) \\ &= O\left[\{\phi(4t)\log t\}^{\frac{1}{2}} \frac{\xi}{\sigma-\frac{1}{2}}\right] + O\left(\frac{\phi(4t)}{\xi}\right) + O(1) \end{aligned}$$

for $\sigma - \frac{1}{2} \geq 1/\log \log T$, $4 \leq t \leq T$. Taking

$$\xi = A \left\{ \frac{\phi(4t)}{\log t} \right\}^{\frac{1}{2}} \frac{1}{(\log \log T)^{\frac{1}{2}}},$$

we obtain $\log \zeta(s) = O[(\log T)^{\frac{1}{2}}(\log \log T)^{\frac{1}{2}}\{\phi(4T)\}^{\frac{1}{2}}]$.

Hence by Lemma β , for $\sigma - \frac{1}{2} \geq 1/\log \log T$,

$$\log \zeta(s) = O[(\log T)^{\frac{1}{2}}(\log \log T)^{\frac{1}{2}}\{\phi(4T+4)\}^{\frac{1}{2}}e^{-A(\sigma-\frac{1}{2})\log \log T}].$$

Hence

$$\int_{\frac{1}{2}+1/\log \log T}^2 \log |\zeta(s)| d\sigma = O[(\log T)^{\frac{1}{2}}(\log \log T)^{-\frac{1}{2}}\{\phi(4T+4)\}^{\frac{1}{2}}]. \quad (14.13.4)$$

Again, the real part of (14.13.3) may be written

$$\log |\zeta(s)| = \int_0^{\xi} \frac{x}{(\sigma-\frac{1}{2})^2+x^2} \{S(t-x)-S(t+x)\} dx + O\left(\frac{\phi(2t)}{\xi}\right) + O(1). \quad (14.13.5)$$

Hence

$$\begin{aligned} \int_{\frac{1}{2}}^{\frac{1}{2}+\mu} \log |\zeta(s)| d\sigma &= \int_0^{\xi} \arctan \frac{\mu}{x} \{S(t-x)-S(t+x)\} dx + \\ &\quad + O\{\mu \phi(2t)/\xi\} + O(\mu) \\ &= O[\xi\{\phi(4t)\log t\}^{\frac{1}{2}}] + O\{\mu \phi(2t)/\xi\} + O(\mu). \end{aligned}$$

Taking $\mu = 1/\log \log T$, and ξ as before,

$$\int_{\frac{1}{2}}^{\frac{1}{2}+1/\log \log T} \log |\zeta(s)| d\sigma = O[(\log T)^{\frac{1}{2}}(\log \log T)^{-\frac{1}{2}}\{\phi(4T)\}^{\frac{1}{2}}]. \quad (14.13.6)$$

Now (14.13.4), (14.13.6), and Theorem 9.9 give

$$S_1(t) = O[(\log T)^{\frac{1}{2}}(\log \log T)^{-\frac{1}{2}}\{\phi(5T)\}^{\frac{1}{2}}] \quad (4 \leq t \leq T). \quad (14.13.7)$$

Varying t and taking the maximum,

$$\phi(T) = O[(\log T)^{\frac{1}{2}}(\log \log T)^{-\frac{1}{2}}\{\phi(5T)\}^{\frac{1}{2}}].$$

Let

$$\psi(T) = \max_{4 \leq t \leq T} \frac{(\log \log t)^2 \phi(t)}{\log t},$$

so that $\psi(T)$ is non-decreasing and

$$\phi(T) \leq \frac{\log T}{(\log \log T)^2} \psi(T).$$

Then (14.13.7) gives

$$\phi(T) = O\left[\frac{\log T}{(\log \log T)^2} \{\psi(5T)\}^{\frac{1}{2}}\right]$$

$$\text{or } \frac{\phi(T)(\log \log T)^2}{\log T} = O[\{\psi(5T)\}^{\frac{1}{2}}] = O[\{\psi(5T_1)\}^{\frac{1}{2}}] \quad (T \leq T_1).$$

Varying T and taking the maximum,

$$\psi(T_1) = O[\{\psi(5T_1)\}^{\frac{1}{2}}].$$

But $\psi(5T_1) < 5\psi(T_1)$ for some arbitrarily large T_1 ; for otherwise

$$\psi(5^n t_0) \geq 5^n \psi(t_0),$$

i.e. $\psi(T) > AT$ for some arbitrarily large T , which is not so, since in fact $\phi(T) = O(\log T)$, $\psi(T) = O((\log \log T)^2)$. Hence

$$\psi(T_1) < A\{\psi(T_1)\}^{\frac{1}{2}}, \quad \psi(T_1) < A,$$

for some arbitrarily large T_1 , and so for all T_1 , since ϕ is non-decreasing.

Hence
$$\phi(T) = O\left\{\frac{\log T}{(\log \log T)^2}\right\}.$$

This proves (14.13.2), and (14.13.1) then follows from Lemma α .

The argument can be extended to show that, if $S_n(t)$ is the n th integral of $S(t)$, then

$$S_n(t) = O\left\{\frac{\log t}{(\log \log t)^{n+1}}\right\}. \quad (14.13.8)$$

14.14. Theorem 14.13 also enables us to prove inequalities for $\zeta(s)$ in the immediate neighbourhood of $\sigma = \frac{1}{2}$, a region not touched by previous arguments. We obtain first

THEOREM 14.14 (A).

$$\zeta\left(\frac{1}{2} + it\right) = O\left\{\exp\left(A \frac{\log t}{\log \log t}\right)\right\}. \quad (14.14.1)$$

We have

$$S(t+x) - S(t) = \{N(t+x) - N(t)\} - \{L(t+x) - L(t)\} - \{f(t+x) - f(t)\},$$

where $f(t)$ is the $O(1/t)$ of (9.3.2), and arises from the asymptotic formula for $\log \Gamma(s)$. Thus $f'(t) = O(1/t^2)$, and since $N(t+x) \geq N(t)$

$$S(t+x) - S(t) > -Ax \log t + O(x/t^2) > -Ax \log t.$$

Hence, by (14.13.5),

$$\begin{aligned} \log |\zeta(s)| &< A \int_0^\xi \frac{x^2 \log t}{(\sigma - \frac{1}{2})^2 + x^2} dx + O\left\{\frac{\log t}{\xi(\log \log t)^2}\right\} + O(1) \\ &< A\xi \log t + O\left\{\frac{\log t}{\xi(\log \log t)^2}\right\} + O(1) \end{aligned}$$

uniformly for $\sigma > \frac{1}{2}$, and so by continuity for $\sigma = \frac{1}{2}$. Taking

$$\xi = 1/\log \log t$$

the result follows.

THEOREM 14.14 (B). We have

$$-\frac{A \log t}{\log \log t} \log \left\{ \frac{2}{(\sigma - \frac{1}{2}) \log \log t} \right\} < \log |\zeta(s)| < \frac{A \log t}{\log \log t} \quad \left(\frac{1}{2} < \sigma \leq \frac{1}{2} + A/\log \log t\right), \quad (14.14.2)$$

$$\arg \zeta(s) = O\left\{\frac{\log t}{\log \log t}\right\} \quad \left(\frac{1}{2} \leq \sigma \leq \frac{1}{2} + A/\log \log t\right). \quad (14.14.3)$$

By (14.13.1) and (14.13.3),

$$\log \zeta(s) = O\left\{\frac{\log t}{\log \log t} \int_0^\xi \frac{dx}{\sqrt{((\sigma - \frac{1}{2})^2 + x^2)}}\right\} + O\left\{\frac{\log t}{\xi(\log \log t)^2}\right\} + O(1).$$

Now
$$\int_0^\xi \frac{du}{\sqrt{((\sigma - \frac{1}{2})^2 + u^2)}} = \int_0^{\xi(\sigma - \frac{1}{2})} \frac{dx'}{\sqrt{(1+x'^2)}},$$

which is less than 1 if $\xi \leq \sigma - \frac{1}{2}$, and otherwise is less than

$$1 + \int_1^{\xi(\sigma - \frac{1}{2})} \frac{dx'}{x'} = 1 + \log \frac{\xi}{\sigma - \frac{1}{2}}.$$

Taking $\xi = 1/\log \log t$, the lower bound in (14.14.2) follows. The upper bound follows from the argument of the previous section. Lastly, taking imaginary parts in (14.13.3),

$$\begin{aligned} \arg \zeta(s) &= \int_0^\xi \frac{\sigma - \frac{1}{2}}{x^2 + (\sigma - \frac{1}{2})^2} \{S(t+x) - S(t-x)\} dx + \\ &\quad + O\left\{\frac{\log t}{\xi(\log \log t)^2}\right\} + O(1) \\ &= O\left\{\frac{\log t}{\log \log t} \int_0^\xi \frac{\sigma - \frac{1}{2}}{x^2 + (\sigma - \frac{1}{2})^2} dx\right\} + O\left\{\frac{\log t}{\xi(\log \log t)^2}\right\} + O(1). \end{aligned}$$

Now
$$\int_0^\xi \frac{\sigma - \frac{1}{2}}{x^2 + (\sigma - \frac{1}{2})^2} dx < \int_0^\infty \frac{\sigma - \frac{1}{2}}{x^2 + (\sigma - \frac{1}{2})^2} dx = \frac{1}{2}\pi.$$

Hence, taking $\xi = 1$, (14.14.3) follows uniformly for $\sigma > \frac{1}{2}$, and so by continuity for $\sigma = \frac{1}{2}$.

In particular

$$\log \zeta(s) = O\left\{\frac{\log t}{\log \log t}\right\} \quad \left(\sigma = \frac{1}{2} + \frac{A}{\log \log t}\right). \quad (14.14.4)$$

From (14.14.4), (14.5.2), and a Phragmén-Lindelöf argument it follows that

$$\log \zeta(s) = O\left\{\frac{(\log t)^{2-2\sigma}}{\log \log t}\right\} \quad (14.14.5)$$

uniformly for
$$\frac{1}{2} + \frac{A}{\log \log t} \leq \sigma \leq 1 - \delta.$$

14.15. Another result in the same order of ideas is an approximate formula for $\log \zeta(s)$, which should be compared with Theorem 9.6 (B).

THEOREM 14.15. For $\frac{1}{2} \leq \sigma \leq 2$,

$$\log \zeta(s) = \sum_{|\mu| < \log \log t} \log(s - \rho) + O\left\{\frac{(\log t \log \log \log t)}{\log \log t}\right\}, \quad (14.15.1)$$

where $\rho = \frac{1}{2} + i\gamma$ runs through zeros of $\zeta(s)$.

In Lemma α of § 3.9, let

$$f(s) = \zeta(s), \quad s_0 = \frac{1}{2} + \frac{1}{\sqrt{3}}\delta + iT, \quad r = \frac{4}{\sqrt{3}}\delta,$$

where $\delta = 1/\log \log T$. By (14.14.4)

$$\left| \frac{1}{\zeta(s_0)} \right| \leq \exp \left(\frac{A \log T}{\log \log T} \right).$$

The upper bound in (14.14.2) gives

$$|\zeta(s)| < \exp \left(\frac{A \log T}{\log \log T} \right)$$

for $|s - s_0| \leq r$, $\sigma \geq \frac{1}{2}$; and for $|s - s_0| \leq r$, $\sigma < \frac{1}{2}$, the functional equation gives

$$|\zeta(s)| < A t^{\frac{1}{2}-\sigma} |\zeta(1-s)| < A t^{\frac{1}{2}} \exp \left(\frac{A \log T}{\log \log T} \right) < \exp \left(\frac{A \log T}{\log \log T} \right).$$

It therefore follows from (3.9.1) that

$$\begin{aligned} \log \zeta(s) - \log \zeta(s_0) - \sum_{|s_0 - \rho| \leq 2\delta/\sqrt{3}} \log(s - \rho) + \\ + \sum_{|s_0 - \rho| \leq 2\delta/\sqrt{3}} \log(s_0 - \rho) = O \left(\frac{\log T}{\log \log T} \right) \end{aligned}$$

for $|s - s_0| \leq \frac{1}{2}r$, and so in particular for $\frac{1}{2} \leq \sigma \leq \frac{1}{2} + \delta$, $t = T$.

$$\text{Now} \quad \log \zeta(s_0) = O \left(\frac{\log T}{\log \log T} \right).$$

$$\text{Also} \quad s_0 - \rho = \frac{1}{\sqrt{3}}\delta + iT - \rho.$$

$$\text{Hence} \quad \frac{1}{\sqrt{3}}\delta \leq |s_0 - \rho| < A,$$

and so, if the logarithm has its principal value,

$$\log(s_0 - \rho) = O \left(\log \frac{1}{\delta} \right) = O(\log \log \log T).$$

Also the number of values of ρ in the above sums does not exceed

$$N \left(T + \frac{2\delta}{\sqrt{3}} \right) - N \left(T - \frac{2\delta}{\sqrt{3}} \right) = O(\delta \log T) + O \left(\frac{\log T}{\log \log T} \right) = O \left(\frac{\log T}{\log \log T} \right),$$

by Theorem 14.13. Hence

$$\sum_{|s_0 - \rho| \leq 2\delta/\sqrt{3}} \log(s_0 - \rho) = O \left(\frac{\log T \log \log \log T}{\log \log T} \right).$$

Since $|T - \gamma| \leq \delta$ if $|s_0 - \rho| \leq 2\delta/\sqrt{3}$, the result follows, with T for t and $\frac{1}{2} \leq \sigma \leq \frac{1}{2} + \delta$. It is also true for $\frac{1}{2} + \delta < \sigma \leq 2$, since in this region

$$\log \zeta(s) = O \left(\frac{\log T}{\log \log T} \right),$$

and, as in the case of the other sum,

$$\sum_{|s_0 - \rho| \leq 2\delta/\sqrt{3}} \log(s - \rho) = O \left(\frac{\log T \log \log \log T}{\log \log T} \right).$$

This proves the theorem.

For $\zeta'(s)/\zeta(s)$ we obtain similarly from Lemma α of § 3.9

$$\frac{\zeta'(s)}{\zeta(s)} = \sum_{|t - \gamma| \leq 1/\log \log t} \frac{1}{s - \rho} + O(\log t). \quad (14.15.2)$$

14.16. THEOREM 14.16. *Each interval $[T, T+1]$ contains a value of t such that*

$$|\zeta(s)| > \exp \left(-A \frac{\log t}{\log \log t} \right) \quad \left(\frac{1}{2} \leq \sigma \leq 2 \right). \quad (14.16.1)$$

Let $\delta = 1/\log \log T$. Then the lower bound (14.16.1) holds automatically for $\sigma \geq \frac{1}{2} + \delta$, by (14.14.4). We therefore assume that $\frac{1}{2} \leq \sigma \leq \frac{1}{2} + \delta$. If $s = \sigma + it$ and $s_0 = \frac{1}{2} + \delta + iT$ then, on integrating (14.15.2), we find

$$\log \frac{\zeta(s)}{\zeta(s_0)} = \sum_{|t - \gamma| \leq \delta} \log \left(\frac{s - \rho}{s_0 - \rho} \right) + O \left(\frac{\log T}{\log \log T} \right).$$

Moreover $\log \zeta(s_0) = O \left(\frac{\log T}{\log \log T} \right)$ by (14.14.4) so that, on taking real parts

$$\begin{aligned} \log |\zeta(s)| &= \sum_{|t - \gamma| \leq \delta} \log \left| \frac{s - \rho}{s_0 - \rho} \right| + O \left(\frac{\log T}{\log \log T} \right) \\ &\geq \sum_{|t - \gamma| \leq \delta} \log \frac{|t - \gamma|}{2\delta} + O \left(\frac{\log T}{\log \log T} \right), \end{aligned}$$

since $|s - \rho| \geq |t - \gamma|$ and $|s_0 - \rho| \leq 2\delta$. We now observe that

$$\begin{aligned} \int_T^{T+1} \sum_{|t - \gamma| \leq \delta} \log \frac{|t - \gamma|}{2\delta} dt &= \sum_{T - \delta \leq \gamma \leq T + 1 + \delta} \int_{\max(\gamma - \delta, T)}^{\min(\gamma + \delta, T + 1)} \log \frac{|t - \gamma|}{2\delta} dt \\ &\geq \sum_{T - \delta \leq \gamma \leq T + 1 + \delta} \int_{T - \delta}^{\gamma + \delta} \log \frac{|t - \gamma|}{2\delta} dt \\ &= \sum_{T - \delta \leq \gamma \leq T + 1 + \delta} (-2\delta - 2\delta \log 2) \\ &\geq -A\delta \log T, \end{aligned}$$

as there are $O(\log T)$ terms in the sum. Hence there is a t for which

$$\sum_{|t-\gamma| \leq \delta} \log \frac{|t-\gamma|}{2\delta} \geq -A\delta \log T$$

and the result follows.

In particular, if ϵ is any positive number, each $(T, T+1)$ contains a t such that

$$\frac{1}{\zeta(s)} = O(\epsilon) \quad \left(\frac{1}{2} \leq \sigma \leq 2\right). \quad (14.16.2)$$

14.17. Mean-value theorems† for $S(t)$ and $S_1(t)$. We consider first $S_1(t)$. We begin by proving

THEOREM 14.17. For $\frac{1}{2}T \leq t \leq T$, $\delta = T^{-\frac{1}{2}}$,

$$\pi S_1(t) = C - \sum_{n=2}^{\infty} \frac{\Lambda_1(n) \cos(t \log n)}{n^{\frac{1}{2}} \log n} e^{-\delta n} + O\left(\frac{1}{\log \log T}\right) \quad (14.17.1)$$

where

$$C = \int_{\frac{1}{2}}^{\infty} \log |\zeta(\sigma)| d\sigma.$$

Making $\beta \rightarrow \infty$ in (9.9.4), we have

$$\pi S_1(t) = C - \int_{\frac{1}{2}}^{\infty} \log |\zeta(\sigma + it)| d\sigma. \quad (14.17.2)$$

Now, integrating (14.4.1) from s to $\frac{1}{2} + it$,

$$\begin{aligned} \log \zeta(s) - \log \zeta\left(\frac{1}{2} + it\right) &= \sum_{n=2}^{\infty} \frac{\Lambda_1(n)}{n^s} e^{-\delta n} - \sum_{n=2}^{\infty} \frac{\Lambda_1(n)}{n^{\frac{1}{2} + it}} e^{-\delta n} \\ &\quad + \sum_{\rho} \int_{\sigma}^{\frac{1}{2}} \delta^{\sigma-\rho} \Gamma(\rho - s_1) d\sigma_1 + O(\delta^{\sigma-\frac{1}{2}} \log t) \quad \left(\frac{1}{2} \leq \sigma \leq \frac{3}{2}\right). \end{aligned}$$

Also, if $\sigma \geq \frac{3}{2}$,

$$\begin{aligned} \log \zeta(\sigma + it) - \sum_{n=2}^{\infty} \frac{\Lambda_1(n)}{n^{\sigma}} e^{-\delta n} &= \sum_{n=2}^{\infty} \frac{\Lambda_1(n)}{n^{\sigma}} (1 - e^{-\delta n}) \\ &= O\left(\sum_{n=2}^{\infty} n^{-\sigma} (1 - e^{-\delta n})\right) = O\left(\sum_{2 \leq n \leq 1/\delta} n^{1-\sigma} \delta + \sum_{n > 1/\delta} n^{-\sigma}\right) \\ &= O(\delta^{\sigma-1} + 2^{-\sigma} \delta \log t). \end{aligned} \quad (14.17.3)$$

Hence, for $\frac{1}{2} \leq \sigma \leq \frac{3}{2}$,

$$\log \zeta(s) = \sum_{n=2}^{\infty} \frac{\Lambda_1(n)}{n^s} e^{-\delta n} + \sum_{\rho} \int_{\sigma}^{\frac{1}{2}} \delta^{\sigma-\rho} \Gamma(\rho - s_1) d\sigma_1 + O(\delta^{\frac{1}{2}}) + O(\delta^{\sigma-\frac{1}{2}} \log t),$$

† Littlewood (5), Titchmarsh (2).

and integrating over $\frac{1}{2} \leq \sigma \leq \frac{3}{2}$,

$$\int_{\frac{1}{2}}^{\frac{3}{2}} \log \zeta(s) d\sigma = \int_{\frac{1}{2}}^{\frac{3}{2}} \left(\sum_{n=2}^{\infty} \frac{\Lambda_1(n)}{n^s} e^{-\delta n} \right) d\sigma + \sum_{\rho} \int_{\frac{1}{2}}^{\frac{3}{2}} (\sigma_1 - \frac{1}{2}) \delta^{\sigma_1-\rho} \Gamma(\rho - s_1) d\sigma_1 + O(\delta^{\frac{1}{2}}) + O(\delta^{\frac{1}{2}} \log t).$$

Also, by (14.17.3),

$$\int_{\frac{1}{2}}^{\infty} \log \zeta(s) d\sigma = \int_{\frac{1}{2}}^{\infty} \left(\sum_{n=2}^{\infty} \frac{\Lambda_1(n)}{n^s} e^{-\delta n} \right) d\sigma + O(\delta^{\frac{1}{2}}),$$

$$\text{and} \quad \int_{\frac{1}{2}}^{\infty} \sum_{n=2}^{\infty} \frac{\Lambda_1(n)}{n^s} e^{-\delta n} d\sigma = \sum_{n=2}^{\infty} \frac{\Lambda_1(n)}{n^{\frac{1}{2} + it}} e^{-\delta n} \log n,$$

the inversion being justified by absolute convergence. Hence

$$\begin{aligned} \pi S_1(t) &= C - \sum_{n=2}^{\infty} \frac{\Lambda_1(n) \cos(t \log n)}{n^{\frac{1}{2}} \log n} e^{-\delta n} \\ &\quad + O\left(\sum_{\rho} \int_{\frac{1}{2}}^{\frac{3}{2}} (\sigma_1 - \frac{1}{2}) \delta^{\sigma_1-\frac{1}{2}} |\Gamma(\rho - s_1)| d\sigma_1\right) + O(\delta^{\frac{1}{2}}) + O(\delta^{\frac{1}{2}} \log t). \end{aligned}$$

Now

$$\Gamma(\rho - s_1) = O(e^{-A|\gamma-t|}) \quad (|\gamma-t| \geq 1),$$

$$\Gamma(\rho - s_1) = O\left(\frac{1}{|\rho - s_1|}\right) = O\left(\frac{1}{((\sigma_1 - \frac{1}{2})^2 + |\gamma - t|^2)^{\frac{1}{2}}}\right) \quad (|\gamma - t| < 1).$$

Hence

$$\begin{aligned} \sum_{\rho} &= \sum_{|\gamma-t| < 1/\log \log t} + \sum_{1/\log \log t \leq |\gamma-t| < 1} + \sum_{|\gamma-t| \geq 1} \\ &= O\left(\sum_{|\gamma-t| < 1/\log \log t} \int_{\frac{1}{2}}^{\frac{3}{2}} \delta^{\sigma_1-\frac{1}{2}} d\sigma_1\right) + \\ &\quad + O\left(\sum_{1/\log \log t \leq |\gamma-t| < 1} \frac{1}{|\gamma-t|} \int_{\frac{1}{2}}^{\frac{3}{2}} (\sigma_1 - \frac{1}{2}) \delta^{\sigma_1-\frac{1}{2}} d\sigma_1\right) + \\ &\quad + O\left(\sum_{|\gamma-t| \geq 1} e^{-A|\gamma-t|} \int_{\frac{1}{2}}^{\frac{3}{2}} (\sigma_1 - \frac{1}{2}) \delta^{\sigma_1-\frac{1}{2}} d\sigma_1\right). \end{aligned}$$

Now

$$\int_{\frac{1}{2}}^{\frac{3}{2}} \delta^{\sigma_1 - \frac{1}{2}} d\sigma_1 < \int_0^{\infty} e^{-x \log 1/\delta} dx = \log^{-1}(1/\delta),$$

$$\int_{\frac{1}{2}}^{\frac{3}{2}} (\sigma_1 - \frac{1}{2}) \delta^{\sigma_1 - \frac{1}{2}} d\sigma_1 < \int_0^{\infty} x e^{-x \log 1/\delta} dx = \log^{-2}(1/\delta).$$

As in § 14.5,

$$\sum_{|y-t| \geq 1} e^{-\Delta |y-t|} = O(\log t).$$

Also, by (14.13.1), for $t-1 \leq t' \leq t+1$

$$N\left(t' + \frac{1}{\log \log t}\right) - N(t') = O\left(\frac{\log t}{\log \log t}\right).$$

Hence

$$\sum_{|y-t| \leq 1/\log \log t} 1 = O\left(\frac{\log t}{\log \log t}\right),$$

and

$$\sum_{t+1/\log \log t < \gamma \leq t+1} \frac{1}{\gamma-t} = \sum_{m < \log \log t} \sum_{t+m/\log \log t < \gamma < (m+1)/\log \log t} \frac{1}{\gamma-t}$$

$$= \sum_{m < \log \log t} O\left(\frac{1}{m/\log \log t} \frac{\log t}{\log t}\right) = O(\log t \log \log \log t).$$

Hence

$$\sum_p = O\left(\frac{\log t}{\log 1/\delta \log \log t}\right) + O\left(\frac{\log t \log \log \log t}{\log^2 1/\delta}\right) + O\left(\frac{\log t}{\log^2 1/\delta}\right) = O\left(\frac{1}{\log \log t}\right)$$

for the given δ and t . This proves the theorem.**14.18. LEMMA 14.18.** *If $a_n = O(1)$, $\delta < \frac{1}{2}$, then*

$$\frac{2}{T} \int_{\frac{1}{2}T}^T \left| \sum_{n=1}^{\infty} \frac{a_n}{n^{\sigma}} e^{-\delta n} \right|^2 dt = \sum_{n=1}^{\infty} \frac{|a_n|^2}{n^{2\sigma}} e^{-2\delta n} + O\left(\frac{1}{T\delta} \log \frac{1}{\delta}\right)$$

uniformly for $\sigma \geq \frac{1}{2}$. Similarly, if $a_n = O(\log n)$, the formula holds with a remainder term

$$O\left(\frac{1}{T\delta} \log^3 \frac{1}{\delta}\right).$$

The left-hand side is

$$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{2}{T} \int_{\frac{1}{2}T}^T \frac{a_m \bar{a}_n}{(mn)^{\sigma}} e^{-(m+n)\delta} \left(\frac{n}{m}\right)^{\sigma} dt = \sum_{m=n} + \sum_{m \neq n}.$$

Clearly

$$\sum_{m=n} = \sum_{n=1}^{\infty} \frac{|a_n|^2}{n^{2\sigma}} e^{-2\delta n}.$$

Also

$$\sum_{m \neq n} = O\left(\frac{1}{T}\right) \sum_{m < n} \frac{e^{-n\delta}}{(mn)^{\frac{1}{2}} \log n/m}.$$

Now

$$\sum_{n=m+1}^{2m} \frac{e^{-n\delta}}{(mn)^{\frac{1}{2}} \log n/m} = O\left(\sum_{n=m+1}^{2m} \frac{e^{-m\delta}}{m \log\{1+(n-m)/m\}}\right)$$

$$= O\left(e^{-m\delta} \sum_{n=m+1}^{2m} \frac{1}{n-m}\right) = O(e^{-m\delta} \log m),$$

$$\sum_{n=2m+1}^{\infty} \frac{e^{-n\delta}}{(mn)^{\frac{1}{2}} \log n/m} = O\left(\frac{1}{m} \sum_{n=2m+1}^{\infty} e^{-n\delta}\right) = O\left(\frac{e^{-m\delta}}{m\delta}\right).$$

$$\text{Hence } \sum_n = O\left(\frac{1}{T}\right) \sum_{m=1}^{\infty} \left(\log m + \frac{1}{m\delta}\right) e^{-m\delta} = O\left(\frac{1}{T\delta} \log \frac{1}{\delta}\right).$$

This proves the first part. In the second part we have a pair of logarithms running throughout the remainder terms, and this is easily seen to produce the extra $\log^2 1/\delta$ in the result.**14.19. THEOREM 14.19.** *As $T \rightarrow \infty$,*

$$\frac{1}{T} \int_0^T \{S_1(t)\}^2 dt \sim \frac{C^2}{\pi^2} + \frac{1}{2\pi^2} \sum_{n=2}^{\infty} \frac{\Lambda_1^2(n)}{n \log^2 n}.$$

Let

$$f(t) = C e^{-\delta} - \sum_{n=2}^{\infty} \frac{\Lambda_1(n) \cos(t \log n)}{n^{\frac{1}{2}} \log n} e^{-\delta n}.$$

Then, as in the lemma,

$$\frac{2}{T} \int_{\frac{1}{2}T}^T \{f(t)\}^2 dt = C^2 e^{-2\delta} + \frac{1}{2} \sum_{n=2}^{\infty} \frac{\Lambda_1^2(n)}{n \log^2 n} e^{-2\delta n} + O\left(\frac{\log 1/\delta}{T\delta}\right),$$

and we can replace δ by 0 in the first two terms on the right with error

$$O(\delta) + O\left(\sum_{n=2}^{\infty} \frac{1-e^{-2\delta n}}{n \log^2 n}\right) = O(\delta) + O\left(\sum_{n \leq 1/\delta} \frac{\delta}{n \log^2 n}\right) + O\left(\sum_{n > 1/\delta} \frac{1}{n \log^2 n}\right)$$

$$= O(1/\log(1/\delta)).$$

Hence, taking $\delta = T^{-\frac{1}{2}}$,

$$\frac{2}{T} \int_{\frac{1}{2}T}^T \{f(t)\}^2 dt = C^2 + \frac{1}{2} \sum_{n=2}^{\infty} \frac{\Lambda_1^2(n)}{n \log^2 n} + O\left(\frac{1}{\log T}\right).$$

Hence

$$\frac{2\pi^2}{T} \int_{\frac{1}{2}T}^T \{S_1(t)\}^2 dt = \frac{2}{T} \int_{\frac{1}{2}T}^T \left\{f(t) + O\left(\frac{1}{\log \log T}\right)\right\}^2 dt$$

$$= \frac{2}{T} \int_{\frac{1}{2}T}^T \{f(t)\}^2 dt + O\left(\frac{1}{T \log \log T} \int_{\frac{1}{2}T}^T |f(t)| dt\right) + O\left(\frac{1}{(\log \log T)^2}\right),$$

and, since $\int_{\frac{1}{2}T}^T |f(t)| dt \leq \left[\int_{\frac{1}{2}T}^T dt \int_{\frac{1}{2}T}^T \{f(t)\}^2 dt \right]^{\frac{1}{2}} = O(T)$,

it follows that

$$\frac{2}{T} \int_{\frac{1}{2}T}^T \{S_1(t)\}^2 dt = \frac{C^2}{\pi^2} + \frac{1}{2\pi^2} \sum_{n=1}^{\infty} \frac{\Lambda_1^2(n)}{n \log^2 n} + O\left(\frac{1}{\log \log T}\right).$$

Replacing T by $\frac{1}{2}T, \frac{1}{4}T, \dots$ and adding, we obtain the result.

14.20. The corresponding problem involving $S(t)$ is naturally more difficult, but it has recently been solved by A. Selberg (4). The solution depends on the following formula for $\zeta'(s)/\zeta(s)$

THEOREM 14.20. *Without any hypothesis*

$$\begin{aligned} \frac{\zeta'(s)}{\zeta(s)} = & - \sum_{n < x^2} \frac{\Lambda_x(n)}{n^s} + \frac{x^{2(1-s)} - x^{1-s}}{(1-s)^2 \log x} + \\ & + \log x \sum_{q=1}^{\infty} \frac{x^{-2q-s} - x^{-2(2q+s)}}{(2q+s)^2} + \log x \sum_{\rho} \frac{x^{\rho-s} - x^{2(\rho-s)}}{(s-\rho)^2}, \end{aligned}$$

where

$$\Lambda_x(n) = \Lambda(n) \quad (1 \leq n \leq x), \quad \frac{\Lambda(n) \log(x^2/n)}{\log x} \quad (x \leq n \leq x^2).$$

Let $\alpha = \max(2, 1+\sigma)$. Then

$$\begin{aligned} & \frac{1}{2\pi i} \int_{\alpha-i\infty}^{\alpha+i\infty} \frac{x^z - x^{2(2z-s)}}{(z-s)^2} \frac{\zeta'(z)}{\zeta(z)} dz \\ &= -\frac{1}{2\pi i} \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s} \int_{\alpha-i\infty}^{\alpha+i\infty} \frac{x^{z-s} - x^{2(2z-s)}}{(z-s)^2 n^z} dz \\ &= -\frac{1}{2\pi i} \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s} \int_{\alpha-\sigma-i\infty}^{\alpha-\sigma+i\infty} \frac{x^{w-s} - x^{2w}}{w^2 n^w} dw \\ &= -\frac{1}{2\pi i} \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s} \left\{ \int_{\alpha-\sigma-i\infty}^{\alpha-\sigma+i\infty} \left(\frac{x}{n}\right)^w \frac{dw}{w^2} - \int_{\alpha-\sigma-i\infty}^{\alpha-\sigma+i\infty} \left(\frac{x^2}{n}\right)^w \frac{dw}{w^2} \right\} \\ &= - \sum_{n < x} \frac{\Lambda(n)}{n^s} \left(\log \frac{x}{n} - \log \frac{x^2}{n} \right) - \sum_{x < n \leq x^2} \frac{\Lambda(n)}{n^s} \left(-\log \frac{x^2}{n} \right) \\ &= \log x \sum_{n < x^2} \frac{\Lambda_x(n)}{n^s}. \end{aligned}$$

Now consider the residues obtained by moving the line of integration to the left. The residue at $z = s$ is $-\log x \zeta'(s)/\zeta(s)$; that at $z = 1$ is

$$-\frac{x^{1-s} - x^{2(1-s)}}{(1-s)^2};$$

those at $z = -2q$ and $z = \rho$ are

$$\frac{x^{-2q-s} - x^{2(-2q-s)}}{(-2q-s)^2}, \quad \frac{x^{\rho-s} - x^{2(\rho-s)}}{(\rho-s)^2},$$

respectively. The result now easily follows.

14.21. THEOREM 14.21. *For $t > 2$, $4 \leq x \leq t^2$,*

$$\sigma_1 = \frac{1}{2} + \frac{1}{\log x},$$

we have

$$S(t) = -\frac{1}{\pi} \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1}} \frac{\sin(t \log n)}{\log n} + O\left\{ \frac{1}{|\log x|} \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1+t}} \right\} + O\left(\frac{\log t}{\log x}\right).$$

By the previous theorem,

$$\frac{\zeta'(\sigma+it)}{\zeta(\sigma+it)} = - \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma+it}} + O\left\{ \frac{x^{2(1-\sigma)} + x^{1-\sigma}}{t^2 \log x} \right\} + \frac{2\omega x^{\frac{1}{2}-\sigma}}{\log x} \sum_{\gamma} \frac{1}{(\sigma_1 - \frac{1}{2})^2 + (t-\gamma)^2} \quad (14.21.1)$$

for $\sigma \geq \sigma_1$, where $|\omega| < 1$. Now

$$\frac{x^{2(1-\sigma)} + x^{1-\sigma}}{t^2 \log x} \leq \frac{x^{1-2\sigma} + x^{-\sigma}}{\log x} < 2x^{\frac{1}{2}-\sigma}.$$

Hence

$$\frac{\zeta'(\sigma+it)}{\zeta(\sigma+it)} = - \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma+it}} + 2\omega x^{\frac{1}{2}-\sigma} \sum_{\gamma} \frac{\sigma_1 - \frac{1}{2}}{(\sigma_1 - \frac{1}{2})^2 + (t-\gamma)^2} + O(x^{\frac{1}{2}-\sigma}). \quad (14.21.2)$$

Now by (2.12.7)

$$\frac{\zeta'(s)}{\zeta(s)} = \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho} \right) + O(\log t).$$

Hence

$$\begin{aligned} \Re \frac{\zeta'(\sigma+it)}{\zeta(\sigma+it)} &= \sum_{\gamma} \left\{ \frac{\sigma - \frac{1}{2}}{(\sigma - \frac{1}{2})^2 + (t-\gamma)^2} + \frac{\frac{1}{2}}{\frac{1}{2} + \gamma^2} \right\} + O(\log t) \\ &= \sum_{\gamma} \frac{\sigma - \frac{1}{2}}{(\sigma - \frac{1}{2})^2 + (t-\gamma)^2} + O(\log t). \end{aligned}$$

Taking real parts in (14.21.2), substituting this on the left, and taking $\sigma = \sigma_1$,

$$\begin{aligned} & \sum_{\gamma} \frac{\sigma_1 - \frac{1}{2}}{(\sigma_1 - \frac{1}{2})^2 + (t - \gamma)^2} + O(\log t) \\ &= -R \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u}} + \frac{2\omega'}{e} \sum_{\gamma} \frac{\sigma_1 - \frac{1}{2}}{(\sigma_1 - \frac{1}{2})^2 + (t - \gamma)^2} \quad (|\omega'| < 1). \end{aligned}$$

Hence

$$\left(1 - \frac{2\omega'}{e}\right) \sum_{\gamma} \frac{\sigma_1 - \frac{1}{2}}{(\sigma_1 - \frac{1}{2})^2 + (t - \gamma)^2} = -R \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u}} + O(\log t).$$

Here

$$1 - \frac{2\omega'}{e} > 1 - \frac{2}{e} > \frac{1}{4}.$$

Hence

$$\sum_{\gamma} \frac{\sigma_1 - \frac{1}{2}}{(\sigma_1 - \frac{1}{2})^2 + (t - \gamma)^2} = O \left| \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u}} \right| + O(\log t). \quad (14.21.3)$$

Inserting this in (14.21.2), we get

$$\frac{\zeta'(\sigma + it)}{\zeta(\sigma + it)} = - \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma + u}} + O \left(x^{\frac{1}{2} - \sigma} \left| \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u}} \right| \right) + O(x^{\frac{1}{2} - \sigma} \log t). \quad (14.21.4)$$

Now

$$\begin{aligned} \arg \zeta \left(\frac{1}{2} + it \right) &= - \int_{\frac{1}{2}}^{\infty} \mathbf{I} \left\{ \frac{\zeta'(\sigma + it)}{\zeta(\sigma + it)} \right\} d\sigma \\ &= - \int_{\sigma_1}^{\infty} \mathbf{I} \left\{ \frac{\zeta'(\sigma + it)}{\zeta(\sigma + it)} \right\} d\sigma - (\sigma_1 - \frac{1}{2}) \mathbf{I} \left\{ \frac{\zeta'(\sigma_1 + it)}{\zeta(\sigma_1 + it)} \right\} \\ &\quad + \int_{\frac{1}{2}}^{\sigma_1} \mathbf{I} \left\{ \frac{\zeta'(\sigma_1 + it)}{\zeta(\sigma_1 + it)} - \frac{\zeta'(\sigma + it)}{\zeta(\sigma + it)} \right\} d\sigma \\ &= J_1 + J_2 + J_3. \end{aligned}$$

By (14.21.4)

$$\begin{aligned} J_1 &= \mathbf{I} \int_{\sigma_1}^{\infty} \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma + u}} d\sigma + O \left(\left| \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u}} \right| \int_{\sigma_1}^{\infty} x^{1 - \sigma} d\sigma \right) + O \left(\log t \int_{\sigma_1}^{\infty} x^{1 - \sigma} d\sigma \right) \\ &= \mathbf{I} \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u} \log n} + O \left(\frac{1}{\log x} \left| \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u}} \right| \right) + O \left(\frac{\log t}{\log x} \right). \end{aligned}$$

Also, by (14.21.4) with $\sigma = \sigma_1$,

$$\begin{aligned} |J_2| &\leq (\sigma_1 - \frac{1}{2}) \left| \frac{\zeta'(\sigma_1 + it)}{\zeta(\sigma_1 + it)} \right| \\ &= O \left((\sigma_1 - \frac{1}{2}) \left| \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u}} \right| \right) + O((\sigma_1 - \frac{1}{2}) \log t) \\ &= O \left(\frac{1}{\log x} \left| \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u}} \right| \right) + O \left(\frac{\log t}{\log x} \right). \end{aligned}$$

It remains to estimate J_3 . For $\frac{1}{2} < \sigma \leq \sigma_1$,

$$\begin{aligned} \mathbf{I} \left\{ \frac{\zeta'(\sigma_1 + it)}{\zeta(\sigma_1 + it)} - \frac{\zeta'(\sigma + it)}{\zeta(\sigma + it)} \right\} &= \sum_{\rho} \left(\frac{1}{\sigma_1 + it - \rho} - \frac{1}{\sigma + it - \rho} \right) + O(\log t) \\ &= \sum_{\gamma} \frac{(t - \gamma) \{ (\sigma - \frac{1}{2})^2 - (\sigma_1 - \frac{1}{2})^2 \}}{(\sigma - \frac{1}{2})^2 + (t - \gamma)^2 \{ (\sigma_1 - \frac{1}{2})^2 + (t - \gamma)^2 \}} + O(\log t). \end{aligned}$$

Hence

$$\begin{aligned} & \left| \mathbf{I} \left\{ \frac{\zeta'(\sigma_1 + it)}{\zeta(\sigma_1 + it)} - \frac{\zeta'(\sigma + it)}{\zeta(\sigma + it)} \right\} \right| \\ &\leq \sum_{\gamma} \frac{|t - \gamma| (\sigma_1 - \frac{1}{2})^2}{\{ (\sigma - \frac{1}{2})^2 + (t - \gamma)^2 \} \{ (\sigma_1 - \frac{1}{2})^2 + (t - \gamma)^2 \}} + O(\log t). \end{aligned}$$

Hence

$$\begin{aligned} |J_3| &\leq \sum_{\gamma} \frac{(\sigma_1 - \frac{1}{2})^2}{(\sigma_1 - \frac{1}{2})^2 + (t - \gamma)^2} \int_{\frac{1}{2}}^{\sigma_1} \frac{|t - \gamma| d\sigma}{(\sigma - \frac{1}{2})^2 + (t - \gamma)^2} + O((\sigma_1 - \frac{1}{2}) \log t) \\ &\leq \frac{1}{2} \pi (\sigma_1 - \frac{1}{2}) \sum_{\gamma} \frac{(\sigma_1 - \frac{1}{2})}{(\sigma_1 - \frac{1}{2})^2 + (t - \gamma)^2} + O((\sigma_1 - \frac{1}{2}) \log t) \\ &= O \left(\frac{1}{\log x} \left| \sum_{n < x^2} \frac{\Lambda_x(n)}{n^{\sigma_1 + u}} \right| \right) + O \left(\frac{\log t}{\log x} \right). \end{aligned}$$

by (14.21.3). The theorem follows from these results.

Theorem 14.21 leads to an alternative proof of Theorem 14.13; for taking $x = \sqrt{(\log t)}$ we obtain

$$\begin{aligned} S(t) &= O \left(\sum_{n < x} \frac{1}{n^{\frac{1}{2}}} \right) + O \left(\frac{1}{\log x} \sum_{n < x^2} \frac{\log n}{n^{\frac{1}{2}}} \right) + O \left(\frac{\log t}{\log x} \right) \\ &= O(x) + O(x) + O \left(\frac{\log t}{\log x} \right) \\ &= O \left(\frac{\log t}{\log \log t} \right). \end{aligned}$$

14.22. THEOREM 14.22. For

$$T^a \leq x \leq T^{\frac{1}{2}} \quad (0 < a \leq \frac{1}{2})$$

$$\int_{\frac{1}{2}T}^T \left\{ S(t) + \frac{1}{\pi} \sum_{p < x^2} \frac{\sin(t \log p)}{\sqrt{p}} \right\}^2 dt = O(T).$$

We have

$$\begin{aligned} S(t) + \frac{1}{\pi} \sum_{p < x^2} \frac{\sin(t \log p)}{\sqrt{p}} &= \frac{1}{\pi} \sum_{p < x^2} \frac{\Lambda(p) - \Lambda_x(p) p^{\frac{1}{2}-\sigma_1}}{p^{\frac{1}{2}} \log p} \sin(t \log p) + \\ &+ O \left\{ \frac{1}{\log x} \left| \sum_{p < x^2} \frac{\Lambda_x(p)}{p^{\sigma_1 + i t}} \right| \right\} + O \left(\left| \sum_{p < x^2} \frac{\Lambda_x(p^2)}{p^{2(\sigma_1 + i t)} \log p} \right| \right) + \\ &+ O \left(\frac{1}{\log x} \left| \sum_{p < x^2} \frac{\Lambda_x(p^2)}{p^{2(\sigma_1 + i t)}} \right| \right) + O \left(\sum_{p > \frac{1}{2}} \frac{1}{p^{\frac{1}{2}}} \right) + O \left(\frac{\log t}{\log x} \right). \end{aligned}$$

The last term is bounded if $\frac{1}{2}T \leq t \leq T$, $x \geq T^a$, where a is a fixed positive constant. The last term but one is

$$O \left(\sum_{r=3}^{\infty} \frac{1}{p^{\frac{1}{2}r}} \right) = O \left(\sum_p \frac{p^{-\frac{3}{2}}}{1-p^{-\frac{1}{2}}} \right) = O(1).$$

Now consider the first term on the right. If $p \leq x$,

$$\begin{aligned} \Lambda(p) - \Lambda_x(p) p^{\frac{1}{2}-\sigma_1} &= (1 - p^{\frac{1}{2}-\sigma_1}) \log p \\ &= (1 - p^{-1/\log x}) \log p = (1 - e^{-\log p / \log x}) \log p = O \left(\frac{\log^2 p}{\log x} \right), \end{aligned}$$

and, if $x < p \leq x^2$, it is

$$O\{\Lambda_x(p)\} = O \left(\log p \frac{\log x^2/p}{\log x} \right) = O \left(\frac{\log^2 p}{\log x} \right).$$

Hence the first term is the imaginary part of

$$\sum_{p < x^2} \frac{\alpha_p}{p^{\frac{1}{2} + i t}},$$

where

$$\alpha_p = \frac{\Lambda(p) - \Lambda_x(p) p^{\frac{1}{2}-\sigma_1}}{\log p} = O(\log p / \log x).$$

Now

$$\begin{aligned} \int_{\frac{1}{2}T}^T \left| \sum_{p < x^2} \frac{\alpha_p}{p^{\frac{1}{2} + i t}} \right|^2 dt &= \sum_{p < x^2} \sum_{q < x^2} \frac{\alpha_p \bar{\alpha}_q}{p^{\frac{1}{2}} q^{\frac{1}{2}}} \int_{\frac{1}{2}T}^T \left(\frac{q}{p} \right)^{it} dt \\ &= O \left(T \sum_{p < x^2} \frac{\alpha_p^2}{p} \right) + O \left(\sum_{p \neq q} \sum_{p < x^2} \frac{|\alpha_p \bar{\alpha}_q|}{p^{\frac{1}{2}} q^{\frac{1}{2}}} \frac{1}{|\log p/q|} \right). \end{aligned}$$

Since

$$\alpha_p^2 = O \left(\frac{\log^2 p}{\log^2 x} \right) = O \left(\frac{\log p}{\log x} \right),$$

the first term is

$$O \left(T \frac{1}{\log x} \sum_{p < x^2} \frac{\log p}{p} \right) = O(T),$$

by (3.14.3). The second term is

$$\begin{aligned} &O \left(\sum_{p < x^2} \sum_{q < \frac{1}{2}p} \frac{|\alpha_p \bar{\alpha}_q|}{p^{\frac{1}{2}} q^{\frac{1}{2}}} \right) + O \left(\sum_{p < x^2} \sum_{\frac{1}{2}p < q < p} \frac{|\alpha_p \bar{\alpha}_q|}{p^{\frac{1}{2}} (p-q)/p} \right) \\ &= O \left(\sum_{p < x^2} \frac{\log p}{p^{\frac{1}{2}} \log x} \right) + O \left(\sum_{p < x^2} \frac{\log^2 p}{\log^2 x} \cdot \log p \right) \\ &= O \left(\sum_{p < x^2} \frac{1}{p^{\frac{1}{2}}} \right) + O \left(\sum_{p < x^2} \log p \right) \\ &= O(x^2) + O(x^2) = O(T) \end{aligned}$$

if $x \leq \sqrt{T}$.

A similar argument clearly applies to the second term. In the third term, the sum is of the form

$$\sum_{p < x} \frac{\alpha'_p}{p^{\frac{1}{2} + i t}},$$

where $\alpha'_p = O(1)$; and

$$\begin{aligned} \int_{\frac{1}{2}T}^T \left| \sum_{p < x} \frac{\alpha'_p}{p^{\frac{1}{2} + i t}} \right|^2 dt &= \sum_{p < x} \sum_{q < x} \frac{\alpha'_p \bar{\alpha}'_q}{pq} \int_{\frac{1}{2}T}^T \left(\frac{q}{p} \right)^{it} dt \\ &= O \left(T \sum_{p < x} \frac{1}{p^3} \right) + O \left(\sum_{p \neq q} \sum_{p < x} \frac{1}{pq |\log p/q|} \right) \\ &= O(T) + O \left(\sum_{p < x} \sum_{q < \frac{1}{2}p} \frac{1}{pq} \right) + O \left(\sum_{p < x} \sum_{\frac{1}{2}p < q < p} \frac{1}{p^2 \cdot (p-q)/p} \right) \\ &= O(T) + O(\log^2 x) + O(\log^2 x) = O(T). \end{aligned}$$

Similarly for the fourth term, and the result follows.

14.23. THEOREM 14.23. If $T^a \leq x \leq T^{\frac{1}{2}}$,

$$\int_{\frac{1}{2}T}^T \left\{ \sum_{p < x^2} \frac{\sin(t \log p)}{\sqrt{p}} \right\}^2 dt = \frac{1}{2} T \log \log T + O(T).$$

This is

$$\sum_{p < x^2} \sum_{q < x^2} \frac{1}{p^{\frac{1}{2}} q^{\frac{1}{2}}} \int_{\frac{1}{2}T}^T \sin(t \log p) \sin(t \log q) dt \\ = \sum_{p < x^2} \frac{1}{p} \left\{ \frac{1}{2} T + O\left(\frac{1}{\log p}\right) \right\} + O\left(\sum_{p \neq q} \sum \frac{1}{p^{\frac{1}{2}} q^{\frac{1}{2}} |\log p/q|} \right).$$

Now, by (3.14.5),

$$\sum_{p < x^2} \frac{1}{p} = \log \log x^2 + O(1) = \log \log T + O(1)$$

and (since $p_n > A n \log n$)

$$\sum \frac{1}{p \log p} = O(1).$$

Hence the first term is

$$\frac{1}{2} T \log \log T + O(T).$$

Also

$$|\log p/q| > A/p > A/x^2.$$

Hence the remainder is

$$O\left\{x^2 \left(\sum_{p < x^2} \frac{1}{p^{\frac{1}{2}}} \right)^2\right\} = O(x^2 \cdot x^2) = O(x^4),$$

and the result follows if $x \leq T^{\frac{1}{2}}$.

14.24. THEOREM 14.24.

$$\text{For } \int_0^T \{S(t)\}^2 dt \sim \frac{1}{2\pi^2} T \log \log T.$$

$$\begin{aligned} \int_{\frac{1}{2}T}^T \{S(t)\}^2 dt &= \int_{\frac{1}{2}T}^T \left\{ S(t) + \frac{1}{\pi} \sum_{p < x^2} \frac{\sin(t \log p)}{\sqrt{p}} - \frac{1}{\pi} \sum_{p < x^2} \frac{\sin(t \log p)}{\sqrt{p}} \right\}^2 dt \\ &= \int_{\frac{1}{2}T}^T \left\{ S(t) + \frac{1}{\pi} \sum_{p < x^2} \frac{\sin(t \log p)}{\sqrt{p}} \right\}^2 dt - \\ &\quad - \frac{2}{\pi} \int_{\frac{1}{2}T}^T \left\{ S(t) + \frac{1}{\pi} \sum_{p < x^2} \frac{\sin(t \log p)}{\sqrt{p}} \right\} \sum_{p < x^2} \frac{\sin(t \log p)}{\sqrt{p}} dt + \\ &\quad + \frac{1}{\pi^2} \int_{\frac{1}{2}T}^T \left\{ \sum_{p < x^2} \frac{\sin(t \log p)}{\sqrt{p}} \right\}^2 dt \\ &= O(T) + O\{T(\log \log T)^{\frac{1}{2}}\} + \frac{1}{4\pi^2} T \log \log T + O(T) \end{aligned}$$

(using Schwarz's inequality on the middle term). The result then follows by addition.

It can be proved in a similar way that

$$\int_0^T \{S(t)\}^{2k} dt \sim \frac{(2k)!}{k!(2\pi)^{2k}} T (\log \log T)^k \quad (14.24.1)$$

for every positive integer k .

14.25. The Dirichlet series for $1/\zeta(s)$. It was proved in § 3.13 that the formula

$$\frac{1}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\mu(n)}{n^s},$$

which is elementary for $\sigma > 1$, holds also for $\sigma = 1$. On the Riemann hypothesis we can go much farther than this.†

THEOREM 14.25 (A). *The series*

$$\sum_{n=1}^{\infty} \frac{\mu(n)}{n^s} \quad (14.25.1)$$

is convergent, and its sum is $1/\zeta(s)$, for every s with $\sigma > \frac{1}{2}$.

In the lemma of § 3.12, take $a_n = \mu(n)$, $f(s) = 1/\zeta(s)$, $c = 2$, and x half an odd integer. We obtain

$$\begin{aligned} \sum_{n < x} \frac{\mu(n)}{n^s} &= \frac{1}{2\pi i} \int_{2-iT}^{2+iT} \frac{1}{\zeta(s+w)} \frac{x^w}{w} dw + O\left(\frac{x^2}{T}\right) \\ &= \frac{1}{2\pi i} \left(\int_{2-iT}^{\frac{1}{2}-\sigma+\delta-iT} + \int_{\frac{1}{2}-\sigma+\delta-iT}^{\frac{1}{2}-\sigma+\delta+iT} + \int_{\frac{1}{2}-\sigma+\delta+iT}^{2+iT} \right) \frac{1}{\zeta(s+w)} \frac{x^w}{w} dw + \frac{1}{\zeta(s)} + O\left(\frac{x^2}{T}\right), \end{aligned}$$

where $0 < \delta < \sigma - \frac{1}{2}$.

By (14.2.5), the first and third integrals are

$$O\left(T^{-1+\epsilon} \int_{\frac{1}{2}-\sigma+\delta}^2 x^u du\right) = O(T^{-1+\epsilon} x^2),$$

and the second integral is

$$O\left\{x^{\frac{1}{2}-\sigma+\delta} \int_{-T}^T (1+|t|)^{-1+\epsilon} dt\right\} = O(x^{\frac{1}{2}-\sigma+\delta} T^{\epsilon}).$$

† Littlewood (1).

$$\text{Hence } \sum_{n \leq x} \frac{\mu(n)}{n^s} = \frac{1}{\zeta(s)} + O(T^{-1+\epsilon}x^2) + O(T^\epsilon x^{\frac{1}{2}-\sigma+3}).$$

Taking, for example, $T = x^2$, the O -terms tend to zero as $x \rightarrow \infty$, and the result follows.

Conversely, if (14.25.1) is convergent for $\sigma > \frac{1}{2}$, it is uniformly convergent for $\sigma \geq \sigma_0 > \frac{1}{2}$, and so in this region represents an analytic function, which is $1/\zeta(s)$ for $\sigma > 1$ and so throughout the region. Hence the Riemann hypothesis is true. We have in fact

THEOREM 14.25 (B). *The convergence of (14.25.1) for $\sigma > \frac{1}{2}$ is a necessary and sufficient condition for the truth of the Riemann hypothesis.*

We shall write

$$M(x) = \sum_{n \leq x} \mu(n).$$

Then we also have

THEOREM 14.25 (C). *A necessary and sufficient condition for the Riemann hypothesis is* $M(x) = O(x^{\frac{1}{2}+\epsilon}).$ (14.25.2)

The lemma of § 3.12 with $s = 0$, x half an odd integer, gives

$$\begin{aligned} M(x) &= \frac{1}{2\pi i} \int_{2-iT}^{2+iT} \frac{1}{\zeta(w)} \frac{x^w}{w} dw + O\left(\frac{x^2}{T}\right) \\ &= \frac{1}{2\pi i} \left(\int_{2-iT}^{\frac{1}{2}+\delta-iT} + \int_{\frac{1}{2}+\delta-iT}^{\frac{1}{2}+\delta+iT} + \int_{\frac{1}{2}+\delta+iT}^{2+iT} \right) \frac{1}{\zeta(w)} \frac{x^w}{w} dw + O\left(\frac{x^2}{T}\right) \\ &= O\left(\int_{-T}^T (1+|v|)^{-1+\epsilon} x^{\frac{1}{2}+\delta} dv\right) + O(T^{\epsilon-1}x^2) + O\left(\frac{x^2}{T}\right) \\ &= O(T^\epsilon x^{\frac{1}{2}+\delta}) + O(x^2 T^{\epsilon-1}), \end{aligned} \quad (14.25.3)$$

by (14.2.6). Taking $T = x^2$, (14.25.2) follows if x is half an odd integer, and so generally.

Conversely, if (14.25.2) holds, then by partial summation (14.25.1) converges for $\sigma > \frac{1}{2}$, and the Riemann hypothesis follows.

14.26. The finer theory of $M(x)$ is extremely obscure, and the results are not nearly so precise as the corresponding ones in the prime-number problem. The best O -result known is

THEOREM 14.26 (A).[†]

$$M(x) = O\left\{x^{\frac{1}{2}} \exp\left(A \frac{\log x}{\log \log x}\right)\right\}. \quad (14.26.1)$$

To prove this, take

$$\delta = \frac{1}{2} + \frac{1}{\log \log T}, \quad T = x^2,$$

in the formula (14.25.3). By (14.14.2),

$$\left| \frac{1}{\zeta(w)} \right| \leq \exp\left(A \frac{\log T}{\log \log T}\right)$$

on the horizontal sides of the contour. The contribution of these is therefore

$$O\left\{x^2 \frac{1}{T} \exp\left(A \frac{\log T}{\log \log T}\right)\right\} = O\left\{\exp\left(A \frac{\log x}{\log \log x}\right)\right\}.$$

On the vertical side, (14.14.2) gives

$$\left| \frac{1}{\zeta(\frac{1}{2} + \delta + iv)} \right| \leq \exp\left(A \frac{\log v}{\log \log v} \log \frac{2 \log \log T}{\log \log v}\right)$$

for $v_0 \leq v \leq T$. Now it is easily seen that the right-hand side is a steadily increasing function of v in this interval. Hence

$$\left| \frac{1}{\zeta(w)} \right| \leq \exp\left(A \frac{\log T}{\log \log T}\right) \quad (v_0 \leq v \leq T).$$

Hence the integral along the vertical side is of the form

$$\begin{aligned} O(x^{\frac{1}{2}+\delta}) + O\left\{x^{\frac{1}{2}+\delta} \exp\left(A \frac{\log T}{\log \log T}\right) \int_{v_0}^T \frac{dv}{v}\right\} \\ = O\left\{x^{\frac{1}{2}+\delta} \exp\left(A \frac{\log T}{\log \log T}\right) \log T\right\} = O\left\{x^{\frac{1}{2}} \exp\left(A \frac{\log x}{\log \log x}\right)\right\}. \end{aligned}$$

This proves the theorem.

THEOREM 14.26 (B).

$$M(x) = \Omega(x^{\frac{1}{2}}). \quad (14.26.2)$$

This is true without any hypothesis. For if the Riemann hypothesis is false, Theorem 14.25 (C) shows that

$$M(x) = \Omega(x^a)$$

[†] Landau (13), Titchmarsh (3).

with some a greater than $\frac{1}{2}$. On the other hand, if the Riemann hypothesis is true, then for $\sigma > \frac{1}{2}$

$$\frac{1}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\mu(n)}{n^s} = \sum_{n=1}^{\infty} M(n) \left\{ \frac{1}{n^s} - \frac{1}{(n+1)^s} \right\} = s \int_1^{\infty} \frac{M(x)}{x^{s+1}} dx. \quad (14.26.3)$$

Suppose that

$$|M(x)| \leq M_0 \quad (1 \leq x < x_0), \quad \leq \delta x^{\frac{1}{2}} \quad (x \geq x_0).$$

Then

$$\begin{aligned} \left| \frac{1}{\zeta(s)} \right| &\leq |s| M_0 \int_1^{x_0} \frac{dx}{x^{\frac{1}{2}}} + |s| \delta \int_{x_0}^{\infty} \frac{dx}{x^{\sigma+\frac{1}{2}}} \\ &< |s| M_0 \int_1^{\infty} \frac{dx}{x^{\frac{1}{2}}} + |s| \delta \int_1^{\infty} \frac{dx}{x^{\sigma+\frac{1}{2}}} \\ &= 2|s| M_0 + \frac{|s| \delta}{\sigma - \frac{1}{2}}. \end{aligned} \quad (14.26.4)$$

But if $\rho = \frac{1}{2} + i\gamma$ is a simple zero of $\zeta(s)$, and $s = \sigma + i\gamma$, $\sigma \rightarrow \frac{1}{2}$, then

$$\frac{1}{\zeta(s)} = \frac{1}{\zeta(s) - \zeta(\rho)} \sim \frac{1}{(\sigma - \frac{1}{2}) \zeta'(\rho)}.$$

We therefore obtain a contradiction if

$$\delta < \frac{1}{|\rho \zeta'(\rho)|}.$$

This proves the theorem.

14.27. Formulae connecting the functions of prime-number theory with series of the form

$$\sum x^{\rho}, \quad \sum_{\rho} \frac{x^{\rho}}{\rho},$$

etc., are well known, and are discussed in the books of Landau and Ingham. Here we prove a similar formula for the function $M(x)$.

THEOREM 14.27. *There is a sequence T_v , $v \leq T_v \leq v+1$, such that*

$$M(x) = \lim_{v \rightarrow \infty} \sum_{|v| < T_v} \frac{x^{\rho}}{\rho \zeta'(\rho)} - 2 + \sum_{n=1}^{\infty} \frac{(-1)^{n-1} (2\pi/x)^{2n}}{(2n)! n \zeta(2n+1)} \quad (14.27.1)$$

if x is not an integer. If x is an integer, $M(x)$ is to be replaced by

$$M(x) - \frac{1}{2} \mu(x).$$

In writing the series we have supposed for simplicity that all the zeros of $\zeta(s)$ are simple; obvious modifications are required if this is not so.

For a fixed non-integral x , (3.12.1), with $a_n = \mu(n)$, $s = 0$, $c = 2$, and v replaced by s , gives

$$M(x) = \frac{1}{2\pi i} \int_{2-iT}^{2+iT} \frac{x^s}{s} \frac{1}{\zeta(s)} ds + O\left(\frac{1}{T}\right).$$

If x is an integer, $\frac{1}{2} \mu(x)$ is to be subtracted from the left-hand side. By the calculus of residues, the first term on the right is equal to

$$\begin{aligned} \sum_{|v| < T} \frac{x^{\rho}}{\rho \zeta'(\rho)} - 2 + \sum_{n=1}^N \frac{(-1)^{n-1} (2\pi/x)^{2n}}{(2n)! n \zeta(2n+1)} + \\ + \frac{1}{2\pi i} \left(\int_{2-iT}^{2N-1-iT} + \int_{2N-1-iT}^{-2N-1-iT} + \int_{-2N-1-iT}^{2+iT} \right) \frac{x^s}{s \zeta(s)} ds, \end{aligned}$$

where T is not the ordinate of a zero. Now

$$\begin{aligned} \int_{-2N-1-iT}^{2N-1-iT} \frac{x^s}{s \zeta(s)} ds &= \int_{2N+2-iT}^{2N+2+iT} \frac{x^{1-s}}{(1-s) \zeta(1-s)} ds \\ &= \int_{2N+2-iT}^{2N+2+iT} \frac{x^{1-s}}{1-s} \frac{2^{s-1} \pi^s}{\cos \frac{1}{2} \pi s \Gamma(s)} \frac{1}{\zeta(s)} ds. \end{aligned}$$

Here

$$\begin{aligned} \frac{1}{\Gamma(s)} &= O(|e^{-(s-\frac{1}{2}) \log s}|) = O(e^{-(\sigma-\frac{1}{2}) \log s + \frac{1}{2} \pi \gamma}) \\ &= O(e^{-(\sigma-\frac{1}{2}) \log s + \frac{1}{2} \pi \gamma}). \end{aligned}$$

Hence the integral is

$$O\left(\int_{-T}^T \frac{1}{T} \left(\frac{2\pi}{x}\right)^{2N+2} e^{2N+2-(2N+\frac{1}{2}) \log(2N+2)} dt\right),$$

which tends to zero as $N \rightarrow \infty$, for a fixed T . Hence we obtain

$$\sum_{|v| < T} \frac{x^{\rho}}{\rho \zeta'(\rho)} - 2 + \sum_{n=1}^{\infty} \frac{(-1)^{n-1} (2\pi/x)^{2n}}{(2n)! n \zeta(2n+1)} + \frac{1}{2\pi i} \left(\int_{2-iT}^{-\infty-iT} + \int_{-\infty+iT}^{2+iT} \right) \frac{x^s}{s \zeta(s)} ds.$$

Also

$$\begin{aligned} \int_{-\infty+iT}^{-1+iT} \frac{x^s}{s \zeta(s)} ds &= \int_{2+iT}^{\infty+iT} \frac{x^{1-s}}{(1-s) \zeta(1-s)} ds = \int_{2+iT}^{\infty+iT} \frac{x^{1-s}}{1-s} \frac{2^{s-1} \pi^s}{\cos \frac{1}{2} \pi s \Gamma(s)} \frac{1}{\zeta(s)} ds \\ &= O\left(\int_{\frac{1}{2}}^{\infty} \frac{1}{T} \left(\frac{2\pi}{x}\right)^{\sigma} e^{-(\sigma-\frac{1}{2}) \log \sigma} d\sigma\right) = O\left(\frac{1}{T}\right). \end{aligned}$$

Also by (14.16.2) we can choose $T = T_\nu$ ($\nu \leq T_\nu \leq \nu+1$) such that

$$\frac{1}{\zeta(s)} = O(t^\epsilon) \quad \left(\frac{1}{2} \leq \sigma \leq 2, t = T_\nu\right).$$

Hence for $-1 \leq \sigma \leq \frac{1}{2}$, $t = T_\nu$

$$\frac{1}{\zeta(s)} = O\left(\frac{|t|^{\sigma-\frac{1}{2}}}{|\zeta(1-s)|}\right) = O(t^\epsilon).$$

Hence

$$\int_{-1-iT_\nu}^{2+iT_\nu} \frac{x^s}{s \zeta(s)} ds = O(T_\nu^{\epsilon-1}).$$

Similarly for the integral over $(2-iT, -\infty-iT)$, and the result stated follows.

It follows from the above theorem that

$$\sum \frac{1}{|\rho \zeta'(\rho)|}$$

is divergent; if it were convergent,

$$\sum \frac{x^\rho}{\rho \zeta'(\rho)}$$

would be uniformly convergent over any finite interval, and $M(x)$ would be continuous.

14.28. The Mertens hypothesis.† It was conjectured by Mertens, from numerical evidence, that

$$|M(n)| < \sqrt{n} \quad (n > 1). \quad (14.28.1)$$

This has not been proved or disproved. It implies the Riemann hypothesis, but is not apparently a consequence of it. A slightly less precise hypothesis would be

$$M(x) = O(x^{\frac{1}{2}}). \quad (14.28.2)$$

The problem has a certain similarity to that of the function $\psi(x)-x$ in prime-number theory, where

$$\psi(x) = \sum_{n \leq x} \Lambda(n).$$

On the Riemann hypothesis, $\psi(x)-x = O(x^{\frac{1}{2}+\epsilon})$, but it is not of the form $O(x^{\frac{1}{2}})$, and in fact

$$\psi(x)-x = \Omega(x^{\frac{1}{2}} \log \log x). \quad (14.28.3)$$

The influence of the factor $\log \log x$ is quite inappreciable as far as

† See references in Landau's *Handbuch*, and von Sterneck (1).

the calculations go, and it might be conjectured that (14.28.2) could be disproved similarly. We shall show, however, that there is an essential difference between the two problems, and that the proof of (14.28.3) cannot be extended to the other case, at any rate in any obvious way.

The proof of (14.28.3) depends on the fact that the real part of

$$\sum_{\gamma > 0} \frac{e^{i\rho s}}{\rho}$$

is unbounded in the neighbourhood of $z = 0$. To deal with $M(x)$ in the same way, we should have to prove that the real part of

$$f(z) = \sum_{\gamma > 0} \frac{e^{i\rho z}}{\rho \zeta'(\rho)} \quad (\Re(z) > 0)$$

is unbounded in the neighbourhood of $z = 0$. This, however, is not the case. For consider the integral

$$\frac{1}{2\pi i} \int \frac{e^{isz}}{s \zeta(s)} ds$$

taken round the rectangle $(-1, 2, 2+iT_n, -1+iT_n)$, where the T_n are those of the previous section, and an indentation is made above $s = 0$. The integral along the upper side of the contour tends to 0 as $n \rightarrow \infty$, and we calculate that

$$f(z) = \frac{1}{2\pi i} \int_{\frac{1}{2}}^{2+i\infty} \frac{e^{isz}}{s \zeta(s)} ds - \frac{1}{2\pi i} \int_{-1}^{-1+i\infty} \frac{e^{isz}}{s \zeta(s)} ds + \frac{1}{2\pi i} \int_{-1}^{\frac{1}{2}} \frac{e^{isz}}{s \zeta(s)} ds.$$

The last term tends to a finite limit as $z \rightarrow 0$. Also

$$|e^{isz}| = e^{\nu-xt} \leq e^\nu \quad (s = -1+it, z = x+it, x > 0)$$

and $1/\zeta(-1+it) = O(t^{-\frac{1}{2}})$. The second term is therefore bounded for $\Re(z) > 0$.

The first term is equal to

$$\frac{1}{2\pi i} \sum_{n=1}^{\infty} \mu(n) \int_{\frac{1}{2}}^{2+i\infty} \frac{e^{isz}}{s n^s} ds.$$

Now, if $n > 1$,

$$\int_{\frac{1}{2}}^{2+i\infty} \frac{e^{s(is-\log n)}}{s} ds = \left[\frac{e^{s(is-\log n)}}{s} \right]_{\frac{1}{2}}^{2+i\infty} + \frac{1}{iz-\log n} \int_{\frac{1}{2}}^{2+i\infty} \frac{e^{s(is-\log n)}}{s^2} ds$$

and $|e^{(2+i\infty)(is-\log n)}| = e^{-2\nu-2 \log n - t\infty} \leq n^{-2} e^{-2\nu}$.

Hence

$$\int_{\frac{1}{2}}^{2+i\infty} \frac{e^{s(is-\log n)}}{s} ds = O\left(\frac{1}{n^2 \log n}\right)$$

uniformly in the neighbourhood of $z = 0$. Hence

$$\sum_{n=2}^{\infty} \mu(n) \int_2^{2+i\infty} \frac{e^{i\lambda z}}{s^{\sigma}} ds = O(1).$$

If $z = re^{i\theta}$, we have

$$\begin{aligned} \int_2^{2+i\infty} \frac{e^{i\lambda z}}{s} ds &= \int_2^{e^{i(1-\theta)}} + \int_{e^{i(1-\theta)}}^{re^{i(1-\theta)}} \\ &= O(1) + \int_1^{\infty} \frac{e^{-r\lambda}}{\lambda} d\lambda \\ &= O(1) + \int_r^{\infty} \frac{e^{-x}}{x} dx \\ &= O(1) + \int_r^1 \frac{dx}{x} + \int_r^1 \frac{e^x - 1}{x} dx + \int_1^{\infty} \frac{e^{-x}}{x} dx \\ &= \log \frac{1}{r} + O(1). \end{aligned}$$

Hence

$$f(z) = \frac{1}{2\pi i} \log \frac{1}{r} + O(1),$$

and consequently $Rf(z)$ is bounded.

14.29. In this section we shall investigate the consequences of the hypothesis that

$$\int_1^X \left\{ \frac{M(x)}{x} \right\}^2 dx = O(\log X). \quad (14.29.1)$$

This is less drastic than the Mertens hypothesis, since it clearly follows from (14.28.2). The corresponding formula with $M(x)$ replaced by $\psi(x) - x$ is a consequence of the Riemann hypothesis.†

THEOREM 14.29 (A). *If (14.29.1) is true, all the zeros of $\zeta(s)$ on the critical line are simple.*

By (14.26.3),

$$\begin{aligned} \left| \frac{1}{\zeta(s)} \right| &\leq |s| \int_1^{\infty} \frac{|M(x)|}{x^{\sigma+1}} dx = |s| \int_1^{\infty} \frac{|M(x)|}{x^{\frac{1}{2}\sigma+\frac{1}{2}}} \frac{1}{x^{\frac{1}{2}\sigma+\frac{1}{2}}} dx \\ &\leq |s| \left(\int_1^{\infty} \frac{M^2(x)}{x^{\sigma+\frac{1}{2}}} dx \int_1^{\infty} \frac{dx}{x^{\sigma+\frac{1}{2}}} \right)^{\frac{1}{2}} = \frac{|s|}{(\sigma-\frac{1}{2})^{\frac{1}{2}}} \left(\int_1^{\infty} \frac{M^2(x)}{x^{\sigma+\frac{1}{2}}} dx \right)^{\frac{1}{2}}. \end{aligned}$$

† Cramér (5).

Let

$$f(X) = \int_1^X \left\{ \frac{M(x)}{x} \right\}^2 dx.$$

Then

$$\begin{aligned} \int_1^{\infty} \frac{M^2(x)}{x^{\sigma+\frac{1}{2}}} dx &= \int_1^{\infty} \frac{f'(x)}{x^{\sigma+\frac{1}{2}}} dx = (\sigma-\frac{1}{2}) \int_1^{\infty} \frac{f(x)}{x^{\sigma+\frac{1}{2}}} dx \\ &= O\left((\sigma-\frac{1}{2}) \int_1^{\infty} \frac{\log x}{x^{\sigma+\frac{1}{2}}} dx\right) = O\left(\int_1^{\infty} \frac{dx}{x^{\sigma+\frac{1}{2}}}\right) = O\left(\frac{1}{\sigma-\frac{1}{2}}\right). \end{aligned}$$

Hence

$$\frac{1}{\zeta(s)} = O\left(\frac{|s|}{\sigma-\frac{1}{2}}\right).$$

Let ρ be a zero and $s = \rho + h$, where $h > 0$. Then $\sigma = \frac{1}{2} + h$, and hence

$$\frac{1}{\zeta(\rho+h)} = O\left(\frac{1}{h}\right). \quad (14.29.2)$$

This would be false for $h \rightarrow 0$ if ρ were a zero of order higher than the first, so that the result follows.

Multiplying each side of (14.29.2) by h , and making $h \rightarrow 0$, we obtain

$$\frac{1}{\zeta'(\rho)} = O(|\rho|). \quad (14.29.3)$$

We can, however, prove more than this.

THEOREM 14.29 (B). *If (14.29.1) is true,*

$$\sum \frac{1}{|\rho \zeta'(\rho)|^2} \quad (14.29.4)$$

is convergent.

This follows from an argument of the 'Bessel's inequality' type. We have

$$\begin{aligned} 0 &\leq \int_1^X \left\{ \frac{M(x)}{x} - \sum_{|\gamma| < T} \frac{x^{\rho-1}}{\rho \zeta'(\rho)} \right\}^2 dx \\ &= \int_1^X \left\{ \frac{M(x)}{x} \right\}^2 dx + \sum_{|\gamma| < T} \sum_{|\gamma'| < T} \frac{1}{\rho \rho' \zeta'(\rho) \zeta'(\rho')} \int_1^X x^{\rho+\rho'-2} dx - \\ &\quad - 2 \sum_{|\gamma| < T} \frac{1}{\rho \zeta'(\rho)} \int_1^X M(x) x^{\rho-2} dx. \end{aligned}$$

In the first sum, the terms with $\rho' = 1 - \rho$ are

$$\sum_{|\gamma| < x} \frac{1}{\rho(1-\rho)\zeta'(\rho)\zeta'(1-\rho)} \int_1^x \frac{dx}{x} = \log X \sum_{|\gamma| < x} \frac{1}{|\rho\zeta'(\rho)|^2},$$

since $1-\rho$ is the conjugate of ρ . In the remaining terms, $\rho = \frac{1}{2} + i\gamma$, $\rho' = \frac{1}{2} + i\gamma'$, where $\gamma' \neq -\gamma$. Hence

$$\int_1^x x^{\rho+\rho'-2} dx = \frac{X^{\rho+\rho'-1} - 1}{\rho+\rho'-1} = O\left(\frac{1}{|\gamma+\gamma'|}\right).$$

Hence the sum of these terms is less than $K_1 = K_1(T)$.

In the last sum we write

$$\int_1^X M(x)x^{\rho-2} dx = \int_1^X M(x)x^{\rho-2} \left(1 - \frac{x}{X}\right) dx + \frac{1}{X} \int_1^X M(x)x^{\rho-1} dx.$$

The last term is

$$\begin{aligned} O\left(\frac{1}{X} \int_1^X |M(x)|x^{\frac{1}{2}} dx\right) &= O\left[\frac{1}{X} \left(\int_1^X \frac{M^2(x)}{x^2} dx \int_1^X x dx\right)^{\frac{1}{2}}\right] \\ &= O\left(\int_1^X \frac{M^2(x)}{x^2} dx\right)^{\frac{1}{2}} = O(\log^{\frac{1}{2}} X), \end{aligned}$$

by (14.29.1). Also

$$\int_1^X M(x)x^{\rho-2} \left(1 - \frac{x}{X}\right) dx = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \frac{X^{w+\rho-1} - 1}{\zeta(w)w(w+\rho)(w+\rho-1)} dw. \quad (14.29.5)$$

To prove this, insert the Dirichlet series for $1/\zeta(w)$ on the right-hand side and integrate term by term. This is justified by absolute convergence. We obtain

$$\sum_{n=1}^{\infty} \frac{\mu(n)}{2\pi i} \int_{2-i\infty}^{2+i\infty} \frac{1}{n^w} \frac{X^{w+\rho-1} - 1}{w(w+\rho)(w+\rho-1)} dw.$$

Evaluating the integral in the usual way by the calculus of residues, we obtain

$$\begin{aligned} \sum_{n \leq X} \mu(n) \left\{ \frac{X^{\rho-1} - n^{\rho-1}}{\rho-1} - \frac{X^{\rho} - n^{\rho}}{X^{\rho}} \right\} &= \sum_{n \leq X} \mu(n) \int_n^X \left(x^{\rho-2} - \frac{x^{\rho-1}}{X} \right) dx \\ &= \int_1^X \sum_{n \leq X} \mu(n)x^{\rho-2} \left(1 - \frac{x}{X}\right) dx = \int_1^X M(x)x^{\rho-2} \left(1 - \frac{x}{X}\right) dx, \end{aligned}$$

and (14.29.5) follows.

Let U be not the ordinate of a zero. Then the right-hand side of (14.29.5) is equal to

$$\frac{1}{2\pi i} \left(\int_{2-i\infty}^{2-iU} + \int_{2-iU}^{\frac{1}{2}-iU} + \int_{\frac{1}{2}-iU}^{\frac{1}{2}+iU} + \int_{\frac{1}{2}+iU}^{2+iU} + \int_{2+iU}^{2+i\infty} \right) + \text{sum of residues in } -U < \Im(w) < U.$$

Let ρ^* run through zeros of $\zeta(s)$ with imaginary parts between $-U$ and U . Let $U > T$. Then there is a pole at $w = 1 - \rho^*$, with residue

$$\frac{\log X}{(1-\rho^*)\zeta'(1-\rho^*)}.$$

At the other ρ^* the residues are

$$\frac{X^{\rho^*+\rho-1} - 1}{\zeta'(\rho^*)\rho^*(\rho^*+\rho-1)(\rho^*+\rho)} = O\left(\frac{1}{(|\rho^*+\rho-1|)(\rho^*+\rho)}\right) = O\left(\frac{1}{|\gamma^*+\gamma|}\right),$$

by (14.29.3), and

$$-\sum_{\substack{-U < \gamma^* < -\gamma \\ \gamma^* \neq -\gamma}} \frac{1}{|\gamma^*+\gamma|^2} \leq \sum_{\gamma^* \neq -\gamma} \frac{1}{|\gamma^*+\gamma|^2} < K_2,$$

where K_2 depends on T , if $|\gamma| < T$, but not on U .

Again

$$\begin{aligned} \int_{2-iU}^{2+i\infty} \frac{X^{w+\rho-1} - 1}{\zeta(w)w(w+\rho)(w+\rho-1)} dw &= O\left(X^{\frac{1}{2}} \int_U^{\infty} \frac{dv}{v(v+\gamma)^2}\right) \\ &= O\left(\frac{X^{\frac{1}{2}}}{U(v+\gamma)}\right) = O\left(\frac{X^{\frac{1}{2}}}{U(U-T)}\right), \end{aligned}$$

and similarly for the integral over $(2-i\infty, 2-iU)$. Also by (14.2.6) and the functional equation

$$\frac{1}{\zeta(\frac{1}{2}+it)} = O\left(\frac{|t|^{-\frac{1}{2}}}{|\zeta(\frac{1}{2}-it)|}\right) = O(|t|^{-\frac{1}{2}}).$$

Hence, since $|w+\rho| \geq \frac{1}{2}$, $|w+\rho-1| \geq \frac{1}{2}$,

$$\int_{\frac{1}{2}-iU}^{\frac{1}{2}+iU} \frac{X^{w+\rho-1} - 1}{\zeta(w)w(w+\rho)(w+\rho-1)} dw = O\left(\int_{-U}^U \frac{|v|^{-\frac{1}{2}}}{(\frac{1}{4}+v^2)^{\frac{1}{2}}} dv\right) = O(1).$$

Finally, by Theorem 14.16, we can choose a sequence of values of U such that

$$\frac{1}{\zeta(w)} = O(|w|) \quad (t = U, \frac{1}{2} \leq \sigma \leq 2).$$

By the functional equation the same result then holds for $\frac{1}{2} \leq \sigma \leq \frac{1}{2} + iU$ also. Hence

$$\int_{\frac{1}{2}+iU}^{2+iU} \frac{X^{w+\rho-1}-1}{\zeta(w)(w+\rho)(w+\rho-1)} dw = O\left(\frac{X^{\frac{1}{2}}}{U^{1-\epsilon}(U+\gamma)^2}\right) = O\left(\frac{X^{\frac{1}{2}}}{U^{1-\epsilon}(U-T)^2}\right),$$

and similarly for the integral over $(2-iU, \frac{1}{2}-iU)$. Making $U \rightarrow \infty$, it follows that

$$\int_1^X M(x)x^{\rho-2}\left(1-\frac{x}{X}\right)dx = \frac{\log X}{(1-\rho)\zeta'(1-\rho)} + R,$$

where $|R| < K_3 = K_3(T)$ if $|\gamma| < T$.

Hence we obtain

$$0 \leq A \log X + \log X \sum_{|\gamma| < T} \frac{1}{|\rho \zeta'(\rho)|^2} - 2 \log X \sum_{|\gamma| < T} \frac{1}{|\rho \zeta'(\rho)|^2} + A \log^{\frac{1}{2}} X + K_4(T),$$

$$\sum_{|\gamma| < T} \frac{1}{|\rho \zeta'(\rho)|^2} \leq A + \frac{A}{\log^{\frac{1}{2}} X} + \frac{K_4(T)}{\log X}.$$

Making $X \rightarrow \infty$,

$$\sum_{|\gamma| < T} \frac{1}{|\rho \zeta'(\rho)|^2} \leq A.$$

Since the right-hand side is now independent of T , the result follows.

In particular $\frac{1}{\zeta'(\rho)} = o(|\rho|)$.

14.30. If (14.29.1) is true,†

$$\frac{1}{\zeta'(\frac{1}{2}+it)} = O\left(\exp\left(\frac{A \log^2 t}{\log \log t}\right)\right).$$

Suppose that the interval $(t-t^3, t+t^3)$ contains γ , the ordinate of a zero. By differentiating (2.1.4) twice,

$$\zeta''(\frac{1}{2}+it) = O(t).$$

Using this and (14.29.3), we obtain

$$\begin{aligned} |\zeta'(\frac{1}{2}+it)| &= \left| \zeta'(\frac{1}{2}+i\gamma) + \int_{\frac{1}{2}+i\gamma}^{\frac{1}{2}+it} \zeta''(s) ds \right| \\ &> \frac{A}{\gamma} - At|t-\gamma| \\ &> \frac{A}{t} - \frac{A}{t^2} > \frac{A}{t}. \end{aligned}$$

† Cramér and Landau (1).

Suppose on the contrary that $(t-t^3, t+t^3)$ is free from ordinates of zeros. Theorem 14.15 gives

$$\log |\zeta(\frac{1}{2}+it)| = \sum_{|t-\gamma| \leq \frac{1}{2} \log \log t} \log |t-\gamma| + O\left(\frac{\log t \log \log \log t}{\log \log t}\right).$$

There are $O(\log t / \log \log t)$ terms in the sum, each being now $O(\log t)$. Hence

$$\log |\zeta(\frac{1}{2}+it)| = O\left(\frac{\log^2 t}{\log \log t}\right), \quad \frac{1}{\zeta(\frac{1}{2}+it)} = O\left(\exp\left(\frac{A \log^2 t}{\log \log t}\right)\right).$$

Now $\pi^{-\frac{1}{2}} \Gamma(\frac{1}{2}+s) \zeta(s)$ is real on $\sigma = \frac{1}{2}$. Hence

$$-\frac{1}{2} \log \pi + \frac{1}{2} \frac{\Gamma'(\frac{1}{2}+s)}{\Gamma(\frac{1}{2}+s)} + \frac{\zeta'(s)}{\zeta(s)}$$

is purely imaginary on $\sigma = \frac{1}{2}$. Hence, on $\sigma = \frac{1}{2}$,

$$\left| \frac{\zeta'(s)}{\zeta(s)} \right| \geq -\Re \frac{\zeta'(s)}{\zeta(s)} = -\frac{1}{2} \log \pi + \frac{1}{2} \Re \frac{\Gamma'(\frac{1}{2}+s)}{\Gamma(\frac{1}{2}+s)} = \frac{1}{2} \log t + O(1) \rightarrow \infty.$$

Hence (without any hypothesis)

$$|\zeta'(\frac{1}{2}+it)| \geq |\zeta(\frac{1}{2}+it)| \quad (t > t_0).$$

This proves the theorem.

14.31. Let $\frac{1}{2}+i\gamma$, $\frac{1}{2}+i\gamma'$ be consecutive complex zeros of $\zeta(s)$. If (14.29.1) is true

$$\gamma' - \gamma > \frac{A}{\gamma} \exp\left(-A \frac{\log \gamma}{\log \log \gamma}\right).$$

We have

$$0 = \int_{\gamma}^{\gamma'} \zeta'(\frac{1}{2}+it) dt = (\gamma' - \gamma) \zeta'(\frac{1}{2}+i\gamma) + \int_{\gamma}^{\gamma'} (\gamma' - t) \zeta''(\frac{1}{2}+it) dt.$$

Hence by (14.29.3)

$$\begin{aligned} \frac{\gamma' - \gamma}{\gamma} &< A \left| \int_{\gamma}^{\gamma'} (\gamma' - t) \zeta''(\frac{1}{2}+it) dt \right| \\ &< A \max_{\gamma < t < \gamma'} |\zeta''(\frac{1}{2}+it)| \int_{\gamma}^{\gamma'} (\gamma' - t) dt = A(\gamma' - \gamma)^2 \max_{\gamma < t < \gamma'} |\zeta''(\frac{1}{2}+it)|. \end{aligned}$$

Now

$$\begin{aligned} \zeta''(\frac{1}{2}+it) &= \frac{1}{\pi i} \int_0^{2\pi} \frac{\zeta(\frac{1}{2}+it+re^{i\theta})}{(re^{i\theta})^3} ire^{i\theta} d\theta = O\left(\frac{1}{r^2} \int_0^{2\pi} |\zeta(\frac{1}{2}+it+re^{i\theta})| d\theta\right) \\ &= O\left(\frac{1}{r^2} \exp\left(A \frac{\log t}{\log \log t}\right) (1+t^{A\epsilon})\right) \end{aligned}$$

by (14.14.1) and the functional equation. Taking $r = 1/\log t$,

$$\zeta^r(\tfrac{1}{2} + it) = O\left(\exp\left(A \frac{\log t}{\log \log t}\right)\right)$$

and the result follows.

14.32. Necessary and sufficient conditions for the Riemann hypothesis. Two such conditions have been given in § 14.25. Other similar conditions occur in the prime-number problem.†

A different kind of condition was stated by M. Riesz.‡ Let

$$F(x) = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} x^k}{(k-1)! \zeta(2k)}. \quad (14.32.1)$$

Then a simple application of the calculus of residues gives

$$F(x) = \frac{i}{2} \int_{a-i\infty}^{a+i\infty} \frac{x^s}{\Gamma(s) \zeta(2s) \sin \pi s} ds = \frac{i}{2\pi} \int_{a-i\infty}^{a+i\infty} \frac{\Gamma(1-s)}{\zeta(2s)} x^s ds,$$

where $\frac{1}{2} < a < 1$. Taking a just greater than $\frac{1}{2}$, it clearly follows that

$$F(x) = O(x^{\frac{1}{2}+\epsilon}).$$

On the Riemann hypothesis we could move the line of integration to $a = \frac{1}{2} + \epsilon$ (using (14.2.5)) and obtain similarly

$$F(x) = O(x^{\frac{1}{2}+\epsilon}). \quad (14.32.2)$$

Conversely, by Mellin's inversion formula,

$$\frac{\Gamma(1-s)}{\zeta(2s)} = - \int_0^{\infty} F(x) x^{-1-s} ds.$$

If (14.32.2) holds, the integral converges uniformly for $\sigma \geq \sigma_0 > \frac{1}{2}$; the analytic function represented is therefore regular for $\sigma > \frac{1}{2}$, and the truth of the Riemann hypothesis follows. Hence (14.32.2) is a necessary and sufficient condition for the Riemann hypothesis.

A similar condition stated by Hardy and Littlewood§ is

$$\sum_{k=1}^{\infty} \frac{(-x)^k}{k! \zeta(2k+1)} = O(x^{-\frac{1}{2}}). \quad (14.32.3)$$

These conditions have a superficial attractiveness since they depend explicitly only on values taken by $\zeta(s)$ at points in $\sigma > 1$; but actually no use has ever been made of them.

† Landau, *Vorlesungen*, ii. 108-56.

‡ M. Riesz (1).

§ Hardy and Littlewood (2).

Conditions for the Riemann hypothesis also occur in the theory of Farey series. Let the fractions h/k with $0 < h \leq k$, $(h, k) = 1$, $k \leq N$, arranged in order of magnitude, be denoted by r_ν ($\nu = 1, 2, \dots$, $\Phi(N)$), where $\Phi(N) = \phi(1) + \dots + \phi(N)$. Let

$$\delta_\nu = r_\nu - \nu/\Phi(N)$$

be the distance between r_ν and the corresponding fraction obtained by dividing up the interval $(0, 1)$ into $\Phi(N)$ equal parts. Then a necessary and sufficient condition for the Riemann hypothesis is†

$$\sum_{\nu=1}^{\Phi(N)} \delta_\nu^2 = O\left(\frac{1}{N^{1-\epsilon}}\right). \quad (14.32.4)$$

An alternative necessary and sufficient condition is§

$$\sum_{\nu=1}^{\Phi(N)} |\delta_\nu| = O(N^{\frac{1}{2}+\epsilon}). \quad (14.32.5)$$

Details are given in Landau's *Vorlesungen*, ii. 167-77.

Still another condition|| can be expressed in terms of the formulae of § 10.1. If $\Xi(t)$ and $\Phi(u)$ are related by (10.1.3), a necessary and sufficient condition that all the zeros of $\Xi(t)$ should be real is that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(x) \Phi(\beta) e^{i(\alpha+\beta)x} e^{(\alpha-\beta)u} (\alpha-\beta)^2 d\alpha d\beta \geq 0 \quad (14.32.6)$$

for all real values of x and y . But no method has been suggested of showing whether such criteria are satisfied or not.

A sufficient condition†† for the Riemann hypothesis is that the partial sums $\sum_{\nu=1}^n \nu^{-s}$ of the series for $\zeta(s)$ should have no zeros in $\sigma > 1$.

NOTES FOR CHAPTER 14

14.33. The argument of § 14.5 may be extended to the strip $\frac{1}{2} \leq \sigma_0 \leq \sigma < \frac{3}{2}$, giving

$$\frac{\zeta'(s)}{\zeta(s)} = O\left(\frac{\delta^{\sigma-1}-1}{1-\delta}\right) + O(\delta^{\sigma-\frac{1}{2}} \log t).$$

The choice $\delta = (\log t)^{-2}$ then yields

$$\frac{\zeta'(s)}{\zeta(s)} \ll \frac{(\log t)^{2-2\sigma}-1}{1-\sigma}$$

† Franel (1).

§ Landau (16).

|| See Pólya (3), § 7.

†† Turán (3).

uniformly for $\sigma_0 \leq \sigma \leq \frac{3}{8}$ and $t \geq 2$, and hence

$$\log \zeta(s) \ll \begin{cases} \log \frac{1}{\sigma-1} & \text{if } 1 + \frac{1}{\log \log t} \leq \sigma \leq \frac{3}{8}, \\ \frac{(\log t)^{2-2\sigma} - 1}{(1-\sigma) \log \log t} + \log \log \log t & \text{if } \sigma_0 \leq \sigma \leq 1 + \frac{1}{\log \log t}. \end{cases}$$

These results, together with those of § 14.14 are the sharpest conditional order-estimates available at present.

14.34. The Ω -result given by Theorem 14.12(A) has been sharpened by Montgomery [3], to give

$$S(t) = \Omega_{\pm} \left(\frac{(\log t)^{\frac{1}{2}}}{(\log \log t)^{\frac{1}{2}}} \right)$$

on the Riemann hypothesis. A minor modification of his method also yields

$$S_1(t) = \Omega_{\pm} \left(\frac{(\log t)^{\frac{1}{2}}}{(\log \log t)^{\frac{3}{2}}} \right).$$

It may be conjectured that these are best possible.

Mueller [2] has shown, on the Riemann hypothesis, that if c is a suitable constant, then $S(t)$ changes sign in any interval $[T, T + c \log \log T]$.

Further results and conjectures on the vertical distribution of the zeros are given by Montgomery [2], who investigated the pair correlation function

$$F(\alpha, T) = \frac{1}{N(T)} \sum_{0 < \gamma, \gamma' \leq T} T^{\alpha(\gamma-\gamma')} w(\gamma-\gamma'),$$

where $w(u) = 4/(4+u^2)$. This is a real-valued, even, non-negative function of α , and satisfies

$$F(\alpha, T) = \alpha + T^{-2\alpha} \log T + O\left(\frac{1}{\log T}\right) + O(\alpha T^{\alpha-1}) + O(T^{-\frac{1}{2}\alpha}) \quad (14.34.1)$$

for $\alpha \geq 0$, whence $F(\alpha, T) \rightarrow \alpha$ as $T \rightarrow \infty$, uniformly for $0 < \delta \leq \alpha \leq 1 - \delta$. Montgomery conjectured that in general

$$F(\alpha, T) \rightarrow \min(\alpha, 1) \quad (14.34.2)$$

uniformly for $0 \leq \delta \leq \alpha \leq A$. This is related to a number of conjectures on the distribution of prime numbers. (See Gallagher and Mueller [1], Heath-Brown [10], and joint work of Goldston and Montgomery in the course of publication.) From (14.34.2) one may deduce that

$$\begin{aligned} & \#\left\{ \gamma, \gamma' \in [0, T] : \frac{2\pi\alpha}{\log T} \leq \gamma - \gamma' \leq \frac{2\pi\beta}{\log T} \right\} \\ & \sim N(T) \left\{ \delta(\alpha, \beta) + \int_{\alpha}^{\beta} 1 - \left(\frac{\sin \pi u}{u} \right)^2 du \right\} \end{aligned}$$

for fixed α, β , as $T \rightarrow \infty$. Here $\delta(\alpha, \beta) = 1$ or 0 according as $\alpha \leq 0 \leq \beta$ or not.

Using (14.34.1), Montgomery showed that

$$\sum_{0 < \gamma \leq T} m(\rho)^2 \leq \left\{ \frac{3}{2} + o(1) \right\} N(T),$$

where $m(\rho)$ is the multiplicity of ρ , and Σ' counts zeros without regard to multiplicity. One may deduce, in the notation of § 10.29, that

$$N^{(1)}(T) \geq \left\{ \frac{3}{2} + o(1) \right\} N(T), \quad (14.34.3)$$

on the Riemann hypothesis. The conjecture (14.34.2) would indeed yield $N^{(1)}(T) \sim N(T)$, i.e. 'almost all' the zeros would be simple. Montgomery also used (14.34.1) to show that

$$\liminf_{n \rightarrow \infty} \frac{\gamma_{n+1} - \gamma_n}{(2\pi/\log \gamma_n)} \leq 0.68; \quad (14.34.4)$$

here $2\pi/\log \gamma_n$ is the average spacing between zeros.

By using a different method, Conrey, Ghosh, and Gonek (in work in the course of publication) have improved (14.34.3). Their starting point is the observation that

$$\left| \sum_{0 < \gamma \leq T} M\left(\frac{1}{2} + i\gamma\right) \zeta'\left(\frac{1}{2} + i\gamma\right) \right|^2 \leq N^{(1)}(T) \sum_{0 < \gamma \leq T} |M\left(\frac{1}{2} + i\gamma\right) \zeta'\left(\frac{1}{2} + i\gamma\right)|^2, \quad (14.34.5)$$

by Cauchy's inequality. The function $M(s)$ is taken to be a mollifier

$$M(s) = \sum_{n \leq y} \mu(n) P\left(\frac{\log y/n}{\log y}\right) n^{-s}, \quad y = T^{1-\varepsilon},$$

where the polynomial $P(x)$ is chosen optimally as $\frac{1}{2}x - \frac{1}{2}x^2$. One may

write the sums occurring in (14.34.5) as integrals

$$\frac{1}{2\pi i} \int_P \frac{\zeta'(s)}{\zeta(s)} M(s) \zeta'(s) ds$$

and

$$\frac{1}{2\pi i} \int_P \frac{\zeta'(s)}{\zeta(s)} M(s) M(1-s) \zeta'(s) \zeta'(1-s) ds,$$

taken around an appropriate rectangular path P . The estimation of these is long and complicated, but leads ultimately to the lower bound

$$N^{(1)}(T) \geq \left\{ \frac{3}{4} + o(1) \right\} N(T).$$

The estimate (14.34.4) has also been improved, firstly by Montgomery and Odlyzko [1], and then by Conrey, Ghosh and Gonek [1]. The latter work produces the constant 0.95172. The corresponding lower bound

$$\limsup_{n \rightarrow \infty} \frac{\gamma_{n+1} - \gamma_n}{2\pi / \log \gamma_n} \geq \lambda > 1 \quad (14.34.6)$$

has been considered by Mueller [1], as well as in the two papers just cited. Here the best result known is that of Conrey, Ghosh, and Gonek [1], which has $\lambda = 2.337$. Indeed, further work by Conrey, Ghosh, and Gonek, which is in the course of publication at the time of writing, yields $\lambda = 2.68$ subject to the generalized Riemann hypothesis (i.e. a Riemann hypothesis for $\zeta(s)$ and all Dirichlet L -functions $L(s, \chi)$). Moreover it seems likely that this condition may be relaxed to the ordinary Riemann hypothesis with further work.

If one asks for bounds of the form (14.34.4) and (14.34.6) which are satisfied by a positive proportion of zeros (as in §9.25) then one may take constants 0.77 and 1.33 (Conrey, Ghosh, Goldston, Gonek, and Heath-Brown [1]).

14.35. It should be remarked in connection with §14.24 that Selberg (4) proved Theorem 14.24 with error term $O(T)$, while the method here yields only $O\{T(\log \log T)^{\frac{1}{2}}\}$. Moreover he obtained the error term $O\{T(\log \log T)^{k-1}\}$ for (14.24.1).

14.36. The argument of the final paragraph of §14.27 may be quantified, and then yields

$$\sum_{|t| \leq T} |\zeta'(\frac{1}{2} + it)|^{-1} \gg T,$$

uniformly for $T \gg T_0$, assuming the Riemann hypothesis and that all the zeros are simple. However a slightly better result comes from combining

the asymptotic formula

$$\sum_{0 < \gamma \leq T} |\zeta'(\frac{1}{2} + i\gamma)|^2 \sim \frac{1}{2} N(T) (\log T)^2$$

of Gonek [2] with the bound (14.34.3). Using Hölder's inequality one may then derive the estimate

$$\sum_{cT < \gamma \leq T} \frac{1}{|\zeta'(\frac{1}{2} + i\gamma)|} \gg T,$$

where Σ^* counts simple zero only, and $c > 0$ is a suitable numerical constant.

14.37. The Mertens hypothesis has been disproved by Odlyzko and te Riele [1], who showed that

$$\limsup_{x \rightarrow \infty} \frac{M(x)}{\sqrt{x}} > 1.06$$

and

$$\liminf_{x \rightarrow \infty} \frac{M(x)}{\sqrt{x}} < -1.009.$$

Their treatment is indirect, and produces no specific x for which $|M(x)| > x^{\frac{1}{2}}$. The method used is computational, and depends on solving numerically the inequalities occurring in Kronecker's theorem, so as to make the first few terms of (14.27.1) pull in the same direction. To this extent Odlyzko and te Riele follow the earlier work of Jurkat and Peyerimhoff [1], but they use a much more efficient algorithm for solving the Diophantine approximation problem.

14.38. Turán (3) conjectured that

$$\sum_{n \leq x} \frac{\lambda(n)}{n} \geq 0 \quad (14.38.1)$$

for all $x > 0$, where $\lambda(n)$ is the Liouville function, given by (1.2.11). He showed that his condition, given in §14.32, implies the above conjecture, which in turn implies the Riemann hypothesis. However Haselgrove [2] proved that (14.38.1) is false in general, thereby showing that Turán's condition does not hold. Later Spira [1] found by calculation that

$$\sum_{n=1}^{19} n^{-s}$$

has a zero in the region $\sigma > 1$.

XV

CALCULATIONS RELATING TO THE ZEROS

15.1. It is possible to verify by means of calculation that all the complex zeros of $\zeta(s)$ up to a certain point lie exactly (not merely approximately) on the critical line. As a simple example we shall find roughly the position of the first complex zero in the upper half-plane, and show that it lies on the critical line.

We consider the function $Z(t) = e^{i\theta} \zeta(\frac{1}{2} + it)$ defined in § 4.17. This is real for real values of t , so that, if $Z(t_1)$ and $Z(t_2)$ have opposite signs, $Z(t)$ vanishes between t_1 and t_2 , and so $\zeta(s)$ has a zero on the critical line between $\frac{1}{2} + it_1$ and $\frac{1}{2} + it_2$.

It follows from (2.2.1) that $\xi(\frac{1}{2}) < 0$, then from (2.1.12) that $\xi(\frac{1}{2}) > 0$, i.e. that $\Xi(0) > 0$; and then from (4.17.3) that $Z(0) < 0$.

We shall next consider the value $t = 6\pi$. Now the argument of § 4.14 shows that, if x is half an odd integer,

$$\left| \zeta(s) - \sum_{n < x} \frac{1}{n^s} \right| \leq \frac{x^{1-\sigma}}{|1-s|} + \frac{2x^{-\sigma}}{2\pi - |t|/x}. \quad (15.1.1)$$

Hence, taking $t > 0$,

$$\left| Z(t) - \sum_{n < x} \frac{\cos(t \log n - \theta)}{n^{\frac{1}{2}}} \right| \leq \frac{x^{\frac{1}{2}}}{t} + \frac{2x^{\frac{1}{2}}}{2\pi x - t}. \quad (15.1.2)$$

For $x = \frac{9}{2}$, $t = 6\pi$, the right-hand side is about 0.6.

We next require an approximation to θ . We have

$$e^{-2i\theta} = \chi(\frac{1}{2} + it) = \pi^{it} \frac{\Gamma(\frac{1}{2} - \frac{1}{2}it)}{\Gamma(\frac{1}{2} + \frac{1}{2}it)},$$

so that

$$\begin{aligned} \theta &= -\frac{1}{2}t \log \pi + \text{Ilog} \Gamma(\frac{1}{2} + \frac{1}{2}it) \\ &= \frac{1}{2}t \log \frac{t}{2\pi} - \frac{1}{2}t - \frac{1}{2}\pi + O\left(\frac{1}{t}\right). \end{aligned}$$

It may be verified that the term $O(1/t)$ is negligible in the calculations.

Writing $\theta = 2\pi K$, and using the values

$$\log 2 = 0.6931, \quad \log 3 = 1.0986,$$

it is found that

$$K = 0.1166, \quad 3 \log 3 - K = 3.179$$

$$3 \log 2 - K = 1.963, \quad 3 \log 4 - K = 4.042,$$

approximately. Hence the cosines in (15.1.2) are all positive, and $\cos 2\pi K = 0.74\dots$. Hence $Z(6\pi) > 0$.

There is therefore one zero at least on the critical line between $t = 0$ and $t = 6\pi$.

Again, the formulae of § 9.3 give

$$N(T) = 1 + 2K + \frac{1}{\pi} \Delta \arg \zeta(s),$$

where Δ denotes variation along $(2, 2 + iT, \frac{1}{2} + iT)$. Now $R \zeta(s) > 0$ on $\sigma = 2$, and an argument similar to that already used, but depending on (15.1.1), shows that $R \zeta(s) > 0$ on $(2 + iT, \frac{1}{2} + iT)$, if $T = 6\pi$. Hence $|\Delta \arg \zeta(s)| < \frac{1}{2}\pi$, and

$$N(6\pi) < \frac{3}{2} + 2K < 2.$$

Hence there is at most one complex zero with imaginary part less than 6π , and so in fact just one, namely the one on the critical line.

15.2. It is plain that the above process can be continued as long as the appropriate changes of sign of the function $Z(t)$ occur. Defining $K = K(t)$, as before, let t_ν be such that

$$K(t_\nu) = \frac{1}{2}\nu - 1 \quad (\nu = 1, 2, \dots). \quad (15.2.1)$$

Then (15.1.2) gives

$$Z(t_\nu) \sim (-1)^\nu \sum_{n < x} \frac{\cos(t_\nu \log n)}{n^{\frac{1}{2}}}.$$

If the sum is dominated by its first term, it is positive, and so $Z(t_\nu)$ has the sign of $(-1)^\nu$. If this is true for ν and $\nu+1$, $Z(t)$ has a zero in the interval $(t_\nu, t_{\nu+1})$.

The value $t = 6\pi$ in the above argument is a rough approximation to t_2 .

The ordinates of the first six zeros are

$$14.13, 21.02, 25.01, 30.42, 32.93, 37.58$$

to two decimal places.† Some of these have been calculated with great accuracy.

15.3. The calculations which the above process requires are very laborious if t is at all large. A much better method is to use the formula (4.17.5) arising from the approximate functional equation. Let us write $t = 2\pi u$,

$$\alpha_n = \alpha_n(u) = n^{-\frac{1}{2}} \cos 2\pi(K - u \log n),$$

and

$$h(\xi) = \frac{\cos 2\pi(\xi^2 - \xi - \frac{1}{16})}{\cos 2\pi\xi}.$$

† See the references Gram (6), Lindelöf (3), in Landau's *Handbuch*.

Then (4.17.5) gives

$$Z(2\pi u) = 2 \sum_{n=1}^m \alpha_n(u) + (-1)^{m-1} u^{-\frac{1}{2}} h(\sqrt{u} - m) + R(u),$$

where $m = [\sqrt{u}]$, and $R(u) = O(u^{-\frac{1}{2}})$. The $\alpha_n(u)$ can be found, for given values of u , from a table of the function $\cos 2\pi x$. In the interval $0 \leq \xi \leq \frac{1}{2}$, $h(\xi)$ decreases steadily from 0.92388 to 0.38268, and $h(1-\xi) = h(\xi)$.

For the purpose of calculation we require a numerical upper bound for $R(u)$. A rather complicated formula of this kind is obtained in Titchmarsh (17), Theorem 2. For values of u which are not too small it can be much simplified, and in fact it is easy to deduce that

$$|R(u)| < \frac{3}{2u^{\frac{1}{2}}} \quad (u > 125).$$

This inequality is sufficient for most purposes.

Occasionally, when $Z(2\pi u)$ is too small, a second term of the Riemann-Siegel asymptotic formula has to be used.

The values of u for which the calculations are performed are the solutions of (15.2.1), since they make α_1 alternately 1 and -1 . In the calculations described in Titchmarsh (17), I began with

$$u = 1.6, \quad K = -0.04865$$

and went as far as

$$u = 62.785, \quad K = 98.5010.$$

The values of u were obtained in succession, and are rather rough approximations to the u_n , so that the K 's are not quite integers or integers and a half.

It was shown in this way that the first 198 zeros of $\zeta(s)$ above the real axis all lie on the line $\sigma = \frac{1}{2}$.

The calculations were carried a great deal farther by Dr. Comrie.† Proceeding on the same lines, it was shown that the first 1,041 zeros of $\zeta(s)$ above the real axis all lie on the critical line, in the interval $0 < t < 1,468$.

One interesting point which emerges from these calculations is that $Z(t_n)$ does not always have the same sign as $(-1)^n$. A considerable number of exceptional cases were found; but in each of these cases there is a neighbouring point t_0 such that $Z(t_0)$ has the sign of $(-1)^n$, and the succession of changes of sign of $Z(t)$ is therefore not interrupted.

15.4. As far as they go, these calculations are all in favour of the truth of the Riemann hypothesis. Nevertheless, it may be that they do

† See Titchmarsh (18).

not go far enough to reveal the real state of affairs. At the end of the table constructed by Dr. Comrie there are only fifteen terms in the series for $Z(t)$, and this is a very small number when we are dealing with oscillating series of this kind. Indeed there is one feature of the table which may suggest a change in its character farther on. In the main, the result is dominated by the first term α_1 , and later terms more or less cancel out. Occasionally (e.g. at $K = 435$) all, or nearly all, the numbers α_n have the same sign, and $Z(t)$ has a large maximum or minimum. As we pass from this to neighbouring values of t , the first few α_n undergo violent changes, while the later ones vary comparatively slowly. The term α_n appears when $u = n^2$, and here

$$\begin{aligned} \cos 2\pi(K - u \log n) &= \cos \pi(u \log(u/n^2) - u - \frac{1}{2} + \dots) \\ &= \cos \pi(n^2 + \frac{1}{2} + \dots) = (-1)^n \cos \frac{1}{2}\pi + \dots, \end{aligned}$$

and

$$\frac{d}{du}(K - u \log n) = \frac{1}{2} \log u - \log n - \frac{1}{192\pi^2 u^2} + \dots \sim -\frac{1}{192\pi^2 u^2}.$$

At its first appearance in the table α_n will therefore be approximately $(-1)^n n^{-\frac{1}{2}} \cos \frac{1}{2}\pi$, and it will vary slowly for some time after its appearance.

It is conceivable that if t , and so the number of terms, were large enough, there might be places where the smaller slowly varying terms would combine to overpower the few quickly varying ones, and so prevent the graph of $Z(t)$ from crossing the zero line between successive maxima. There are too few terms in the table already constructed to test this possibility.

There are, of course, relations between the numbers α_n which destroy any too simple argument of this kind. If the Riemann hypothesis is true, there must be some relation, at present hidden, which prevents the suggested possibility from ever occurring at all.

No doubt the whole matter will soon be put to the test of modern methods of calculation. Naturally the Riemann hypothesis cannot be proved by calculation, but, if it is false, it could be disproved by the discovery of exceptions in this way.

NOTES FOR CHAPTER 15

15.5. A number of workers have checked the Riemann hypothesis over increasingly large ranges. At the time of writing the most extensive calculation is that of van de Lune and te Riele (as reported in Odlyzko and te Riele [1]), who have found that the first 1.5×10^9 non-trivial zeros are simple and lie on the critical line.

ORIGINAL PAPERS

[This list includes that given in my Cambridge tract; it does not include papers referred to in Landau's Handbuch der Lehre von der Verteilung der Primzahlen, 1909.]

ABBREVIATIONS

A.M.	Acta Mathematica.
C.R.	Comptes rendus de l'Académie des sciences (Paris).
J.L.M.S.	Journal of the London Mathematical Society.
J.M.	Journal für die reine und angewandte Mathematik.
M.A.	Mathematische Annalen.
M.Z.	Mathematische Zeitschrift.
P.C.P.S.	Proceedings of the Cambridge Philosophical Society.
P.L.M.S.	Proceedings of the London Mathematical Society.
Q.J.O.	Quarterly Journal of Mathematics (Oxford Series).

ANANDA-RAU, K.

- (1) The infinite product for $(s-1)\zeta(s)$, *M.Z.* 20 (1924), 156-64.

ARWIN, A.

- (1) A functional equation from the theory of the Riemann $\zeta(s)$ -function, *Annals of Math.* (2), 24 (1923), 359-66.

ATKINSON, F. V.

- (1) The mean value of the zeta-function on the critical line, *Q.J.O.* 10 (1939) 123-8.
- (2) The mean value of the zeta-function on the critical line, *P.L.M.S.* (2), 47 (1941), 174-200.
- (3) A divisor problem, *Q.J.O.* 12 (1941), 193-200.
- (4) A mean value property of the Riemann zeta-function, *J.L.M.S.* 23 (1948), 123-35.
- (5) The Abel summation of certain Dirichlet series, *Q.J.O.* 19 (1948), 59-64.
- (6) The Riemann zeta-function, *Duke Math. J.* 17 (1950), 63-8.

BABINI, J.

- (1) Über einige Eigenschaften der Riemannschen $\zeta(z)$ -Funktion, *An. Soc. Cl. Argent.* 118 (1934), 209-15.

BACKLUND, R.

- (1) Einige numerische Rechnungen, die Nullpunkte der Riemannschen ζ -Funktion betreffend, *Öfversigt Finska Vetensk. Soc. (A)*, 54 (1911-12), No. 3.
- (2) Sur les zéros de la fonction $\zeta(s)$ de Riemann, *C.R.* 158 (1914), 1979-81.
- (3) Über die Nullstellen der Riemannschen Zetafunktion, *A.M.* 41 (1918), 345-75.
- (4) Über die Beziehung zwischen Anwachsen und Nullstellen der Zetafunktion, *Öfversigt Finska Vetensk. Soc.* 61 (1918-19), No. 9.

BEAUFAIN, J.

- (1) Sur la fonction $\zeta(s, w)$ et la fonction $\zeta(s)$ de Riemann, *Acad. Royale de Belgique* (2), 3 (1909), No. 1.

BELLMAN, R.

- (1) The Dirichlet divisor problem, *Duke Math. J.* 14 (1947), 411-17.
- (2) An analog of an identity due to Wilton, *Duke Math. J.* 16 (1949), 539-45.
- (3) Wigert's approximate functional equation and the Riemann zeta-function, *Duke Math. J.* 16 (1949), 547-52.

BOHR, H.

- (1) En Sætning om ζ -Funktioner, *Nyt. Tidss. for Math.* (B), 21 (1910), 61-6.
- (2) Über das Verhalten von $\zeta(s)$ in der Halbebene $\sigma > 1$, *Göttinger Nachrichten* (1911), 400-28.
- (3) Sur l'existence de valeurs arbitrairement petites de la fonction $\zeta(s) = \zeta(\sigma+it)$ de Riemann pour $\sigma > 1$, *Öfversigt Vidensk. Selsk. København* (1911), 201-8.
- (4) Sur la fonction $\zeta(s)$ dans le demi-plan $\sigma > 1$, *C.R.* 154 (1912), 1078-81.
- (5) Über die Funktion $\zeta'(s)/\zeta(s)$, *J.M.* 141 (1912), 217-34.
- (6) En nyt Bevis for, at den Riemann'ske Zetafunktion $\zeta(s) = \zeta(\sigma+it)$ har uendelig mange Nulpunkter indenfor Parallel-strimlerne $0 \leq t \leq 1$, *Nyt. Tidss. for Math.* (B) 23 (1912), 81-5.
- (7) Om de Vaerdier, den Riemann'ske Funktion $\zeta(\sigma+it)$ antager i Halvplanen $\sigma > 1$, 2. *Skand. Math. Kongr.* (1912), 113-21.
- (8) Note sur la fonction zeta de Riemann $\zeta(s) = \zeta(\sigma+it)$ sur la droite $\sigma = 1$, *Öfversigt Vidensk. Selsk. København* (1913), 3-11.
- (9) Lösung des absoluten Konvergenzproblems einer allgemeinen Klasse Dirichletscher Reihen, *A.M.* 36 (1913), 197-240.
- (10) Sur la fonction $\zeta(s)$ de Riemann, *C.R.* 158 (1914), 1986-8.
- (11) Zur Theorie der Riemannschen Zetafunktion im kritischen Streifen, *A.M.* 40 (1915), 67-100.
- (12) Die Riemannsche Zetafunktion, *Deutsche Math. Ver.* 24 (1915), 1-17.
- (13) Über eine quasi-periodische Eigenschaft Dirichletscher Reihen mit Anwendung auf die Dirichletschen L -Funktionen, *M.A.* 85 (1922), 115-22.
- (14) Über diophantische Approximationen und ihre Anwendungen auf Dirichletsche Reihen, besonders auf die Riemannsche Zetafunktion, 5. *Skand. Math. Kongr.* (1923), 131-54.
- (15) Another proof of Kronecker's theorem, *P.L.M.S.* (2), 21 (1922), 315-16.
- (16) Again the Kronecker theorem, *J.L.M.S.* 9 (1934), 5-6.

BOHR, H., and COUBRANT, R.

- (1) Neue Anwendungen der Theorie der Diophantischen Approximationen auf die Riemannsche Zetafunktion, *J.M.* 144 (1914), 249-74.

BOHR, H., and JESSEN, B.

- (1), (2) Über die Wertverteilung der Riemannschen Zetafunktion, *A.M.* 54 (1930), 1-35 and *ibid.* 53 (1932), 1-55.
- (3) One more proof of Kronecker's theorem, *J.L.M.S.* 7 (1932), 274-5.
- (4) Mean-value theorems for the Riemann zeta-function, *Q.J.O.* 5 (1934), 43-7.
- (5) On the distribution of the values of the Riemann zeta-function, *Amer. J. Math.* 58 (1936), 35-44.

BOHR, H., and LANDAU, E.

- (1) Über das Verhalten von $\zeta(s)$ und $\zeta'(s)$ in der Nähe der Geraden $\sigma = 1$, *Göttinger Nachrichten* (1910), 303-30.
- (2) Über die Zetafunktion, *Rend. di Palermo*, 32 (1911), 278-85.
- (3) Beiträge zur Theorie der Riemannschen Zetafunktion, *M.A.* 74 (1913), 3-30.
- (4) Ein Satz über Dirichletsche Reihen mit Anwendung auf die ζ -Funktion und die L -Funktionen, *Rend. di Palermo*, 37 (1914), 269-72.
- (5) Sur les zéros de la fonction $\zeta(s)$ de Riemann, *C.R.* 158 (1914), 106-10.
- (6) Über das Verhalten von $1/\zeta(s)$ auf der Geraden $\sigma = 1$, *Göttinger Nachrichten* (1923), 71-80.
- (7) Nachtrag zu unseren Abhandlungen aus den Jahrgängen 1910 und 1923, *Göttinger Nachrichten* (1924), 168-72.

BOHR, H., LANDAU, E., and LITTLEWOOD, J. E.

- (1) Sur la fonction $\zeta(s)$ dans le voisinage de la droite $\sigma = \frac{1}{2}$, *Bull. Acad. Belgique*, 15 (1913), 1144-75.

BORCHSENUS, V., and JESSEN, E.

- (1) Mean motions and values of the Riemann zeta-function, *A.M.* 80 (1948), 97-166.

BOWKAMP, C. J.

- (1) Über die Riemannsche Zetafunktion für positive, gerade Werte des Argumentes, *Nieuw Arch. Wisk.* 19 (1936), 50-8.

BRIKA, M.

- (1) Über eine Gestalt der Riemannschen Reihe $\zeta(s)$ für $s =$ gerade ganze Zahl, *Bull. Soc. Math. Grèce*, 14 (1933), 36-8.

BRUN, V.

- (1) On the function $[x]$, *P.C.P.S.* 20 (1920), 299-303.
(2) Deux transformations élémentaires de la fonction zêta de Riemann, *Revista Ci. Lima*, 41 (1939), 517-25.

BURRAU, C.

- (1) Numerische Lösung der Gleichung $\frac{2-p \log 2}{1-2^{-2p}} = \sum_{n=2}^{\infty} \frac{p_n^{-p} \log p_n}{1-p_n^{-2p}}$ wo p_n die Reihe der Primzahlen von 3 an durchläuft, *J.M.* 142 (1912), 51-3.

CARLSON, F.

- (1) Über die Nullstellen der Dirichletschen Reihen und der Riemannschen ζ -Funktion, *Arkiv för Mat. Astr. och Fysik*, 15 (1920), No. 20.
(2), (3) Contributions à la théorie des séries de Dirichlet, *Arkiv för Mat. Astr. och Fysik*, 16 (1922), No. 18, and 19 (1926), No. 25.

CHOWLA, S. D.

- (1) On some identities involving zeta-functions, *Journal Indian Math. Soc.* 17 (1928), 153-63.

CORPUT, J. G. VAN DER

- (1) Zahlentheoretische Abschätzungen, *M.A.* 84 (1921), 53-79.
(2) Verschärfung der Abschätzung beim Teilerproblem, *M.A.* 87 (1922), 39-65.
(3) Neue zahlentheoretische Abschätzungen, erste Mitteilung, *M.A.* 89 (1923), 215-54.
(4) Zum Teilerproblem, *M.A.* 98 (1928), 697-716.
(5) Zahlentheoretische Abschätzungen, mit Anwendung auf Gitterpunktprobleme, *M.Z.* 28 (1928), 301-10.
(6) Neue zahlentheoretische Abschätzungen, zweite Mitteilung, *M.Z.* 29 (1929), 397-426.
(7) Über Weylsche Summen, *Mathematica B* (1936-7), 1-30.

CORPUT, J. G. VAN DER, and KOKSMA, J. F.

- (1) Sur l'ordre de grandeur de la fonction $\zeta(s)$ de Riemann dans la bande critique, *Annales de Toulouse* (3), 22 (1930), 1-39.

CRAIG, C. F.

- (1) On the Riemann ζ -Function, *Bull. Amer. Math. Soc.* 29 (1923), 337-40.

CHAMÉR, H.

- (1) Über die Nullstellen der Zetafunktion, *M.Z.* 2 (1918), 237-41.
(2) Studien über die Nullstellen der Riemannschen Zetafunktion, *M.Z.* 4 (1919), 104-30.

- (3) Bemerkung zu der vorstehenden Arbeit des Herrn E. Landau, *M.Z.* 6 (1920), 155-7.

- (4) Über das Teilerproblem von Piltz, *Arkiv för Mat. Astr. och Fysik*, 16 (1922), No. 21.

- (5) Ein Mittelwertsatz in der Primzahltheorie, *M.Z.* 12 (1922), 147-53.

CHAMÉR, H., and LANDAU, E.

- (1) Über die Zetafunktion auf der Mittellinie des kritischen Streifens, *Arkiv för Mat. Astr. och Fysik*, 16 (1920), No. 28.

CRUM, M. M.

- (1) On some Dirichlet series, *J.L.M.S.* 15 (1940), 10-15.

DAVENPORT, H.

- (1) Note on mean-value theorems for the Riemann zeta-function, *J.L.M.S.* 10 (1935), 136-8.

DAVENPORT, H., and HEILBRONN, H.

- (1), (2) On the zeros of certain Dirichlet series I, II, *J.L.M.S.* 11 (1936), 181-5 and 307-12.

DENJOY, A.

- (1) L'hypothèse de Riemann sur la distribution des zéros de $\zeta(s)$, reliée à la théorie des probabilités, *C.R.* 192 (1931), 656-8.

DEURING, M.

- (1) Imaginäre quadratische Zahlkörper mit der Klassenzahl 1, *M.Z.* 37 (1933), 405-15.
(2) On Epstein's zeta-function, *Annals of Math.* (2) 38 (1937), 684-93.

ESTERMANN, T.

- (1) On certain functions represented by Dirichlet series, *P.L.M.S.* (2), 27 (1928), 435-48.
(2) On a problem of analytic continuation, *P.L.M.S.* 27 (1928), 471-82.
(3) A proof of Kronecker's theorem by induction, *J.L.M.S.* 8 (1933), 18-20.

FAVARD, J.

- (1) Sur la répartition des points où une fonction presque périodique prend une valeur donnée, *C.R.* 194 (1932), 1714-16.

FEJÉR, L.

- (1) Nombre de changements de signe d'une fonction dans un intervalle et ses moments, *C.R.* 158 (1914), 1328-31.

FEKETE, M.

- (1) Sur une limite inférieure des changements de signe d'une fonction dans un intervalle, *C.R.* 158 (1914), 1256-8.
(2) The zeros of Riemann's zeta-function on the critical line, *J.L.M.S.* 1 (1926), 15-19.

FLETT, T. M.

- (1) On the function $\sum_{n=1}^{\infty} \frac{1}{n} \sin \frac{t}{n}$, *J.L.M.S.* 25 (1950), 5-19.

- (2) On a coefficient problem of Littlewood and some trigonometrical sums, *Q.J.O.* (2) 2 (1951), 26-52.

FRANKE, J.

- (1) Les suites de Farey et le problème des nombres premiers, *Göttinger Nachrichten* (1924), 198-201.

GABRIEL, R. M.

- (1) Some results concerning the integrals of moduli of regular functions along certain curves, *J.L.M.S.* 2 (1927), 112-17.

GRAM, J. P.

- (1) Tafeln für die Riemannsche Zetafunktion, *Skriften København* (8), 9 (1925), 311-25.

GRONWALL, T. H.

- (1) Sur la fonction $\zeta(s)$ de Riemann au voisinage de $\sigma = 1$, *Rend. di Palermo*, 35 (1913), 95-102.
- (2) Über das Verhalten der Riemannschen Zeta-funktion auf der Geraden $\sigma = 1$, *Arch. der Math. u. Phys.* (3) 21 (1913), 231-8.

GROSSMAN, J.

- (1) Über die Nullstellen der Riemannschen Zeta-funktion und der Dirichlet'schen L -Funktionen, *Dissertation*, Göttingen (1913).

GUINAND, A. P.

- (1) A formula for $\zeta(s)$ in the critical strip, *J.L.M.S.* 14 (1939), 97-106.
- (2) Some Fourier transforms in prime-number theory, *Q.J.O.* 18 (1947), 53-64.
- (3) Some formulae for the Riemann zeta-function, *J.L.M.S.* 22 (1947), 14-18.
- (4) Fourier reciprocities and the Riemann zeta-function, *P.L.M.S.* 51 (1949), 401-14.

HADAMARD, J.

- (1) Une application d'une formule intégrale relative aux séries de Dirichlet, *Bull. Soc. Math. de France*, 56, ii (1927), 43-4.

HAMBURGER, H.

- (1), (2), (3) Über die Riemannsche Funktionalgleichung der ζ -Funktion, *M.Z.* 10 (1921), 240-54; 11 (1922), 224-45; 13 (1922), 283-311.
- (4) Über einige Beziehungen, die mit der Funktionalgleichung der Riemannschen ζ -Funktion äquivalent sind, *M.A.* 85 (1922), 129-40.

HARDY, G. H.

- (1) Sur les zéros de la fonction $\zeta(s)$ de Riemann, *C.R.* 158 (1914), 1012-14.
- (2) On Dirichlet's divisor problem, *P.L.M.S.* (2), 15 (1916), 1-25.
- (3) On the average order of the arithmetical functions $F(n)$ and $\Delta(n)$, *P.L.M.S.* (2), 15 (1915), 192-213.
- (4) On some definite integrals considered by Mellin, *Messenger of Math.* 49 (1919), 85-91.
- (5) Ramanujan's trigonometrical function $c_k(n)$, *P.C.P.S.* 20 (1920), 263-71.
- (6) On the integration of Fourier series, *Messenger of Math.* 51 (1922), 186-92.
- (7) A new proof of the functional equation for the zeta-function, *Mat. Tidskrift*, B (1922), 71-3.
- (8) Note on a theorem of Mertens, *J.L.M.S.* 2 (1926), 70-2.

HARDY, G. H., INGHAM, A. E., and PÓLYA, G.

- (1) Theorems concerning mean values of analytic functions, *Proc. Royal Soc. (A)*, 113 (1936), 542-69.

HARDY, G. H., and LITTLEWOOD, J. E.

- (1) Some problems of Diophantine approximation, *Internat. Congress of Math., Cambridge* (1912), 1, 223-9.
- (2) Contributions to the theory of the Riemann zeta-function and the theory of the distribution of primes, *A.M.* 41 (1918), 119-96.

- (3) The zeros of Riemann's zeta-function on the critical line, *M.Z.* 10 (1921), 283-317.
- (4) The approximate functional equation in the theory of the zeta-function, with applications to the divisor problems of Dirichlet and Piltz, *P.L.M.S.* (2), 21 (1922), 39-74.
- (5) On Lindelöf's hypothesis concerning the Riemann zeta-function, *Proc. Royal Soc. (A)*, 103 (1923), 403-12.
- (6) The approximate functional equations for $\zeta(s)$ and $\zeta'(s)$, *P.L.M.S.* (2), 29 (1928), 81-97.

HARTMAN, P.

- (1) Mean motions and almost periodic functions, *Trans. Amer. Math. Soc.* 46 (1939), 66-81.

HASELGRÖVE, C. B.

- (1) A connexion between the zeros and the mean values of $\zeta(s)$, *J.L.M.S.* 24 (1949), 215-22.

HASSE, H.

- (1) Beweis des Analogons der Riemannschen Vermutung für die Artinschen und F.K. Schmidtschen Kongruenzzetafunktionen in gewissen elliptischen Fällen, *Göttinger Nachrichten*, 42 (1933), 253-62.

HAVILAND, E. K.

- (1) On the asymptotic behavior of the Riemann ξ -function, *Amer. J. Math.* 67 (1945), 411-16.

HECKE, E.

- (1) Über die Lösungen der Riemannschen Funktionalgleichung, *M.Z.* 16 (1923), 301-7.
- (2) Über die Bestimmung Dirichletscher Reihen durch ihre Funktionalgleichung, *M.A.* 112 (1936), 664-99.
- (3) Herleitung des Euler-Produktes der Zetafunktion und einiger L -Reihen aus ihrer Funktionalgleichung, *M.A.* 119 (1944), 266-87.

HEILBRONN, H.

- (1) Über den Primzahlssatz von Herrn Hoheisel, *M.Z.* 36 (1933), 394-423.

HILLE, E.

- (1) A problem in 'Factorisatio Numerorum', *Acta Arith.* 2 (1936), 134-44.

HOHEISEL, G.

- (1) Normalfolgen und Zetafunktion, *Jahresber. Schles. Gesell.* 100 (1927), 1-7.
- (2) Eine Illustration zur Riemannschen Vermutung, *M.A.* 99 (1928), 150-61.
- (3) Über das Verhalten des reziproken Wertes der Riemannschen Zeta-Funktion, *Sitzungsber. Preuss. Akad. Wiss.* (1929), 219-23.
- (4) Nullstellenanzahl und Mittelwerte der Zetafunktion, *Sitzungsber. Preuss. Akad. Wiss.* (1930), 72-82.
- (5) Primzahlprobleme in der Analysis, *Sitzungsber. Preuss. Akad. Wiss.* (1930), 580-8.

HÖLDER, O.

- (1) Über gewisse der Möbiusschen Funktion $\mu(n)$ verwandte zahlentheoretische Funktionen, der Dirichletschen Multiplikation und eine Verallgemeinerung der Umkehrformeln, *Ber. Verh. sächs. Akad. Leipzig*, 85 (1933), 3-28.

HUA, L. K.

- (1) An improvement of Vinogradov's mean-value theorem and several applications, *Q.J.O.* 20 (1949), 48-61.

HUTCHINSON, J. I.

- (1) On the roots of the Riemann zeta-function, *Trans. Amer. Math. Soc.* 27 (1925), 49-60.

INGHAM, A. E.

- (1) Mean-value theorems in the theory of the Riemann zeta-function, *P.L.M.S.* (2), 27 (1926), 273-300.
- (2) Some asymptotic formulae in the theory of numbers, *J.L.M.S.* 2 (1927), 202-8.
- (3) Notes on Riemann's ζ -function and Dirichlet's L -functions, *J.L.M.S.* 5 (1930), 107-12.
- (4) Mean-value theorems and the Riemann zeta-function, *Q.J.O.* 4 (1933), 279-90.
- (5) On the difference between consecutive primes, *Q.J.O.* 8 (1937), 255-66.
- (6) On the estimation of $N(\alpha, T)$, *Q.J.O.* 11 (1940), 291-2.
- (7) On two conjectures in the theory of numbers, *Amer. J. Math.* 64 (1942), 313-19.

JARNÍK, V., and LANDAU, E.

- (1) Untersuchungen über einen van der Corputschen Satz, *M.Z.* 39 (1935), 745-67.

JESSEN, B.

- (1) Eine Integrationstheorie für Funktionen unendlich vieler Veränderlichen, mit Anwendung auf das Werteverteilungsproblem für fastperiodische Funktionen, insbesondere für die Riemannsche Zetafunktion, *Mat. Tidsskrift, B* (1932), 59-65.
- (2) Mouvement moyen et distribution des valeurs des fonctions presque-périodiques, *10. Skand. Math. Kongr.* (1946), 301-12.

JESSEN, B., and WINTNER, A.

- (1) Distribution functions and the Riemann zeta-function, *Trans. Amer. Math. Soc.* 38 (1935), 48-88.

KAC, M., and STEINHAUS, H.

- (1) Sur les fonctions indépendantes (N), *Studia Math.* 7 (1938), 1-15.

KAMPEN, E. R. VAN

- (1) On the addition of convex curves and the densities of certain infinite convolutions, *Amer. J. Math.* 59 (1937), 679-95.

KAMPEN, E. R. VAN, and WINTNER, A.

- (1) Convolutions of distributions on convex curves and the Riemann zeta-function, *Amer. J. Math.* 59 (1937), 175-204.

KERSHNER, R.

- (1) On the values of the Riemann ζ -function on fixed lines $\sigma > 1$, *Amer. J. Math.* 59 (1937), 167-74.

KERSHNER, R., and WINTNER, A.

- (1) On the boundary of the range of values of $\zeta(s)$, *Amer. J. Math.* 58 (1936), 421-5.
- (2) On the asymptotic distribution of $\zeta'/\zeta(s)$ in the critical strip, *Amer. J. Math.* 59 (1937), 673-8.

KIENAST, A.

- (1) Über die Dirichletschen Reihen für $\zeta^2(s)$, $L^2(s)$, *Comment. Math. Helv.* 8 (1936), 359-70.

KLOOSTERMAN, H. D.

- (1) Een integraal voor de ζ -functie van Riemann, *Christiaan Huygens Math. Tijdschrift*, 2 (1922), 172-7.

KLUYVER, J. C.

- (1) On certain series of Mr. Hardy, *Quart. J. of Math.* 50 (1924), 185-92.

KOBER, H.

- (1) Transformationen einer bestimmten Besselschen Reihe sowie von Potenzen der Riemannschen ζ -Funktion und von verwandten Funktionen, *J.M.* 173 (1935), 65-78.
- (2) Eine der Riemannschen verwandte Funktionalgleichung, *M.Z.* 39 (1935), 630-3.
- (3) Funktionen, die den Potenzen der Riemannschen Zetafunktion verwandt sind, und Potenzreihen, die über den Einheitskreis nicht fortsetzbar sind, *J.M.* 174 (1936), 206-25.
- (4) Eine Mittelwertformel der Riemannschen Zetafunktion, *Compositio Math.* 3 (1936), 174-89.

KOCH, H. VON

- (1) Contribution à la théorie des nombres premiers, *A.M.* 33 (1910), 293-320.

KOSLIAKOV, N.

- (1) Some integral representations of the square of Riemann's function $\Xi(s)$, *C.R. Acad. Sci. U.R.S.S.* 2 (1934), 401-4.
- (2) Integral for the square of Riemann's function, *C.R. Acad. Sci. U.R.S.S.* n.s. 2 (1936), 87-90.
- (3) Some formulae for the functions $\zeta(s)$ and $\zeta_r(s)$, *C.R. Acad. Sci. U.R.S.S.* (2), 25 (1939), 567-9.

KRAMASCHKE, L.

- (1) Nullstellen der Zetafunktion, *Deutsche Math.* 2 (1937), 107-10.

KUSMIN, R.

- (1) Sur les zéro de la fonction $\zeta(s)$ de Riemann, *C.R. Acad. Sci. U.R.S.S.* 2 (1934), 398-400.

LANDAU, E.

- (1) Zur Theorie der Riemannschen Zetafunktion, *Vierteljahrsschr. Naturf. Ges. Zürich.* 56 (1911), 125-48.
- (2) Über die Nullstellen der Zetafunktion, *M.A.* 71 (1911), 548-64.
- (3) Ein Satz über die ζ -Funktion, *Nyt. Tidsk.* 22 (B), (1911), 1-7.
- (4) Über einige Summen, die von den Nullstellen der Riemannschen Zetafunktion abhängen, *A.M.* 35 (1912), 271-94.
- (5) Über die Anzahl der Gitterpunkte in gewissen Bereichen, *Göttinger Nachrichten* (1912), 687-771.
- (6) Gelöste und ungelöste Probleme aus der Theorie der Primzahlverteilung und der Riemannschen Zetafunktion, *Jahresber. der Deutschen Math. Ver.* 21 (1912), 208-28.
- (7) Gelöste und ungelöste Probleme aus der Theorie der Primzahlverteilung und der Riemannschen Zetafunktion, *Proc. 5 Internat. Math. Congr.* (1913), 1, 93-108.
- (8) Über die Hardy'sche Entdeckung unendlich vieler Nullstellen der Zetafunktion mit reellem Teil $\frac{1}{2}$, *M.A.* 76 (1915), 212-43.
- (9) Über die Wigersche asymptotische Funktionalgleichung für die Lambersche Reihe, *Arch. d. Math. u. Phys.* (3), 27 (1916), 144-6.
- (10) Neuer Beweis eines Satzes von Herrn Valiron, *Jahresber. der Deutschen Math. Ver.* 29 (1920), 239.
- (11) Über die Nullstellen der Zetafunktion, *M.Z.* 6 (1920), 151-4.

- (12) Über die Nullstellen der Dirichletschen Reihen und der Riemannschen ζ -Funktion, *Arkiv för Mat. Astr. och Fysik*, 16 (1921), No. 7.
- (13) Über die Möbiussche Funktion, *Rend. di Palermo*, 48 (1924), 277-80.
- (14) Über die Wurzeln der Zetafunktion, *M.Z.* 20 (1924), 98-104.
- (15) Über die ζ -Funktion und die L -Funktionen, *M.Z.* 20 (1924), 105-25.
- (16) Bemerkung zu der vorstehenden Arbeit von Herrn Franel, *Göttinger Nachrichten* (1924), 202-6.
- (17) Über die Riemannsche Zetafunktion in der Nähe von $\sigma = 1$, *Rend. di Palermo*, 50 (1926), 423-7.
- (18) Über die Zetafunktion und die Hadamardsche Theorie der ganzen Funktionen, *M.Z.* 26 (1927), 170-5.
- (19) Über das Konvergenzgebiet einer mit der Riemannschen Zetafunktion zusammenhängenden Reihe, *M.A.* 97 (1927), 251-90.
- (20) Bemerkung zu einer Arbeit von Hrn. Hoheisel über die Zetafunktion, *Sitzungsber. Preuss. Akad. Wiss.* (1929), 271-5.
- (21) Über die Fareyreihe und die Riemannsche Vermutung, *Göttinger Nachrichten* (1932), 347-52.
- (22) Über den Wertevorrat von $\zeta(s)$ in der Halbebene $\sigma > 1$, *Göttinger Nachrichten* (1933), 81-91.

LANDAU, E., and WALFISZ, A.

- (1) Über die Nichtfortsetzbarkeit einiger durch Dirichletsche Reihen definierter Funktionen, *Rend. di Palermo* 44 (1919), 82-6.

LERCH, M.

- (1) Über die Bestimmung der Koeffizienten in der Potenzreihe für die Funktion $\zeta(s)$, *Časopis*, 43 (1914), 611-22.

LETTENMEYER, F.

- (1) Neuer Beweis des allgemeinen Kroneckerschen Approximationssatzes, *P.L.M.S.* (2) 21 (1923), 306-14.

LEVINSON, N.

- (1) On Hardy's theorem on the zeros of the zeta function, *J. Math. Phys. Mass. Inst. Tech.* 19 (1940), 159-60.

LITTLEWOOD, J. E.

- (1) Quelques conséquences de l'hypothèse que la fonction $\zeta(s)$ de Riemann n'a pas de zéros dans le demi-plan $\Re(s) > \frac{1}{2}$, *C.R.* 154 (1912), 263-6.
- (2) Researches in the theory of the Riemann ζ -function, *P.L.M.S.* (2) 20 (1922), Records xxii-xxviii.
- (3) Two notes on the Riemann zeta-function, *P.C.P.S.* 22 (1924), 234-42.
- (4) On the zeros of the Riemann zeta-function, *P.C.P.S.* 22 (1924), 295-318.
- (5) On the Riemann zeta-function, *P.L.M.S.* (2), 24 (1925), 175-201.
- (6) On the function $1/\zeta(1+it)$, *P.L.M.S.* (2), 27 (1928), 349-57.

MAIER, W.

- (1) Gitterfunktionen der Zahlentheorie, *M.A.* 113 (1936), 363-79.

MALURKAR, S. L.

- (1) On the application of Herr Mellin's integrals to some series, *Journal Indian Math. Soc.* 16 (1925), 130-8.

MATTSO, R.

- (1) Eine neue Darstellung der Riemann'schen Zetafunktion, *Arkiv för Mat. Astr. och Fysik*, 19 (1926), No. 26.

MELLIN, H.

- (1) Über die Nullstellen der Zetafunktion, *Annales Acad. Scientiarum Fennicae* (A), 10 (1917), No. 11.

MEULENBELD, B.

- (1) Een approximatieve Functionaalbetrekking van de Zetafunctie van Riemann, *Dissertation, Groningen*, 1936.

MIKOLÁS, M.

- (1) Sur l'hypothèse de Riemann, *C.R.* 228 (1949), 633-6.

MIN, S. H.

- (1) On the order of $\zeta(\frac{1}{2} + it)$, *Trans. Amer. Math. Soc.* 65 (1949), 448-72.

MIYATAKE, O.

- (1) On Riemann's ξ -function, *Tôhoku Math. Journal*, 46 (1939), 180-72.

MORDELL, L. J.

- (1) Some applications of Fourier series in the analytic theory of numbers, *P.C.P.S.* 34 (1928), 585-96.
- (2) Poisson's summation formulae and the Riemann zeta-function, *J.L.M.S.* 4 (1929), 285-91.
- (3) On the Riemann hypothesis and imaginary quadratic fields with a given class number, *J.L.M.S.* 9 (1934), 289-98.

MÜNTZ, C. H.

- (1) Beziehungen der Riemannschen ζ -Funktion zu willkürlichen reellen Funktionen, *Mat. Tidskrift*, B (1922), 39-47.

MUTATEKER, V. L.

- (1) On some formulae in the theory of the zeta-function, *Journal Indian Math. Soc.* 19 (1932), 220-4.

NEVANLINNA, F. and R.

- (1), (2) Über die Nullstellen der Riemannschen Zetafunktion, *M.Z.* 20 (1924), 253-63, and 23 (1925), 159-60.

OSTROWSKI, A.

- (1) Notiz über den Wertevorrat der Riemannschen ζ -Funktion am Rande des kritischen Streifens, *Jahresbericht Deutsch. Math. Verein.* 43 (1933), 58-64.

PALEY, R. E. A. C., and WIENER, N.

- (1) Notes on the theory and application of Fourier transforms V, *Trans. Amer. Math. Soc.* 35 (1933), 768-81.

PHILLIPS, ERIC

- (1) The zeta-function of Riemann; further developments of van der Corput's method, *Q.J.O.* 4 (1933), 208-25.
- (2) A note on the zeros of $\zeta(s)$, *Q.J.O.* 6 (1935), 137-45.

POL, B. VAN DER,

- (1) An electro-mechanical investigation of the Riemann zeta-function in the critical strip, *Bull. Amer. Math. Soc.* 53 (1947), 976-81.

PÓLYA, G.

- (1) Bemerkung über die Integraldarstellung der Riemannschen ξ -Funktion, *A.M.* 48 (1926), 305-17.
- (2) On the zeros of certain trigonometric integrals, *J.L.M.S.* 1 (1926), 98-9.
- (3) Über die algebraisch-funktiontheoretischen Untersuchungen von J. L. W. V. Jensen, *Kgl. Danske Videnskabsn. Selskab.* 7 (1927), No. 17.
- (4) Über trigonometrische Integrale mit nur reellen Nullstellen, *J.M.* 158 (1927), 6-18.

POPOV, A. I.

- (1) Several series containing primes and roots of $\zeta(s)$, *C.R. Acad. Sci. U.R.S.S.*, N.S. 41 (1943), 362-3.

POTTER, H. S. A., and TITCHMARSH, E. C.

- (1) The zeros of Epstein's zeta-functions, *P.L.M.S.* (2) 39 (1935), 372-84.

RADEMACHER, H.

- (1) Ein neuer Beweis für die Funktionalgleichung der ζ -Funktion. *M.Z.* 31 (1930), 39-44.

RAMANUJAN, S.

- (1) New expressions for Riemann's functions $\xi(s)$ and $\Xi(t)$, *Quart. J. of Math.* 46 (1915), 253-61.
 (2) Some formulae in the analytic theory of numbers, *Messenger of Math.* 45 (1915), 81-4.
 (3) On certain trigonometrical sums and their applications in the theory of numbers, *Trans. Camb. Phil. Soc.* 22 (1918), 259-76.

RAMASWAMI, V.

- (1) Notes on Riemann's ζ -function, *J.L.M.S.* 9 (1934), 165-9.

RIESZ, M.

- (1) Sur l'hypothèse de Riemann, *A.M.* 40 (1916), 185-90.

SCHNEE, W.

- (1) Die Funktionalgleichung der Zetafunktion und der Dirichletschen Reihen mit periodischen Koeffizienten, *M.Z.* 31 (1930), 378-90.

SELBERG, A.

- (1) On the zeros of Riemann's zeta-function on the critical line, *Arch. for Math. og Naturv.* B, 45 (1942), 101-14.
 (2) On the zeros of Riemann's zeta-function, *Skr. Norske Vid. Akad. Oslo* (1942), no. 10.
 (3) On the normal density of primes in small intervals, and the difference between consecutive primes, *Arch. for Math. og Naturv.* B, 47 (1943), No. 6.
 (4) On the remainder in the formula for $N(T)$, the number of zeros of $\zeta(s)$ in the strip $0 < t < T$, *Avhandlingar Norske Vid. Akad. Oslo* (1944), no. 1.
 (5) Contributions to the theory of the Riemann zeta-function, *Arch. for Math. og Naturv.* B, 48 (1946), no. 5.
 (6) The zeta-function and the Riemann hypothesis, *10. Skand. Math. Kongr.* (1946), 187-200.
 (7) An elementary proof of the prime-number theorem, *Ann. of Math.* (2) 50 (1949), 305-13.

SELBERG, S.

- (1) Bemerkung zu einer Arbeit von Viggo Brun über die Riemannsche Zetafunktion, *Norske Vid. Selekt. Forh.* 13 (1940), 17-19.

SIEGEL, C. L.

- (1) Bemerkung zu einem Satz von Hamburger über die Funktionalgleichung der Riemannschen Zetafunktion, *M.A.* 86 (1922), 276-9.
 (2) Über Riemanns Nachlass zur analytischen Zahlentheorie, *Quellen und Studien zur Geschichte der Math. Astr. und Physik, Abt. B: Studien*, 2 (1932), 45-80.
 (3) Contributions to the theory of the Dirichlet L -series and the Epstein Zeta-functions, *Annals of Math.* 44 (1943), 143-72.

SPEISER, A.

- (1) Geometrisches zur Riemannschen Zetafunktion, *M.A.* 110 (1934), 514-21.

STEEN, E. W. P.

- (1) A linear transformation connected with the Riemann zeta-function, *P.L.M.S.* (2), 41 (1936), 151-75.

STERNECK, R. VON

- (1) Neue empirische Daten über die zahlentheoretische Funktion $\sigma(n)$, *Internat. Congress of Math., Cambridge* (1912), 1, 341-3.

SZÁSZ, O.

- (1) Introduction to the theory of divergent series, Cincinnati, 1944 (*Math. Rev.* 6, 45).

TAYLOR, P. R.

- (1) On the Riemann zeta-function, *Q.J.O.* 16 (1945), 1-21.

TOHDAKOFF, N. G.

- (1) Sur les zéros de la fonction $\zeta(s)$, *C.R.* 202 (1936), 191-3.
 (2) On zeros of the function $\zeta(s)$, *C.R. Acad. Sci. U.R.S.S.* 1 (x) (1936), 201-4.
 (3) On zeros of Dirichlet's L -functions, *Mat. Sbornik* (1) 43 (1936), 591-602.
 (4) On Weyl's sums, *Mat. Sbornik* (2) 44 (1937), 17-35.
 (5) On the functions $\zeta(s)$ and $\pi(x)$, *C.R. Acad. Sci. U.R.S.S.* 21 (1938), 421-2.

THIRUVENKATACHARYA, V.

- (1) On some properties of the zeta-function, *Journal Indian Math. Soc.* 19 (1931), 92-6.

TITCHMARSH, E. C.

- (1) The mean-value of the zeta-function on the critical line, *P.L.M.S.* (2) 27 (1928), 137-60.
 (2) On the remainder in the formula for $N(T)$, the number of zeros of $\zeta(s)$ in the strip $0 < t < T$, *P.L.M.S.* (2), 27 (1928), 449-58.
 (3) A consequence of the Riemann hypothesis, *J.L.M.S.* 2 (1927), 247-54.
 (4) On an inequality satisfied by the zeta-function of Riemann, *P.L.M.S.* (2), 28 (1928), 70-80.
 (5) On the zeros of the Riemann zeta-function, *P.L.M.S.* (2) 30 (1929), 319-21.
 (6) Mean value theorems in the theory of the Riemann zeta-function, *Messenger of Math.* 58 (1929), 125-9.
 (7) The zeros of Dirichlet's L -functions, *P.L.M.S.* (2) 32 (1931), 488-500.
 (8)-(12) On van der Corput's method and the zeta-function of Riemann, *Q.J.O.* 2 (1931), 161-73; 2 (1931), 313-20; 3 (1932), 133-41; 5 (1934), 98-105; 5 (1934), 195-210.
 (13) On the Riemann zeta-function, *P.C.P.S.* 28 (1932), 273-4.
 (14) On the function $1/\zeta(1+it)$, *Q.J.O.* 4 (1933), 64-70.
 (15) On Epstein's zeta-function, *P.L.M.S.* (2) 36 (1934), 485-500.
 (16) The lattice-points in a circle *P.L.M.S.* (2) 38 (1935), 96-115.
 (17), (18) The zeros of the Riemann zeta-function, *Proc. Royal Soc. (A)*, 151 (1935), 234-55, and *ibid.* 157 (1936), 261-3.
 (19) The mean value of $|\zeta(\frac{1}{2}+it)|^4$, *Q.J.O.* 8 (1937), 107-12.
 (20) On $\zeta(s)$ and $\pi(x)$, *Q.J.O.* 9 (1938), 97-108.
 (21) The approximate functional equation for $\zeta^2(s)$, *Q.J.O.* 9 (1938), 109-14.
 (22) On divisor problems, *Q.J.O.* 9 (1938), 210-20.
 (23) A convexity theorem, *J.L.M.S.* 13 (1938), 196-7.
 (24) On the order of $\zeta(\frac{1}{2}+it)$, *Q.J.O.* 13 (1942), 11-17.
 (25) Some properties of the Riemann zeta-function, *Q.J.O.* 14 (1943), 16-26.
 (26) On the zeros of the Riemann zeta-function, *Q.J.O.* 18 (1947), 4-16.

TORELLI, G.

(1) Studio sulla funzione $\zeta(s)$ di Riemann, *Napoli Rend.* (3) 19 (1913), 212-16.

TSUJII, M.

(1) On the zeros of the Riemann zeta-function, *Proc. Imp. Acad. Tokyo*, 18 (1942), 631-4.

TURÁN, P.

(1) Über die Verteilung der Primzahlen I, *Acta Szeged*, 10 (1941), 81-104.(2) On Riemann's hypothesis, *Bull. Acad. Sci. U.R.S.S.* 11 (1947), 197-262.(3) On some approximative Dirichlet-polynomials in the theory of the zeta-function of Riemann, *Danske Vidensk. Selskab*, 24 (1948), Nr. 17.

TURNING, A. M.

(1) A method for the calculation of the zeta-function, *P.L.M.S.* (2), 48 (1943), 180-97.

UTZINGER, A. A.

(1) Die reellen Züge der Riemannschen Zetafunktion, *Dissertation, Zürich*, 1934 (see *Zentralblatt für Math.* 10, 163).

VALIRON, G.

(1) Sur les fonctions entières d'ordre nul et d'ordre fini, *Annales de Toulouse* (3), 5 (1914), 117-257.

VALLÉE POUSSIN, C. DE LA

(1), (2) Sur les zéros de $\zeta(s)$ de Riemann, *C.R.* 163 (1916), 418-21 and 471-3.

VINOGRADOV, I. M.

(1) On Weyl's sums, *Mat. Sbornik*, 42 (1935), 521-30.(2) A new method of resolving of certain general questions of the theory of numbers, *Mat. Sbornik* (1), 43 (1936), 9-19.(3) A new method of estimation of trigonometrical sums, *Mat. Sbornik* (1), 43 (1936), 175-188.(4) The method of trigonometrical sums in the theory of numbers, *Trav. Inst. Math. Stekloff* 23 (1947).

VORONÓV, G.

(1) Sur un problème du calcul des fonctions asymptotiques, *J.M.* 126 (1903), 241-82.(2) Sur une fonction transcendante et ses applications à la sommation de quelques séries, *Annales de l'École Normale* (3) 21 (1904), 207-68 and 459-534.

WALFISZ, A.

(1) Zur Abschätzung von $\zeta(\frac{1}{2}+it)$, *Göttinger Nachrichten* (1924), 155-8.(2) Über Gitterpunkte in mehrdimensionalen Ellipsoiden VIII, *Travaux de l'Institut Math. de Pbilisi*, 5 (1938), 181-96.

WALTHER, A.

(1) Über die Extrema der Riemannschen Zetafunktion bei reellem Argument, *Jahresbericht Deutsch. Math. Verein.* 34 (1925), 171-7.(2) Anschauliches zur Riemannschen Zetafunktion, *A.M.* 48 (1926), 393-400.

WANG, F. T.

(1) A remark on the mean-value theorem of Riemann's zeta-function, *Science Reports Tôhoku Imperial Univ.* (1) 25 (1936), 381-91.(2) On the mean-value theorem of Riemann's zeta-function, *Science Reports Tôhoku Imperial Univ.* (1) 25 (1936), 392-414.(3) A note on zeros of Riemann zeta-function, *Proc. Imp. Acad. Tokyo*, 12, (1937), 305-6.(4) A formula on Riemann zeta-function, *Ann. of Math.* (2) 46 (1945), 89-92.(5) A note on the Riemann zeta-function, *Bull. Amer. Math. Soc.* 52 (1946), 319-21.(6) A mean-value theorem of the Riemann zeta function, *Q.J.O.* 18 (1947), 1-3.

WATSON, G. N.

(1) Some properties of the extended zeta-function, *P.L.M.S.* (2) 12 (1913), 288-96.

WENNBERG, S.

(1) Zur Theorie der Dirichletschen Reihen, *Dissertation, Upsala*, 1920.

WEYL, H.

(1) Über die Gleichverteilung von Zahlen mod. Eins, *M.A.* 77 (1916), 313-52.(2) Zur Abschätzung von $\zeta(1+it)$, *M.Z.* 10 (1921), 88-101.

WHITTAKER, J. M.

(1) An inequality for the Riemann zeta-function, *P.L.M.S.* (2), 41 (1936), 544-52.(2) A mean-value theorem for analytic functions, *P.L.M.S.* (2), 42 (1936), 186-95.

WIGERT, S.

(1) Sur la série de Lambert et son application à la théorie des nombres, *A.M.* 41 (1916), 197-218.(2) Sur la théorie de la fonction $\zeta(s)$ de Riemann, *Arkiv for Mat. Astr. och Fysik* (1919), No. 12.(3) On a problem concerning the Riemann ζ -function, *P.C.P.S.* 21 (1921), 17-21.

WILSON, B. M.

(1) Proofs of some formulae enunciated by Ramanujan, *P.L.M.S.* (2) 21 (1922), 235-55.

WILTON, J. R.

(1) Note on the zeros of Riemann's ζ -function, *Messenger of Math.* 45 (1915), 180-3.(2) A proof of Burnside's formula for $\log \Gamma(x+1)$ and certain allied properties of Riemann's ζ -function, *Messenger of Math.* 52 (1922), 90-3.(3) A note on the coefficients in the expansion of $\zeta(s, x)$ in powers of $s-1$, *Quart. J. of Math.* 50 (1927), 329-32.(4) An approximate functional equation for the product of two ζ -functions, *P.L.M.S.* (2), 31 (1930), 11-17.(5) The mean value of the zeta-function on the critical line, *J.L.M.S.* 5 (1930), 28-32.

WINTNER, A.

(1) A note on the distribution of the zeros of the zeta-function, *Amer. J. Math.* 57 (1935), 101-2.(2) A note on the Riemann ζ -function, *J.L.M.S.* 10 (1935), 82-3.(3) The almost periodic behavior of the function $1/\zeta(1+it)$, *Duke Math. J.* 2 (1936), 443-6.(4) Riemann's hypothesis and almost periodic behavior, *Revista Ci. Lima*, 41 (1939), 575-85.(5) On the asymptotic behavior of the Riemann zeta-function on the line $\sigma = 1$, *Amer. J. Math.* 63 (1941), 575-80.(6) Riemann's hypothesis and harmonic analysis, *Duke Math. J.* 10 (1943), 99-105.(7) The behavior of Euler's product on the boundary of convergence, *Duke Math. J.* 10 (1943), 429-40.(8) Random factorizations and Riemann's hypothesis, *Duke Math. J.* 11 (1944), 267-75.

FURTHER REFERENCES

ANDERSON, R. J.

- [1] Simple zeros of the Riemann zeta-function, *J. Number Theory*, 17 (1983), 176-182.

ATKINSON, F. V.

- [1] The mean value of the Riemann zeta-function, *Acta Math.* 81 (1949), 353-376.

BAGCHI, B.

- [1] A joint universality theorem for Dirichlet L-functions, *Math. Zeit.* 181 (1982), 319-335.

BALASUBRAMANIAN, R.

- [1] An improvement of a theorem of Titchmarsh on the mean square of $\zeta(\frac{1}{2} + it)$, *Proc. London Math. Soc.* (3), 36 (1978), 540-576.

BALASUBRAMANIAN, R., CONREY, J. B., and HEATH-BROWN, D. R.

- [1] Asymptotic mean square of the product of the Riemann zeta-function and a Dirichlet polynomial, *J. Reine Angew. Math.* 357 (1985), 161-181.

BALASUBRAMANIAN, R., and RAMACHANDRA, K.

- [1] The place of an identity of Ramanujan in prime number theory, *Proc. Indian Acad. Sci.* 83 A (1976), 156-165.
- [2] On the frequency of Titchmarsh's phenomenon for $\zeta(s)$. III, *Proc. Indian Acad. Sci.* 86 A (1977), 341-351.
- [3] On the zeros of the Riemann Zeta function and L-series -II, *Hardy-Ramanujan J.* 5 (1982), 1-30.

BERNDT, B. C.

- [1] The number of zeros for $\zeta^{(k)}(s)$, *J. London Math. Soc.* (2), 2 (1970), 577-580.

BURGESS, D. A.

- [1] On character sums and L-series. II, *Proc. London Math. Soc.* (3), 13 (1963), 524-536.

CHEN, J.-R.

- [1] On the order of $\zeta(\frac{1}{2} + it)$, *Chinese Math. Acta*, 6 (1965), 463-478.

CHIH, T.-T.

- [1] A divisor problem, *Acad. Sinica Sci. Record*, 3 (1950), 177-182.

CONREY, J. B.

- [1] Zeros of derivatives of Riemann's xi-function on the critical line, *J. Number Theory*, 16 (1983), 49-74.

CONREY, J. B., and GHOSH, A.

- [1] On mean values of the zeta-function, *Mathematika*, 31 (1984), 159-161.

CONREY, J. B., GHOSH, A., GOLDSTON, D., GONEK, S. M., and HEATH-BROWN, D. R.

- [1] On the distribution of gaps between zeros of the zeta-function, *Quart. J. Math. Oxford* (2), 36 (1985), 43-51.

CONREY, J. B., GHOSH, A., and GONEK, S. M.

- [1] A note on gaps between zeros of the zeta function, *Bull. London Math. Soc.* 16 (1984), 421-424.

CORRADI, K., and KÁTAI, I.

- [1] A comment on K. S. Gangadharan's paper entitled "Two classical lattice point problems", *Magyar Tud. Akad. Mat. Fiz. Oszt. Kézl.* 17 (1967), 89-97.

FURTHER REFERENCES

407

DESJOUILLERS, J.-M., and IWANEC, H.

- [1] Kloosterman sums and Fourier coefficients of cusp forms, *Invent. Math.* 70 (1982), 219-288.

- [2] Power mean values of the Riemann zeta-function, *Mathematika*, 29 (1982), 202-212.

- [3] Power mean values of the Riemann zeta-function II, *Acta Arith.* 48 (1984), 305-312.

DIAMOND, H.

- [1] Elementary methods in the study of the distribution of prime numbers, *Bull. Amer. Math. Soc.* 7 (1982), 553-589.

ERDŐS, P.

- [1] On a new method in elementary number theory which leads to an elementary proof of the prime number theorem, *Proc. Nat. Acad. Sci. USA*, 35 (1949), 374-384.

ESTERMANN, T.

- [1] On the representation of a number as the sum of two products, *Proc. London Math. Soc.* (2), 31 (1930), 123-133.

- [2] On Kloosterman's sum, *Mathematika*, 8 (1961), 83-86.

FUJII, A.

- [1] On the distribution of the zeros of the Riemann Zeta function in short intervals, *Bull. Amer. Math. Soc.* 81 (1975), 139-142.

- [2] On the difference between r consecutive ordinates of the Riemann Zeta function, *Proc. Japan Acad.* 51 (1975), 741-743.

- [3] On the problem of divisors, *Acta Arith.* 31 (1976), 355-360.

GALLAGHER, P. X., and MUELLER, J. H.

- [1] Primes and zeros in short intervals, *J. Reine Angew. Math.* 303/304 (1978), 205-220.

GANGADHARAN, K. S.

- [1] Two classical lattice point problems, *Proc. Camb. Phil. Soc.* 57 (1961), 699-721.

GHOSH, A.

- [1] On Riemann's zeta function—sign changes of $S(T)$, *Recent progress in analytic number theory*, Vol I, 25-46, (Academic Press, London, 1981).

- [2] On the Riemann Zeta function—Mean value theorems and the distribution of $|S(T)|$, *J. Number Theory*, 17 (1983), 93-102.

GONEK, S. M.

- [1] Analytic properties of zeta and L-functions, Thesis, Univ. Michigan, Ann Arbor, 1979.

- [2] Mean values of the Riemann zeta-function and its derivatives, *Invent. Math.* 75 (1984), 123-141.

GOOD, A.

- [1] Ein Ω -Resultat für das quadratische Mittel der Riemannschen Zetafunktion auf der kritische Linie, *Invent. Math.* 41 (1977), 233-251.

HAFNER, J. L.

- [1] New omega theorems for two classical lattice point problems, *Invent. Math.* 63 (1981), 181-186.

- [2] On the average order of a class of arithmetic functions, *J. Number Theory*, 15 (1982), 36-76.

- HALASZ, G.
 [1] Über die Mittelwerte multiplikativer zahlentheoretischer Funktionen, *Acta Math. Acad. Sci. Hungar.* 19 (1968), 365–403.
- HALASZ, G., and TURÁN, P.
 [1] On the distribution of Roots of Riemann zeta and allied functions, I, *J. Number Theory*, 1 (1969), 121–137.
- HANKE, W.
 [1] Verschärfung der Abschätzung von $\zeta(\frac{1}{2} + it)$, *Acta Arith.* 8 (1962–63), 357–430.
- HARDY, G. H.
 [1] On Dirichlet's divisor problem, *Proc. London Math. Soc.* (2), 15 (1916), 1–25.
- HASELIGROVE, C. B.
 [1] A connection between the zeros and the mean values of $\zeta(s)$, *J. London Math. Soc.* 24 (1949), 215–222.
 [2] A disproof of a conjecture of Pólya, *Mathematika*, 5 (1958), 141–145.
- HEATH-BROWN, D. R.
 [1] The mean square of the Riemann Zeta-function, *Mathematika*, 25 (1978), 177–184.
 [2] The twelfth power moment of the Riemann Zeta-function, *Quart. J. Math. Oxford* (2), 29 (1978), 443–462.
 [3] Hybrid bounds for Dirichlet L-functions, *Invent. Math.* 47 (1978), 149–170.
 [4] The fourth power moment of the Riemann Zeta-function, *Proc. London Math. Soc.* (3), 38 (1979), 385–422.
 [5] Simple zeros of the Riemann Zeta-function on the critical line, *Bull. London Math. Soc.* 11 (1979), 17–18.
 [6] Zero density estimates for the Riemann Zeta-function and Dirichlet L-functions, *J. London Math. Soc.* (2), 20 (1979), 221–232.
 [7] Fractional moments of the Riemann Zeta-function, *J. London Math. Soc.* (2), 24 (1981), 65–78.
 [8] Mean values of the Zeta-function and divisor problems, *Recent progress in analytic number theory*, Vol I, 115–119, (Academic Press, London, 1981).
 [9] Hybrid bounds for L-functions: a q-analogue of van der Corput's method and a t-analogue of Burgess's method, *Recent progress in analytic number theory*, Vol I, 121–126, (Academic Press, London, 1981).
 [10] Gaps between primes, and the pair correlation of zeros of the zeta-function, *Acta Arith.* 41 (1982), 85–99.
 [11] The Pjateckiĭ-Sapiro prime number theorem, *J. Number Theory*, 16 (1983), 242–266.
- HUXLEY, M. N.,
 [1] On the difference between consecutive primes, *Invent. Math.* 15 (1972), 155–164.
- INGHAM, A. E.
 [1] On two classical lattice point problems, *Proc. Camb. Phil. Soc.* 36 (1940), 131–138.
- IVIĆ, A.
 [1] Large values of the error term in the divisor problem, *Invent. Math.* 71 (1983), 513–520.
 [2] A zero-density theorem for the Riemann zeta-function, *Trudy Mat. Inst. Steklov.* 163 (1984), 85–89.

- [3] *The Riemann zeta-function*, (Wiley-Interscience, New York, 1985).
- IWANIEC, H.
 [1] Fourier coefficients of cusp forms and the Riemann Zeta-function, *Exposé No. 18, Sémin. Théor. Nombres*, Université Bordeaux, (1979/80).
- JURKAT, W., and PEYERIMHOFF, A.
 [1] A constructive approach to Kronecker approximations and its applications to the Mertens conjecture, *J. Reine Angew. Math.* 286/287 (1976), 322–340.
- JUTILA, M.
 [1] Zero-density estimates for L-functions, *Acta Arith.* 32 (1977), 52–62.
 [2] Zeros of the zeta-function near the critical line, *Studies in pure mathematics, to the memory of Paul Turán*, 385–394, (Birkhäuser, Basel-Stuttgart, 1982).
 [3] Riemann's zeta-function and the divisor problem, *Arkiv för Mat.* 21 (1983), 75–96.
 [4] On the value distribution of the zeta-function on the critical line, *Bull. London Math. Soc.* 15 (1983), 513–518.
- KARATSUBA, A. A.
 [1] Estimates of trigonometric sums by Vinogradov's method, and some applications, *Proc. Steklov. Inst. Math.* 119 (1971), 241–255.
 [2] *Principles of analytic number theory*, (Russian), (Izdat. 'Nauka', Moscow, 1975).
- KOLESNIK, G.
 [1] The improvement of the error term in the divisor problem, *Mat. Zametki*, 6 (1969), 545–554.
 [2] On the estimation of certain trigonometric sums, *Acta Arith.* 25 (1973), 7–30.
 [3] On the estimation of multiple exponential sums, *Recent progress in analytic number theory*, Vol I, 231–246, (Academic Press, London, 1981).
 [4] On the order of $\zeta(\frac{1}{2} + it)$ and $\Delta(R)$, *Pacific J. Math.* 82 (1982), 107–122.
 [5] On the method of exponent pairs, *Acta Arith.* 45 (1985), 115–143.
- KOROBOV, N. M.
 [1] Estimates of trigonometric sums and their applications, *Uspehi Mat. Nauk*, 13 (1958), 185–192.
- KUBOTA, T., and LEOPOLDT, H. W.
 [1] Eine p-adische Theorie der Zetawerte. I. Einführung der p-adischen Dirichletschen L-funktionen, *J. Reine Angew. Math.* 214/215 (1964), 328–339.
- LAVRIK, A. F.
 [1] The functional equation for Dirichlet L-functions and the problem of divisors in arithmetic progressions, *Izv. Akad. Nauk SSSR Ser. Mat.* 30 (1966), 433–448.
- LAVRIK, A. F., and SOBOLEV, A. Š.
 [1] On the remainder term in the elementary proof of the prime number theorem, *Dokl. Akad. Nauk SSSR*, 211 (1973), 534–536.
- LEVINSON, N.
 [1] Ω -theorems for the Riemann zeta-function, *Acta Arith.* 20 (1972), 319–332.
 [2] More than one third of the zeros of Riemann's zeta-function are on $\sigma = \frac{1}{2}$, *Adv. Math.* 13 (1974), 383–436.

- [3] Zeros of derivative of Riemann's ζ -function, *Bull. Amer. Math. Soc.* 80 (1974), 951-954.
- [4] A simplification of the proof that $N_0(T) > \frac{1}{2}N(T)$ for Riemann's zeta-function, *Adv. Math.* 18 (1975), 239-242.
- [5] Deduction of semi-optimal mollifier for obtaining lower bounds for $N_0(T)$ for Riemann's zeta-function, *Proc. Nat. Acad. Sci. USA*, 72 (1975), 294-297.
- [6] Almost all roots of $\zeta(s) = a$ are arbitrarily close to $\sigma = \frac{1}{2}$, *Proc. Nat. Acad. Sci. USA*, 72 (1975), 1322-1324.
- LEVINSON, N., and MONTGOMERY, H. L.
- [1] Zeros of the derivative of the Riemann zeta-function, *Acta Math.* 133 (1974), 49-65.
- LOU, S.-T.
- [1] A lower bound for the number of zeros of Riemann's zeta-function on $\sigma = \frac{1}{2}$, *Recent progress in analytic number theory*, 319-324, Vol. I, (Academic press, London, 1981).
- MONTGOMERY, H. L.
- [1] *Topics in multiplicative number theory*, Lecture Notes in Math. 227 (Springer, Berlin, 1971).
- [2] The pair correlation of zeros of the zeta-function, *Analytic number theory* (Proc. Symp. Pure Math. Vol XXIV), 181-193, (Amer. Math. Soc., Providence, R. I., 1973).
- [3] Extreme values of the Riemann zeta-function, *Comment. Math. Helv.* 52 (1977), 511-518.
- MONTGOMERY, H. L., and ODLYZKO, A. M.
- [1] Gaps between zeros of the zeta-function, *Topics in classical number theory*, (Coll. Math. Soc. János Bolyai, 34), 1079-1106, (Budapest, 1981).
- MONTGOMERY, H. L., and VAUGHAN, R. C.
- [1] Hilbert's inequality, *J. London Math. Soc.* (2), 8 (1974), 73-82.
- MOTOHASHI, Y.
- [1] An elementary proof of Vinogradov's zero-free region for the Riemann zeta-function, *Recent progress in analytic number theory*, Vol. I, 257-267, (Academic Press, London, 1981).
- [2] A note on the approximate functional equation for $\zeta^2(s)$, *Proc. Japan Acad.* 59A (1983), 392-396.
- [3] A note on the approximate functional equation for $\zeta^2(s)$. II, *Proc. Japan Acad.* 59A (1983), 469-472.
- MUELLER, J. H.
- [1] On the difference between the consecutive zeros of the Riemann zeta-function, *J. Number Theory*, 14 (1982), 327-331.
- [2] On the Riemann Zeta-function $\zeta(s)$ - gaps between sign changes of $S(t)$, *Mathematika*, 29 (1983), 264-269.
- ODLYZKO, A. M., and TRIEDEL, H. J. J.
- [1] Disproof of Mertens conjecture, *J. Reine Angew. Math.* 357 (1985), 138-160.
- OSTROWSKI, A.
- [1] Über Dirichletsche Reihen und algebraische Differentialgleichungen, *Math. Zeit.* 8 (1920), 115-143.

- PINTZ, J.
- [1] On the remainder term of the prime number formula and the zeros of Riemann's zeta-function, *Number theory, Noordwijkerhout 1983*, 186-197, Lecture Notes in Math. 1068, (Springer, Berlin, 1984).
- RAMACHANDRA, K.
- [1] On the zeros of the Riemann zeta-function and L-series, *Acta Arith.* 34 (1978), 211-218.
- [2] Some remarks on a theorem of Montgomery and Vaughan, *J. Number Theory*, 11 (1980), 465-471.
- [3] Some remarks on the mean value of the Riemann Zeta-function and other Dirichlet series - II, *Hardy-Ramanujan J.* 3 (1980), 1-24.
- [4] Some remarks on the mean value of the Riemann Zeta-function and other Dirichlet series - III, *Ann. Acad. Sci. Fenn. Ser. A1 Math.* 5 (1980), 145-158.
- [5] Mean-value of the Riemann Zeta-function and other remarks - II, *Trudy Mat. Inst. Steklov.* 163 (1984), 200-204.
- RANKIN, R. A.
- [1] Van der Corput's method and the theory of exponent pairs, *Quart. J. Math. Oxford* (2), 6 (1955), 147-153.
- REICH, A.
- [1] Zetafunktion und Differenzen-Differentialgleichungen, *Arch. Math. (Basel)*, 38 (1982), 226-235.
- RIECHT, H.-E.
- [1] Verschärfung der Abschätzung beim Dirichletschen Teilerproblem, *Math. Zeit.* 58 (1953), 204-218.
- [2] Einführung in die Theorie der starken Riesz'schen Summierbarkeit von Dirichletreihen, *Nachr. Akad. Wiss. Göttingen (Math.-Phys.) Kl. II*, 1960, 17-75.
- [3] Zur Abschätzung der Riemannschen Zetafunktion in der Nähe der Vertikalen $\sigma = 1$, *Math. Ann.* 169 (1967), 97-101.
- SELBERG, A.
- [1] The zeta-function and the Riemann Hypothesis, *Skandinaviske Matematikerkongres*, 10 (1946), 187-200.
- [2] An elementary proof of the prime number theorem, *Ann. Math.* (2), 50 (1949), 305-313.
- [3] Discontinuous groups and harmonic analysis, *Proc. Internat. Congr. Mathematicians (Stockholm, 1962)*, 177-189 (Inst. Mittag-Leffler, Djursholm, 1963).
- SPIRA, R.
- [1] Zeros of sections of the zeta function. II, *Math. Comp.* 22 (1968), 163-173.
- SRINIVASAN, B. R.
- [1] Lattice point problems of many-dimensional hyperboloids III, *Math. Ann.* 160 (1965), 280-311.
- STEČKIN, S. B.
- [1] Mean values of the modulus of a trigonometric sum, *Trudy Mat. Inst. Steklov.* 134 (1975), 283-309, 411.
- TATE, J. T.
- [1] Fourier analysis in number fields, and Hecke's zeta-functions, *Algebraic Number Theory* (Proc. Instructional Conf., Brighton, 1965), 305-347, (Thompson, Washington DC, 1967).