

or, because

$$2\psi(x) + 1 = x^{-1/2} \left[2\psi\left(\frac{1}{x}\right) + 1 \right] \quad (\text{Jacobi, Fund., p. 184}),$$

that

$$\begin{aligned} \Pi\left(\frac{s}{2} - 1\right) \pi^{-s/2} \zeta(s) &= \int_1^\infty \psi(x) x^{(s/2)-1} dx + \int_0^1 \psi\left(\frac{1}{x}\right) x^{(s-3)/2} dx \\ &\quad + \frac{1}{2} \int_0^1 (x^{(s-3)/2} - x^{(s/2)-1}) dx \\ &= \frac{1}{s(s-1)} + \int_1^\infty \psi(x) (x^{(s/2)-1} + x^{-(1+s/2)}) dx. \end{aligned}$$

I now set $s = \frac{1}{2} + ti$ and

$$\Pi\left(\frac{s}{2}\right) (s-1) \pi^{-s/2} \zeta(s) = \xi(t)$$

so that

$$\xi(t) = \frac{1}{2} - (tt + \frac{1}{4}) \int_1^\infty \psi(x) x^{-3/4} \cos(\frac{1}{2}t \log x) dx$$

or also

$$\xi(t) = 4 \int_1^\infty \frac{d[x^{3/2}\psi'(x)]}{dx} x^{-1/4} \cos\left(\frac{1}{2}t \log x\right) dx.$$

This function is finite for all finite values of t and can be developed as a power series in tt which converges very rapidly. Now since for values of s with real part greater than 1, $\log \zeta(s) = -\sum \log(1 - p^{-s})$ is finite and since the same is true of the other factors of $\xi(t)$, the function $\xi(t)$ can vanish only when the imaginary part of t lies between $\frac{1}{2}i$ and $-\frac{1}{2}i$. The number of roots of $\xi(t) = 0$ whose real parts lie between 0 and T is about

$$= \frac{T}{2\pi} \log \frac{T}{2\pi} - \frac{T}{2\pi}$$

because the integral $\int d \log \xi(t)$ taken in the positive sense around the domain consisting of all values whose imaginary parts lie between $\frac{1}{2}i$ and $-\frac{1}{2}i$ and whose real parts lie between 0 and T is (up to a fraction of the order of magnitude of $1/T$) equal to $[T \log (T/2\pi) - T]i$ and is, on the other hand, equal to the number of roots of $\xi(t) = 0$ in the domain multiplied by $2\pi i$. One finds in fact about this many real roots within these bounds and it is very likely that all of the roots are real. One would of course like to have a rigorous proof of this, but I have put aside the search for such a proof after some fleeting vain attempts because it is not necessary for the immediate objective of my investigation.

If one denotes by α the roots of the equation $\xi(\alpha) = 0$, then one can express $\log \xi(t)$ as

$$\sum \log \left(1 - \frac{tt}{\alpha\alpha} \right) + \log \xi(0)$$

because, since the density of roots of size t grows only like $\log(t/2\pi)$ as t grows, this expression converges and for infinite t is only infinite like $t \log t$; thus it differs from $\log \xi(t)$ by a function of tt which is continuous and finite for finite t and which, when divided by tt , is infinitely small for infinite t . This difference is therefore a constant, the value of which can be determined by setting $t = 0$.

With these preparatory facts, the number of primes less than x can now be determined.

Let $F(x)$, when x is not exactly equal to a prime, be equal to this number, but when x is a prime let it be greater by $\frac{1}{2}$ so that for an x where $F(x)$ jumps

$$F(x) = \frac{F(x+0) + F(x-0)}{2}.$$

If one sets

$$p^{-s} = s \int_p^\infty x^{-s-1} dx, \quad p^{-2s} = s \int_{p^2}^\infty x^{-s-1} dx, \quad \dots$$

in the formula

$\log \zeta(s) = -\sum \log(1 - p^{-s}) = \sum p^{-s} + \frac{1}{2} \sum p^{-2s} + \frac{1}{3} \sum p^{-3s} + \dots$,
one finds

$$\frac{\log \zeta(s)}{s} = \int_1^\infty f(x) x^{-s-1} dx$$

when one denotes

$$F(x) + \frac{1}{2}F(x^{1/2}) + \frac{1}{3}F(x^{1/3}) + \dots$$

by $f(x)$.

This equation is valid for every complex value $a + bi$ of s provided $a > 1$. But when in such circumstances

$$g(s) = \int_0^\infty h(x) x^{-s} d \log x$$

is valid, the function h can be expressed in terms of g by means of Fourier's theorem. The equation splits when h is real and when $g(a + bi) = g_1(b) + ig_2(b)$ into the two equations

$$g_1(b) = \int_0^\infty h(x) x^{-a} \cos(b \log x) d \log x,$$

$$ig_2(b) = -i \int_0^\infty h(x) x^{-a} \sin(b \log x) d \log x.$$

When both equations are multiplied by $[\cos(b \log y) + i \sin(b \log y)] db$ and integrated from $-\infty$ to $+\infty$, one finds in both cases that the right side is $\pi h(y) y^{-a}$ so that when they are added and multiplied by iy^a

$$2\pi i h(y) = \int_{a-\infty i}^{a+\infty i} g(s) y^s ds,$$

where the integration is to be carried out in such a way that the real part of s remains constant.†

The integral represents, for a value of y where the function $h(y)$ has a jump, the middle value between the two values of h on either side of the jump. The function f was defined in such a way that it too has this property, so one has in full generality

$$f(y) = \frac{1}{2\pi i} \int_{a-\infty i}^{a+\infty i} \frac{\log \zeta(s)}{s} y^s ds.$$

For $\log \zeta$ one can now substitute the expression

$$\begin{aligned} & \frac{s}{2} \log \pi - \log(s-1) - \log \Pi\left(\frac{s}{2}\right) \\ & + \sum_{\alpha} \log \left[1 + \frac{(s-\frac{1}{2})^2}{\alpha\alpha} \right] + \log \xi(0) \end{aligned}$$

found above; the integrals of the individual terms of this expression will not converge, however, when they are taken to infinity, so it is advantageous to reformulate the equation as

$$f(x) = -\frac{1}{2\pi i} \frac{1}{\log x} \int_{a-\infty i}^{a+\infty i} \frac{d \frac{\log \zeta(s)}{s}}{ds} x^s ds$$

by integration by parts.

Since

$$-\log \Pi\left(\frac{s}{2}\right) = \lim \left[\sum_{n=1}^m \log \left(1 + \frac{s}{2n} \right) - \frac{s}{2} \log m \right]$$

for $m = \infty$ and therefore,

$$-\frac{d \frac{1}{s} \log \Pi\left(\frac{s}{2}\right)}{ds} = \sum_1^{\infty} \frac{d \frac{1}{s} \log \left(1 + \frac{s}{2n} \right)}{ds},$$

all of the terms in the expression for $f(x)$ except for the term

$$\frac{1}{2\pi i} \frac{1}{\log x} \int_{a-\infty i}^{a+\infty i} \frac{1}{ss} \log \xi(0) x^s ds = \log \xi(0)$$

take the form

$$\pm \frac{1}{2\pi i} \frac{1}{\log x} \int_{a-\infty i}^{a+\infty i} \frac{d \left[\frac{1}{s} \log \left(1 - \frac{s}{\beta} \right) \right]}{ds} x^s ds.$$

But

$$\frac{d \left[\frac{1}{s} \log \left(1 - \frac{s}{\beta} \right) \right]}{d\beta} = \frac{1}{(\beta-s)\beta}$$

†This argument is not quite correct. See the relevant note in Riemann's collected works [R1] (translator's note).

and, when the real part of s is greater than the real part of β ,

$$-\frac{1}{2\pi i} \int_{a-\infty i}^{a+\infty i} \frac{x^s ds}{(\beta - s)\beta} = \frac{x^\beta}{\beta} = \int_{\infty}^x x^{\beta-1} dx$$

or

$$= \int_0^x x^{\beta-1} dx$$

depending on whether† the real part of β is negative or positive. Thus

$$\begin{aligned} & \frac{1}{2\pi i} \frac{1}{\log x} \int_{a-\infty i}^{a+\infty i} \frac{d\left[\frac{1}{s} \log\left(1 - \frac{s}{\beta}\right)\right]}{ds} x^s ds \\ &= -\frac{1}{2\pi i} \int_{a-\infty i}^{a+\infty i} \frac{1}{s} \log\left(1 - \frac{s}{\beta}\right) x^s ds \\ &= \int_{\infty}^x \frac{x^{\beta-1}}{\log x} dx + \text{const} \end{aligned}$$

in the first case and

$$= \int_0^x \frac{x^{\beta-1}}{\log x} dx + \text{const}$$

in the second case.

In the first case the constant of integration can be determined by taking β to be negative and infinite. In the second case the integral from 0 to x takes on two values which differ by $2\pi i$ depending on whether the path of integration is in the upper halfplane or in the lower halfplane; if the path of integration is in the upper halfplane, the integral will be infinitely small when the coefficient of i in β is infinite and positive, and if the path is in the lower halfplane, the integral will be infinitely small when the coefficient of i in β is infinite and negative. This shows how to determine the values of $\log[1 - (s/\beta)]$ on the left side in such a way that the constants of integration drop out.

By setting these values in the expression for $f(x)$ one finds

$$\begin{aligned} f(x) &= \text{Li}(x) - \sum_{\alpha} [\text{Li}(x^{(1/2)+\alpha i}) + \text{Li}(x^{(1/2)-\alpha i})] \\ &\quad + \int_x^{\infty} \frac{1}{x^2 - 1} \frac{dx}{x \log x} + \log \xi(0), \end{aligned}$$

where‡ the sum \sum_{α} is over all positive roots (or all roots with positive real parts) of the equation $\xi(\alpha) = 0$, ordered according to their size. It is possible, by means of a more exact discussion of the function ξ , easily to show that with this ordering of the roots the sum of the series

$$\sum_{\alpha} [\text{Li}(x^{(1/2)+\alpha i}) + \text{Li}(x^{(1/2)-\alpha i})] \log x$$

†Note that this excludes the possibility $\text{Re } \beta = 0$ and therefore does not apply to roots, if any, on the imaginary axis (translator's note).

‡Concerning the erroneous value of $\log \xi(0)$ in this formula, see Chapter 1 (translator's note).

is the same as the limiting value of

$$\frac{1}{2\pi i} \int_{a-bi}^{a+bi} \frac{d \frac{1}{s} \sum \log \left[1 + \frac{(s - \frac{1}{2})^2}{\alpha\alpha} \right]}{ds} x^s ds$$

as b grows without bound; by a different ordering, however, it can approach any arbitrary real value.

From $f(x)$ one can find $F(x)$ by inverting

$$f(x) = \sum \frac{1}{n} F(x^{1/n})$$

to find

$$F(x) = \sum (-1)^\mu \frac{1}{m} f(x^{1/m}),$$

where m ranges over all positive integers which are not divisible by any square other than 1 and where μ denotes the number of prime factors of m .

If \sum_α is restricted to a finite number of terms, then the derivative of the expression for $f(x)$ or, except for a part which decreases very rapidly as x increases,

$$\frac{1}{\log x} - 2 \sum_\alpha \frac{\cos(\alpha \log x) x^{-1/2}}{\log x}$$

gives an approximate expression for the density of primes + half the density of prime squares + $\frac{1}{3}$ the density of prime cubes, etc., of magnitude x .

Thus the known approximation $F(x) = \text{Li}(x)$ is correct only to an order of magnitude of $x^{1/2}$ and gives a value which is somewhat too large, because the nonperiodic† terms in the expression of $F(x)$ are, except for quantities which remain bounded as x increases,

$$\begin{aligned} \text{Li}(x) - \frac{1}{2} \text{Li}(x^{1/2}) - \frac{1}{3} \text{Li}(x^{1/3}) - \frac{1}{5} \text{Li}(x^{1/5}) \\ + \frac{1}{6} \text{Li}(x^{1/6}) - \frac{1}{7} \text{Li}(x^{1/7}) + \dots \end{aligned}$$

In fact the comparison of $\text{Li}(x)$ with the number of primes less than x which was undertaken by Gauss and Goldschmidt and which was pursued up to $x =$ three million shows that the number of primes is already less than $\text{Li}(x)$ in the first hundred thousand and that the difference, with minor fluctuations, increases gradually as x increases. The thickening and thinning of primes which is represented by the periodic terms in the formula has also been observed in the counts of primes, without, however, any possibility of establishing a law for it having been noticed. It would be interesting in a future count to examine the influence of individual periodic terms in the formula for the density of primes. More regular than the behavior of $F(x)$ is the behavior of $f(x)$ which already in the first hundred is on average very nearly equal to $\text{Li}(x) + \log \xi(0)$.

†Strictly speaking, the terms $\text{Li}(x^{(1/2)+\alpha i})$ are not periodic but merely oscillatory (translator's note).

References

- A1 Abel, N. H., Solution de quelques problemes a l'aide d'integrales définies. *Mag. Naturvidenskaberne* 2 (1823). (Also "Oeuvres," 1, pp. 11–27.)
- A2 Abel, N. H., Letter to Holmboe dated 4 August 1823. In "Niels Henrik Abel, Memorial, Publié à L'Occasion du Centenaire de sa Naissance."
- A3 Ahlfors, L. V., "Complex Analysis." McGraw-Hill, New York, 1953.
- B1 Backlund, R., Sur les zéros de la fonction $\zeta(s)$ de Riemann. *C. R. Acad. Sci. Paris* 158, 1979–1982 (1914).
- B2 Backlund, R., Über die Nullstellen der Riemannschen Zetafunktion. Dissertation, Helsingfors, 1916.
- B3 Backlund, R., Über die Nullstellen der Riemannschen Zetafunktion. *Acta Math.* 41, 345–375 (1918).
- B4 Backlund, R., Über die Beziehung zwischen Anwachsen und Nullstellen der Zetafunktion. *Öfversigt Finska Vetensk. Soc.* 61, No. 9 (1918–1919).
- B5 Bochner, S., "Vorlesungen über Fouriersche Integrale," Akademische Verlagsgesellschaft M. B. H., Leipzig, 1932. (Reprinted by Chelsea, Bronx, New York, 1948.)
- B6 Bochner, S., Ein Satz von Landau und Ikehara. *Math. Z.* 37, 1–9 (1933).
- B6a Bohr, H., and Landau, E., Über das Verhalten von $\zeta(s)$ und $\zeta_K(s)$ in der Nähe der Geraden $\sigma = 1$. *Göttinger Nachr.* 303–330 (1910).
- B7 Bohr, H., and Landau, E., Beiträge zur Theorie der Riemannschen Zetafunktion. *Math. Ann.* 74, 3–30 (1913). (Also Bohr's "Collected Works," Vol. I, B11.)
- B8 Bohr, H., and Landau, E., Ein Satz über Dirichletsche Reihen mit Anwendung *Rend. Circ. Mat. Palermo* 37, 269–272 (1914). (Also Bohr's "Collected Works," Vol. I, B13.)
- B9 Bombieri, E., Sulle formule di A. Selberg generalizzate per classi di funzioni aritmetiche e le applicazioni al del resto nel "Primzahlsatz." *Riv. Mat. Univ. Parma* [2] 3, 393–440 (1962).
- C1 Cantor, M., "Vorlesungen über Geschichte der Mathematik," Vol. 3, p. 663. Teubner, Leipzig, 1898.
- C2 Chebyshev, P. L., Sur la fonction qui détermine la totalité des nombres premiers inférieurs à une limite donnée. *J. Math. Pures Appl.* [1] 17 (1852). (First published 1848. Available in both French and Russian Editions of Collected Works.)
- C3 Chebyshev, P. L., Mémoire sur les nombres premiers. *J. Math. Pures Appl.* [1] 17 (1852). (First published 1850. Available in both French and Russian Editions of Collected Works.)

- C4 Chebyshev, P., "Oeuvres." St. Petersburg. (Republished by Chelsea, Bronx, New York, 1962 (?).)
- C5 Chebyshev, P., "Polnoe Sobranie Sochineniy." Press of the USSR Acad. of Sci., Moscow and Leningrad, 1947.
- D1 Debye, P. J. W., Näherungsformeln für die Zylinderfunktionen . . . *Math. Ann.* **67**, 535–558 (1909). (Engl. Transl. in "The Collected Papers of Peter J. W. Debye." Wiley (Interscience), New York, 1954.)
- D2 Denjoy, A., L'Hypothèse de Riemann sur la distribution des zéros de $\zeta(s)$, reliée à la théorie des probabilités. *C. R. Acad. Sci. Paris* **192**, 656–658 (1931).
- D3 Dirichlet, P. G. Lejeune, Sur l'usage des series infinies dans la théorie des nombres. *J. Reine Angew. Math.* **18**, 257–274 (1838). (Also "Werke," Vol. I, pp. 359–374.)
- E1 Edwards, H. M., "Advanced Calculus." Houghton, Boston, Massachusetts, 1969.
- E2 Euclid, "Elements," Book 9, Prop. 20.
- E3 Euler, L., De progressionibus transcendentibus . . . *Comm. Acad. Sci. Petropolitanae* **5**, 36–57 (1730). (Also "Opera" (1), Vol. 14, pp. 1–24.)
- E4 Euler, L., Variarum observationes circa series infinitas. *Comm. Acad. Sci. Petropolitanae* **9**, 222–236 (1737). ("Opera" (1), Vol. 14, pp. 216–244.)
- E5 Euler, L., "Introductio in Analysin Infinitorum," Chapter 15. Bousquet and Socios., Lausanne, 1748. ("Opera" (1), Vol. 8.)
- E6 Euler, L., "Institutiones Calculi Differentialis," Pt. 2, Chapters 5 and 6. Acad. Imp. Sci. Petropolitanae, St. Petersburg, 1755. ("Opera" (1), Vol. 10.)
- E7 Euler, L., Remarques sur un beau rapport entre les séries des puissances tant directes que réciproques. *Mem. Acad. Sci. Berlin* **17**, 83–106 (1761). ("Opera" (1), Vol. 15, pp. 70–90.)
- E8 Euler, L., Survey of Euler's work on the factorial function (German). Introduction to Euler's "Opera" (1), Vol. 19, 1932.
- F1 Franel, J., and Landau, E., Les suites de Farey et le problème des nombres premiers. *Göttinger Nachr.*, 198–206 (1924).
- F2 Fuss, P.-H., "Correspondance Mathématique et Physique." Imp. Acad. Sci., St. Petersburg, 1843. (Reprinted, Johnson Reprint Corp., 1968.)
- G1 Gauss, C. F., Circa seriem infinitam

$$1 + \frac{\alpha\beta}{1\cdot\gamma}x + \frac{\alpha(\alpha+1)\beta(\beta+1)}{1\cdot2\cdot\gamma(\gamma+1)}x^2 + \dots,$$

- Comm. Soc. Reg. Sci. Göttingensis* **2** (1813). (Also "Werke," Vol. III, pp. 125–160.)
- G2 Gauss, C. F., Letter to Enke dated 24 December 1849. "Werke," Vol. II, pp. 444–447. Königlich Gesellschaft der Wissenschaften zu Göttingen.
- G3 Gauss, C. F., "Werke," Vol. XI, p. 11. Königlich Gesellschaft der Wissenschaften zu Göttingen.
- G4 Genocchi, A., Formule per determinare quanti siano i numeri primi fino ad un dato limite (Résumé of Riemann's Paper). *Ann. Mat. Pura Appl.* [1] **3**, 52–59 (1860).
- G5 Gram, J.-P., Sur les Zéros de la Fonction $\zeta(s)$ de Riemann. *Acta Math.* **27**, 289–304 (1903).
- H1 Hadamard, J., Étude sur les Propriétés des Fonctions Entières et en Particulier d'une Fonction Considérée par Riemann. *J. Math. Pures Appl.* [4] **9**, 171–215 (1893). (Also in "Oeuvres.")
- H2 Hadamard, J., Sur la distribution des zeros de la fonction $\zeta(s)$ et ses conséquences arithmétiques. *Bull. Soc. Math. France* **24**, 199–220 (1896). (Also in "Oeuvres.")
- H3 Hardy, G. H., Sur les Zéros de la Fonction $\zeta(s)$ de Riemann. *C. R. Acad. Sci. Paris* **158**, 1012–1014 (1914). (Also in "Collected Papers.")

- H3a Hardy, G. H., Prime numbers, *Brit. Ass. Rep.*, 1915, 350–354. [“Collected Papers,” Vol. 2, pp. 14–19.]
- H4 Hardy, G. H., “Ramanujan.” Cambridge Univ. Press, London and New York, 1940. (Reprinted by Chelsea, Bronx, New York.)
- H5 Hardy, G. H., “Divergent Series.” Oxford Univ. Press, London and New York, 1956. (First Edition, 1949).
- H6 Hardy, G. H., and Littlewood, J. E., The zeros of Riemann’s zeta-function on the critical line. *Math. Z.* **10**, 283–317 (1921). (Also in Hardy’s “Collected Papers.”)
- H7 Hardy, G. H., and Wright, E. M., “An Introduction to the Theory of Numbers.” Oxford Univ. Press, London and New York, 1938.
- H8 Haselgrove, C. B., “Tables of the Riemann Zeta Function,” Roy. Soc. Math. Tables, Vol. 6. Cambridge Univ. Press, London and New York, 1960.
- H9 Hilbert, D. Problèmes futures des mathématiques. *C. R. 2nd Congr. Int. Math., Paris*, 1902, p. 85.
- H10 Holmgren, E., On Primtalens Fördelning. *Öfversigt Kgl. Vetensk. Acad. Stockholm* **59**, 221–225 (1902–1903).
- H11 Hutchinson, J. I., On the roots of the Riemann zeta-function. *Trans. Amer. Math. Soc.* **27**, 49–60 (1925).
- I1 Ikehara, S., An extension of Landau’s theorem in the analytic theory of numbers. *J. Math. and Phys.* **10**, 1–12 (1931).
- I2 Ingham, A. E., “The Distribution of Prime Numbers,” Cambridge Tracts in Math. and Math. Phys., No. 30. Cambridge Univ. Press, London and New York, 1932. (Republished by Stechert-Hafner, New York and London, 1964.)
- J1 Jacobi, C. G. J., Suite des notices sur les fonctions elliptiques. *J. Reine Angew. Math.* **3**, 303–310 (1828). (Also “Werke,” Vol. 1, pp. 255–263.)
- J2 Jeffreys, H. J., and Jeffreys, B. S., “Methods of Mathematical Physics.” Cambridge Univ. Press, London and New York, 1946.
- J3 Jensen, J. L. V. W., Sur un nouvel et important théorème de la théorie des fonctions. *Acta Math.* **22**, 359–364 (1899).
- K1 Von Koch, H., Sur la distribution des nombres premiers. *Acta Math.* **24**, 159–182 (1901).
- K2 Kuzmin, R., Contribution to the theory of a class of Dirichlet series (Russian). *Izv. Akad. Nauk SSSR, Ser. Math. Nat. Sci.* **7**, 115–124 (1930).
- K3 Kuzmin, R., On the roots of Dirichlet series (Russian). *Izv. Akad. Nauk SSSR Ser. Math. Nat. Sci.* **7**, 1471–1491 (1934).
- L1 Lakatos, I., Proofs and refutations (I–IV). *British J. Philos. Sci.* (1963).
- L2 Landau, E., Nouvelle démonstration pour la formule de Riemann *Ann. Sci. École Norm. Sup.* [3] **25**, 399–442 (1908).
- L3 Landau, E., “Handbuch der Lehre von der Verteilung der Primzahlen.” Teubner, Leipzig, 1909. (Reprinted by Chelsea, Bronx, New York.)
- L3a Landau, E., Zur Theorie der Riemannschen Zetafunktion, *Vierteljahrsschr. Naturf. Ges. Zurich.* **56**, 125–148 (1911).
- L4 Legendre, A. M., “Théorie des Nombres,” 4th ed., Vol. 2, 4th Pt., §VIII Paris, 1830. (Reprinted, Librairie Sci. Tech., A. Blanchard, Paris, 1955.)
- L5 Lehman, R. S., Separation of zeros of the Riemann zeta function, I. *Math. Comp.* **20**, 542–550 (1966).
- L6 Lehman, R. S., On the difference $\pi(x) - \text{Li}(x)$. *Acta Arith.* **11**, 397–410 (1966).
- L7 Lehmer, D. H., On the roots of the Riemann zeta-function. *Acta Math.* **95**, 291–298 (1956).

- L8 Lehmer, D. H., Extended computation of the Riemann zeta-function. *Mathematika* 3, 102–108 (1956).
- L9 Lehmer D. N., List of prime numbers from 1 to 10,006,721. Publ. No. 165. Carnegie Inst. of Washington, Washington, D.C. 1913. (Reprinted, Hafner, New York, 1956.)
- L10 Levinson, N., A motivated account of an elementary proof of the prime number theorem. *Amer. Math. Monthly* 76, 225–245 (1969).
- L11 Lindelöf, E., Quelques remarques sur la croissance de la fonction $\zeta(s)$. *Bull. Sci. Math.* [2] 32₁, 341–356 (1908).
- L12 Littlewood, J. E., Quelques conséquences de l'hypothèse que la fonction $\zeta(s)$ n'a pas de zéros dans le demi-plan $\operatorname{Re}(s) > \frac{1}{2}$. *C. R. Acad. Sci. Paris* 154, 263–266 (1912).
- L13 Littlewood, J. E., Sur la distribution des nombres premiers. *C. R. Acad. Sci. Paris* 158, 1869–1872 (1914).
- L14 Littlewood, J. E., On the zeros of the Riemann zeta-function. *Proc. Cambridge Philos. Soc.* 22, 295–318 (1924).
- M1 von Mangoldt, H., Zu Riemann's Abhandlung 'Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse'. *J. Reine Angew. Math.* 114, 255–305 (1895).
- M2 von Mangoldt, H., Beweis der Gleichung $\sum_{k=1}^{\infty} u(k)/k = 0$. *S.-B. Kgl. Preuss. Akad. Wiss. Berlin*, 835–852 (1897).
- M3 von Mangoldt, H., Zur Verteilung der Nullstellen der Riemannschen Funktion $\xi(t)$. *Math. Ann.* 60, 1–19 (1905).
- M4 Mellin, H., Eine Formel für den Logarithmus transzendenter Funktionen von endlichem Geschlecht. *Acta Soc. Sci. Fenn.* 29 (1900).
- M5 Mertens, F., Ein Beitrag zur Analytischen Zahlentheorie. *J. Reine Angew. Math.* 78, 46–62 (1874).
- M6 Mertens, F., Über eine Eigenschaft der Riemann'schen ζ -Function. *S.-B. Akad. Wiss. Wien Math.-Natur. Kl. Abt. 2A* 107, 1429–1434 (1898).
- P1 Polya, G., Über die Nullstellen gewisser ganzer Funktionen. *Math. Z.* 2, 352–383 (1918).
- P2 Polya, G., Über trigonometrische Integrale mit nur reellen Nullstellen. *J. Reine Angew. Math.* 158, 6–18 (1927).
- R1 Riemann, B., "Gesammelte Werke." Teubner, Leipzig, 1892. (Reprinted by Dover Books, New York, 1953.)
- R1a Riemann, B., Unpublished Papers, Handschriftenabteilung Niedersächsische Staats- und Universitätsbibliothek, Göttingen.
- R2 Riemann, B., "Partielle Differentialgleichungen" (Lectures edited and prepared for publication by K. Hattendorf). Vieweg, Braunschweig, 1876.
- R3 Rosser, J. B., Yohe, J. M., and Schoenfeld, L., Rigorous computation and the zeros of the Riemann zeta-function. *Cong. Proc. Int. Federation Information Process.* 1968, pp. 70–76, Spartan, Washington, D.C. and Macmillan, New York, 1969.
- S1 Selberg, A., On the zeros of Riemann's zeta-function. *Skr. Norske Vid.-Akad. Oslo* No. 10 (1942).
- S2 Selberg, A., The zeta-function and the Riemann hypothesis. *Proc. Skand. Math. Kongr., 10th*, pp. 187–200, Jul. Giellerups Forlag, Kobenhavn, 1947.
- S3 Selberg, A., An elementary proof of the prime number theorem. *Ann. of Math.* [2] 50, 305–313 (1949).
- S4 Siegel, C. L., Über Riemanns Nachlaß zur analytischen Zahlentheorie. *Quellen Studien zur Geschichte der Math. Astron. und Phys. Abt. B: Studien* 2, 45–80 (1932). (Also in "Gesammelte Abhandlungen," Vol. 1. Springer-Verlag, Berlin and New York, 1966.)

- S5 Stieltjes, T. J., Sur le developpement de $\log \Gamma(a)$. *J. Math. Pures Appl.* [4] **5**, 425–444 (1889). (Also in “Oeuvres.”)
- S6 Stieltjes, T. J., Letters in “Correspondance d’Hermite et de Stieltjes,” 2 Vols. Gauthier-Villars, Paris, 1905. (Especially Letter No. 79 and No. 4 in the Appendix.)
- S7 Stirling, J., “Methodus Differentialis Sive Tractatus de Summatione et Interpolatione Serierum Infinitarum.” London, 1730.
- T1 Tauber, A., Ein Satz aus der Theorie der unendlichen Reihen. *Monatsh. Math.* **8**, 273–277 (1897).
- T2 Taylor, A. E., “Advanced Calculus.” Ginn (Blaisdell), Boston, Massachusetts, 1955.
- T3 Titchmarsh, E. C., “The Zeta-Function of Riemann,” Cambridge Tracts in Math. and Math. Phys. No. 26. Cambridge Univ. Press, London and New York, 1930.
- T4 Titchmarsh, E. C., On van der Corput’s method and the zeta-function of Riemann (IV). *Quart. J. Math. Oxford Ser.* **5**, 98–105 (1934).
- T5 Titchmarsh, E. C., The zeros of the Riemann zeta-function. *Proc. Roy. Soc. Ser. A* **151**, 234–255 (1935).
- T6 Titchmarsh, E. C., The zeros of the Riemann zeta-function. *Proc. Roy. Soc. Ser. A* **157**, 261–263 (1936).
- T7 Titchmarsh, E. C., “Introduction to the Theory of Fourier Integrals.” Oxford Univ. Press, London and New York, 1937.
- T8 Titchmarsh, E. C., “The Theory of the Riemann Zeta-Function.” Oxford Univ. Press, London and New York, 1951.
- T9 Turing, A. M., Some calculations of the Riemann zeta-function. *Proc. London Math. Soc.* [3] **3**, 99–117 (1953).
- V1 de la Vallée Poussin, C.-J., Recherches analytiques sur la théorie des nombres (première partie). *Ann. Soc. Sci. Bruxelles* [I] **20**₂, 183–256 (1896).
- V2 de la Vallée Poussin, C.-J., Sur la fonction $\zeta(s)$ de Riemann et le nombre des nombres premiers inférieurs a une limite donnée. *Mém. Couronnés et Autres Mém. Publ. Acad. Roy. Sci., des Lettres Beaux-Arts Belg.* **59** (1899–1900).
- W1 Walfisz, A., “Weylsche Exponentialsummen in der Neueren Zahlentheorie.” VEB Deutscher Verlag der Wiss., Berlin, 1963.
- W2 Weil, A., “Courbes Algébriques et Variétés Abéliennes.” Hermann, Paris, 1948.
- W3 Wiener, N., Tauberian theorems, *Ann. of Math.* **33**, 1–100 (1932).
- W4 Weyl, H., Zur Abschätzung von $\zeta(1 + it)$. *Math. Z.* **10**, 88–101 (1921).
- W5 Wirsing, E., Elementare Beweise des Primzahlsatzes mit Restglied I. *J. Reine Angew. Math* **211**, 205–214 (1962).

Index

A

- ABEL
 integral formula for $\zeta(s)$, 9, 221
 Abel's theorem, 278
 Alpha notation for roots $\rho = \frac{1}{2} + i\alpha$, 36
 Analytic continuation
 of a particular integral, 276–277
 Riemann's conception of, 9
 of zeta function, 11, 16n, 115
 Approximate functional equation, *see*
 Hardy and Littlewood
 Asymptotic expansions, 87, *see also*
 Euler–Maclaurin summation for
 evaluation of zeta, Logarithmic
 integral, Riemann–Siegel asymp-
 totic formula, Stirling's series

B

- BACKLUND, 97
 determination of $N(T)$ for $T = 200$,
 128–129
 estimation of $N(T)$, 132–134
 theorem on Lindelöf hypothesis, 182,
 188–190
 Bernoulli numbers, 11, 103
 $B_{2n+1} = 0$, 11n, 14n, 103
 Bernoulli polynomials $B_n(x)$, 100–103
 periodified $\tilde{B}_n(x)$, 104
 BOHR and LANDAU
 Riemann hypothesis implies S un-
 bounded, 181, 201–202
 theorem on roots near $\text{Re } s = \frac{1}{2}$, 19,
 193–195
 BOMBIERI, 297

C

- Cesaro averages, 279
 CHEBYSHEV
 estimates of π from estimates of ψ ,
 68, 76–77
 estimation of $\pi(x)$, 3–4, 281–284
 mentioned by Riemann, 5
 Chebyshev's identity, 281–284
 Conditional convergence
 in Fourier analysis, 23
 of sums over ρ , 20–21, 30, 35, 49–50

D

- DEBYE, 139
 Definite integrals, *see* Riemann, evalua-
 tion of definite integrals
 DENJOY, 268–269
 DIRICHLET, 298, 299
 acquaintanceship with Chebyshev, 4
 use of Euler product formula, 7

E

- ERDÖS, 282, 288
 EUCLID, 1
 EULER
 and factorial function, 7–8
 formula $\sum_{n=-\infty}^{\infty} x^n = 0$, 212
 and functional equation of zeta, 12
 ϕ -function, 250–251
 product formula, 1, 6–7, 22–23, 50n
 statement of $\sum \mu(n)/n = 0$, 92
 $\sum p^{-1}$ diverges, 1

Euler–Maclaurin summation, 97
 for evaluation of zeta, 114–115
 statement of method, 98–106
 summary of method, 106
see also Stirling's series
 Euler product formula, *see* Euler
 Euler's constant, 67, 106n
 Explicit formulas, *see* J function, psi
 function

F

Factorial function, 7–9, *see also* Stirling's
 series
 Farey series, 263–264
 related to Riemann hypothesis,
 264–267
 Fourier analysis, 23–25, 203–225
 adjoint of an operator, 205
 interpretation of Chebyshev's identity,
 282–283
 inversion formulas, 24, 27, 51, 54–56,
 205, 213–215
 transform of an (invariant) operator,
 204
 Fourier series of $\bar{B}_1(x)$, 196
 Fourier transform, 209, 211, *see also*
 Fourier analysis, transform of an
 operator
 FRANEL and LANDAU
 Farey series and the Riemann hy-
 pothesis, 263–267

G

G function, 206
 functional equation 209–210
 Gamma function, *see* Factorial function
 GAUSS, 299, 305
 counts of primes, 2, 305
 on density of primes, 2
 notation Π for factorial, 8
 GRAM
 computation of 15 roots ρ , 96–97
 Gram points, 125–126
 Gram's law, 127, 171
 exceptions, 126–127, 176n
 statement of, 126

H

H function, 207
 HADAMARD
 proof of prime number theorem, 6,
 38, 68
 proof of product formula for ξ , 18,
 21, 39
 proof that $\zeta(1+it) \neq 0$, 69–72
 publication of three circles theorem, 187
 HARDY, 136
 infinitely many roots on $\text{Re } s = \frac{1}{2}$, 19,
 226–229
 HARDY and LITTLEWOOD
 approximate functional equation,
 201n, 229n
 average of $|\zeta(s)|^2$ on $\text{Re } s = \text{const}$, 195
 estimate of $\zeta(\frac{1}{2}+it)$, 201
 KT roots on $\text{Re } s = \frac{1}{2}$, 19, 226, 229–
 237
 reformulation of Lindelöf hypothesis,
 201
 Tauberian theorems for Cesaro
 averages, 279
 use of Tauberian theorem to prove
 prime number theorem, 280
 HASELGROVE, 96, 121, 157, 161, 178
 on exceptions to Gram's law, 126n
 excerpts from tables, 122–123, 158
 HILBERT, 6, 298
 HUTCHINSON, 97
 numerical analysis of roots ρ , 126–127,
 129–132

I

Ikehara's theorem, 281

J

J function, 22
 Riemann's formula for, 33, 48, 61–65
 in terms of $\pi(x)$, 33
 JACOBI, 15
 Jensen's theorem, 39–41

K

KUZMIN
 proof of Riemann–Siegel integral
 formula, 273–278

L

- LANDAU, 62, 62n, 136
 average of $|\zeta(s)|^2$ on $\text{Re} = \text{const}$, 195
o, *O* notation, 200
see also Bohr and Landau, Franel and Landau
- LEGENDRE
 on density of primes, 3
 notation for gamma function, 8
- Legendre relation for factorial function, 9
- LEHMAN, 269
 verification of Riemann hypothesis to g_{250000} , 172
- LEHMER, D. H.
 computations of roots ρ , 175–179
 verification of Riemann hypothesis to g_{250000} , 172
- LEHMER, D. N.
 counts of primes, 3
 on Riemann's formula for $\pi(x)$, 35
- Lehmer's phenomenon, 179
- LEVINSON, 288
- LINDELÖF
 estimates of growth of $\zeta(s)$, 182–186
- Lindelöf hypothesis, 186, 177n, 188, 201
 as consequence of Riemann hypothesis, 188
- Lindelöf's theorem, 184
 modified, 186
- LITTLEWOOD
 improvement of $\beta < 1$, 200
 improvement of Bohr–Landau theorem, 195
 improvement of Tauber's theorem, 279
 $\int_0^T S(t) dt = O(\log T)$, 173
 $\pi(x) < \text{Li}(x)$ fails, 269
 Riemann hypothesis and growth of M , 261
 use of three circles theorem, 187
see also Hardy and Littlewood
- $\text{Li}(x)$, *see* Logarithmic integral
- Logarithmic integral $\text{Li}(x)$, 26
 asymptotic formula for, 86
 estimate of $\text{Li}(x^\beta)$ as $x \rightarrow \infty$, 90
 value at x^β for complex β , 30

M

- M function (sum of Möbius μ), 260
- MACLAURIN, *see* Euler–Maclaurin summation
- VON MANGOLDT
 estimate of $N(T)$, 173
 proof of formula for ψ , 50–61
 proof of Riemann's estimate of $N(T)$, 133
 proof of Riemann's formula for $J(x)$, 48, 61–65
 proof that $\sum \mu(n)/n = 0$, 92
 statement of formula for ψ , 49, 54, 66
- MELLIN, 25n
 estimate of $\zeta(1 + it)$, 183
- MERTENS
 proof that $\zeta(1 + it) \neq 0$, 79–80
- Mertens's theorem, 6
- Möbius inversion, 34, 217–218, 283, 285
- Mu function of Lindelöf $\mu(\sigma)$, 186
- Mu function of Möbius $\mu(n)$, 34, 91–92, 217

N

- N function, 128
 Backlund's verification of Riemann's estimate, 132–134
 evaluated by Turing's method, 172–175
 Riemann's estimate, 18–19, 301
- Nonsense, 212, 217

O

- o*, *O* notation, 200

P

- Parseval's equation, 215–216
- Pi function $\pi(x)$, 4, 33
 approximations to, 84–91
 in terms of $J(x)$, 34
see also Prime number theorem
- Poisson summation formula, 209–210
- POLYA
 theorems on functions with zeros on $\text{Re } s = \frac{1}{2}$, 269–273

- Prime number theorem, 4
 - improved remainder, 84, 200
 - proof, 68–77
- Product formula for sine, 9, 18, 47, 224
- Product formula for $\xi(s)$, 20–21
 - proof, 46–47
- Psi function $\psi(x)$ of Chebyshev, 49
 - von Mangoldt's formula for, 49–61
 - see also* Chebyshev

R

- Ramanujan's formula, 218–225
- Rho, *see* Roots ρ
- RIEMANN
 - analytic functions treated globally, 20
 - comments on $\pi(x) \sim \text{Li}(x)$, 34–36, 269, 305
 - computations of roots, 159–162
 - error involving $\xi(0)$, 31
 - estimate of $N(T)$, 18–19, 301
 - evaluation of definite integrals, 12–13, 19, 26–33, 146–148
 - "everywhere valid" formula for $\zeta(s)$, 9–11
 - explicit formula for $J(x)$ ($= f(x)$), 33, 304
 - introduction of function ξ , 16
 - manuscript with statement of asymptotic formula, 156–157
 - paper on $\pi(x)$, 1–38
 - proofs of the functional equation, 12–16, 166–170, 274, 300–301
 - questions unresolved by, 37–38
 - skill as analyst, 136
 - statement of Riemann hypothesis, 19, 30n, 301
 - translation of paper on $\pi(x)$, 299–305
 - use of "Fouriers theorem", 23–25, 302–303
 - use of saddle point method, 139n
 - view of analytic continuation, 9, 20
- Riemann hypothesis, 6, 19
 - implies Lindelöf hypothesis, 188
 - in light of Riemann–Siegel formula, 164–166
 - probabilistic interpretation, 268–269
 - related to error in prime number theorem, 88–91

- related to Farey series, 263–267
- related to growth of $M(x)$, 260–263
- verified to $T = 200$, 129
- verified to $T = 300$, 129–132
- verified to $T = g_{1040}$, 171
- verified to $T = g_{250000}$, 172
- verified to $T = g_{2500000}$, 172
- verified to $T = g_{35000000}$, 172, 179–180
- Riemann–Siegel asymptotic formula, 136–164
 - error estimates, 162–164
 - manuscript, 156–157
 - statement, 154
- Riemann–Siegel integral formula, 137, 166–170
 - proof by Kuzmin, 273–278
- Roots ρ of $\xi(s) = 0$, 18–19
 - computations of, 96, 157–162, 178
 - crude estimate of density, 42–43
 - von Mangoldt's estimate of density, 56–58
 - real parts < 1 , 70–72, 79–81, 200
 - Riemann's estimate of density 21, 43, 302
 - see also* Conditional convergence, Bohr and Landau
- Rosser's rule, 180–181
- ROSSER, YOHE, and SHOENFELD
 - error estimate for Riemann–Siegel formula, 163
 - use of Turing's method, 175, 180
 - verification of Riemann hypothesis to $T = g_{35000000}$, 179–180

S

- S function $S(T)$, 173
 - estimates of 174, 190–193, 201–202
- Saddle point method, 139–140
- SELBERG
 - elementary proof of prime number theorem, 282, 288–297
 - $KT \log T$ roots on $\text{Re } s = \frac{1}{2}$, 19, 226, 237–259
- Selberg's inequality, 284–288
- Self-reciprocal operators, 210–211
- SIEGEL
 - discovery of Riemann–Siegel formula, 136

proof that Riemann–Siegel formula is asymptotic, 163
 Slit plane, 111
 Steepest descent, method of, 139–140
 STIELTJES
 alleged proof of Riemann hypothesis, 262–263
 estimate of remainder in Stirling's series, 112
 evaluation of the constant in Stirling's series, 113
 STIRLING, 109
 Stirling's formula, *see* Stirling's series
 Stirling's series 106–114
 for $\Pi'(s)/\Pi(s)$, 113
 statement, 109
 Stieltjes's estimate of the remainder, 112

T

TAUBER, 278
 Tauberian theorems, 278–281
 Theta function of Jacobi, 15, 170, 227
 Theta function θ of Chebyshev, 76, 288
 Theta function $\vartheta(t)$ in $\zeta(\frac{1}{2} + it) = Z(t) \exp i \vartheta(t)$, 119
 asymptotic formula for, 120–121
 Three circles theorem, 187
 TITCHMARSH
 average of $\zeta(\frac{1}{2} + ig_n)$ is 2, 229
 error estimate for Riemann–Siegel formula, 162–163
 as secondary source, 199, 226
 verification of Riemann hypothesis to $T = g_{1040}$, 171
 Turing's method of evaluating $N(T)$, 172–175

V

DE LA VALLÉE POUSSIN

estimate of error in prime number theorem, 78–95
 proof of prime number theorem, 68
 proof that $\sum \mu(n)/n = 0$, 91–95
 Vinogradov's estimate of $\zeta(1 + it)$, 200

W

WEIL, 298
 Weyl's estimates of $\zeta(1 + it)$, 200
 Wiener's Tauberian theorem, 280–281
 WIRSING, 288, 294, 297

X

Xi function $\xi(s)$, 16–18
 rate of growth, 41
 as transform of self-adjoint operator, 206–213

Z

Z function $Z(t)$, 119
 asymptotic formula for, *see* Riemann–Siegel asymptotic formula
 Zeta function
 analytic continuation of, 9–11, 16n
 evaluation by Euler–Maclaurin summation, 114–118
 as Fourier transform of summation operator, 204
 functional equation, 12–16, 222, 224–225, *see also* Euler, Riemann
 growth related to zeros, 182, 188, 190, 193, 200
 growth in strip $0 \leq \operatorname{Re} s \leq 1$, 182–201
 value of $\zeta'(0)/\zeta(0)$, 66–67, 134–135
 values at integers, 11–12
 values at negative integers, 216–217

This Page Intentionally Left Blank