

✓ Congratulations! You passed!

[Go to next item](#)

Grade received 100% Latest Submission Grade 100% To pass 70% or higher

1.

1 / 1 point



Which of the following is a valid step used during feature scaling?


- ☐ Add the mean (average) from each value and then divide by the (max - min).
- ☒ Subtract the mean (average) from each value and then divide by the (max - min).

✓ Correct

This is called mean normalization.

 2. Suppose a friend ran gradient descent three separate times with three choices of the learning rate α and plotted the learning curves for each (cost J for each iteration).

1 / 1 point


 For which case, A or B, was the learning rate α likely too large?

- ☐ Neither Case A nor B
- ☐ case A only
- ☐ Both Cases A and B
- ☒ case B only

✓ Correct

The cost is increasing as training continues, which likely indicates that the learning rate alpha is too large.

3. Of the circumstances below, for which one is feature scaling particularly helpful?

1 / 1 point

- ☐ Feature scaling is helpful when all the features in the original data (before scaling is applied) range from 0 to 1.
- ☒ Feature scaling is helpful when one feature is much larger (or smaller) than another feature.

✓ Correct

For example, the “house size” in square feet may be as high as 2,000, which is much larger than the feature “number of bedrooms” having a value between 1 and 5 for most houses in the modern era.

4.

1 / 1 point

You are helping a grocery store predict its revenue, and have data on its items sold per week, and price per item. What could be a useful engineered feature?

- ☐ For each product, calculate the number of items sold divided by the price per item.
- ☒ For each product, calculate the number of items sold times price per item.

✓ Correct

This feature can be interpreted as the revenue generated for each product.

5. True/False? With polynomial regression, the predicted values $f_{w,b}(x)$ does not necessarily have to be a straight line (or linear) function of the input feature x .

1 / 1 point

- ☒ True
- ☐ False

✓ Correct

A polynomial function can be non-linear. This can potentially help the model to fit the training data better.