

✔ **Congratulations! You passed!**

Grade received **85.71%** To pass 80% or higher

Go to next item

1. In this quiz you will diagonalise some matrices and apply this to simplify calculations.

0 / 1 point

Given the matrix  $T = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$  and change of basis matrix  $C = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$  (whose columns are eigenvectors of  $T$ ), calculate the diagonal matrix  $D = C^{-1}TC$ .

- ☐  $\begin{bmatrix} 5 & 0 \\ 0 & 4 \end{bmatrix}$
- ☐  $\begin{bmatrix} 6 & 0 \\ 0 & 3 \end{bmatrix}$
- ☒  $\begin{bmatrix} 9 & 0 \\ 0 & 20 \end{bmatrix}$
- ☐  $\begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$

✘ **Incorrect**

Be careful when multiplying matrices together.

2. Given the matrix  $T = \begin{bmatrix} 2 & 7 \\ 0 & -1 \end{bmatrix}$  and change of basis matrix  $C = \begin{bmatrix} 7 & 1 \\ -3 & 0 \end{bmatrix}$  (whose columns are eigenvectors of  $T$ ), calculate the diagonal matrix  $D = C^{-1}TC$ .

1 / 1 point

- ☒  $\begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix}$
- ☐  $\begin{bmatrix} 1 & 0 \\ 0 & -2 \end{bmatrix}$
- ☐  $\begin{bmatrix} 7 & 0 \\ 0 & 0 \end{bmatrix}$
- ☐  $\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$

✔ **Correct**

Well done!

3. Given the matrix  $T = \begin{bmatrix} 1 & 0 \\ 2 & -1 \end{bmatrix}$  and change of basis matrix  $C = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  (whose columns are eigenvectors of  $T$ ), calculate the diagonal matrix  $D = C^{-1}TC$ .

1 / 1 point

- ☐  $\begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix}$
- ☒  $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$
- ☐  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
- ☐  $\begin{bmatrix} 0 & 0 \\ 0 & -1 \end{bmatrix}$

✔ **Correct**

Well done!

4. Given a diagonal matrix  $D = \begin{bmatrix} a & 0 \\ 0 & a \end{bmatrix}$ , and a change of basis matrix  $C = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$  with inverse  $C = \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix}$ , calculate  $T = CDC^{-1}$ .

1 / 1 point

- ☐  $\begin{bmatrix} a & 0 \\ 0 & -a \end{bmatrix}$
- ☒  $\begin{bmatrix} a & 0 \\ 0 & a \end{bmatrix}$
- ☐  $\begin{bmatrix} -a & 0 \\ 0 & a \end{bmatrix}$
- ☐  $\begin{bmatrix} -a & 0 \\ 0 & 0 \end{bmatrix}$

$$\begin{bmatrix} 0 & -a \end{bmatrix}$$



Correct

Well done! As it turns out, because  $D$  is a special type of diagonal matrix, where all entries on the diagonal are the same, this matrix is just a scalar multiple of the identity matrix. Hence, given any change of co-ordinates, this matrix remains the same.

5. Given that  $T = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 5 & 0 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ -1 & 1 \end{bmatrix}$ , calculate  $T^3$ .

1 / 1 point

- ☐  $\begin{bmatrix} -61 & 3 \\ 122 & 186 \end{bmatrix}$
- ☐  $\begin{bmatrix} 122 & 186 \\ -61 & 3 \end{bmatrix}$
- ☒  $\begin{bmatrix} 186 & -61 \\ 122 & 3 \end{bmatrix}$
- ☐  $\begin{bmatrix} 3 & 122 \\ 186 & -61 \end{bmatrix}$



Correct

Well done!

6. Given that  $T = \begin{bmatrix} 2 & 7 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 7 & 1 \\ -3 & 0 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 0 & -1/3 \\ 1 & 7/3 \end{bmatrix}$ , calculate  $T^3$ .

1 / 1 point

- ☐  $\begin{bmatrix} -1 & 21 \\ 8 & 0 \end{bmatrix}$
- ☐  $\begin{bmatrix} 21 & 8 \\ 0 & -1 \end{bmatrix}$
- ☒  $\begin{bmatrix} 8 & 21 \\ 0 & -1 \end{bmatrix}$
- ☐  $\begin{bmatrix} 0 & -1 \\ 21 & 8 \end{bmatrix}$



Correct

Well done!

7. Given that  $T = \begin{bmatrix} 1 & 0 \\ 2 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$ , calculate  $T^5$ .

1 / 1 point

- ☐  $\begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}$
- ☒  $\begin{bmatrix} 1 & 0 \\ 2 & -1 \end{bmatrix}$
- ☐  $\begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$
- ☐  $\begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}$



Correct

Well done!