

TOPICS IN GRAVITATIONAL-WAVE ASTRONOMY

By
Prayush Kumar

A THESIS SUBMITTED IN
PARTIAL FULFILLMENT OF THE
REQUIREMENTS FOR THE DEGREE OF

DOCTOR OF PHILOSOPHY

IN
PHYSICS

at
Syracuse University
August 2014

TOPICS IN GRAVITATIONAL-WAVE ASTRONOMY

By
Prayush Kumar

A THESIS SUBMITTED IN
PARTIAL FULFILLMENT OF THE
REQUIREMENTS FOR THE DEGREE OF

DOCTOR OF PHILOSOPHY

IN
PHYSICS

at
Syracuse University
August 2014

Duncan Brown

Date

Graduate School Approval

Date

TOPICS IN GRAVITATIONAL-WAVE ASTRONOMY

By
Prayush Kumar

Syracuse University, 2014

Under the Supervision of Professor Duncan Brown

ABSTRACT

Gravitational waves are a consequence of the general theory of relativity. Direct detection of such waves will provide a wealth of information about physics, astronomy, and cosmology. A worldwide effort is currently underway to make the first direct detection of gravitational waves. The global network of detectors includes the Laser Interferometer Gravitational-wave Observatory (LIGO), which recently completed its sixth science run.

A particularly promising source of gravitational waves is a binary system consisting of two neutron stars and/or black holes. As the objects orbit each other they emit gravitational radiation, lose energy, and spiral inwards. This produces a characteristic “chirp” signal for which we can search in the LIGO data. Currently this is done using matched-filter techniques, which correlate the detector data against analytic models of the emitted gravitational waves. Several choices must be made in constructing a search for signals from such binary coalescences.

Any discrepancy between the signals and the models used will reduce the effectiveness of the matched filter. However, the analytic models are based on approximations which are not valid through the entire evolution of the binary. In recent years numerical relativity has had impressive success in simulating the final phases of the coalescence of binary black holes. While numerical relativity is too computationally expensive to use directly in the search, this progress has made it possible to perform realistic tests of the LIGO searches. The results of such tests can be used to improve the efficiency of searches.

Conversely, noise in the LIGO and Virgo detectors can reduce the efficiency. This must be addressed by characterizing the quality of the data from the detectors, and removing from the analysis times that will be detrimental to the search.

In this thesis we utilize recent results from numerical relativity to study both the degree to which analytic models match realistic waveforms and the ability of LIGO searches to make detections. We also apply the matched-filter search to the problem of removing times of excess noise from the search.

Duncan Brown

Date

© Copyright 2014

by

Prayush Kumar

*to
my parents*

Contents

Preface	x
Acknowledgments	xii
Conventions	xiii
List of Tables	xvii
List of Figures	xxxiii
1 Introduction	1
2 Gravitational waves from compact binaries	7
3 Search template banks for low-mass binary black holes in the Advanced gravitational-wave detector era	8
3.1 Introduction	8
3.2 Waveforms and Template Bank Placement	11
3.2.1 Waveform Approximants	11
3.2.2 Bank Placement metric	17
3.3 Results	19
3.3.1 EOBNRv2 templates placed using TaylorF2 metric	21
3.3.2 Effectualness of TaylorF2 templates	24
3.3.3 Effect of sub-dominant modes	27
3.4 Conclusions	33

4 Binary black hole search template banks with Numerical Relativity waveforms	35
4.1 Introduction	35
4.2 Waveforms	39
4.2.1 Numerical Relativity simulations	39
4.2.2 Post-Newtonian waveforms	40
4.2.3 PN-NR hybrid waveforms	41
4.2.4 Effective-One-Body model	42
4.3 Quantifying waveform accuracy & bank effectualness	43
4.4 Constructing a template bank for NR waveforms	48
4.5 Constructing a template bank for NR-PN hybrids	53
4.6 Complete NR-PN hybrid bank for non-spinning BBH	60
4.7 Conclusions	66
5 NINJA-2: Detecting gravitational waveforms modelled using numerical binary black hole simulations	71
5.1 Introduction	72
5.2 PN-NR Hybrid Waveforms	78
5.3 Modified Detector Noise	82
5.4 Injection Parameters	87
5.5 Search Pipelines	90
5.5.1 Coherent WaveBurst	91
5.5.2 <i>ihope</i>	91
5.5.3 Parameter estimation	94
5.6 Blind Injection Challenge Results	97
5.6.1 Coherent WaveBurst	97
5.6.2 <i>ihope</i>	98
5.7 Parameter Estimation Results	103
5.8 Sensitivity evaluation	105
5.9 Conclusion	110
6 Modeling of intermediate mass-ratio inspirals: Exploring the form of the self-force in the intermediate mass-ratio regime	119
6.1 Introduction	119

6.2	Effective One Body model	122
6.2.1	EOB dynamics	122
6.2.2	Modelling of the inspiral and plunge evolution	126
6.2.3	Merger and ring-down calibration of the EOB model	128
6.3	Self-force corrections in the intermediate-mass-ratio regime	130
6.3.1	IMRI self-force model	130
6.3.2	Radial evolution prescription	133
6.4	ISCO shift: connecting the extreme, intermediate and comparable-mass ratio regimes	144
6.5	Conclusions	150
6.6	Appendix	152
7	Self-forced evolutions of an implicit rotating source: a natural framework to model comparable and intermediate mass-ratio systems from inspiral through ringdown	155
7.1	Introduction	155
7.2	Modeling	160
7.2.1	Nomenclature	160
7.2.2	Inspiral evolution	161
7.2.3	Dissipative dynamics	166
7.2.4	Transition and plunge phases	172
7.2.5	Ringdown Waveform	177
7.2.6	Ringdown waveform construction in the context of an Implicit Rotating Source	179
7.2.7	Summary	181
7.3	Conclusions	185
8	Conclusions	188
	Bibliography	226

Preface

The work presented in this thesis stems from my participation in the LIGO Scientific Collaboration (LSC) and the NINJA-2 Collaboration. This work does not reflect the scientific opinion of the LSC and it was not reviewed by the collaboration.

Chapter 3 is based on material from

D. A. Brown, P. Kumar and A. H. Nitz, “Template banks to search for low-mass binary black holes with Advanced Gravitational-wave detectors,” *Phys. Rev. D* **87**, 082004 (2013)

Chapter 4 is based on material from

P. Kumar, I. MacDonald, D. A. Brown, H. P. Pfeiffer, K. Cannon, M. Boyle, L. E. Kidder, A. H. Mroue, B. Szilagyi and A. Zeninoglu, “Template banks for binary black hole searches with numerical relativity waveforms,” *Phys. Rev. D* **89**, 042002 (2014)

Chapter 5 is based on material from

The LIGO Scientific Collaboration, the Virgo Collaboration, the NINJA-2 Collaboration, “The NINJA-2 project: Detecting and characterizing gravitational waveforms modelled using numerical binary black hole simulations,” *arXiv:1401.0939* (2014)
Submitted to Classical and Quantum Gravity

Chapter 6 is based on material from

E. A. Huerta, P. Kumar, D. A. Brown, “Accurate modeling of intermediate-mass-ratio inspirals: Exploring the form of the self-force in the intermediate-mass-ratio regime,”

Phys. Rev. D **86**, 024024 (2012)

Chapter 7 is based on material from

E. A. Huerta, P. Kumar, J. R. Gair, S. T. McWilliams, “Self-forced evolutions of an implicit rotating source: a natural framework to model comparable and intermediate mass-ratio systems from inspiral through ringdown,” *arXiv:1403.0561* (2014)

Submitted to Phys. Rev. D

Acknowledgments

I have been fortunate to work with many wonderful people through my membership in the LIGO and NINJA collaborations. Sadly space makes it impossible to thank them all individually, and I hope the vast majority I must omit will forgive me.

I would first and foremost like to thank my advisor, Duncan Brown..

Conventions

We adopt the Einstein summation convention where repeated indices are summed over. Parenthesis are shorthand for the fully symmetric sum

$$A_{(\alpha\beta)} = \frac{1}{2} (A_{\alpha\beta} + A_{\beta\alpha})$$

and square brackets are fully antisymmetric

$$A_{[\alpha\beta]} = \frac{1}{2} (A_{\alpha\beta} - A_{\beta\alpha})$$

We take the signature of spacetime to be $(-1, +1, +1, +1)$.

Except where otherwise noted we work in geometric units where $G = c = 1$. We will often measure masses in multiples of the mass of the sun $1M_\odot \approx 1.99 \times 10^{30}$ kg

We define the Fourier transform of a function of time $g(t)$ to be $\tilde{g}(f)$, where

$$\tilde{g}(f) = \int_{-\infty}^{\infty} g(t) e^{-2\pi ift} dt$$

and the inverse Fourier transform is

$$g(t) = \int_{-\infty}^{\infty} \tilde{g}(f) e^{2\pi ift} dt$$

This convention differs from that used in some gravitational-wave literature, but is the adopted convention in the LIGO Scientific Collaboration.

The time-stamps of interferometer data are measured in Global Positioning System (GPS) seconds: seconds since 00:00.00 UTC January 6, 1980 as measured by an atomic clock.

List of Tables

1	SpEC BBH simulations used in this study. Given are symmetric mass-ratio η , mass-ratio $q = m_1/m_2$, and the length in orbits of the simulation. The last column gives the lowest total masses for which the NR simulations cover the entire coalescence process within the sensitive band of aLIGO, starting at 15 Hz.	40
2	List of mass-ratios, a template bank restricted to which will be effectual over the region of the non-spinning BBH mass space where $m_1 + m_2 \gtrsim 12M_\odot$ and $1 \leq q \leq 10$. The fraction of optimal SNR recovered by such a bank, accounting for discreteness losses, remains above 98%. This is shown in Fig. 17.	63
3	Submissions to NINJA-2	83
4	Submissions to NINJA-2	84

7	The <code>ihope</code> search results. The Event IDs correspond to the Event ID of each blind injection given in Table 5; this association is based on the time of the candidates relative to the time of the injections. The cFARs are calculated from all possible 5 s time shifts over the entire two-month dataset. M and q give the total mass and mass ratio respectively that were recovered in each detector. The recovered SNR (ρ_{rec}) and re-weighted SNR ($\hat{\rho}$) are reported separately for each detector. To calculate the cFARs, the quadrature sum of $\hat{\rho}$ was used. Unless noted, the cFARs were calculated after category 1-3 vetoes were applied.	99
8	Summary of the mass and spin estimates for events 1-7 for each of the models used. Median values are quoted along with the upper and lower bounds of the 95% credible intervals. The injected values for each event are included for reference.	106
9	Summary of the sky location estimates for events 1-7 for each of the models used. Median values are quoted along with the upper and lower bounds of the 95% credible intervals, as well as the area of the 95% credible regions in the sky. The injected values for each event are included for reference.	107
10	Ω_{ansatz} fit coefficients	133
11	\dot{L}_z fit coefficients	134
12	$z_{\text{SF}}(x)$ fit coefficients	135

13	The Table shows the value of the angular momentum L_z as a function of the gauge invariant object $x = (M\Omega)^{2/3}$ evaluated at the ISCO radius (see Eq. (6.28)). The SF (self-force) values are computed using the prescription given in Eq. (6.32) to compute the orbital frequency shift, and Eq. (6.28) with the coefficients b_i quoted in [280], i.e., evaluated in the context of extreme-mass-ratio inspirals. Incomplete stands for a prescription in which we use the prescription for the orbital frequency given by Eq. (6.38), and the angular momentum prescription given by Eq. (6.28) with the coefficients b_i quoted in [280]. The b_i values of the IMRI model are obtained using a model that reproduces the dynamical evolution of intermediate-mass-ratio inspirals, as compared with the EOBNRv2 model, and for which we have derived a new prescription for the red-shift observable $z_{\text{SF}}(x)$ which actually reproduces the features of true inspirals.	147
14	The table summarizes the nomenclature we will use throughout our analysis.	161

List of Figures

1	This figure shows the effectualness of a bank of EOBNRv2 templates, placed using the 2PN accurate hexagonal template placement of Ref. [63], to search for a population of BBH signals simulated with EOBNRv2 waveforms. The masses of the BBH population are chosen from a uniform distribution of component masses between 3 and $25 M_{\odot}$. For each injection, we plot the component masses of the injection, and the fitting factor (\mathcal{FF}).	22
2	This figure shows a cumulative histogram of the fraction of the BBH signal space (on the y-axis), where the bank of EOBNRv2 waveforms has \mathcal{FF} less than the respective values on the x-axis. The EOBNRv2 bank has a fitting factor \mathcal{FF} below the 0.97 for less than $\sim 1.5\%$ of all simulated signals with component-masses m_1, m_2 between $3 M_{\odot}$ and $25 M_{\odot}$	23
3	The fitting factor \mathcal{FF} of a bank of TaylorF2 waveforms, constructed with $MM = 0.97$, for a population of BBH systems which are modeled using EOBNRv2 signals.	25
4	The (blue) curve shows the upper-bound on total-mass for the sub-region over which the TaylorF2 bank has a minimal-fitting-factor as given on the x-axis. We observe that the TaylorF2 bank has a minimal-fitting-factor of 0.965 (0.947) for the region with total masses below $\sim 11.4 M_{\odot}$ ($19 M_{\odot}$). The minimal-fitting-factor is the fitting-factor value which is less than the fitting-factors of the TaylorF2 bank for $\geq 99.75\%$ of the points sampled in the sub-region.	26

- 7 (left) This figure shows $\epsilon_V(\theta_1 = \{m_1, m_2\})$ in the component-mass space (see Eq. 3.29). This gives the fraction of total observable volume that is visible to a search which uses the leading order $l = m = 2$ EOBNRv2 waveform template bank, placed with the 2PN accurate TaylorF2 bank placement metric. For a population of signals, that is distributed uniformly in spacial volume, this is equivalent to the fraction of the maximum possible event observation rate that we get with the use of a discrete bank of matched-filters. We observe that the loss in event observation rate, averaged over all parameters (uniformly distributed) but $\theta_1 = \{m_1, m_2\}$, does not exceed $\sim 11\%$ for any region of the component-mass space. (right) This figure shows $\epsilon_V(\theta_1 = \{q, \iota\})$ over the $q - \iota$ plane. We note that the maximum averaged loss in the detection rate is for systems with high mass ratios *and* $\iota \in [1.08, 2.02]$, and can go as high as $\sim 20\%$ for such systems.

8 We show the *true* (upper) and the *hybrid* (lower) waveform manifolds here, with the signal residing in the former, and a discrete bank of templates placed along lines of constant mass-ratio in the latter. Both manifolds are embedded in the same space of all possible waveforms. The true signal waveform is denoted as h_x^{tr} , while the templates in the bank are labelled h_b^M . The hybrid waveform that matches the signal H_x^{tr} best is shown as h_{\perp}^M . Also shown is the “distance” between the signal and the hybrid template that has the highest overlap with it. This figure is qualitatively similar to Fig. 3 of Ref. [91].

9 The color at each point gives the number of waveform cycles N_{cyc} , for that particular binary, which contain 99% of the signal power in the aLIGO sensitivity band. The figure is truncated to exclude the region where $N_{\text{cyc}} > 40$. The solid curve shows the lower bounding edge of the region with $\mathcal{M}_c = 27M_{\odot}$

- 10 The color at each point in the figure gives the value of $\mathcal{FF} \simeq 1 - \Gamma_{\text{bank}}$ of the bank for that binary, for the NR bank restricted to $\mathcal{S}_q = \{1, 2, 3, 4, 6, 8\}$. This is the same as the fraction of the optimal SNR, for the binary, that the template bank recovers. The black dots show the location of the templates in the bank. We note that they all lie along straight lines of constant q passing through the origin. The region shaded light-grey (towards the bottom of the figure) is where the \mathcal{FF} drops sharply below 97%.

11 This figure is similar to Fig. 10. The color at each point gives the value of $\mathcal{FF} \simeq 1 - \Gamma_{\text{bank}}$ of the bank for that binary, for the NR bank restricted to $\mathcal{S}_q = \{1, 2, 3, 4, 6, 9.2\}$. The black dots show the location of the templates in the bank. The region shaded light-grey (towards the bottom of the figure) is where the \mathcal{FF} drops below 97%. We note that with an additional NR waveform for mass ratio $q = 9.2$, the coverage of the bank is extended to include binaries with $10 \leq q \leq 11$

12 Bounds on mismatches of PN-NR hybrid waveforms, for the currently existing NR simulations. The PN error is for hybrids matched at $M\omega_m = 0.025$ for $q = 1$, $M\omega_m = 0.038$ for $q = 2$, and $M\omega_m = 0.042$ for $q = 3, 4, 6, 8$. The black circles indicate the lower bound of the template bank in Table 1. The black square show the lower bound with a hybrid error of 1.5%. The inset shows these lower bounds as a function of mass ratio.

- 13 These figures show fitting factors \mathcal{FF} obtained when using a discrete mass-ratio template bank for $q = 1, 2, 3, 4, 6, 8$. For each mass-ratio, the templates are extended down to a total mass where the NR-PN hybridization mismatch becomes 3%. The bank is placed using the stochastic algorithm, similar to Ref. [126, 127, 128]. The black dots show the location of the templates. The fitting factor on the left plot does *not* take into account the hybridization error, and therefore shows the effect of the sparse placement of the templates alone. The right plot accounts for the hybridization error and gives the actual fraction of the optimal SNR that would be recovered with this bank of NR-PN hybrid templates. The region bounded by the magenta (solid) line in both plots indicates the lower end of the coverage of the bank of un-hybridized NR waveforms. Lastly, the shaded grey dots show the points where the fitting factor was below 96.5% 56
- 14 These figures are similar to Fig. 13. The figures show fitting factors \mathcal{FF} obtained when using a discrete mass-ratio template bank for $q = 1, 2, 3, 4, 6, 8$. For each mass-ratio, the templates are extended down to a total mass where the NR-PN hybridization mismatch becomes 3%. Templates are placed independently for each mass-ratio, and span the full range of total masses. For each mass-ratio, neighboring templates are required to have an overlap of 97%. The union of the six single- q one-dimensional banks is taken as the final bank. The black dots show the location of the templates. The fitting factor on the left plot does *not* take into account the hybridization error, and therefore shows the effect of the sparse placement of the templates alone. The right plot accounts for the hybridization error and gives the actual fraction of the optimal SNR that would be recovered with this bank of NR-PN hybrid templates. The region bounded by the magenta (solid) line in both plots indicates the lower end of the coverage of the bank of un-hybridized NR waveforms. Lastly, the shaded grey dots show the points where the fitting factor was below 96.5% 57

- 15 This figure is similar to Fig. 14. The figures show fitting factors \mathcal{FF} obtained when using a discrete mass-ratio template bank for $q = 1, 2, 3, 4, 6, 8$. Templates are placed independently for each mass-ratio, and span the range of total masses, down to the region where the hybrid errors become 3%. For each mass-ratio, neighboring templates are required to have an overlap of 97%. The union of the six single- q one-dimensional banks is taken as the final bank. The black dots show the location of the templates. The GW signals are modeled using the EOBNRv2 approximant [38], while TaylorT4+NR hybrids are used as templates. The fitting factor on the left plot shows the combined effect of the sparse placement of the templates, and the (relatively small) disagreement between the hybrid and EOBNRv2 waveforms. The right plot explicitly accounts for the hybridization error and gives the (conservative) actual fraction of the optimal SNR that would be recovered with this bank of NR-PN hybrid templates. The region bounded by the magenta (solid) line in both plots indicates the lower end of the coverage of the bank of un-hybridized NR waveforms. Lastly, the shaded grey dots show the points where the fitting factor was below 96.5%.
- 16 These figures show the coverage of template banks restricted to single mass-ratios, i.e. (from left to right) $q = 1, 4, 8$. We note that at higher total masses, the templates are correlated to simulated signals for considerably different mass-ratios, than at lower total masses. This agrees with what we expect as with decreasing total mass, the number of cycles in the sensitive frequency band of Advanced LIGO increases.

58

61

17	This figure shows fitting factors for a hybrid template bank which samples from the 26 mass ratios $q = 1, 1.5, 1.75, 2, \dots, 9.6$, and allows coverage to masses down to $m_1 + m_2 = 12M_\odot$ and $1 \leq q \leq 10$, with a minimal-match of 98% at the lowest masses. The left and right panel show the same on $M - q$ and $m_1 - m_2$ axes, respectively. The magenta lines, in both panels, shows the upper bound in total mass, below which frequency-domain PN waveforms can be used to construct template banks for aLIGO searches [73, 123]. The dash-dotted line in the right panel shows the lower mass limit on the smaller component object, to which a bank of currently available NR-PN hybrids can cover, i.e. $\min(m_1, m_2) = 12M_\odot$ (see Sec. 4.5). The blue (solid) curve in the right panel gives the lower mass limit to which a bank of currently available NR waveforms can cover (see Sec. 4.4). Thus, between the simulations listed in Table 2, and frequency domain PN waveforms, we can search for the entire range of BBH masses.	62
18	The maximum mismatch between different PN approximants for hybrid waveforms plotted against the total mass for at different matching frequencies ($M\omega_m$). The dotted lines indicate a mismatch of 1.5% and a lower total mass limit, $12M_\odot$ for $q = 1$, and $M_2 = 3M_\odot$ for $q = 4, 8$. The thick dashed lines indicate the currently possible matching frequency for hybrids based on the length of NR waveforms. The numbers next to each line indicate the number of orbits before merger where the PN and NR (or EOB) waveforms were stitched together.	64
19	This plot shows the lower mass limit of a template bank constructed with hybrid waveforms in terms of the number of NR orbits (left panel) and initial gravitational wave frequency (right panel) needed to have a PN error below 1.5% (solid curves) or 3% (dashed curves). The dotted line indicates the lower total mass limit when one component mass is $3M_\odot$	65
20	Parameters of the NINJA-2 submissions	79

21	Left: predicted sensitivity curves for aLIGO and AdV. Shown are both the design curves and predicted 2015 – 2016 <i>early</i> sensitivity curves. Also shown is the early AdV noise curve rescaled such that the horizon distance for a ($10 M_{\odot}$, $10 M_{\odot}$) binary system is equal to that obtained with the early aLIGO noise curve. Right: Horizon distance as a function of observed total mass for the early aLIGO and rescaled early AdV sensitivity curves. This plot is made considering only equal mass, non-spinning systems and calculated using the <code>EOBNRv2</code> [166] waveform approximant. Results in this paper are generated from the early aLIGO noise curve and the rescaled early AdV curve.	85
22	Sensitivity curves of the recolored data for the LIGO Hanford detector (left) and the Virgo detector (right). In both cases the black dashed line shows the predicted 2015 – 2016 sensitivity curve (with the scaling factor added for Virgo). The dark colored region indicates the range between the 10 % and 90 % quantiles of the PSD over time. The lighter region shows the range between the minimum and maximum of the PSD over time.	87
23	SNR time series in a 20 s window around a known glitch in the original data (left) and in the recolored data (right). While the SNR time series clearly change, the primary features of the glitch are preserved across the recoloring procedure. These SNR time series were obtained by matched-filtering a short stretch of recolored and original data against a ($23.7, 1.3$) M_{\odot} template.	88
24	SNR distribution	102
25	The 95% credible regions for the <code>EOBNRv2</code> (blue), <code>PhenomB$_{\chi=0}$</code> (red), and <code>PhenomB</code> (green) models. Injected values are indicated by the “x” and the neutron star and mass gap regions are indicated where relevant. A strong degeneracy between spin and mass ratio results in systematic biases and artificially strong constraints on mass estimates when spin is ignored (i.e. <code>EOBNRv2</code> , <code>PhenomB$_{\chi=0}$</code>). By accounting for spin the <code>PhenomB</code> model produces estimates consistent with the injected sources.	113

26	The 90% (dashed), 95% (solid), and 99% (dotted) credible regions of the two-dimensional marginalized posterior for the mass ratio and effective spin χ using the <code>PhenomB</code> model. The strong correlation between mass ratio and spin is responsible for the systematically weaker constraints placed on component masses when analyzing signals with an aligned-spin model.	114
27	(a) shows a spectrogram of the noise in Hanford centered on the end time of the injection (before the blind injection is added). A non-Gaussian feature is present with peak energy slightly after the end time of the injection. (b) shows the 95% credible regions for the <code>EOBNRv2</code> analysis of the event 1 injection in real noise (solid) and noiseless data (dashed). Injected values are indicated by the “x”. The fact that the injected values are well outside of the 95% credible region when real noise is present, and not in noiseless data, leads to the conclusion the non-Gaussian feature in the noise led to significant systematic biases in parameter estimates.	115
28	(Top) Evidences for the non-spinning (<code>EOBNRv2</code> and <code>PhenomB$_{\chi=0}$</code>) and spin-aligned (<code>PhenomB</code>) models. (Bottom) Bayes factors showing the support for the spin-aligned model over the non-spinning <code>PhenomB$_{\chi=0}$</code> model. Events 3 and 4 are found to be spinning with high certainty. Event 7, with the largest spin magnitude, has little support for spin, though this is likely due to its low SNR.	116
29	95% credible regions for the sky position of all seven events, using the <code>EOBNRv2</code> (blue), <code>PhenomB$_{\chi=0}$</code> (red), and <code>PhenomB</code> (green) models. Despite the substantial differences between these models, their sky localization ability is very consistent.	117

- 30 Volume-weighted average sensitive distance as a function of total mass, for a variety of NR waveforms. The sensitive distance is defined as the volume-weighted average distance up to which a CBC signal with a given waveform and total mass can be detected, where we average over the source extrinsic parameters and over the varying detector noise levels in the data set via Monte Carlo integration. *Top left*—Waveforms with equal component masses and various spins. Here the dotted lines represent predicted performances if an aligned-spin search had been performed. Details of this prediction are given in section 5.8. *Top right*—Waveforms with zero component spins and various mass ratios. *Bottom left*—Waveforms with equal-mass, non-spinning components. The dashed black line represents the predicted sensitivity curve of non-spinning waveforms with the aLIGO sensitivity curve, as described in 5.8. The dotted black line represents the predicted sensitivity with the AdV sensitivity curve. *Bottom right*—Waveforms with unequal spins, or unequal component masses and non-zero spins. The waveform designations are given in Table 3. These plots were generated using the data described in section 5.3 and the distribution of signals described in section 5.4. 118
- 31 The panels show the orbital frequency, as a function of the radial coordinate r , obtained using the fit described in the text (solid red line), and the orbital frequency predicted by the EOB model (solid black line). The systems shown in the panels correspond to binaries of component masses $17M_{\odot} + 100M_{\odot}$ (top-left panel), $10M_{\odot} + 100M_{\odot}$ (top-right) and $1M_{\odot} + 100M_{\odot}$ (bottom-left panel). The bottom-right panel demonstrates the accuracy with which our IMRI model reproduces the orbital frequency predicted by the EOBNRv2 for the systems 17:100 (dashed blue), 10:100 (dashed-dot red) and 1:100 (solid black). Note that the spikes are due to artifacts in the interpolation function used to plot the EOBNRv2 orbital frequency and the numerical fit used to reproduce it. Also notice that the discrepancy between the data and the fit is always smaller than one part in a thousand. 138

- 32 We compare the calibrated flux of angular momentum described in Eq. (6.26) against the prediction of the EOBNRv2 model for binaries of component masses $17M_{\odot} + 100M_{\odot}$ (top left panel), $10M_{\odot} + 100M_{\odot}$ (top right panel), and $1M_{\odot} + 100M_{\odot}$ (bottom-left panel). The bottom-right panel shows the accuracy with which the IMRI model reproduces the EOBNRv2 angular momentum flux, $\dot{L} = dL_z/dt$, for the systems: 17:100 (dashed blue), 10:100 (dashed-dot red) and 1:100 (solid black). The spikes in the bottom-right panel are due to numerical artifacts of the interpolating function used to plot the EOBNRv2 angular momentum flux and the numerical fit to reproduce it. The fit is such that its discrepancy to the EOBNRv2 is always smaller than one part in a thousand. 139
- 33 The left panel shows the angular momentum flux for binaries of mass ratio 1:100 computed using the EOBNRv2 model and the fit to Teukolsky data introduced in [282]. Both formalisms present a remarkable agreement all the way down to the ISCO. Note that the angular momentum flux fit to Teukolsky data is valid only from early inspiral until the ISCO. The right panel shows the relative difference between the two prescriptions for the angular momentum flux $\dot{L} = dL_z/dt$ 140
- 34 The panels show the radial and azimuthal evolution for a $17M_{\odot} + 100M_{\odot}$ system (top panels), $10M_{\odot} + 100M_{\odot}$ system (middle panels), and $1M_{\odot} + 100M_{\odot}$ system (bottom panels), obtained using the IMRI model compared against the EOBNRv2 model. 141
- 35 The panels show the accuracy with which the IMRI model proposed in the paper reproduces the radial (right panel) and azimuthal (left-panel) time evolution predicted by the EOBNRv2 model for the systems $17M_{\odot} + 100M_{\odot}$ (dashed blue), $10M_{\odot} + 100M_{\odot}$ (dashed-dot red), and $1M_{\odot} + 100M_{\odot}$ (solid black). Notice that our model reproduces the EOBNRv2 orbital and azimuthal evolution point to point with an accuracy better than one part in a thousand. 142

- 36 The panels show the radial evolution using the EOBNRv2 model, the IMRI model described in the text, and a model (Unfitted) which is the same as the IMRI model described in text except for the fact that the coefficients used in Eq. (6.29) for the function $z_{\text{SF}}(x)$ were derived using available self-force data from extreme-mass ratio calculations [280]. The plots correspond to binaries of component masses $17M_{\odot} + 100M_{\odot}$ (top-left panel), $10M_{\odot} + 100M_{\odot}$ (top-right panel) and $1M_{\odot} + 100M_{\odot}$ (bottom-left panel). The bottom-right panel shows that the orbital evolution predicted by a model that incorporates self-force corrections from extreme-mass-ratio inspirals (EMRIs) deviates from the orbital evolution predicted by the EOBNRv2 model at late inspiral. This discrepancy is more noticeable for systems with mass-ratios 17:100 (dashed blue) and 10:100 (dashed-dot red). Furthermore, for smaller mass-ratios, 1:100 (solid black), the discrepancy becomes comparatively smaller, as expected. 143
- 37 ISCO shift using various approximations as described in the text. The ‘Numerical Data’ has been obtained using calculations in the extreme-mass ratio regime [272] and the EOBNRv2. The prescription that encapsulates results from the extreme, intermediate and comparable mass-ratio regime is labeled as ‘Numerical Fit’ and is given by Eq. (6.38) in the main text. 146

38 The panels show the invariant angular momentum $L_z(x)$, with $x = (M\Omega)^{2/3}$, from early inspiral to the ISCO using two different prescriptions. The ‘IMRI’ prescription reproduces accurately the expected inspiral evolution, as compared with results from the EOBNRv2 model —which has been calibrated to NR simulations. The ‘Self-Force’ prescription encapsulates self-force corrections derived in the context of EMRIs [278, 280]. Note that for the three binary systems studied, $17M_\odot + 100M_\odot$ (top-left panel), $10M_\odot + 100M_\odot$ (top-right panel), and $1M_\odot + 100M_\odot$ (bottom-left panel), the ‘Self Force’ prescription for the angular momentum presents a deviation from its expected value that becomes more noticeable near the ISCO. The bottom-right panel shows the fractional accuracy between the two different prescriptions for the angular momentum for the systems 17:100 (dashed blue), 10:100 (dashed-dot red) and 1:100 (solid black). Notice also that, as expected, for binaries with small mass-ratio the ‘Self Force’ and ‘IMRI’ prescriptions are fairly consistent all the way down to the ISCO. As in the previous plots, the spikes in the bottom-right panel are due to numerical artifacts.

149

39 The panels show the accuracy with which our model can reproduce the dynamics of an equal mass (EM) binary system with masses ($10M_\odot + 10M_\odot$), as compared with the EOBNRv2 model. The top panels show that our model can reproduce the orbital angular evolution and the flux of angular momentum predicted by the EOBNRv2 model point to point with an accuracy better than one part in a thousand. The middle panels show that our model can reproduce with a similar accuracy the radial and azimuthal evolution of a $10M_\odot + 10M_\odot$ binary system. The bottom panel shows that the expression for the gauge-invariant angular momentum $L_z(x)$ for an EM mass deviates clearly from the EMRI prescription even before nearing the ISCO.

154

- 40 The panel shows the expected sensitivity for two configurations of the Einstein Telescope (ET), namely, ETD (black), ETB (blue) and LIGO’s Zero Detuned High Power (ZDHP) configuration (red). The vertical axis measures $F_{\text{normalized}} = (f/f_{\max})^{-7/6} \sqrt{S_n(f_{\max})/S_n(f)}$, where f_{\max} is the maximum of the corresponding power spectral density, $S_n(f)$ 157
- 41 The phase discrepancy in radians between the PN approximant TaylorT4, and the Effective One Body model, shown as a function of time from $r = 30M$ to the point when the TaylorT4 model reaches the ISCO. The systems have mass-ratio, q , total mass, M_{tot} , and final phase discrepancy, $\Delta\Phi$: $(q, M_{\text{tot}}, \Delta\Phi) = (1/6, 7M_{\odot}, 21.5 \text{ rads})$ (top-left), $(1/8, 9M_{\odot}, 30.2 \text{ rads})$ (top-right), $(1/10, 11M_{\odot}, 70.1 \text{ rads})$ (bottom-left) and $(1/15, 16M_{\odot}, 83.2 \text{ rads})$ (bottom-right) respectively. 159
- 42 The panels show the energy and angular momentum given by Eqs. (7.13)-(7.14), respectively. We show the functional form of these parameters for binary systems with mass-ratio values, from top to bottom, $q \in [0, 1/100, 1/20, 1/10, 1/6, 1/5, 1]$ 164
- 43 The location of the innermost stable circular orbit is determined by the condition $dE/dx = 0$. The panel shows dE/dx as a function of the gauge invariant quantity $x = (M\Omega)^{2/3}$. The various curves represent binary systems with mass-ratios, from top to bottom, $q \in [0, 1/100, 1/20, 1/10, 1/6, 1/5, 1]$. Note that binaries with mass-ratios $q \gtrsim 1/6$ do not have an ISCO in this model. 165
- 44 The (top, bottom) panels show the results of the optimization runs that were used to constrain the values of the b_i coefficients given in Eq. (7.21). The panels show the results for binaries of mass-ratio $q \in [1/6, 1/8]$, and total mass $M_{\text{tot}} = [7M_{\odot}, 9M_{\odot}]$. The ‘optimal’ value for the coefficients has been chosen by ensuring that the flux prescription minimizes the phase discrepancy between the EOB model and our self-force model. The color bar shows the phase difference squared between both models, which is integrated from $r = 20M$ all the way to the light-ring. 170

45	The panels show the orbital phase evolution of a self-force model making use of optimized PN energy flux given by Eq. (7.20) and the phase evolution as predicted by the EOB model. The [top/bottom] panels exhibit this evolution for a compact binary with mass–ratio $q = [(1/6, 1/8), (1/10, 1/15)]$, and total mass $M_{\text{tot}} = [(7M_{\odot}, 9M_{\odot}), (11M_{\odot}, 16M_{\odot})]$, respectively.	171
46	Left panel: The object sits at the minimum of the effective potential, Eq. (7.32), which corresponds to the case $\xi = L - L_{\text{ISCO}} \gg 0$. Right panel. Blue (top) curve: radial geodesic motion, which corresponds to $\xi = L - L_{\text{ISCO}} \gg 0$; Red (middle) curve: the object nears the ISCO and the orbit shrinks due to radiation reaction. Note that the minimum of the potential has moved inwards ($\xi = 0.35$). Yellow (bottom) curve: body’s inertia prevents it from staying at the minimum of the potential, and adiabatic inspiral breaks down ($\xi = 0$). At this point the transition regime takes over the late inspiral evolution [288]. Note: this plot is based on Figure 1 of [288].	173
47	The left panel shows the effective potential for a Schwarzschild BH without including finite mass–ratio corrections. Note that the minimum of the potential takes place at the ISCO, which can be determined using Eq. (7.18). The right panel exhibits the influence of finite mass–ratio corrections on the effective potential used to modify Ori and Thorne transition regime [288]. The curves represent binaries, from top to bottom, with mass–ratios $q \in [0, 1/100, 1/20, 1/10, 1/6, 1/5, 1]$	174
48	(Top, bottom) panels: the left panel shows the inspiral, transition and plunge radial evolution for a BH binary of mass–ratio $q = (1/6, 1/8)$ — and total mass $M_{\text{tot}} \in (7M_{\odot}, 9M_{\odot})$ — using the coordinate transformation given by Eq. (7.15). The right panel shows the orbital frequency $M\Omega$ from late inspiral all the way to the light ring. The evolution starts from an initial radial value $r = 30M$	176
49	As in Figure 48, but with the (top, bottom) panels showing the radial and orbital frequency evolution for binaries with mass–ratios $q = (1/10, 1/15)$, and total mass $M_{\text{tot}} \in (11M_{\odot}, 16M_{\odot})$, respectively. As before, the evolution starts from an initial radial value $r = 30M$	177

- 50 The panels show sample waveforms from inspiral to ringdown for systems with mass-ratios $q \in [1/6, 1/8]$ —and total mass $M_{\text{tot}} \in (7M_{\odot}, 9M_{\odot})$ — (top panels—from left to right) and $q \in [1/10, 1/15]$ —and total mass $M_{\text{tot}} \in (11M_{\odot}, 16M_{\odot})$ — (bottom panels —from left to right). The inspiral evolution for the [top, bottom] panels starts from $r = [30M, 25M]$ 180
- 51 The panels show sample the late-time evolution of waveforms whose ringdown phase is modeled using the implicit rotating source (IRS) model and a sum of quasinormal modes (QNMs). The systems shown correspond to binaries with mass-ratios $q \in [1/6, 1/8]$ —and total mass $M_{\text{tot}} \in (7M_{\odot}, 9M_{\odot})$ — (top panels—from left to right) and $q \in [1/10, 1/15]$ —and total mass $M_{\text{tot}} \in (11M_{\odot}, 16M_{\odot})$ — (bottom panels —from left to right). The inspiral evolution for the [top, bottom] panels starts from $r = [30M, 25M]$ 181

Chapter 1

Introduction

Gravitational waves, as predicted by General Relativity, are ripples in the curvature of spacetime that propagate at the speed of light. Like water waves on the ocean, the concept of gravitational wave requires an ideal, i.e. a smooth and unperturbed background on which the waves propagate. Unlike water waves, however, gravitational waves are not the motion of a material medium, but are ripples in the fabric of spacetime itself. The phenomena can be intuitively described with an analogy to electromagnetism. When an electromagnetic wave is incident upon a charged particle, it accelerates the particle in a direction transverse to its own direction of propagation. This acceleration depends on the electric charge and the inertial mass of the particle. Incident gravitational waves, similarly, produce an acceleration that is transverse to their direction of propagation; with the difference that the acceleration produced is independent of the mass of the particle experiencing it and will be the same for any particle at that point. Free falling particles also experience the same acceleration, and an observer sitting in their frame, also known as a local inertial frame, will not be able to measure the local acceleration. It is only through non-local measurements between points that are separated in spacetime that one would be able to discern the effect of a passing gravitational wave.

Gravitational waves carry energy away from their source. They couple minimally with matter, which allows for the preservation of information from distant sources. This will enable us to do precision observation and characterization of astrophysical sources. On the other hand, the relatively weak coupling with matter also makes

them demonstrably quite difficult to detect directly in a laboratory. The first indirect detection of gravitational waves was made by Russell Hulse and Joseph Taylor, when they discovered a binary system with a pulsating neutron star PSR 1913+16 in 1974 [1]. They observed pulses of radio waves emitted by the pulsar, and through precise measurement of their timing, they were able to determine the rate at which the binary orbit was shrinking. Orbital energy is directly co-related with the separation of the objects in the binary, and their measurement allowed them to calculate the rate at which the binary was losing energy. It was soon shown that this rate agreed within a percent with that predicted by General Relativity [2, 3]. Hulse and Taylor were awarded the Nobel prize in Physics in 1993 for this discovery. In the following years, there have been similar pulsar binaries have been discovered, allowing for more precise measurements of the effect of gravitational waves as predicted by General Relativity [4].

Felix Pirani first suggested the use of an interferometer for precise measurement of the distance between two test masses in 1956 [5]. In 1971, Moss, Miller and Forward built and tested the first laser interferometric gravitational wave transducer [6]. The design of the modern interferometric detectors are built on the contributions of Weiss [7] and Drever [8] from 1970s. The core of such detectors is a Michelson interferometer, with two laser cavities oriented perpendicular to each other. The end mirrors of each cavity is held vertically, and are practically freely falling in the transverse direction. Incoming gravitational waves will cause different changes in the length of each cavity, and these instruments measure with great precision the change in the difference in the length of both cavities [9]. There is currently a concerted effort worldwide towards the direct detection of gravitational waves, and several ground based gravitational wave observatories are being planned and built. These include:

1. Laser Interferometer Gravitational-wave Observatory (LIGO): This United States funded project comprises of two independent detectors, one located in Hanford, Washington and the other in Livingston, Louisiana.
2. Virgo observatory: A French-Italian project based in Cascina, Italy.
3. GEO 600, near Sarstedt, Germany.
4. The Kamioka Gravitational Wave Detector (KAGRA), formerly the Large-scale

Cryogenic Gravitational-wave Telescope (LCGT), is a Japanese project, and has begun construction in the tunnels of Kamioka mines in Japan.

5. LIGO-India: This is a tentative project between LIGO and several institutions in India, aimed at having a third LIGO-like instrument in the eastern hemisphere.

The first-generation LIGO and Virgo detectors were constructed in stages, and began observation in 2002 and 2007, respectively. There were 6 observation runs performed with LIGO, called Science run 1 (S1) to Science run 6 (S6). In 2007, during S5, LIGO reached its design sensitivity, at which time it was sensitive to gravitational waves from systems similar to the Hulse-Taylor binary pulsar up to distances of FILLME Mpc . For comparison, the Virgo supercluster, of which our own galaxy is a part, has a diameter of 33 Mpc. In the sixth Science run, the sensitivity of the instruments was improved by increasing the laser power in its cavities, adding an output-mode cleaner and implementing other technological upgrades. This *enhanced* LIGO operated from July 2009 to October 2010, being sensitive to similar fiducial systems out to FILLME Mpc . Similarly, the Virgo detector observed in 3 Virgo Science Runs (VSR1-3), extending over late 2007 to 2011. At the peak of their combined sensitivity, the three detectors were sensitive to Hulse-Taylor type binaries out to FILLME Mpc . While no detection was made, the LIGO-Virgo Scientific Collaboration (LVC) was able to put upper limits on the rate of compact object binary mergers within its range. Results from these observation periods have been published since, e.g. [10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24].

The second generation Advanced LIGO (aLIGO) and Virgo detectors are currently under construction and commissioning [25, 26]. The two aLIGO detectors are expected to begin operation late next year, and plan to achieve their design sensitivity by 2019 [27]. The Virgo detector is also being upgraded on a similar timeline. These upgrades will increase the sensitivity of both detectors in two important ways, (i) provide a factor of 10 improvement in sensitivity across the entire LIGO frequency band, and (ii) lower the lower frequency bound from 40 Hz down to about 15 Hz [25]. These enhancements would mean a thousandfold increase in the volume of the universe that we would be able to probe for source of gravitational waves. Astrophysical estimates and numerical simulations suggest that the Advanced detectors would detect about

40 mergers a year of Hulse-Taylor type compact binaries [28].

Gravitational wave astronomy can probe regions that the electromagnetic spectrum cannot, like black holes, interiors of neutron stars, supernovae mechanism etc. They also offer the unique avenue of probing gravity in the strong-field regime, providing a valuable test of General Relativity. There are several astrophysical sources of gravitational waves, such as supernovae explosions, cosmic strings, black hole ring-downs, etc. Binaries of stellar-mass compact objects are of special interest for ground based detectors like LIGO and Virgo as they emit gravitational radiation in the operating frequency range of the detectors, which extends from about 15 Hz to a few kHz. As binary stars evolve and undergo supernovae, a significant fraction of them get disrupted. The ones that survive form binaries containing neutron stars and/or black holes. While the mass of neutron stars are observationally constrained between $1 - 3M_{\odot}$, the upper bound on the mass of stellar-collapse black holes is something that can also be better determined with gravitational wave observations.

Searches for gravitational waves from compact binaries operate by matched-filtering the detector data to dig out the weak gravitational wave signatures that are otherwise buried in the instrumental noise. They take advantage of the fact that we can theoretically model the expected form of the gravitational wave signatures, called *waveforms*, and use those as filtering templates. While an exact analytic solution in General Relativity for the dynamics of compact binary systems has not been found, several perturbative formalisms exist that are capable of describing the inspiral dynamics. In the slow-motion large-separation post-Newtonian (PN) approximation [29], various orbital quantities are calculated as Taylor series in the binary velocity parameter v (normalized by the speed of light). There exist careful resummation techniques, e.g. within the Effective-One-Body (EOB) framework [30], that are built on PN theory results and extend them to the strong-field fast-motion regime which the binary enters close to its merger. For binaries where one body is substantially more massive than the other, the recently developed self-force (SF) formalism proves to be accurate [31, 32]. On the other hand, there has been tremendous progress recently in Numerical Relativity, which includes high-accuracy numerical simulations of binary black hole mergers in non-perturbative General Relativity. The field saw the first breakthrough simulations in 2005 by Frans Pretorius [33, 34], the Goddard group [35] and the research group at the University of Texas – Brownville and Florida Atlantic

University [35]. More recently, the Simulating eXtreme Spacetimes (SXS) collaboration [36] between California Institute of Technology, Cornell University, Canadian Institute of Theoretical Astrophysics, and the California State University at Fullerton has made rapid progress in increasing the stability and efficiency of the numerical methods involved in simulating mergers of black holes [37]. Owing to the computational complexity of these simulations, they are available only for a restricted set of binary configurations. The information from these simulations is traditionally used to calibrate semi-analytic waveform models [38], as well as purely phenomenological closed-form models [39]. There has also been concerted progress in including the effect of the internal matter structure of the neutron stars in binaries, e.g. [40], and in simulating other matter systems, e.g. supernovae [41]. In this dissertation, we will restrict ourselves to systems in vacuum, neglecting possible matter effects.

With its increased sensitivity, the Advanced detectors will need more accurate waveform templates as well as better methods for efficiently using those to filter out the noise from true gravitational wave signals. The goal of the research presented in this dissertation is to construct, and improve the techniques of using waveform templates in gravitational wave searches. The rest of the dissertation is organized as follows:

1. In chapter 2, I describe the production of gravitational waves from compact binaries, and its effect on the interferometric detectors.
2. In chapter 3 I study the importance of the accuracy of the PN theory in the predictive modeling of the late-inspiral of binaries. We estimate the range of binary parameters where the weak-field slow-motion PN theory is accurate and sufficient for aLIGO filter templates.
3. In chapter 4, I study the possibility of using a restricted set of accurate Numerical Relativity simulations as filter template in aLIGO searches.
4. In chapter 5, I describe work done as part of the NINJA-2 project. This involved using accurate NR waveforms as simulated signals embedded in LIGO noise that is recolored to aLIGO sensitivity, in order to test different search algorithms that are being developed by the LIGO-Virgo collaboration.

5. In chapter 6, I describe a waveform model that we developed using recent calculations in the self-force picture. This model suffices to describe the inspiral dynamics of compact binaries with $m_1/m_2 \gtrsim 6$.
6. In chapter 7, I describe a waveform model that completes the one described in the chapter 6 by extending to and accurately capturing the late-inspiral, plunge and merger phases of compact binary coalescence.
7. Finally, chapter 8 is a summary of the conclusions from the research presented in this work.

Chapter 2

Gravitational waves from compact binaries

Chapter 3

Search template banks for low-mass binary black holes in the Advanced gravitational-wave detector era

3.1 Introduction

Over the last decade, there has been tremendous progress towards the first direct detection of gravitational waves. Construction of the Advanced Laser Interferometer Gravitational-Wave Observatory (aLIGO) is underway, with completion scheduled for 2014 [25]. Similar upgrades to the French-Italian Virgo detector [26] have commenced and construction of the Japanese KAGRA detector has begun [42]. When these second-generation gravitational-wave detectors reach design sensitivity, they will increase the observable volume of the universe by a thousandfold or more [43], compared to the first-generation detectors.

The inspiral and merger of binary black holes (BBHs) are expected to be an important source for detection by aLIGO [44]. The rate of BBH coalescences that will be observed by aLIGO at design sensitivity is estimated to be between 0.2 yr^{-1} and 1000 yr^{-1} [28]. Accurate knowledge of the gravitational-wave signals generated by BBHs is crucial for detecting and extracting information about these sources. To provide such waveforms, the effective one body (EOB) model [30] has been calibrated

to numerical simulations of black hole mergers [45, 46, 47, 48, 49, 50, 51, 38]. A new EOB waveform family (called EOBNRv2) has been recently proposed that incorporates information from several non-spinning BBH simulations, with black hole ring-down quasi-normal modes [52, 53] attached to provide a complete BBH waveform [38]. The EOBNRv2 waveform is believed to be sufficiently accurate to search for non-spinning BBH signals in the aLIGO sensitive band (10-1000 Hz).

Past searches for BBHs [14, 15, 16, 17, 10] used matched-filtering [54, 55] to search for coalescing compact binaries. These searches divided the BBH mass space into a *low-mass* region with $M = m_1 + m_2 \lesssim 25 M_\odot$ and a *high-mass* region with $M \gtrsim 25 M_\odot$. In this chapter, we focus attention on BBH systems with component masses between $3 M_\odot \lesssim m_1, m_2 \lesssim 25 M_\odot$, which encompasses mass distribution of black hole candidates observed in low-mass X-ray binaries [56]. aLIGO will be able to detect coalescing BBH systems with component masses $m_1 = m_2 = 25 M_\odot$ to a maximum distance of up to ~ 3.6 Gpc. Since we do not know *a priori* the masses of BBHs that gravitational-wave detectors will observe, searches use a *bank* of template waveforms which covers the range of BBH component masses of interest [57, 58]. This technique is sensitive to the accuracy of the waveform templates that are used as filters and the algorithm used to place the template waveforms [59]. An accurate template bank is required as input for matched filter searches in the Fourier domain [55], as well as newer search algorithms such as the singular value decomposition [60].

In this chapter, we investigate three items of importance to advanced-detector BBH searches: First, we study the accuracy of template placement algorithms for BBH searches using EOBNRv2 waveforms. Optimal template placement requires a metric for creating a grid of waveforms in the desired region of parameter space [61], however no analytic metric exists for the EOBNRv2 waveform. In the absence of such a metric, we construct a template bank using the second-order post-Newtonian hexagonal placement algorithm [62, 63, 64, 65]. This metric is used to place template grid points for the aLIGO zero-detuning high power sensitivity curve [66] and we use EOBNRv2 waveforms at these points as search templates. We find that the existing algorithm works well for BBHs with component masses in the range $3 M_\odot \leq m_1, m_2 \leq 25 M_\odot$. For a template bank constructed with a minimal match of 97% less than 1.5% of non-spinning BBH signals have a mismatch greater than 3%. We therefore conclude that the existing bank placement algorithm is sufficiently

accurate for non-spinning BBH searches in this mass region. Second, we investigate the mass range in which the (computationally less expensive) third-and-a-half-order TaylorF2 post-Newtonian waveforms [57, 67, 68, ?, ?, 69, 70, 71, ?, 72] can be used without significant loss in event rate, and where full inspiral-merger-ringdown EOBNRv2 waveforms are required. We construct a TaylorF2 template bank designed to lose no more than 3% of the matched filter signal-to-noise ratio and use the EOBNRv2 model as signal waveforms. We find that for non-spinning BBHs with $M \lesssim 11.4 M_{\odot}$, the TaylorF2 search performs as expected, with a loss of no more than 10% in the event rate. For higher masses larger event rate losses are observed. A similar study was performed in Ref. [73] using an older version of the EOB model and our results are quantitatively similar. We therefore recommend that this limit is used as the boundary between TaylorF2 and EOBNRv2 waveforms in Advanced LIGO searches. Finally, we investigate the effect of modes other than the dominant $l = m = 2$ mode on BBH searches in aLIGO. The horizon distance of aLIGO (and hence the event rate) is computed considering only the dominant mode of the emitted gravitational waves, since current searches only filter for this mode [28]. However, the inclusion of sub-dominant modes in gravitational-wave template could increase the reach of aLIGO [74, 75]. If we assume that BBH signals are accurately modeled by the EOBNRv2 waveform including the five leading modes, we find that for systems with $(m_1/m_2) \leq 1.68$ or inclination angle: $\iota \geq 2.68$ or $\iota \leq 0.31$ radians, there is no significant loss in the total possible signal-to-noise ratio due to neglecting modes other than $l = m = 2$ in the template waveforms, if one uses a 97% minimal-match bank placed using the hexagonal bank placement algorithm [62, 63, 64, 65]. However, for systems with mass-ratio $(q) \geq 4$ and $1.08 \leq \iota \leq 2.02$, including higher order modes could increase the signal-to-noise ratio by as much as 8% in aLIGO. This increase in amplitude may be offset by the increase in false alarm rate from implementing searches which also include sub-dominant waveform modes in templates, so we encourage the investigation of such algorithms in real detector data.

The remainder of this chapter is organized as follows: In Sec. 3.2 we review the gravitational waveform models used in this study. In Sec. 3.3 we present the results of large-scale Monte Carlo signal injections to test the effectualness of the template banks under investigation. Finally in Sec. 3.4 we review our findings and reccomendations for future work.

3.2 Waveforms and Template Bank Placement

3.2.1 Waveform Approximants

The dynamics of a BBH system can be broadly divided into three regimes: (i) The early inspiral, when the separation between the black holes is large and their velocity is small, can be modeled using results from post-Newtonian (PN) theory [29]. The gravitational-wave phasing of non-spinning binaries is available up to 3.5PN order [?, ?, 69, 70, 71, ?, 72]. (ii) Accurately modeling the late-inspiral and merger requires the numerical solution of the Einstein equations [33, 34, ?, 76, 77, 78, 79]. (iii) The final ring-down phase can be modeled using a super-position of quasi-normal modes (QNMs) which describe the oscillations of the perturbed Kerr black-hole that is formed from the coalescence [52, 53].

Numerical simulation of BBH systems are computationally expensive, and results are only available for a relatively small number of binary systems (see e.g. [80]). The EOB model [30] provides a framework for computing the gravitational waveforms emitted during the inspiral and merger of BBH systems. By attaching a QNM waveform and calibrating the model to numerical relativity (NR) simulations, the EOB framework provides for accurate modeling of complete BBH waveforms (EOBNR). The EOBNR waveforms can be computed at relatively low cost for arbitrary points in the waveform parameter space [45, 46, 47, 48, 49, 50, 51, 38]. In particular the EOB model has recently been tuned against high-accuracy numerical relativity simulations of non-spinning BBHs of mass-ratios $q = \{1, 2, 3, 4, 6\}$, where $q \equiv m_1/m_2$ [38]; we refer to this as the EOBNRv2 model, which we review the major features of below. Throughout, we set $G = c = 1$.

The EOB approach maps the fully general-relativistic dynamics of the two-body system to that of an *effective* mass moving in a deformed Schwarzschild spacetime [30]. The physical dynamics is contained in the deformed-spacetime's metric coefficients, the EOB Hamiltonian [30], and the radiation-reaction force. In polar coordinates (r, Φ) , the EOB metric is written as

$$ds_{\text{eff}}^2 = -A(r)dt^2 + \frac{A(r)}{D(r)}dr^2 + r^2(d\Theta^2 + \sin^2\Theta d\Phi^2). \quad (3.1)$$

The geodesic dynamics of the *effective* mass $\mu = m_1m_2/M$ in the background of

Eq. (3.1) is described by an effective Hamiltonian H^{eff} [30, 81]. The EOBNRv2 model uses Pade-resummations of the third-order post-Newtonian Taylor expansions of the metric coefficients $A(r)$ and $D(r)$, with additional 4PN and 5PN coefficients that are calibrated [46, 47, 48, 49, 38] to ensure that the dynamics agrees closely with NR simulations of comparable mass binaries.

Gravitational waves carry energy and angular momentum away from the binary, and the resulting radiation-reaction force \hat{F}_Φ causes the orbits to shrink. This is related to the energy flux as

$$\hat{F}_\Phi = -\frac{1}{\eta\hat{\Omega}} \frac{dE}{dt} = -\frac{1}{\eta v^3} \frac{dE}{dt}, \quad (3.2)$$

where, $v = (\hat{\Omega})^{1/3} = (\pi M f)^{1/3}$ and f is the instantaneous gravitational-wave frequency. The energy flux dE/dt is obtained by summing over the contribution from each term in the multipole expansion of the waveform, i.e.

$$\frac{dE}{dt} = \frac{\hat{\Omega}^2}{8\pi} \sum_l \sum_m \left| \frac{\mathcal{R}}{M} h_{lm} \right|^2. \quad (3.3)$$

\mathcal{R} is the physical distance to the binary, and h_{lm} are the multipoles of the waveform when it is decomposed in spin weighted spherical harmonic basis as

$$h_+ - i h_\times = \frac{M}{\mathcal{R}} \sum_{l=2}^{\infty} \sum_{m=-l}^{m=l} Y_{-2}^{lm} h_{lm}, \quad (3.4)$$

where Y_{-2}^{lm} are the spin weighted spherical harmonics, and h_+ and h_\times are the two orthogonal gravitational wave polarizations. These waveform multipoles depend on the coordinates and their conjugate momenta, and their Taylor expansions were re-summed as products of individually re-summed factors [82],

$$h_{lm} = h_{lm}^F N_{lm}, \quad (3.5a)$$

$$h_{lm}^F = h_{lm}^{(N,\epsilon)} \hat{S}_{\text{eff}}^{(\epsilon)} T_{lm} e^{i\delta_{lm}} (\rho_{lm})^l; \quad (3.5b)$$

where ϵ is 0 if $(l+m)$ is even, and is 1 otherwise. This factorized-re-summation of the waveform multipoles ensures agreement with NR waveform multipoles [46, 47, 45].

The first factor $h_{lm}^{(N,\epsilon)}$ is the re-summation of the Newtonian order contribution and the second factor $\hat{S}_{\text{eff}}^{(\epsilon)}$ is the source term, given by the mass or the current moments of the binary in the EOB formalism [82, 83]. The tail term T_{lm} is the re-summation of the leading order logarithmic terms that enter into the transfer function of the near-zone multipolar waves to the far-zone [83]. The last term N_{lm} attempts to capture the non-circularity of the quasi-circular orbits. While calculating the energy flux in this study we follow exactly the prescription of Ref. [38], which calibrates the coefficients of the flux so that resulting EOB waveform multipoles reproduce their NR counterparts with high accuracy.

We use the EOBNRv2 Hamiltonian and flux in the equations of motion for the binary, given by

$$\frac{dr}{d\hat{t}} \equiv \frac{\partial \hat{H}^{\text{real}}}{\partial p_r} = \frac{A(r)}{\sqrt{D(r)}} \frac{\partial \hat{H}^{\text{real}}}{\partial p_{r*}}(r, p_{r*}, p_\Phi), \quad (3.6a)$$

$$\frac{d\Phi}{d\hat{t}} \equiv \hat{\Omega} = \frac{\partial \hat{H}^{\text{real}}}{\partial p_\Phi}(r, p_{r*}, p_\Phi), \quad (3.6b)$$

$$\frac{dp_{r*}}{d\hat{t}} = -\frac{A(r)}{\sqrt{D(r)}} \frac{\partial \hat{H}^{\text{real}}}{\partial r}(r, p_{r*}, p_\Phi), \quad (3.6c)$$

$$\frac{dp_\Phi}{d\hat{t}} = \hat{F}_\Phi(r, p_{r*}, p_\Phi); \quad (3.6d)$$

where, $\hat{t} (\equiv t/M)$ is time in dimensionless units.

To obtain the initial values of the coordinates $(r, \Phi, p_{r*}, p_\Phi)$ that the system starts out in, we use the conditions for motion on spherical orbits derived in Ref.[84], where they treat the case of a generic precessing binary. We take their non-spinning limit to define the initial configuration of the binary, requiring

$$\frac{\partial \hat{H}^{\text{real}}}{\partial r} = 0, \quad (3.7a)$$

$$\frac{\partial \hat{H}^{\text{real}}}{\partial p_{r*}} = \frac{1}{\eta} \frac{dE}{dt} \frac{(\partial^2 \hat{H}^{\text{real}} / \partial r \partial p_\Phi)}{(\partial \hat{H}^{\text{real}} / \partial p_\Phi)(\partial^2 \hat{H}^{\text{real}} / \partial r^2)}, \quad (3.7b)$$

$$\frac{\partial \hat{H}^{\text{real}}}{\partial p_\Phi} = \hat{\Omega}_0, \quad (3.7c)$$

where $\hat{\Omega}_0 = \pi M f_0$, with f_0 being the starting gravitational wave frequency. Simplifying Eq.(3.7a), and ignoring the terms involving p_{r*} , as $p_{r*} \ll p_\Phi/r$ in the early

inspiral, we get a relation between p_Φ and r :

$$p_\Phi^2 = \frac{r^3 A'(r)}{2A(r) - rA'(r)}, \quad (3.8)$$

where the prime(') denotes $\partial/\partial r$. Substituting this in Eq.(3.7c), we get the relation:

$$\frac{A'(r)}{2r \left(1 + 2\eta \left(\frac{A(r)}{\sqrt{A(r) - \frac{1}{2}r A'(r)}} - 1 \right) \right)} = \hat{\Omega}_0^2. \quad (3.9)$$

Thus, between Eq.(3.9) and Eq.(3.8), we get the initial values of (r, p_Φ) , corresponding to the initial gravitational wave frequency f_0 , and by substituting these into Eq. 3.7b, we obtain the initial value of p_{r*} . With these values, we integrate the equations of motion to obtain the evolution of the coordinates and momenta $(r(t), \Phi(t), p_r(t), p_\Phi(t))$ over the course of inspiral, until the light-ring is reached. In the EOB model, the light-ring is defined as the local maximum of the orbital frequency $\hat{\Omega}$. From the coordinate evolution, we also calculate $h_{lm}^F(t)$, which is the analytic expression for the waveform multipole without the non-quasi-circular correction factor (defined in Eq. (3.5b)). While generating $h_{lm}^F(t)$ from the dynamics, the values for the free parameters in the expressions for δ_{lm} and ρ_{lm} , are taken from Eqn.[38a-39b] of Ref. [38], where they optimize these parameters to minimize the phase and amplitude discrepancy between the respective EOB waveform multipoles and those extracted from NR simulations.

The EOB ringdown waveform is modeled as a sum of N quasi-normal-modes (QNMs) [46, 47, 49, 52]

$$h_{lm}^{\text{RD}}(t) = \sum_{n=0}^{N-1} A_{lmn} e^{-i\sigma_{lmn}(t-t_{lm}^{\text{match}})}, \quad (3.10)$$

where $N = 8$ for the model we consider. The matching time t_{lm}^{match} is the time at which the inspiral-plunge and the ringdown waveforms are attached and is chosen to be the time at which the amplitude of the inspiral-plunge part of $h_{lm}(t)$ peaks (i.e. t_{peak}^{lm}) [46, 38]. The complex frequencies of the modes σ_{lmn} depend on the mass M_f and spin a_f of the BH that is formed from the coalescence of the binary. We use

the relations of Ref. [38], given by

$$\frac{M_f}{M} = 1 + \left(\sqrt{\frac{8}{9}} - 1 \right) \eta - 0.4333\eta^2 - 0.4392\eta^3, \quad (3.11a)$$

$$\frac{a_f}{M} = \sqrt{12}\eta - 3.871\eta^2 + 4.028\eta^3. \quad (3.11b)$$

Using the mass and spin of the final BH, the complex frequencies of the QNMs can be obtained from Ref. [52], where these were calculated using perturbation theory. The complex amplitudes A_{lmn} are determined by a hybrid-comb numerical matching procedure described in detail in Sec.II C of Ref. [38].

Finally, we combine the inspiral waveform multipole $h_{lm}(t)$ and the ringdown waveform $h^{\text{RD}}(t)$ to obtain the complete inspiral-merger-ringdown EOB waveform $h^{\text{IMR}}(t)$,

$$h_{lm}^{\text{IMR}}(t) = h_{lm}(t)\Theta(t_{lm}^{\text{match}} - t) + h^{\text{RD}}(t)\Theta(t - t_{lm}^{\text{match}}), \quad (3.12)$$

where $\Theta(x) = 1$ for $x \geq 0$, and 0 otherwise. These multipoles are combined to give the two orthogonal polarizations of the gravitational waveform, h_+ and h_\times ,

$$h_+ - i h_\times = \frac{M}{\mathcal{R}} \sum_l \sum_m Y_{-2}^{lm}(\iota, \theta_c) h_{lm}^{\text{IMR}}, \quad (3.13)$$

where ι is the inclination angle that the binary's angular momentum makes with the line of sight, and θ_c is a fiduciary phase angle. To ensure the correctness of our results, we wrote independent code to implement the EOBNRv2 waveform based solely on the content of Ref. [38]. We then validated our code against the EOBNRv2 waveform algorithm in the LSC Algorithm Library (LAL) [85]. We find agreement between these two implementations, giving us confidence in both our results and the correctness of the LAL EOBNRv2 code.

Previous searches for stellar-mass BBHs with total mass $M \lesssim 25M_\odot$ in LIGO and Virgo used the restricted TaylorF2 PN waveforms [57, 67, 68]. Since this waveform is analytically generated in the frequency domain, it has two computational advantages over the EOBNRv2 model: First, the TaylorF2 model does not require either the numerical solution of coupled ODEs or a Fourier transform to generate the frequency domain signal required by a matched filter. We compared the speed of generating and Fourier transforming EOBNRv2 waveforms, to the speed of generating Taylor F2

waveforms in the frequency domain, and found that the former can be $\mathcal{O}(10^2)$ times slower than the latter. Second, the TaylorF2 model can be implemented trivially as a kernel on Graphics Processing Units, allowing search pipelines to leverage significant speed increases due to the fast floating-point performance of GPU hardware. We found the generation of TaylorF2 waveforms using GPUs to be $\mathcal{O}(10^4)$ times faster than generating and Fourier transforming EOBNRv2 waveforms on CPUs. However, use of the TaylorF2 waveform may result in a loss in search efficiency due to inaccuracies of the PN approximation for BBHs. To investigate the loss in search efficiency versus computational efficiency, we use the restricted TaylorF2 waveform described below.

The Fourier transform of a gravitational waveform $h(t)$ is defined by

$$\tilde{h}(f) = \int_{-\infty}^{\infty} e^{-2\pi i f t} h(t) dt. \quad (3.14)$$

Using the stationary phase approximation [86], the Taylor F2 waveform $\tilde{h}(f)$ can be written directly in the frequency domain as

$$\tilde{h}(f) = A f^{-7/6} e^{i\Psi(f)}, \quad (3.15)$$

where we have kept only the leading-order amplitude terms; this is known as the restricted PN waveform. The amplitude $A \propto \mathcal{M}_c^{5/6}/\mathcal{R}$, where \mathcal{M}_c is the *chirp-mass* of the binary, $\mathcal{M}_c = (m_1 + m_2) \eta^{3/5}$, $\eta = m_1 m_2 / (m_1 + m_2)^2$ is the symmetric mass ratio, and \mathcal{R} is the distance to the binary. The Fourier phase of the waveform at

3.5PN order is given by [57, 67, 87, 88, 89, 90, 55]

$$\begin{aligned}
\Psi(f) = & 2\pi f t_c - \phi_c - \frac{\pi}{4} + \frac{3}{128} \frac{1}{\eta} v^{-5} \left[1 + \left(\frac{3715}{756} + \frac{55}{9} \eta \right) v^2 \right. \\
& - 16\pi v^3 + \left(\frac{15293365}{508032} + \frac{27145}{504} \eta + \frac{3085}{72} \eta^2 \right) v^4 \\
& + \left(\frac{38645}{756} - \frac{65}{9} \eta \right) \left(1 + 3 \log \left(\frac{v}{v_{\text{lso}}} \right) \right) \pi v^5 \\
& + \left[\frac{11583231236531}{4694215680} - \frac{640}{3} \pi^2 - \frac{6848}{21} \gamma_E \right. \\
& - \frac{6828}{21} \log(4v) + \left(-\frac{15737765635}{3048192} + \frac{2255}{12} \pi^2 \right) \eta \\
& + \left. \frac{76055}{1728} \eta^2 - \frac{127825}{1296} \eta^3 \right] v^6 \\
& \left. + \left(\frac{77096675}{254016} + \frac{378515}{1512} \eta - \frac{74045}{756} \eta^2 \right) \pi v^7 \right], \tag{3.16}
\end{aligned}$$

where $v = (\pi M f)^{1/3}$ is the characteristic velocity of the binary, and γ is Euler's constant. The initial conditions are set by starting the waveform from a given gravitational-wave frequency $f = f_{\text{low}}$ and the waveform is terminated at the frequency of a test particle at the innermost stable circular orbit (ISCO) of a Schwarzschild black hole ($r = 6M$).

3.2.2 Bank Placement metric

The frequency weighted overlap between two waveforms h_1 and h_2 , can be written as

$$(h_1|h_2) \equiv 2 \int_{f_{\min}}^{f_{\max}} \frac{\tilde{h}_1^*(f) \tilde{h}_2(f) + \tilde{h}_1(f) \tilde{h}_2^*(f)}{S_n(f)} df, \tag{3.17}$$

where $S_n(f)$ is the one-sided power spectral density (PSD) of the detector noise. The normalized overlap between the two waveforms is given by

$$(\hat{h}_1|\hat{h}_2) = \frac{(h_1|h_2)}{\sqrt{(h_1|h_1)(h_2|h_2)}}. \tag{3.18}$$

In addition to the two mass-parameters of the binary, this normalized overlap is also sensitive to the relative phase of coalescence ϕ_c and to the difference in the time of

coalescence between the two waveforms h_1 and h_2 , t_c . These two parameters (ϕ_c , t_c) can be analytically maximized over to get the maximized overlap \mathcal{O}

$$\mathcal{O}(h_1, h_2) = \max_{\phi_c, t_c} \left(\hat{h}_1 | \hat{h}_2 e^{i(2\pi f t_c - \phi_c)} \right), \quad (3.19)$$

which gives a measure of how “close” the two waveforms are in the waveform manifold. The mismatch M between the same two waveforms is written as,

$$M(h_1, h_2) = 1 - \mathcal{O}(h_1, h_2). \quad (3.20)$$

The match (Eq. 4.6) can be regarded as an inner-product on the space of intrinsic template parameters, and thus one can define a *metric* on this space [64, 61] (at the point θ_1) as

$$g_{ij}(\theta_1) = -\frac{1}{2} \left. \frac{\partial^2 \mathcal{O}(h(\theta_1), h(\theta_2))}{\partial \theta_1^i \partial \theta_2^j} \right|_{\theta_1^k = \theta_2^k}, \quad (3.21)$$

where θ_1 is the set of intrinsic parameters (i.e. m_1, m_2 or some combination) of the binary. Thus the mismatch between waveforms produced by systems with nearly equal mass parameters can be given by

$$M(h(\theta), h(\theta + \Delta\theta)) \simeq g_{ij}(\theta) \Delta\theta^i \Delta\theta^j. \quad (3.22)$$

For the TaylorF2 approximant, $h(\theta)$ is given by Eq. (3.15, 3.16), and hence using Eq. (4.5, 4.6) we can get $\mathcal{O}(h(\theta_1), h(\theta_2))$ as an analytic function of θ_1 and θ_2 (albeit involving an integral over frequency). This gives a measure of mismatches between neighbouring points in the manifold of the mass-parameters, and hence a hexagonal 2D lattice placement can be used in the manifold of the mass parameters [64] (and references therein), to construct a geometric lattice based template bank [62, 61, 64].

On the other hand, for the EOBNRv2 approximant, $h(\theta)$ is obtained through numerical solutions of the Hamiltonian equations, Eq.(3.6). In this case, the calculation of the metric would involve derivatives of coordinate evolution obtained from numerically integrated equations of motion, which could introduce numerical instabilities in the metric. So the concept of a metric, as in Eq. (3.21)', cannot be used in a convenient (semi-) analytic form for the construction of a bank with the EOBNRv2 approximant.

3.3 Results

To assess the effectualness of the template banks constructed here, we compute the fitting factors [59] of the template bank, defined as follows. If h_a^e is the waveform emitted by a BBH system then the *Fitting Factor* of a bank of template waveforms (modeled using approximant X) for this waveform, is defined as the maximum value of maximized normalized overlaps between h_a^e and all members h_b^X of the bank of template waveforms [59]; i.e.

$$\mathcal{FF}(a, X) = \max_{b \in \text{bank}} \mathcal{O}(h_a^e, h_b^X). \quad (3.23)$$

This quantity simultaneously quantifies the loss in recovered signal-to-noise ratio (SNR) due to the discreteness of the bank, and the inaccuracy of the template model. The similarly defined quantity MM (minimal match) quantifies the loss in SNR due to only the discreteness of the bank as both the *exact* and the template waveform is modeled with the same waveform model, i.e.

$$\text{MM} = \min_a \max_{b \in \text{bank}} \mathcal{O}(h_a^X, h_b^X), \quad (3.24)$$

where a is any point in the space covered by the bank, and X is the waveform approximant. For a detection search that aims at less than 10% (15%) loss in event detection rate due to the discreteness of the bank and the inaccuracy of the waveform model, we require a bank of template waveforms that has \mathcal{FF} above 0.965 (0.947) [91, 92, 73]. Throughout, we use the aLIGO zero-detuning high power noise curve as the PSD for bank placement and overlap calculations, and set $f_{\min} = 15$ Hz. The waveforms are generated at a sample rate of 8192 Hz, and we set $f_{\max} = 4096$ Hz, i.e. the Nyquist frequency.

The expectation value of the SNR for a signal, ρ , from a source located at a distance D is proportional to $1/D$, which comes from the dependence of the amplitude on the distance. In other words, the range to which a souce can be seen by the detector

$$D_{\text{obs}} = \frac{(g, g)}{\rho^*}, \quad (3.25)$$

where g is the GW strain produced by the same source at the detector, when located

at a unit distance from the detector, and ρ^* is the threshold on SNR required for detection (typically taken as $\rho^* = 8$). For non-precessing binaries, for which the sky-location (θ, ϕ) and polarization angles (ψ) do not change over the course of inspiral, the effective volume in which the same source can be detected is $\propto D_{\text{obs}}^3$ [93], i.e.

$$V_{\text{obs}} = k D_{\text{obs}}^3, \quad (3.26)$$

where the proportionality constant k comes from averaging over various possible sky positions of the binary. The use of discrete template banks, and lack of knowledge of the *true* GW signal model, leads to the observed SNR ρ' being lower than the optimal SNR $\rho = (h, h)$, i.e.

$$\rho' = \mathcal{FF} \rho, \quad (3.27)$$

where \mathcal{FF} is the fitting-factor of the template bank employed in the search for the particular system. The observable volume hence goes down as

$$V_{\text{obs}}^{\text{eff}} = k (\mathcal{FF} \times D_{\text{obs}})^3. \quad (3.28)$$

If we assume that the source population is distributed uniformly in spacial volume in the universe, then the ratio $V_{\text{obs}}^{\text{eff}}/V_{\text{obs}}$ also gives the fraction of systems within the detector's reach that will be seen by the matched-filtering search. For a system with given mass-parameters θ_1 , the ratio of the total $V_{\text{obs}}^{\text{eff}}$ available to it for different inclinations and sky-locations, to the total V_{obs} available to it for the same samples of angles, will give an estimate of the fraction of systems with those masses (marginalized over other parameters - they being uniformly distributed) that will be seen by the matched-filter search. This quantity,

$$\begin{aligned} \epsilon_V(\theta_1) &= \frac{\sum_{\theta_2} V_{\text{obs}}^{\text{eff}}(\theta_1, \theta_2)}{\sum_{\theta_2} V_{\text{obs}}(\theta_1, \theta_2)}, \\ &= \frac{\sum_{\theta_2} \mathcal{FF}^3(\theta_1, \theta_2) V_{\text{obs}}(\theta_1, \theta_2)}{\sum_{\theta_2} V_{\text{obs}}(\theta_1, \theta_2)}, \end{aligned} \quad (3.29)$$

where $\theta_2 = \{\iota, \theta, \phi, \psi\}$ are the parameters being averaged over, we will refer to as the *volume-weighted fitting-factor*. It essentially measures the average of the fractional observable volume loss, weighted by the actual available observable volume, and so simultaneously downweights the loss in the observable volume for binary configurations to which the detector is relatively less sensitive to begin with. We can give the parameter sets θ_1 and θ_2 different elements than the ones shown here, i.e. $\theta_1 \neq \{m_1, m_2\}, \theta_2 \neq \{\iota, \theta, \phi, \psi\}, \theta_1 \cup \theta_2 = \{m_1, m_2, \iota, \theta, \phi, \psi\}$, in order to obtain more information about another set of parameters θ'_1 .

3.3.1 EOBNRv2 templates placed using TaylorF2 metric

In this section we measure the effectualness of the second-order post-Newtonian hexagonal template bank placement metric described in Ref. [63] when used to place EOBNRv2 waveform templates for aLIGO. The same template placement algorithm was used to place a grid of third-and-a-half order post-Newtonian order TaylorF2 waveforms for low-mass BBH detection searches for initial LIGO and Virgo observations [14, 15, 16, 17, 10]. We construct a template bank which has a desired minimal match of 0.97 for waveforms with component masses between $3M_\odot \leq m_1, m_2 \leq 25M_\odot$. This template bank contains 10,753 template grid points in (m_1, m_2) space for the aLIGO noise curve, compared to 373 grid points for the initial LIGO design noise curve. For the template waveforms at each grid point, we use the EOBNRv2 waveforms, rather than TaylorF2 waveforms. Since the metric itself was derived using second-order TaylorF2 waveforms, we do not, *a priori* know if this metric is a good measure to use to place template banks for EOBNRv2 waveforms.

To test the effectualness of this template bank, we perform a Monte-Carlo simulation over the $3M_\odot \leq m_1, m_2 \leq 25M_\odot$ BBH mass space to find regions where the bank placement algorithm leads to under-coverage. We sample 90,000 points uniformly distributed in individual component masses. For each of these points, we generate an EOBNRv2 waveform for the system with component masses given by the coordinates of the point. We record the \mathcal{FF} of the template bank for each of the randomly generated BBH waveforms in the Monte-Carlo simulation. Since we use EOBNRv2 waveforms both to model the true BBH signals and as matched-filter templates, any departure in fitting factor from unity is due to the placement of the template bank grid.




Figure 1 : This figure shows the effectualness of a bank of EOBNRv2 templates, placed using the 2PN accurate hexagonal template placement of Ref. [63], to search for a population of BBH signals simulated with EOBNRv2 waveforms. The masses of the BBH population are chosen from a uniform distribution of component masses between 3 and $25 M_{\odot}$. For each injection, we plot the component masses of the injection, and the fitting factor (\mathcal{FF}).




Figure 2 : This figure shows a cumulative histogram of the fraction of the BBH signal space (on the y-axis), where the bank of EOBNRv2 waveforms has \mathcal{FF} less than the respective values on the x-axis. The EOBNRv2 bank has a fitting factor \mathcal{FF} below the 0.97 for less than $\sim 1.5\%$ of all simulated signals with component-masses m_1, m_2 between $3 M_\odot$ and $25 M_\odot$.

For a bank of template waveforms constructed with a MM of 0.97, Fig. 1 and Fig. 2 show that the \mathcal{FF} of the bank remains above 0.97 for $\sim 98.5\%$ of all simulated BBH signals. Less than $\sim 1.5\%$ of signals have a minimal match of less than 0.97, with the smallest value over the 90,000 sampled points being ~ 0.96 . The diagonal features observed in Fig. 1 are due to the hexagonal bank placement algorithm and are related to the ellipses of constant chirp mass in Fig. 4 of Ref. [63]. From these results, we conclude that the existing template bank placement metric adequately covers the BBH mass space with EOBNRv2 waveform templates; it is not necessary to construct a metric specific to the EOBNRv2 model. aLIGO detection searches can employ the second-order post-Newtonian bank placement metric with the hexagonal placement algorithms [64, 62, 63, 61, 65] to place template banks for EOBNRv2 waveforms without a significant drop in the recovered signal-to-noise ratio.

3.3.2 Effectualness of TaylorF2 templates

We next explore the efficiency of using the computationally cheaper TaylorF2 waveforms to search for a population of BBH signals with component masses between $(3\text{--}25)M_\odot$. The signals from this population are modeled with the full EOBNRv2 waveforms. We use the same template bank placement as above, however now we use the third-and-a-half PN order TaylorF2 model as the template waveforms. This model does not capture the merger and ringdown of BBH signals, as it is terminated at the Schwarzschild test-particle ISCO. Furthermore, it diverges from the true BBH signal in the late inspiral. It is important to determine when these effects become important.

We sample the $(3\text{--}25)M_\odot$ BBH component mass space at 100,000 points by generating an EOBNRv2 waveform to generate the “true” signal waveform. We generate a bank of TaylorF2 template waveforms over the same region, and calculate its \mathcal{FF} for each of the sample points, against the corresponding EOBNRv2 waveform. Fig. 3 shows the distribution of the \mathcal{FF} obtained for the TaylorF2 bank. Clearly the TaylorF2 bank is not effectual for the entire BBH region considered, with mismatches of up to 18% observed. We divide the sampled component mass space into sub-regions which consist of systems with total masses below different thresholds, and compute the minimal-fitting-factor of the bank over those. In Fig. 4, the blue (solid) curve




Figure 3 : The fitting factor \mathcal{FF} of a bank of TaylorF2 waveforms, constructed with $MM = 0.97$, for a population of BBH systems which are modeled using EOBNRv2 signals.




Figure 4 : The (blue) curve shows the upper-bound on total-mass for the sub-region over which the TaylorF2 bank has a minimal-fitting-factor as given on the x-axis. We observe that the TaylorF2 bank has a minimal-fitting-factor of 0.965 (0.947) for the region with total masses below $\sim 11.4M_\odot$ ($19M_\odot$). The minimal-fitting-factor is the fitting-factor value which is less than the fitting-factors of the TaylorF2 bank for $\geq 99.75\%$ of the points sampled in the sub-region.

shows the upper-limit on total mass for different sub-regions against the minimal-fitting-factor of the TaylorF2 bank over those. The minimal-fitting-factor over a sub-region is taken to be the fitting-factor value which is less than the fitting-factors for $\geq 99.75\%$ of the points sampled in the sub-region. We find that the TaylorF2 template bank has \mathcal{FF} above 0.965 (0.947) for the region with total masses below $11.4M_\odot$ ($19M_\odot$). We conclude that the TaylorF2 bank is effectual for BBH signals below $\sim 11.4M_\odot$.

The value of our limit on total-mass is in agreement with the previous study in Ref. [73], however this analysis used the EOBNRv1 model [94] and an older version of the Advanced LIGO noise curve [73]. This agreement provides confidence that this limit will be robust in aLIGO searches and we propose this limit as the upper cutoff for the computationally cheaper TaylorF2 search. To investigate the loss in the \mathcal{FF} due to the mismatch in the template and signal waveform models, we also performed a Monte-Carlo simulation using a denser TaylorF2 bank with $MM = 0.99$. We found that using this dense bank of third-and-a-half order TaylorF2 waveforms, we can relax the limit on the upper mass to $\sim 16.3M_\odot$ ($21.8M_\odot$) and still achieve a \mathcal{FF} above 0.965 (0.947), for over 99.75% of the signals sampled in the region. However, increasing the minimal match increases the size of the template bank from 10,753 to 29,588 templates. This is a significant increase, compared to the cost of filtering with EOBNRv2 templates.

3.3.3 Effect of sub-dominant modes

Having established that the second-order post-Newtonian hexagonal template bank is effectual for placing a bank of EOBNRv2 templates, we now investigate the effect of neglecting sub-dominant modes in BBH searches. The sensitivity reach of the aLIGO detectors is normally computed assuming that the search is only sensitive to the dominant $l = m = 2$ mode of the gravitational waveform. For binary black hole signals, sub-dominant modes may contain significant power [75]. A search that includes these modes could, in principle, have an increased reach (and hence event rate) compared to a search that only uses the dominant mode. The EOBNRv2 model of Ref. [38] has been calibrated against higher order modes from numerical relativity simulations. We investigate the effect of ignoring these modes in a search by modeling the BBH signal as an EOBNRv2 signal containing the dominant and sub-dominant




Figure 5 : (left) The \mathcal{FF} of a bank of EOBNRv2 waveforms, constructed with a minimal match of 0.97 at each point in the stellar-mass BBH component-mass region. While the templates are modeled as the dominant-mode $l = m = 2$ EOBNRv2 waveforms, the signals are modeled including the sub-dominant waveform modes as well (EOBNRv2HM). (right) This figure shows the upper-bound on mass-ratio (q) for the region where a bank of EOBNRv2 templates has a minimal-fitting-factor as given on the x-axis. We observe that for the region with $q \leq 1.68$ (4), the minimum-match of the bank is below 0.965 (0.947). From both the figures, we notice a systematic fall in the coverage of the EOBNRv2 template bank with increasing mass-ratio.




Figure 6 : (left) The \mathcal{FF} of a bank of EOBNRv2 waveforms, constructed with a minimal match of 0.97 at each point in the stellar-mass BBH $q - \iota$ space. While the templates are modeled as the dominant-mode $l = m = 2$ EOBNRv2 waveforms, the signals are modeled including the sub-dominant waveform modes as well (EOBNRv2HM). We observe a loss in fitting-factors, upto $\sim 8\%$, for systems with high mass-ratios (q) and inclination angle (ι) close to $\pi/2$. (right) The \mathcal{FF} for the same population of signals, now shown on the $M - \iota$ plane. We observe the loss in fitting factors to be relatively lesser for more massive binaries.




Figure 7 : (left) This figure shows $\epsilon_V(\theta_1 = \{m_1, m_2\})$ in the component-mass space (see Eq. 3.29). This gives the fraction of total observable volume that is visible to a search which uses the leading order $l = m = 2$ EOBNRv2 waveform template bank, placed with the 2PN accurate TaylorF2 bank placement metric. For a population of signals, that is distributed uniformly in spacial volume, this is equivalent to the fraction of the maximum possible event observation rate that we get with the use of a discrete bank of matched-filters. We observe that the loss in event observation rate, averaged over all parameters (uniformly distributed) but $\theta_1 = \{m_1, m_2\}$, does not exceed $\sim 11\%$ for any region of the component-mass space. (right) This figure shows $\epsilon_V(\theta_1 = \{q, \iota\})$ over the $q - \iota$ plane. We note that the maximum averaged loss in the detection rate is for systems with high mass ratios *and* $\iota \in [1.08, 2.02]$, and can go as high as $\sim 20\%$ for such systems.

multipoles: $h_{lm} = h_{22}, h_{21}, h_{33}, h_{44}, h_{55}$ (which we call EOBNRv2HM) and computing the fitting factor of leading-order EOBNRv2 templates placed using the TaylorF2 metric.

We simulate a population of BBH signals by sampling 100,000 systems uniformly in the $m_1, m_2 \in [3, 25] M_\odot$ component-mass space. These EOBNRv2HM signals are uniformly distributed in sky-location angles and inclination and polarization angles, which appear in the detector's response function to the gravitational-wave signal [95]. The template bank is again placed with a desired minimal match of 0.97 and for each of signal waveforms, we calculate the \mathcal{FF} against the entire bank of EOBNRv2 waveform templates. Fig. 5 (left panel) shows the value of the \mathcal{FF} of the bank of EOBNRv2 waveform templates over the sampled component-mass space. As expected, the highest fitting factors are observed close to the equal mass line, since when the mass ratio is close to unity, the amplitude of the sub-leading waveform modes is several orders of magnitude smaller than the amplitude of the dominant mode. As the mass ratio increases, the relative amplitude of the sub-leading multipoles increases, as illustrated by Fig. 1 of Ref. [38] and the fitting factor decreases. This pattern is brought out further in Fig. 6 (left panel), where we show the \mathcal{FF} values in the mass ratio - inclination angle ($q - \iota$) plane. We observe that when the orbital angular momentum is either parallel or anti-parallel to the line of sight from the detector, the sub-leading multipoles do not contribute significantly to the signal. This is what we would expect from Eq. 3.13, as the spin-weighted harmonics are proportional to $\sin(\frac{\ell}{2})\cos(\frac{\ell}{2})$, except when $l = m = 2$. Similar to Sec. 3.3.2, we divide the sampled component-mass space into sub-regions bounded by $1 \leq q \leq q_{\text{threshold}}$, and compute the minimal-fitting-factor of the EOBNRv2 template bank over those. In Fig. 5 (right panel), the blue (solid) curve shows the value of $q_{\text{threshold}}$ for each restricted sub-region against the minimal-fitting-factor of the bank over the same. For systems with mass-ratio q below 1.68 (4), we find that the \mathcal{FF} of the EOBNRv2 waveform bank is above 0.965 (0.947) over 99.75% of this restricted region. These results demonstrate that the effect of ignoring sub-dominant modes does not cause a significant loss in the total possible signal-to-noise if the mass ratio is less than 1.68. Similar analysis over the range of possible inclination angles shows that the EOBNRv2 waveform bank has fitting factors above 0.965 (0.947) for systems with $2.68 (2.02) \leq \iota \leq \pi$, and $0 \leq \iota \leq 0.31 (1.08)$ (see Fig. 6, left panel).

Fitting factors as low as 0.92 are observed for systems with high mass-ratios *and* inclination angle close to $\pi/2$. As these are also binary configurations to which the detector is relatively less sensitive to [75], fitting-factors alone do not answer the question of where in the parameter space do we lose the most, in terms of detection rate. To address this question, we compute the volume-weighted fitting-factors ϵ_V of the EOBNRv2 template bank, over the sampled BBH parameter space. This gives us an estimate of the expected loss in detection rate, if the source population is distributed uniformly in spacial volume and uniformly in intrinsic and extrinsic source parameters. Fig. 7 (left panel) shows ϵ_V calculated in bins over the component-mass space. In this figure, the color of each bin in the component-mass plane corresponds to, for a population which has all other parameters i.e. the inclination angle and sky/polarization angles uniformly distributed over their possible ranges, the averaged loss in the detection rate incurred due to the use of a bank of leading-order $l = m = 2$ EOBNRv2 templates, placed using the 2PN bank placement metric. We observe that the maximum loss incurred goes up to only $\sim 10\% - 11\%$, which is within our acceptable threshold. Looking at Fig. 6 (left panel), the maximum loss in fitting factor occurs for systems with inclination angles close to $\pi/2$, but (for the same mass-ratio) these gets averaged out with systems with inclinations close to 0 or π , which leads to the low averaged detection-rate losses we observe in Fig. 7 (left panel). The right panel of Fig. 7 shows the same quantity, ϵ_V , calculated over bins in the mass-ratio - inclination angle plane. As expected, we observe that, letting all other parameters be distributed uniformly over their possible ranges, systems with high mass-ratios *and* inclination angles close to $\pi/2$ will incur (averaged) losses in observation volume of up to $\sim 20\%$.

These results suggest that a search that includes higher order modes could achieve a non-trivial increase in sensitivity over leading-order mode templates, only in detecting systems with high q *and* $1.08 \leq \iota \leq 2.02$. However, an algorithm that includes sub-dominant modes could have an increased false-alarm rate (background) over a search that includes only the leading-order mode, and hence the overall gain in search efficiency might not be significant.

3.4 Conclusions

We used the TaylorF2 second-order post-Newtonian hexagonal placement algorithm of Refs. [64, 63, 65] to construct a template bank of EOBNRv2 waveforms with MM of 0.97. We calculated the fitting factor (\mathcal{FF}) of this bank against $\sim 90,000$ simulated EOBNRv2 signals with component masses uniformly distributed between $3M_{\odot} \leq m_1, m_2 \leq 25M_{\odot}$. We find that the \mathcal{FF} of the template bank is greater than 0.97 for 98.5% of the simulated EOBNRv2 signals, assuming the zero-detuning high power noise spectrum for aLIGO sensitivity [66]. We conclude that the existing placement algorithm is effectual for use in aLIGO BBH searches, assuming that EOBNRv2 is an accurate model of BBH signals in this mass region. We then demonstrated that use of the computationally cheaper third-and-a-half order TaylorF2 waveform results in a loss in search efficiency due to inaccuracies of the post-Newtonian approximation, and neglect of merger-ringdown for BBHs with a total mass $M > 11.4 M_{\odot}$. However, below this limit the TaylorF2 model is an acceptable signal for BBH searches. This was done using a bank with a MM of 0.97. By increasing the density of the bank to 0.99MM, the limit on total-mass can be relaxed to $16.3 M_{\odot}$, with an increase in computational cost due to the number of templates increasing by a factor of ~ 2.7 . Finally, we investigated the loss in the SNR incurred by the using template banks constructed using only the leading order mode of EOBNRv2 waveforms. We found that a leading-order $l = m = 2$ EOBNRv2 template bank constructed with a MM of 0.97 is effectual to search for BBHs for which $1 \leq (m_1/m_2) \leq 1.68$ or $\iota \geq 2.68$ or $\iota \leq 0.31$ radians, and there is no significant loss in potential signal-to-noise ratio for systems with q as high as 4 or $2.02 \leq \iota \leq 2.68$ or $0.31 \leq \iota \leq 1.08$. We also observed that the maximum loss in detection-rate, for a binary with given mass parameters, averaging over other parameters - which are taken to be uniformly distributed over their possible ranges, goes only to a maximum of $\sim 10\% - 11\%$. For any given pair of binary masses, the loss is highest when the binary is inclined at $\simeq \pi/2$, and can go up to $\sim 20\%$, and is lower when its angular momentum is close to being parallel or anti-parallel to the line of sight from the detector. These effects average out, and hence for a population which is expected to have a uniform distribution of inclination angles (and uniform distribution in spacial volume), the average loss in detection rate was estimated to be not higher than $\sim 11\%$. Thus, using EOBNRv2HM templates

is unlikely to give a significant increase in the range to which such a population of sources can be detected. For BBHs with $(m_1/m_2) \geq 4$ and $1.08 \leq \iota \leq 2.02$, detection searches could possibly gain sensitivity by the use of EOBNRv2HM waveforms, if they can be implemented without increasing the false alarm rate.

Our results suggest that a significant portion of the non-spinning stellar-mass BBH parameter space can be searched for using LIGO’s existing search algorithms. For systems with total mass below $\sim 11.4M_\odot$ template banks of TaylorF2 can be used without significant loss in event rate. For higher mass systems, neglecting high-order modes in an EOBNRv2 search does not cause a substantial reduction in the maximum possible reach of BBH searches. Finally, we note that our study does not consider BBH systems with BH masses higher than $M = 25 M_\odot$, or the effect of black hole component spins. Future work will extend this study for systems with spinning and/or precessing black holes and consider the effect of non-Gaussian transients in real detector noise.

Chapter 4

Binary black hole search template banks with Numerical Relativity waveforms

4.1 Introduction

Upgrades to the LIGO and Virgo observatories are underway [25, 26], with first observation runs planned for 2015 [27]. The construction of the Japanese detector KAGRA has also begun [42]. The advanced detectors will be sensitive to gravitational waves at frequencies down to $\sim 10\text{Hz}$, with an order of magnitude increase in sensitivity across the band. This is a significant improvement over the lower cutoff of 40Hz for initial LIGO. Estimates for the expected rate of detection have been placed between 0.4 – 1000 stellar-mass binary black hole (BBH) mergers a year [28]. The uncertainty in these estimates comes from the uncertainties in the various factors that govern the physical processes in the BBH formation channels [96, 97]. In sub-solar metallicity environments, stars (in binaries) are expected to lose relatively less mass to stellar winds and form more massive remnants [98, 99, 100]. Population synthesis studies estimate that sub-solar metallicity environments within the horizon of advanced detectors could increase the detection rates to be as high as a few thousand per year [101, 102]. On the other hand, high recoil momenta during core-collapse and merger during the common-envelope phase of the binary star evolution could also decrease the detection rates drastically [100, 101].

Past GW searches have focused on GW bursts [11, 103, 12]; coalescing compact binaries [14, 15, 16, 17, 10, 18, 19], and ringdowns of perturbed black holes [13], amongst others [20, 21, 22, 23, 24]. For coalescing BBHs, detection searches involve matched-filtering [54, 55] of the instrument data using large banks of theoretically modeled waveform templates as filters [57, 64, 61, 63, 62, 65]. The matched-filter is the optimal linear filter to maximize the signal-to-noise ratio (SNR), in the presence of stochastic noise [104]. It requires an accurate modeling of the gravitational waveform emitted by the source binary. Early LIGO-Virgo searches employed template banks of Post-Newtonian (PN) inspiral waveforms [14, 15, 16, 17, 10], while more recent searches targeting high mass BBHs used complete inspiral-merger-ringdown (IMR) waveform templates [18, 19].

Recent developments in Numerical Relativity (NR) have provided complete simulations of BBH dynamics in the strong-field regime, i.e. during the late-inspiral and merger phases [33, 105, 35, 34, 106]. These simulations have contributed unprecedented physical insights to the understanding of BBH mergers (see, e.g., [107, 108, 109, 110] for recent overviews of the field). Due to their high computational cost, fully numerical simulations currently span a few tens of inspiral orbits before merger. For mass-ratios $q = m_1/m_2 = 1, 2, 3, 4, 6, 8$, the multi-domain Spectral Einstein code (SpEC) [111] has been used to simulate 15–33 inspiral merger orbits [79, 112, ?]. These simulations have been used to calibrate waveform models, for example, within the effective-one-body (EOB) formalism [30, 46, 38, 113]. Alternately, inspiral waveforms from PN theory can be joined to numerical BBH inspiral and merger waveforms, to construct longer *hybrid* waveforms [114, 115, 116, 117, 118]. NR-PN hybrids have been used to calibrate phenomenological waveform models [119, 39], and within the NINJA project [120, 80] to study the efficacy of various GW search algorithms towards realistic (NR) signals [121, 122].

Constructing template banks for gravitational wave searches has been a long sought goal for NR. Traditionally, intermediary waveform models are calibrated against numerical simulations and then used in template banks for LIGO searches [18, 19]. In this chapter we explore an alternative to this traditional approach, proposing the use of NR waveforms themselves and hybrids constructed out of them as search templates. For a proof of principle, we focus on the non-spinning BBH space, with the aim of extending to spinning binaries in future work. We investigate exactly where in the mass

space can the existing NR waveforms/hybrids be used as templates, finding that only six simulations are sufficient to cover binaries with $m_{1,2} \gtrsim 12M_\odot$ upto mass-ratio 10. This method can also be used as a guide for the placement of parameters for future NR simulations. Recent work has shown that existing PN waveforms are sufficient for aLIGO searches for $M = m_1 + m_2 \lesssim 12M_\odot$ [73, 123]. To extend the NR/hybrid bank coverage down to $M \simeq 12M_\odot$, we demonstrate that a total of 26 simulations would be sufficient. The template banks are constructed with the requirement that the net SNR recovered for any BBH signal should remain above 96.5% of its optimal value. Enforcing this tells that that these 26 simulations would be required to be ~ 50 orbits long. This goal is achievable, given the recent progress in simulation technology [116, 37, 124]. Our template banks are viable for GW searches with aLIGO, and the framework for using hybrids within the LIGO-Virgo software framework has been demonstrated in the NINJA-2 collaboration [125]. In this chapter, we also derive waveform modeling error bounds which are independent of analytical models. These can be extended straightforwardly to assess the accuracy of such models.

First, we construct a bank for purely-NR templates, restricting to currently available simulations [116, 112, 79, 37, 112]. We use a stochastic algorithm similar to Ref. [126, 127, 128], and place a template bank grid constrained to $q = m_1/m_2 = \{1, 2, 3, 4, 6, 8\}$. The bank placement algorithm uses the EOB model from Ref. [38] (EOBNRv2). As this model was calibrated against NR for most of these mass-ratios, we expect the manifold of EOBNRv2 to be a reasonable approximation for the NR manifold. In Sec. 4.5, we demonstrate that this approximation holds well for NR-PN hybrids as well. To demonstrate the efficacy of the bank, we measure its fitting-factors (FFs) [59] over the BBH mass space. We simulate a population of 100,000 BBH waveforms with masses sampled uniformly over $3M_\odot \leq m_{1,2} \leq 200M_\odot$ and $M = m_1 + m_2 \leq 200M_\odot$, and filter them through the template bank to characterize its SNR recovery. For a bank of NR templates, any SNR loss accrued will be due to the coarseness of the bank grid. We measure this requiring both signals and templates to be in the same manifold, using the EOBNRv2 model for both. We find that for systems with chirp mass $\mathcal{M}_c \equiv (m_1 + m_2)^{-1/5}(m_1 m_2)^{3/5}$ above $\sim 27M_\odot$ and $1 \leq q \leq 10$, this bank has FFs $\geq 97\%$ and is sufficiently accurate to be used in GW searches. We also show that the coverage of the purely-NR bank can be extended to include $10 \leq q \leq 11$, if we instead constrain it to templates with mass-ratios

$$q = \{1, 2, 3, 4, 6, 9.2\}.$$

Second, we demonstrate that currently available PN-NR hybrid waveforms can be used as templates to search for BBHs with much lower masses. The hybrids used correspond to mass-ratios $q = \{1, 2, 3, 4, 6, 8\}$. We use two distinct methods of bank placement to construct a bank with these mass-ratios, and compare the two. The first method is the stochastic algorithm we use for purely-NR templates. The second is a deterministic algorithm, that constructs the two-dimensional bank (in M and q) through a union of six one-dimensional banks, placed separately for each allowed value of mass-ratio. Templates are placed over the total mass dimension by requiring that all pairs of neighboring templates have the same noise weighted overlap. As before, we measure the SNR loss from both banks, due to the discrete placement of the templates, by simulating a population of 100,000 BBH signals, to find the SNR recovered. We measure the intrinsic hybrid errors using the method of Ref. [115, 116], and subsequently account for them in the SNR recovery fraction. We find that the NR-PN hybrid bank is effectual for detecting BBHs with $m_{1,2} \geq 12M_\odot$, with FFs $\geq 96.5\%$. The number of templates required was found to be close to that of a bank constructed using the second-order TaylorF2 hexagonal bank placement algorithm [57, 64, 61, 63, 62, 65]. We note that by pre-generating the template for the least massive binary for each of the mass-ratios that contribute to the bank, we can re-scale it on-the-fly to different total masses in the frequency domain [129]. Used in detection searches, such a bank would be computationally inexpensive to generate relative to a bank of time-domain modeled waveforms.

Finally, we determine the minimal set of NR simulations that we would need to extend the bank down to $M \simeq 12M_\odot$. We find that a bank that samples from the set of 26 mass-ratios listed in Table 2 would be sufficiently dense, even at the lowest masses, for binaries with mass-ratios $1 \leq q \leq 10$. We show that this bank recovers more than 98% of the optimal SNR, not accounting for hybrid errors. To restrict the loss in event detection rate below 10%, we restrict the total SNR loss below 3.5%. This implies the hybrid error mismatches stay below 1.5%, which constrains the length of the NR part for each hybrid. We find that NR simulations spanning about 50 orbits of late-inspiral, merger and ringdown would suffice to reduce the PN truncation error to the desired level. With such a bank of NR-PN hybrids and purely-PN templates for lower masses, we can construct GW searches for stellar-mass BBHs with mass-ratios

$q \leq 10$.

The chapter is organized as follows, in Sec. 4.2.1, we discuss the NR waveforms used in this study, in Sec. 4.2.2 we describe the PN models used to construct the NR-PN hybrids, and in Sec. 4.2.3 we describe the construction of hybrid waveforms. In Sec. 4.2.4 we describe the EOB model that we use to place and test the template banks. In Sec. 4.3 we describe the accuracy measures used in quantifying the loss in signal-to-noise ratio in a matched-filtering search when using a discrete bank of templates and in the construction of hybrid waveforms. In Sec. 4.4 we describe the construction and efficacy of the NR-only banks, while in Sec. 4.5 we discuss the same for the NR-PN hybrid template banks constructed with currently available NR waveforms. In Sec. 4.6, we investigate the parameter and length requirements for future NR simulations in order to cover the entire non-spinning parameter space with $12M_{\odot} \leq M \leq 200M_{\odot}$, $m_{1,2} \geq 3M_{\odot}$, and $1 \leq q \leq 10$. Finally, in Sec. 4.7 we summarize the results.

4.2 Waveforms

In the sections that follow, we will describe the construction of template banks for NR or NR-PN hybrid waveform templates. The NR waveforms that we use correspond to mass ratios $q = \{1, 2, 3, 4, 6, 8\}$, and were simulated using the SpEC code [111]. The construction of hybrid waveforms involves joining a long inspiral portion, modeled using PN theory, to the merger-ringdown waveform from NR. In this section we describe both, the NR waveforms and the PN models used in our study. Measuring the effectualness of these banks involves simulating a population of BBH signals. We use the recently published EOBNRv2 model [38] to obtain waveforms for BBHs with arbitrary masses. This model was calibrated against five out of the six NR simulations we use to construct our banks, and is expected to be faithful at comparable mass ratios [38]. In this section, we briefly summarize the construction of EOBNRv2 waveforms as well.

4.2.1 Numerical Relativity simulations

The numerical relativity waveforms used in this chapter were produced with the SpEC code [111], a multi-domain pseudospectral code to solve Einstein's equations.

η	q	Length (in orbits)	Minimum total mass (M_\odot)
0.25	1	33	49
0.2222	2	15	76
0.1875	3	18	82
0.1600	4	15	87
0.1224	6	20	83
0.0988	8	25	83

Table 1 : SpEC BBH simulations used in this study. Given are symmetric mass-ratio η , mass-ratio $q = m_1/m_2$, and the length in orbits of the simulation. The last column gives the lowest total masses for which the NR simulations cover the entire coalescence process within the sensitive band of aLIGO, starting at 15 Hz.

SpEC uses Generalized Harmonic coordinates, spectral methods, and a flexible domain decomposition, all of which contribute to it being one of the most accurate and efficient codes for computing the gravitational waves from binary black hole systems. High accuracy numerical simulations of the late-inspiral, merger and ring-down of coalescing binary black-holes have been recently performed for mass-ratios $q \equiv m_1/m_2 \in \{1, 2, 3, 4, 6, 8\}$ [79, 130, 131, 112].

The equal-mass, non-spinning waveform covers 33 inspiral orbits and was first discussed in [116, 112]. This waveform was obtained with numerical techniques similar to those of [79]. The unequal-mass waveforms of mass ratios 2, 3, 4, and 6 were presented in detail in Ref. [79]. The simulation with mass ratio 6 covers about 20 orbits and the simulations with mass ratios 2, 3, and 4 are somewhat shorter and cover about 15 orbits. The unequal mass waveform with mass ratio 8 was presented as part of the large waveform catalog in [37, 112]. It is approximately 25 orbits in length. We summarize the NR simulations used in this study in Table 1. In the table, we also give the lowest total masses for which the NR waveforms span the aLIGO band, starting at 15 Hz.

4.2.2 Post-Newtonian waveforms

Post-Newtonian (PN) theory is a perturbative approach to describing the motion of a compact object binary, during the slow-motion and weak-field regime, i.e. the inspiral phase. The conserved energy of a binary in orbit, E , has been calculated to 3PN order in literature [132, 70, 69, 71, 133, 134, 135]. Using the adiabatic approximation, we treat the course of inspiral as a series of radially shrinking circular

orbits. This is valid during the inspiral when the angular velocity of the binary evolves more slowly than the orbital time-scale. The radial separation shrinks as the binary loses energy to gravitational radiation that propagates outwards from the system. The energy flux from a binary F is known in PN theory to 3.5PN order [136, 89, 137, 138]. Combining the energy balance equation, $dE/dt = -F$, with Kepler's law gives a description of the radial and orbital phase evolution of the binary. We start the waveform where the GW frequency enters the sensitive frequency band of advanced LIGO, i.e. at 15Hz. Depending on the way the expressions for orbital energy and flux are combined to obtain the coordinate evolution for the binary, we get different Taylor{T1,T2,T3,T4} time-domain approximants. Using the stationary phase approximation [86], frequency-domain equivalents of these approximants, i.e. TaylorFn, can be constructed. Past GW searches have extensively used the TaylorF2 approximant, as it has a closed form and mitigates the computational cost of generating and numerically fourier-transforming time-domain template [14, 15, 16, 17, 10]. We refer the reader to Ref. [29, 139] for an overview. From the coordinate evolution, we obtain the emitted gravitational waveform; approximating it by the quadrupolar multipole $h_{2,\pm 2}$ which is the dominant mode of the waveform.

4.2.3 PN-NR hybrid waveforms

The hybridization procedure used for this investigation is described in Sec. 3.3 of Ref. [115]: The PN waveform, $h_{\text{PN}}(t)$, is time and phase shifted to match the NR waveform, $h_{\text{NR}}(t)$, and they are smoothly joined together in a GW frequency interval centered at ω_m with width $\delta\omega$:

$$\omega_m - \frac{\delta\omega}{2} \leq \omega \leq \omega_m + \frac{\delta\omega}{2}. \quad (4.1)$$

This translates into a matching interval $t_{\min} < t < t_{\max}$ because the GW frequency continuously increases during the inspiral of the binary. As argued in Ref. [115], we choose $\delta\omega = 0.1\omega_m$ because it offers a good compromise of suppressing residual oscillations in the matching time, while still allowing $h_{\text{PN}}(t)$ to be matched as closely as possible to the beginning of $h_{\text{NR}}(t)$.

The PN waveform depends on a (formal) coalescence time, t_c , and phase, Φ_c . These two parameters are determined by minimizing the GW phase difference in the

matching interval $[t_{\min}, t_{\max}]$ as follows:

$$t'_c, \Phi'_c = \underset{t_c, \Phi_c}{\operatorname{argmin}} \int_{t_{\min}}^{t_{\max}} (\phi_{\text{PN}}(t; t_c, \Phi_c) - \phi_{\text{NR}}(t))^2 dt, \quad (4.2)$$

where t'_c and Φ'_c are the time and phase parameters for the best matching between $h_{\text{PN}}(t)$ and $h_{\text{NR}}(t)$, and $\phi(t)$ is the phase of the (2,2) mode of the gravitational radiation. Since we consider only the (2,2) mode, this procedure is identical to time and phase shifting the PN waveform until it has best agreement with NR as measured by the integral in Eq. (4.2). The hybrid waveform is then constructed in the form

$$h_{\text{H}}(t) \equiv \mathcal{F}(t)h_{\text{PN}}(t; t'_c, \Phi'_c) + [1 - \mathcal{F}(t)]h_{\text{NR}}(t), \quad (4.3)$$

where $\mathcal{F}(t)$ is a blending function defined as

$$\mathcal{F}(t) \equiv \begin{cases} 1, & t < t_{\min} \\ \cos^2 \frac{\pi(t-t_{\min})}{2(t_{\max}-t_{\min})}, & t_{\min} \leq t < t_{\max} \\ 0, & t \geq t_{\max}. \end{cases} \quad (4.4)$$

In this work, we construct all hybrids using the same procedure, Eqs. (4.1)–(4.4), and vary only the PN approximant and the matching frequency ω_m .

4.2.4 Effective-One-Body model

Full numerical simulations are available for a limited number of binary mass combinations. We use a recently proposed EOB model [38], which we refer to as EOBNRv2, as a substitute to model the signal from binaries with arbitrary component masses in this chapter. This model was calibrated to most of the numerical simulations that we use to construct templates banks, which span the range of masses we consider here well. So we expect this approximation to hold. We describe the model briefly here.

The EOB formalism maps the dynamics of a two-body system onto an effective-mass moving in a deformed Schwarzschild-like background [30]. The formalism has evolved to use Padé-resummations of perturbative expansions calculated from PN theory, and allows for the introduction of higher (unknown) order PN terms that are subsequently calibrated against NR simulations of BBHs [50, 51, 48, 82, 46]. The

EOB model proposed recently in Ref. [38] has been calibrated to SpEC NR waveforms for binaries of mass-ratios $q = \{1, 2, 3, 4, 6\}$, where $q \equiv m_1/m_2$, and is the one that we use in this chapter (we will refer to this model as EOBNRv2).

The dynamics of the binary enters in the metric coefficient of the deformed Schwarzschild-like background, the EOB Hamiltonian [30], and the radiation-reaction force. These are known to 3PN order [30, 81] from PN theory. The 4PN & 5PN terms were introduced in the potential $A(r)$, which was Padé resummed and calibrated to NR simulations [46, 47, 48, 49, 38]. We use the resummed potential calibrated in Ref. [38] (see Eq. (5-9)). The geodesic dynamics of the reduced mass $\mu = m_1m_2/M$ in the deformed background is described by the Hamiltonian H^{eff} given by Eq. (3) in [38]. The Hamiltonian describing the conservative dynamics of the binary (labeled the *real* Hamiltonian H^{real}) is related to \hat{H}^{eff} as in Eq. (4) of [38]. The inspiral-merger dynamics can be obtained by numerically solving the Hamiltonian equations of motion for H^{real} , see e.g. Eq.(10) of [38].

The angular momentum carried away from the binary by the outwards propagating GWs results in a radiation-reaction force that causes the orbits to shrink. This is due to the flux of energy from the binary, which is obtained by summing over the contribution from each term in the multipolar decomposition of the inspiral-merger EOB waveform. Complete resummed expressions for these multipoles [82] can be read off from Eq.(13)-(20) of Ref. [38]. In this chapter, as for PN waveforms, we model the inspiral-merger part by summing over the dominant $h_{2,\pm 2}$ multipoles.

The EOB merger-ringdown part is modeled as a sum of N quasi-normal-modes (QNMs), where $N = 8$ for EOBNRv2 [46, 47, 49, 52]. The ringdown frequencies depend on the mass and spin of the BH that is formed from the coalescence of the binary. The inspiral-merger and ringdown parts are attached by matching them at the time at which the amplitude of the inspiral-merger waveform peaks. [46, 38]. The matching procedure followed is explained in detail Sec. II C of Ref. [38]. By combining them, we obtain the complete waveform for a BBH system.

4.3 Quantifying waveform accuracy & bank effectualness

To assess the recovery of SNR from template banks with NR waveforms or NR-PN hybrids as templates, we use the measures proposed in Ref. [59, 57, 58]. The

gravitational waveform emitted during and driving a BBH coalescence is denoted as $h(t)$, or simply h . The inner product between two waveforms h_1 and h_2 is

$$(h_1|h_2) \equiv 4 \int_{f_{\min}}^{f_{\text{Ny}}} \frac{\tilde{h}_1(f)\tilde{h}_2^*(f)}{S_n(f)} df, \quad (4.5)$$

where $S_n(f)$ is the one-sided power spectral density (PSD) of the detector noise, which is assumed to be stationary and Gaussian with zero mean; f_{\min} is the lower frequency cutoff for filtering; f_{Ny} is the Nyquist frequency corresponding to the waveform sampling rate; and $\tilde{h}(f)$ denotes the Fourier transform of $h(t)$. In this chapter, we take $S_n(|f|)$ to be the *zero-detuning high power* noise curve for aLIGO, for both bank placement and overlap calculations [66]; and set the lower frequency cutoff $f_{\min} = 15$ Hz. The peak GW frequency for the lowest binary masses that we consider, i.e. for $m_1 + m_2 \simeq 12M_\odot$, is ~ 2.1 kHz during the ringdown phase. We sample the waveforms at 8192 Hz, preserving the information content up to the Nyquist frequency $f_{\text{Ny}} = 4096$ Hz. A waveform, h , is normalized (made to be a unit vector) by $\hat{h} = h/\sqrt{h|h|}$. In addition to being sensitive to their intrinsic mass parameters, the inner product of two normalized waveforms is sensitive to phase and time shift differences between the two, ϕ_c and t_c . These two parameters (ϕ_c and t_c) can be analytically maximized over to obtain the maximized overlap \mathcal{O} ,

$$\mathcal{O}(h_1, h_2) = \max_{\phi_c, t_c} (\hat{h}_1|\hat{h}_2(\phi_c, t_c)), \quad (4.6)$$

which gives a measure of how “close” the two waveforms are in the waveform manifold, disregarding differences in overall amplitude. The *mismatch* \mathcal{M} between the two waveforms is then

$$\mathcal{M}(h_1, h_2) = 1 - \mathcal{O}(h_1, h_2). \quad (4.7)$$

Matched-filtering detection searches employ a discrete bank of modeled waveforms as filters. The optimal signal-to-noise ratio (SNR) is obtained when the detector strain $s(t) \equiv h^{\text{tr}}(t) + n(t)$ is filtered with the *true* waveform h^{tr} itself, i.e.

$$\rho_{\text{opt}} = \max_{\phi_c, t_c} (h^{\text{tr}}|\hat{h}^{\text{tr}}(\phi_c, t_c)) = ||h^{\text{tr}}||, \quad (4.8)$$

where $||h^{\text{tr}}|| \equiv \sqrt{(h^{\text{tr}}, h^{\text{tr}})}$ is the noise weighted norm of the waveform. With a discrete

bank of filter templates, the SNR we recover

$$\rho \simeq \mathcal{O}(h^{\text{tr}}, h_b) \|h^{\text{tr}}\| = \mathcal{O}(h^{\text{tr}}, h_b) \rho_{\text{opt}}, \quad (4.9)$$

where h_b is the filter template in the *bank* (subscript b) that has the highest overlap with the signal h^{tr} . The furthest distance to which GW signals can be detected is proportional to the matched-filter SNR that the search algorithm finds the signal with. Note that $0 \leq \mathcal{O}(h^{\text{tr}}, h_b) \leq 1$, so the recovered SNR $\rho \leq \rho_{\text{opt}}$ (c.f. Eq. (4.9)). For a BBH population uniformly distributed in spacial volume, the detection rate would decrease as $\mathcal{O}(h^{\text{tr}}, h_b)^3$. Searches that aim at restricting the loss in the detection rate strictly below 10% (or 15%), would require a bank of template waveforms that have \mathcal{O} above 0.965 (or 0.947) with *any* incoming signal [91, 92].

Any template bank has two sources for loss in SNR: (i) the discreteness of the bank grid in the physical parameter space of the BBHs, and, (ii) the disagreement between the actual GW signal h^{tr} and the modeled template waveforms used as filters. We de-coupled these to estimate the SNR loss. Signal waveforms are denoted as h_x^{tr} in what follows, where the superscript tr indicates a *true* signal, and the subscript x indicates the mass parameters of the corresponding binary. Template waveforms are denoted as h_b^M , where M denotes the waveform *model*, and b indicates that it is a member of the discrete *bank*. Fig. 8 shows the signal h_x^{tr} in its manifold, and the bank of templates h_b^M residing in the model waveform manifold, both being embedded in the same space of all possible waveforms. The point h_{\perp}^M is the waveform which has the smallest mismatch in the entire (continuous) model manifold with h_x^{tr} , i.e. $h_{\perp}^M : \mathcal{M}(h_x^{\text{tr}}, h_{\perp}^M) = \min_y \mathcal{M}(h_x^{\text{tr}}, h_y^M)$. The fraction of the optimal SNR recovered at different points x in the binary mass space can be quantified by measuring the fitting factor \mathcal{FF} of the bank [59],

$$\mathcal{FF}(x) = 1 - \min_b \mathcal{M}(h_x^{\text{tr}}, h_b^M). \quad (4.10)$$

For two waveforms h_1 and h_2 close to each other in the waveform manifold: $\|h_1\| \simeq \|h_2\|$, and mutually aligned in phase and time such that the overlap between them is maximized,

$$\|h_1 - h_2\|^2 \simeq 2(h_1|h_1) \left(1 - \frac{(h_1|h_2)}{\sqrt{(h_1|h_1)}\sqrt{(h_1|h_1)}} \right). \quad (4.11)$$




Figure 8 : We show the *true* (upper) and the *hybrid* (lower) waveform manifolds here, with the signal residing in the former, and a discrete bank of templates placed along lines of constant mass-ratio in the latter. Both manifolds are embedded in the same space of all possible waveforms. The true signal waveform is denoted as h_x^{tr} , while the templates in the bank are labelled h_b^M . The hybrid waveform that matches the signal H_x^{tr} best is shown as h_{\perp}^M . Also shown is the “distance” between the signal and the hybrid template that has the highest overlap with it. This figure is qualitatively similar to Fig. 3 of Ref. [91].

The mismatch can, hence, be written as (c.f. Eq. (4.7))

$$\mathcal{M}(h_1, h_2) = \frac{1}{2\|h_1\|^2} \|h_1 - h_2\|^2. \quad (4.12)$$

We note that this equation is an upper bound for Eq. (25) of Ref. [140]. From this relation, and treating the space embedding the true and model waveform manifolds as Euclidean at the scale of template separation, we can separate out the effects of bank coarseness and template inaccuracies as

$$\mathcal{FF}(x) = 1 - \min_b \frac{1}{2\|h_x^{\text{tr}}\|^2} \|h_x^{\text{tr}} - h_b^M\|^2, \quad (4.13a)$$

$$= 1 - \Gamma_{\text{Hyb}}(x) - \Gamma_{\text{bank}}(x); \quad (4.13b)$$

where

$$\Gamma_{\text{Hyb}}(x) \equiv \frac{1}{2\|h_x^{\text{tr}}\|^2} \|h_x^{\text{tr}} - h_{\perp}^M\|^2 = \mathcal{M}(h_x^{\text{tr}}, h_{\perp}^M) \quad (4.14)$$

is the SNR loss from model waveform errors out of the manifold of true signals; and

$$\Gamma_{\text{bank}}(x) \equiv \min_b \frac{1}{2\|h_x^{\text{tr}}\|^2} \|h_{\perp}^M - h_b^M\|^2 = \min_b \mathcal{M}(h_{\perp}^M, h_b^M) \quad (4.15)$$

is the loss in SNR from the distant spacing of templates in the bank. The decomposition in Eq. (4.13b) allows for the measurement of the two effects separately. NR-PN hybrids have the inspiral portion of the waveform, from PN theory, joined to the available late-inspiral and merger portion from NR (as described in Sec. 4.2.3). Towards the late inspiral, the PN waveforms accumulate phase errors, contaminating the hybrids [115, 116]. For each hybrid, we constrain this effect using mismatches between hybrids constructed from the same NR simulation and different PN models, i.e.

$$\Gamma_{\text{Hyb}}(x) \leq \mathcal{M}(h_x^{\text{tr}}, h_x^{\text{Hyb}}) \lesssim \max_{(i,j)} \mathcal{M}(h_x^{M_i}, h_x^{M_j}), \quad (4.16)$$

where $M_i = \text{TaylorT}[1,2,3,4]+\text{NR}$. However, this is only possible for a few values of mass-ratio for which NR simulations are available. We assume Γ_{Hyb} to be a slowly and smoothly varying quantity over the component-mass space at the scale of template grid separation. At any arbitrary point x in the mass space we approximate Γ_{Hyb}

with its value for the “closest” template, i.e.

$$\Gamma_{\text{Hyb}}(x) \leq \max_{(i,j)} \mathcal{M}(h_x^{M_i}, h_x^{M_j}) \simeq \max_{(i,j)} \mathcal{M}(h_b^{M_i}, h_b^{M_j}), \quad (4.17)$$

where h_b^M is the hybrid template in the bank with the highest overlap with the signal at x .

The other contribution to SNR loss comes from the discrete placement of templates in the mass space. In Fig. 8, this is shown in the manifold of the template model. As NR waveforms (or hybrids) are available for a few values of mass-ratio, we measure this in the manifold of EOBNRv2 waveforms. The EOBNRv2 model reproduces most of the NR simulations that were considered here well [38], allowing for this approximation to hold. For the same reason, we expect h_x^{EOBNRv2} to be close to h_b^{EOBNRv2} , with an injective mapping between the two. This allows us to approximate (c.f. Eq. (4.15))

$$\Gamma_{\text{bank}}(x) \simeq \min_b \mathcal{M}(h_x^{\text{EOBNRv2}}, h_b^{\text{EOBNRv2}}). \quad (4.18)$$

In Sec. 4.4, we construct template banks that use purely-NR templates, which have negligible waveform errors. The SNR recovery from such banks is characterized with

$$\mathcal{FF}(x) = 1 - \Gamma_{\text{bank}}(x), \quad (4.19)$$

where the SNR loss from bank coarseness is obtained using Eq. (4.18). In Sec. 4.5, 4.6, we construct template banks aimed at using NR-PN hybrid templates. Their SNR recovery is characterized using Eq. (4.13b), where the additional contribution from the hybrid waveform errors are obtained using Eq. (4.17).

4.4 Constructing a template bank for NR waveforms

In this section we demonstrate the effectualness of a template bank viable for using NR waveforms as templates. The gravitational-wave phase of the dominant waveform multipole extracted from runs at different resolutions was found to converge within ~ 0.3 rad for $q = 3, 4, 6$, and within ~ 0.06 rad for $q = 1, 2$ at merger (see Fig. (6) of Ref. [79], and Fig. (6,7) of Ref. [38] for a compilation). Most of this phase disagreement accumulates over a relatively short duration of $\sim 50M - 100M$ before merger, and




Figure 9 : The color at each point gives the number of waveform cycles N_{cyc} , for that particular binary, which contain 99% of the signal power in the aLIGO sensitivity band. The figure is truncated to exclude the region where $N_{\text{cyc}} > 40$. The solid curve shows the lower bounding edge of the region with $\mathcal{M}_c = 27 M_\odot$.

is significantly lower over the preceding inspiral and plunge. As the matched-filter SNR accumulates secularly over the entire waveform, these numerical phase errors are negligible in terms of mismatches. We set $\Gamma_{\text{Hyb}} = 0$ while computing the fitting factors, so one is left with considering Γ_{bank} to determine the fidelity of the bank (c.f. Eq. (4.13b)).

With NR simulations as templates, the region that the bank can cover is restricted to binaries that have approximately the same number of waveform cycles within the sensitive frequency band of the detectors as the simulations themselves. We take their fiducial length to be ~ 40 GW cycles [141]. For BBHs with $3M_{\odot} \leq m_1, m_2 \leq 200M_{\odot}$ and $m_1 + m_2 \leq 200M_{\odot}$ we map out the region with 99% of the signal power within 40 cycles as the target region of the purely-NR bank. For samples taken over the mass space, we determine the frequency interval $[f_1, f_2]$ for which

$$\int_{f_1}^{f_2} df \frac{|\tilde{h}(f)|^2}{S_n(|f|)} = 0.99 \times \int_{f_{\min}}^{f_{\text{Ny}}} df \frac{|\tilde{h}(f)|^2}{S_n(|f|)}. \quad (4.20)$$

This is done by finding the peak of the integrand in Eq. (4.20) and integrating symmetrically outwards from there, in time, till the interval $[f_1, f_2]$ is found. The number of waveform cycles in this interval is

$$N_{\text{cyc}} = \frac{\Phi(t(f_2)) - \Phi(t(f_1))}{2\pi}, \quad (4.21)$$

where $\Phi(t)$ is the instantaneous phase of the waveform, $h_+(t) - i h_{\times}(t) = A(t) e^{-i\Phi(t)}$, un-wrapped to be a monotonic function of time. We find that for a significant portion of the mass-region, the signal power is contained within 40 waveform cycles. This is shown in Fig. 9, where the color at each point gives N_{cyc} for that system, and the region with $N_{\text{cyc}} > 40$ is excluded. Conservatively, this region is bounded by $\mathcal{M}_c = 27M_{\odot}$, as shown by the solid curve in the figure.

We place a bank over this region, using a stochastic method similar to Ref. [126, 127, 128]. The algorithm begins by taking an empty bank, corresponding to step 0. At step i , a proposal point (q, M) is picked by first choosing a value for q from the restricted set $\mathcal{S}_q = \{1, 2, 3, 4, 6, 8\}$. The total mass M is subsequently sampled from the restricted interval corresponding to the pre-drawn q . The proposal is accepted if the waveform at this point has overlaps $\mathcal{O} < 0.97$ with all the templates in the bank




Figure 10 : The color at each point in the figure gives the value of $\mathcal{FF} \simeq 1 - \Gamma_{\text{bank}}$ of the bank for that binary, for the NR bank restricted to $\mathcal{S}_q = \{1, 2, 3, 4, 6, 8\}$. This is the same as the fraction of the optimal SNR, for the binary, that the template bank recovers. The black dots show the location of the templates in the bank. We note that they all lie along straight lines of constant q passing through the origin. The region shaded light-grey (towards the bottom of the figure) is where the \mathcal{FF} drops sharply below 97%.

from step $i - 1$. This gives the bank at step i . The process is repeated till the fraction of proposals being accepted falls below $\sim 10^{-4}$, and $\gtrsim 99\%$ of the parameter space is covered effectually. To complete the coverage, 100,000 points are sampled over the region of mass space depicted in Fig. 9, and \mathcal{FF} of the bank is computed at each point. With the islands of undercoverage isolated, the points sampled in these regions are added to the bank, pushing their mass-ratios to the two neighboring mass-ratios in \mathcal{S}_q along lines of constant chirp mass. This helps accelerate the convergence of the bank, albeit at the cost of over-populating it, as the algorithm for computing the \mathcal{FF} for the sampled points is parallelizable.

We asses the effectualness of the bank, as discussed in Sec. 4.3, using Eq. (4.19). We draw a population of 100,000 BBH signals, uniformly from the binary mass space, and filter them through the bank. Fig. 10 shows the \mathcal{FF} , or the fraction of the optimal SNR recovered by the bank. The region shown is restricted to binaries with $N_{\text{cyc}} \leq 40$. The black dots in the figure show the position of templates in the bank. The bank




Figure 11 : This figure is similar to Fig. 10. The color at each point gives the value of $\mathcal{FF} \simeq 1 - \Gamma_{\text{bank}}$ of the bank for that binary, for the NR bank restricted to $\mathcal{S}_q = \{1, 2, 3, 4, 6, 9.2\}$. The black dots show the location of the templates in the bank. The region shaded light-grey (towards the bottom of the figure) is where the \mathcal{FF} drops below 97%. We note that with an additional NR waveform for mass ratio $q = 9.2$, the coverage of the bank is extended to include binaries with $10 \leq q \leq 11$.

recovers $\geq 97\%$ of the optimal SNR over the entire region of interest for $q \leq 10$. We propose an additional simulation for $q = 9.2$, to increase the coverage to higher mass-ratios. Substituting this for $q = 8$ in the set of allowed mass-ratios \mathcal{S}_q , we place another bank as before, with $\mathcal{S}_q = \{1, 2, 3, 4, 6, 9.2\}$. The SNR loss from this bank is shown in Fig. 11. This bank recovers $\geq 97\%$ of the SNR for systems with $q \leq 11$ and $N_{\text{cyc}} \leq 40$. The choice of the additional simulation at $q = 9.2$ was made by choosing a value *close-to* the highest possible value of q that does not lead to under-coverage in the region between $q = 6$ and that value. The *exact* highest allowed value was not chosen to reduce the sensitivity of the coverage of the bank to fluctuations in detector sensitivity.

We conclude that with only six NR waveforms for non-spinning BBHs, that are ~ 20 orbits (or 40 GW cycles) in length, a template bank can be constructed that is effectual for detecting binaries with chirp mass above $27M_\odot$ and $1 \leq q \leq 10$. With an additional simulation for $q = 9.2$, this bank can be extended to higher mass-ratios, i.e. to $1 \leq q \leq 11$.

4.5 Constructing a template bank for NR-PN hybrids

The template bank contructed in Sec. 4.4 is effectual for GW detection searches focussed at relatively massive binaries with $\mathcal{M}_c \gtrsim 27M_\odot$. As the NR waveforms are restricted to a small number of orbits, it is useful to consider NR-PN hybrids to bring the lower mass limit down on the template bank. PN waveforms can be generated for an arbitrarily large number of inspiral orbits, reasonably accurately and relatively cheaply. Thus, a hybrid waveform comprised of a long PN early-inspiral and an NR late-inspiral, merger, and ringdown could also be arbitrarily long. There are, however, uncertainties in the PN waveforms, due to the unknown higher-order terms. During the late-inspiral and merger phase, these terms become more important and the PN description becomes less accurate. In addition, when more of the late-inspiral is in the detector's sensitivity frequency range, hybrid waveform mismatches due to the PN errors become increasingly large, and reduce the recovered SNR. Thus, when hybridizing PN and NR waveforms, there must be enough NR orbits that the PN error is sufficiently low for the considered detector noise-curve. In this section, we construct an NR + PN hybrid template bank, for currently available NR waveforms,

and determine the lowest value of binary masses to which it covers.

The hybrids we use are constructed by joining the PN and NR portions, as described in Sec. 4.2.3. The number of orbits before merger at which they are joined depends on the length of the available NR waveforms. We estimate the PN waveform errors using hybridization mismatches Γ_{Hyb} , as discussed in Sec. 4.3. Fig. 12 shows the same for all the hybrids, as a function of total mass. In terms of orbital frequency, these are matched at $M\omega_m = 0.025$ for $q = 1$, $M\omega_m = 0.038$ for $q = 2$, and $M\omega_m = 0.042$ for $q = 3, 4, 6, 8$. In terms of number of orbits before merger, this is 26.9 orbits for $q = 1$, 13.6 orbits for $q = 2$, 12.6 orbits for $q = 3$, 14.3 orbits for $q = 4$, 17.8 orbits for $q = 6$, and 21.4 orbits for $q = 8$. The dotted line indicates a mismatch of 1.5%, a comparatively tight bound that leaves flexibility to accommodate errors due to template bank discreteness. The black circles show the hybrid mismatches at the lower mass bound of the NR-only template bank in Table 1, which are negligible. The inset shows this minimum mass as a function of mass ratio, as well as the minimum attainable mass if we accept a hybrid error of 1.5%. At lower masses, the mismatches increase sharply with more of the PN part moving into the Advanced LIGO sensitivity band. This is due to the nature of the frequency dependence of the detector sensitivity. The detectors will be most sensitive in a comparatively narrow frequency band. As the hybridization frequency sweeps through this band, the hybrid errors rise sharply. They fall again at the lowest masses, for which mostly the PN portion stays within the sensitive band.

We now consider template banks viable for hybrids constructed from currently available NR waveforms at mass ratios $q = 1, 2, 3, 4, 6, 8$. The lower mass limit, in this case, is extended down to masses where the hybridization error exceeds 3%. We demonstrate two independent methods of laying down the bank grid. First, we use the stochastic placement method that proceeds as described in Sec. 4.4. The templates are sampled over the total mass - mass-ratio (M, q) coordinates, sampling q from the restricted set. The total mass M is sampled from the continuous interval between the lower mass limit, which is different for each q , and the upper limit of $200M_\odot$. To assess the SNR loss from the sparse placement of the templates, we simulate a population of 100,000 BBH signal waveforms, with masses sampled with $3M_\odot \leq m_{1,2} \leq 200M_\odot$ and $M \leq 200M_\odot$, and filter them through the bank. This portion of the SNR loss needs to be measured with both signals and templates in the same waveform manifold.




Figure 12 : Bounds on mismatches of PN-NR hybrid waveforms, for the currently existing NR simulations. The PN error is for hybrids matched at $M\omega_m = 0.025$ for $q = 1$, $M\omega_m = 0.038$ for $q = 2$, and $M\omega_m = 0.042$ for $q = 3, 4, 6, 8$. The black circles indicate the lower bound of the template bank in Table 1. The black square show the lower bound with a hybrid error of 1.5%. The inset shows these lower bounds as a function of mass ratio.




Figure 13 : These figures show fitting factors \mathcal{FF} obtained when using a discrete mass-ratio template bank for $q = 1, 2, 3, 4, 6, 8$. For each mass-ratio, the templates are extended down to a total mass where the NR-PN hybridization mismatch becomes 3%. The bank is placed using the stochastic algorithm, similar to Ref. [126, 127, 128]. The black dots show the location of the templates. The fitting factor on the left plot does *not* take into account the hybridization error, and therefore shows the effect of the sparse placement of the templates alone. The right plot accounts for the hybridization error and gives the actual fraction of the optimal SNR that would be recovered with this bank of NR-PN hybrid templates. The region bounded by the magenta (solid) line in both plots indicates the lower end of the coverage of the bank of un-hybridized NR waveforms. Lastly, the shaded grey dots show the points where the fitting factor was below 96.5%.




Figure 14 : These figures are similar to Fig. 13. The figures show fitting factors $\mathcal{F}\mathcal{F}$ obtained when using a discrete mass-ratio template bank for $q = 1, 2, 3, 4, 6, 8$. For each mass-ratio, the templates are extended down to a total mass where the NR-PN hybridization mismatch becomes 3%. Templates are placed independently for each mass-ratio, and span the full range of total masses. For each mass-ratio, neighboring templates are required to have an overlap of 97%. The union of the six single- q one-dimensional banks is taken as the final bank. The black dots show the location of the templates. The fitting factor on the left plot does *not* take into account the hybridization error, and therefore shows the effect of the sparse placement of the templates alone. The right plot accounts for the hybridization error and gives the actual fraction of the optimal SNR that would be recovered with this bank of NR-PN hybrid templates. The region bounded by the magenta (solid) line in both plots indicates the lower end of the coverage of the bank of un-hybridized NR waveforms. Lastly, the shaded grey dots show the points where the fitting factor was below 96.5%.




Figure 15 : This figure is similar to Fig. 14. The figures show fitting factors \mathcal{FF} obtained when using a discrete mass-ratio template bank for $q = 1, 2, 3, 4, 6, 8$. Templates are placed independently for each mass-ratio, and span the range of total masses, down to the region where the hybrid errors become 3%. For each mass-ratio, neighboring templates are required to have an overlap of 97%. The union of the six single- q one-dimensional banks is taken as the final bank. The black dots show the location of the templates. The GW signals are modeled using the EOBNRv2 approximant [38], while TaylorT4+NR hybrids are used as templates. The fitting factor on the left plot shows the combined effect of the sparse placement of the templates, and the (relatively small) disagreement between the hybrid and EOBNRv2 waveforms. The right plot explicitly accounts for the hybridization error and gives the (conservative) actual fraction of the optimal SNR that would be recovered with this bank of NR-PN hybrid templates. The region bounded by the magenta (solid) line in both plots indicates the lower end of the coverage of the bank of un-hybridized NR waveforms. Lastly, the shaded grey

We use the EOBNRv2 approximant [38] to model both, as it has been calibrated to most of the NR waveforms we consider here, and it allows us to model waveforms for arbitrary systems. The left panel of Fig. 13 shows the fraction of the optimal SNR that the bank recovers, accounting for its discreteness alone. We observe that, with just six mass-ratios, the bank can be extended to much lower masses before it is limited by the restricted sampling of mass-ratios for the templates. For binaries with both black-holes more massive than $\sim 12M_{\odot}$, the spacing between mass-ratios was found to be sufficiently dense. The total SNR loss, after subtracting out the hybrid mismatches from Fig. 12, are shown in the right panel of Fig. 13. At the lowest masses, the coverage shrinks between the lines of constant q over which the templates are placed, due to the hybrid errors increasing sharply. We conclude that this bank is viable for hybrid templates for GW searches for BBHs with $m_{1,2} \geq 12M_{\odot}$, $1 \leq q \leq 10$, and $M \leq 200M_{\odot}$. Over this region the bank will recover more than 96.5% of the optimal SNR. This is a significant increase over the coverage allowed for with the purely-NR bank, the region of coverage of which is shown in the right panel of Fig. 14, bounded at lowest masses by the magenta (solid) curve.

Second, we demonstrate a non-stochastic algorithm of bank placement, with comparable results. We first construct six independent bank grids, each restricted to one of the mass-ratios $q = 1, 2, 3, 4, 6, 8$, and spanning the full range of total masses. The template with the lowest total mass is chosen by requiring the hybrid mismatch to be 3% at that point. The spacing between neighboring templates is given by requiring that the overlap between them be 97%. We take the union of these banks as the final two-dimensional bank. As before, we measure the SNR loss due to discreteness of the bank and the waveform errors in the templates separately. To estimate the former, we simulate a population of 100,000 BBH systems, and filter them through the bank. The signals and the templates are both modeled with the EOBNRv2 model. The left panel of Fig. 14 reveals the fraction of SNR recovered over the mass space, accounting for the sparsity of the bank alone, i.e. $1 - \Gamma_{\text{bank}}$. At lower masses, we again start to see gaps between the lines of constant mass ratio which become significant at $m_{1,2} \leq 12M_{\odot}$. The right panel of Fig. 14 shows the final fraction of the optimal SNR recovered, i.e. the \mathcal{FF} as defined in Eq. (4.13b). As before, these are computed by subtracting out the hybrid mismatches Γ_{Hyb} in addition to the discrete mismatches, as described in Sec. 4.3.

The efficacy of both methods of template bank construction can be compared from Fig. 13 and Fig. 14. We observe that the final banks from either of the algorithms have very similar SNR recovery, and are both effectual over the range of masses we consider here. Both were also found to give a very similar number of templates. The uniform-in-overlap method yields a grid with 2,325 templates. The stochastic bank, on the other hand, was placed with a requirement of 98% minimal mismatch, and had 2,457 templates. This however includes templates with $m_{1,2} < 12M_\odot$. Restricted to provide coverage over the region with $m_{1,2} \geq 12M_\odot$, $1 \leq q \leq 10$, and $M \leq 200M_\odot$, the two methods yield banks with 627 and 667 templates respectively. The size of these banks is comparable to one constructed using the second-order post-Newtonian TaylorF2 hexagonal template placement method [62, 63, 64, 65], which yields a grid of 522 and 736 templates, for a minimal match of 97% and 98%, respectively.

Finally, we test the robustness of these results using TaylorT4+NR hybrids as templates. As before, we simulate a population of 100,000 BBH signal waveforms. As we do not have hybrids for arbitrary binary masses, we model the signals as EOBNRv2 waveforms. This population is filtered against a bank of hybrid templates. The SNR recovered is shown in the left panel of Fig. 15. Comparing with the left panels of Fig. 13, 14, we find that the EOBNRv2 manifold is a reasonable approximation for the hybrid manifold; and that, at lower masses, there is a small systematic bias in the hybrids towards EOBNRv2 signals with slightly higher mass-ratios. The right panel of Fig. 15 shows the fraction of optimal SNR recovered after subtracting out the hybrid mismatches from the left panel. The similarity of the \mathcal{FF} distribution between the right panels of Fig. 15 and Fig. 13, 14 is remarkable. This gives us confidence that the EOBNRv2 model is a good approximation for testing NR/hybrid template banks, as we do in this chapter; and that a template bank of NR+PN hybrids is indeed effectual for binaries with $m_{1,2} \geq 12M_\odot$, $M \leq 200M_\odot$ and $q \leq 10$.

4.6 Complete NR-PN hybrid bank for non-spinning BBH

The last sections outlined properties of template banks of NR waveforms (and their hybrids) which are available today. We also investigate the parameter and length requirements for future NR simulations, that would let us construct detection template banks all the way to $M = m_1 + m_2 = 12M_\odot$. This lower limit was chosen following




Figure 16 : These figures show the coverage of template banks restricted to single mass-ratios, i.e. (from left to right) $q = 1, 4, 8$. We note that at higher total masses, the templates are correlated to simulated signals for considerably different mass-ratios, than at lower total masses. This agrees with what we expect as with decreasing total mass, the number of cycles in the sensitive frequency band of Advanced LIGO increases.

Ref. [123, 73] which showed that the region with $M \lesssim 12M_\odot$ can be covered with banks of post-Newtonian inspiral-only waveforms.

Constructing such a bank is a two-step process. First, we pick mass-ratios that allow construction of such a bank given long enough waveforms for these mass-ratios. Second, one needs to determine the necessary length of the NR portion of the waveforms, such that the PN-hybridization error is sufficiently low for all masses of interest.

To motivate the first step, Fig. 16 shows the coverage of banks that sample from a *single* mass-ratio each (from left to right: $q = 1, 4, 8$). We see that the resolution in q required at lower values of M increases sharply below $M \sim 60M_\odot$. This follows from the increase in the number of waveform cycles in aLIGO frequency band as the total mass decreases, which, in turn, increases the discriminatory resolution of the matched-filter along the q axis. To determine the least set of mass-ratios which would sample the q axis sufficiently densely at lower masses, we iteratively add mass-ratios to the allowed set and test banks restricted to sample from it. We find that, constrained to the set \mathcal{S}_q given in Table 2, a template bank can be constructed that has a minimal match of 98% at the lowest masses. The left panel of Fig. 17 shows the loss in SNR




Figure 17 : This figure shows fitting factors for a hybrid template bank which samples from the 26 mass ratios $q = 1, 1.5, 1.75, 2, \dots, 9.6$, and allows coverage to masses down to $m_1 + m_2 = 12M_\odot$ and $1 \leq q \leq 10$, with a minimal-match of 98% at the lowest masses. The left and right panel show the same on $M - q$ and $m_1 - m_2$ axes, respectively. The magenta lines, in both panels, shows the upper bound in total mass, below which frequency-domain PN waveforms can be used to construct template banks for aLIGO searches [73, 123]. The dash-dotted line in the right panel shows the lower mass limit on the smaller component object, to which a bank of currently available NR-PN hybrids can cover, i.e. $\min(m_1, m_2) = 12M_\odot$ (see Sec. 4.5). The blue (solid) curve in the right panel gives the lower mass limit to which a bank of currently available NR waveforms can cover (see Sec. 4.4). Thus, between the simulations listed in Table 2, and frequency domain PN waveforms, we can search for the entire range of BBH masses.

$q (\equiv m_1/m_2)$
1, 1.5, 1.75,
2, 2.25, 2.5, 2.75,
3, 3.25, 3.5, 3.8,
4.05, 4.35, 4.65, 4.95,
5.25, 5.55, 5.85,
6.2, 6.55,
7, 7.5,
8, 8.5,
9, 9.6

Table 2 : List of mass-ratios, a template bank restricted to which will be effectual over the region of the non-spinning BBH mass space where $m_1 + m_2 \gtrsim 12M_\odot$ and $1 \leq q \leq 10$. The fraction of optimal SNR recovered by such a bank, accounting for discreteness losses, remains above 98%. This is shown in Fig. 17.

due to bank grid coarseness, i.e. $1 - \Gamma_{\text{bank}}$. This loss remains below 2% for mass-ratios $1 \leq q \leq 10$, even at $M = 12M_\odot$. This leaves a margin of 1.5% for the hybrid mismatches that would incur due to the hybridization of the NR merger waveforms with long PN inspirals. The right panel of Fig. 17 shows the same data in the m_1 - m_2 plane. In this figure, the region covered by the NR-only bank is above the blue (solid) curve, while that covered by a bank of the currently available NR-PN hybrids is above the line of $m_2 = 12M_\odot$ (with $m_2 \leq m_1$). The region from Ref. [123, 73] that can be covered by PN templates is in the bottom left corner, bounded by the magenta (solid) line. Our bank restricted to the set of 26 mass-ratios, as above, provides additional coverage for binaries with $M \geq 12M_\odot$, $m_2 \leq 12M_\odot$ and $1 \leq q \leq 10$. Thus between purely-PN and NR/NR-PN hybrid templates, we can construct effectual searches for non-spinning BBHs with $q \leq 10$.

Having the set of required mass-ratios \mathcal{S}_q determined, we need to decide on the length requirements for the NR simulations, in order to control the PN hybridization error. For a series of matching frequencies, we construct NR-PN hybrids with Taylor{T1,T2,T3,T4} inspirals, and compute their pairwise mismatch as a function of total mass. The maximum of these mismatches serves a conservative bound on the PN-hybridization error for that hybrid (c.f. Eq. (4.17)). Fig. 18 shows the results of this calculation. Each panel of Fig. 18 focuses on one mass-ratio. Within each panel, each line represents one matching-frequency, with lines moving down toward earlier hybridization with smaller mismatches. Because the hybridization frequency is not






Figure 18 : The maximum mismatch between different PN approximants for hybrid waveforms plotted against the total mass for at different matching frequencies ($M\omega_m$). The dotted lines indicate a mismatch of 1.5% and a lower total mass limit, $12M_\odot$ for $q = 1$, and $M_2 = 3M_\odot$ for $q = 4, 8$. The thick dashed lines indicate the currently possible matching frequency for hybrids based on the length of NR waveforms. The numbers next to each line indicate the number of orbits before merger where the PN and NR (or EOB) waveforms were stitched together.



particularly intuitive, the lines are labeled by the number of orbits of the NR portion of the hybrid-waveform. For a short number of orbits this calculation is indeed done with NR waveforms, whereas for large number of orbits, we substitute EOBNRv2 waveforms. The dashed lines represent the earliest one can match a NR+PN hybrid given the currently available NR waveforms, and are the same as the $q = 1, 4$, and 8 lines in Fig 12. The solid curves show the results using EOB hybrids, while the dotted curves (just barely visible) show the results with NR hybrids. They are virtually identical, which is a confirmation that EOB hybrids can act as a good proxy for NR hybrids in this case. The horizontal dotted line indicates a mismatch of 1.5%, while the vertical dotted line shows a lower mass limit for each mass ratio: $12M_{\odot}$ for $q = 1$, which is the point at which one can construct a template bank with only PN inspirals, $15M_{\odot}$ for $q = 4$, and $27M_{\odot}$ for $q = 8$, which are the lower mass limits if both component masses are $\geq 3M_{\odot}$.

Fig. 19 presents the information obtained in the previous paragraph in a different way. Given NR-PN hybrids with N orbits of NR, the shaded areas in the left panel of Fig. 19 indicate the region of parameter space for which such hybrids have hybridization errors smaller than 1.5%. As before, we see that for high masses, comparatively few NR orbits are sufficient (e.g. the purple $N = 15$ region), whereas lower total masses require increasingly more NR orbits. The dashed lines indicate the region of parameter space with hybrid error below 3%. The black dotted line designates the point where one component mass is greater than $3M_{\odot}$, which is a reasonable lower mass limit for a physical black hole. The right panel shows this same analysis instead with initial GW frequency indicated by the solid and dashed lines. Thus, for the region of parameter space we're interested in, no more than ~ 50 NR orbits, or an initial GW frequency of $M\omega = 0.025$ would be necessary to construct a detection bank with hybrid mismatches below 1.5%.

4.7 Conclusions

The upgrades currently being installed to increase the sensitivity of the ground based interferometric gravitational-wave detectors LIGO and Virgo [25, 26] are scheduled to complete within the next two years. The second generation detectors will have a factor of 10 better sensitivity across the sensitive frequency band, with the lower

frequency limit being pushed from 40Hz down to $\sim 10\text{Hz}$. They will be able to detect GWs from stellar-mass BBHs up to distances of a few Gpc, with the expected frequency of detection between $0.4 - 1000 \text{ yr}^{-1}$ [28].

Gravitational-wave detection searches for BBHs operate by matched-filtering the detector data against a bank of modeled waveform templates [142, 57, 64, 61, 63, 62, 65]. Early LIGO-Virgo searches employed PN waveform template banks that spanned only the inspiral phase of the coalescence [14, 15, 16, 17, 10]. Recent work has shown that a similar bank of PN templates would be effectual for the advanced detectors, to detect non-spinning BBHs with $m_1 + m_2 \lesssim 12M_\odot$ [123, 73]. Searches from the observation period between 2005 – 07 and 2009 – 10 employed templates that also included the late-inspiral, merger and ringdown phases of binary coalescence [18, 19].

Recent advancements in Numerical Relativity have led to high-accuracy simulations of the late-inspiral and mergers of BBHs. The multi-domain SpEC code [111] has been used to perform simulations for non-spinning binaries with mass-ratios $q = 1, 2, 3, 4, 6, 8$ [79, 112, ?]. Owing to their high computational complexity, the length of these simulations varies between 15 – 33 orbits. Accurate modeling of the late-inspiral and merger phases is important for stellar mass BBHs, as they merge at frequencies that the advanced detectors would be sensitive to [123]. Analytic models, like those within the Effective-One-Body formalism, have been calibrated to the NR simulations to increase their accuracy during these phases [30, 46, 38, 113]. Other independent models have also been developed using information from NR simulations and their hybrids [143, 119, 39, 144]. An alternate derived prescription is that of NR+PN hybrid waveforms, that are constructed by joining long PN early-inspirals and late-inspiral-merger simulations from NR [114, 115, 116, 117, 118].

NR has long sought to contribute template banks for gravitational-wave searches. Due to the restrictions on the length and number of NR waveforms, this has been conventionally pursued by calibrating intermediary waveform models, and using those for search templates. In this chapter, we explore the alternative of using NR waveforms and their hybrids directly in template banks. We demonstrate the feasibility of this idea for non-spinning binaries, and extending it to spinning binaries would be the subject of a future work. We find that with only six non-spinning NR simulations, we can cover down to $m_{1,2} \gtrsim 12M_\odot$. We show that with 26 additional NR simulations, we can complete the non-spinning template banks down to $M \simeq 12M_\odot$, below which

existing PN waveforms have been shown to suffice for aLIGO. From template bank accuracy requirements, we are able to put a bound on the required length and initial GW frequencies for the new simulations. This method can therefore be used to lay down the parameters for future simulations.

First, we construct a bank for using pure-NR waveforms as templates, using a stochastic algorithm similar to Ref. [126, 127, 128]. The filter templates are constrained to mass-ratios for which we have NR simulations available, i.e. $q = 1, 2, 3, 4, 6, 8$. We assume that the simulations available to us are ≥ 20 orbits in length. To test the bank, we simulate a population of 100,000 BBH signals and filter them through the bank. The signals and templates are both modeled with the EOBNRv2 model [38]. We demonstrate that this bank is indeed effectual and recovers $\geq 97\%$ of the optimal SNR for GWs from BBHs with mass-ratios $1 \leq q \leq 10$ and chirp-mass $\mathcal{M}_c \equiv (m_1 + m_2)^{-1/5}(m_1 m_2)^{3/5}$ above $27 M_\odot$. Fig. 10 shows this fraction at different simulated points over the binary mass space. With an additional simulation for $q = 9.2$, we are able to extend the coverage to higher mass-ratios. We show that a bank viable for NR waveform templates for $q = 1, 2, 3, 4, 6, 9.2$, would recover $\geq 97\%$ of the optimal SNR for BBHs with $10 \leq q \leq 11$. The SNR recovery fraction from such a bank is shown in Fig. 11.

Second, we construct effectual banks for currently available NR-PN hybrid waveform templates. We derive a bound on waveform model errors, which is independent of analytical models and can be used to independently assess the errors of such models (see Sec. 4.3 for details). This allows us to estimate the hybrid waveform mismatches due to PN error, which are negligible at high masses, and become significant at lower binary masses. We take their contribution to the SNR loss into account while characterizing template banks. For hybrid banks, we demonstrate and compare two independent algorithms of template bank construction. First, we stochastically place a bank grid, as for the purely-NR template bank. Second, we lay down independent sub-banks for each mass-ratio, with a fixed overlap between neighboring templates, and take their union as the final bank. To test these banks, we simulate a population of 100,000 BBH signals and filter them through each. We simulate the GW signals and the templates using the recently developed EOBNRv2 model [38]. The fraction of the optimal SNR recovered by the two banks, before and after accounting for the hybrid errors, are shown in the left and right panels of Fig. 13 and Fig. 14

(respectively). We observe that for BBHs with $m_{1,2} \geq 12M_\odot$ hybrid template banks will recover $\geq 96.5\%$ of the optimal SNR. For testing the robustness of our conclusions, we also test the banks using TaylorT4+NR hybrid templates. The SNR recovery from a bank of these is shown in Fig. 15. We conclude that, the currently available NR+PN hybrid waveforms can be used as templates in a matched-filtering search for GWs from BBHs with $m_{1,2} \geq 12M_\odot$ and $1 \leq q \leq 10$. The number of templates required to provide coverage over this region was found to be comparable to a bank constructed using the second-order post-Newtonian TaylorF2 hexagonal template placement method [62, 63, 64, 65]. The two algorithms we demonstrate yield grids of 667 and 627 templates, respectively; while the metric based placement method yields a grid of 522 and 736 templates, for 97% and 98% minimal match, respectively.

At lower mass, the length of the waveform in the sensitive frequency band of the detectors increases, increasing the resolution of the matched-filter. We therefore see regions of undercoverage between mass ratios for which we have NR/hybrid templates (see, e.g. Fig. 14 at the left edge). For $M \lesssim 12M_\odot$, existing PN waveforms were shown to be sufficient for aLIGO searches. We find the additional simulations that would be needed to extend the hybrid template bank down to $12M_\odot$. We show that a bank of hybrids restricted to the 26 mass-ratios listed in Table 2 would be sufficiently dense at $12M_\odot$. This demonstrates that the method proposed here can be used to decide which NR simulations should be prioritized for the purpose of the GW detection problem. By filtering a population of 100,000 BBH signals through this bank, we show that the SNR loss due to its discreteness stays below 2% over the entire relevant range of masses. The fraction of optimal SNR recovered is shown in Fig. 17. Constraining the detection rate loss below 10% requires that detection template banks recover more than 96.5% of the optimal SNR. Therefore our bank would need hybrids with hybridization mismatches below 1.5%. From this accuracy requirement, we obtain the length requirement for all the 26 simulations. This is depicted in the left panel of Fig. 19, where we show the region of the mass space that can be covered with hybrids, as the length of their NR portion varies. We find that for $1 \leq q \leq 10$ the new simulations should be about 50 orbits in length. In the right panel of Fig. 19 we show the corresponding initial GW frequencies. The requirement of ~ 50 orbit long NR simulations is ambitious, but certainly feasible with the current BBH simulation

technology [124].

In summary, we refer to the right panel of Fig. 17. The region above the dashed (red) line and above the solid (blue) line can be covered with a bank of purely-NR waveforms currently available. The region above the dashed (red) and the dash-dotted (black) line can be covered with the same simulations hybridized to long PN inspirals. With an additional set of NR simulations summarized in Table 2, the coverage of the bank can be extended down to the magenta (solid) line in the lower left corner of the figure. Thus between hybrids and PN waveforms, we can cover the entire non-spinning BBH space. The ability to use hybrid waveforms within the software infrastructure of the LIGO-Virgo collaboration has been demonstrated in the NINJA-2 collaboration [125]. The template banks we present here can be directly used in aLIGO searches. This work will be most useful when extended to aligned spin and precessing binaries [145, 146], which is the subject of a future work.

The detector noise power is modeled using the zero-detuning high-power noise curve for Advanced LIGO [66]. The construction of our template banks is sensitive to the breadth of the frequency range that the detector would be sensitive to. The noise curve we use is the broad-band final design sensitivity estimate. For lower sensitivities at the low/high frequencies, our results would become more conservative, i.e. the template banks would over-cover (and not under-cover).

We finally note that in this chapter we have only considered the dominant (2, 2) mode of the spherical decomposition of the gravitational waveform. For high mass-ratios and high binary masses, other modes would also become important, both for spinning as well as non-spinning black hole binaries [75, 123, 147]. Thus, in future work, it would be relevant to examine the sub-dominant modes of the gravitational waves. Lastly, though we have looked at the feasibility of using this template bank for Advanced LIGO as a single detector, this instrument will be part of a network of detectors, which comes with increased sensitivity and sky localization. For this reason, in subsequent studies it would be useful to consider a network of detectors.

Chapter 5

NINJA-2: Detecting gravitational waveforms modelled using numerical binary black hole simulations

The NINJA-1 project was a huge success in bringing the numerical relativity and gravitational-wave astronomy communities together. The project also resulted in several intriguing qualitative results. Among these were promising indications that the alterations suggested by the studies of chapter ?? could indeed improve search efficiencies.

However, NINJA-1 only began the process of testing detection and parameter estimation pipelines against realistic signals. The follow-up project, NINJA-2, is ongoing as of the time of writing. NINJA-2 aims to remove some of the shortcomings of NINJA-1 and allow quantitative studies of the performance of gravitational-wave searches in various regions of signal parameter space. Specifically, NINJA-2 addresses issues with both the waveform submissions and the noise used to construct the data sets. This chapter describes the contributed waveforms and the studies that have been performed to validate them. The next chapter will discuss the construction of the data sets and will present preliminary results from the low- and high-mass CBC pipelines.

5.1 Introduction

A network of second-generation laser interferometric gravitational-wave (GW) observatories is presently under construction. The US-based Advanced Laser Interferometer Gravitational Wave Observatory (aLIGO) [25] is expected to have its initial observing run in 2015 utilizing observatories in Hanford, Washington and Livingston, Louisiana (denoted “H” and “L”, respectively). aLIGO will then work towards reaching design sensitivity, expected in 2018-20 [27]. The French-Italian Advanced Virgo (AdV) observatory [148, 26] (denoted “V”) is expected to follow shortly after the aLIGO instruments. The cryogenically cooled KAGRA observatory [149, 42] and a India-based aLIGO facility [150, 151] are due to begin operations around 2020, providing a 5-site network to explore the gravitational-wave sky in detail.

These second-generation observatories will have an order of magnitude increase in sensitivity over their first generation counterparts and will be sensitive to a broader range of gravitational-wave frequencies [25, 26, 42]. One of the primary observational targets for this global network is the inspiral, merger and ringdown of a binary system containing two black holes [152]. With aLIGO and AdV operating at their final design sensitivities it is expected that 0.4 - 1000 binary black hole (BBH) coalescences will be observed per year of operation [153]. Directly observing the collision of two black holes will allow gravitational-wave astronomers to understand the physics of black-hole spacetimes and to explore the strong-field conditions of the theory of general relativity [95].

Exploring the underlying mass and spin distributions of stellar-mass black holes can tell us a great deal about the end stages of massive-star evolution. The mass measurements of compact objects made to date suggest a gap between the most massive neutron stars ($\lesssim 3 M_{\odot}$) [154] and the least massive black holes ($\gtrsim 5 M_{\odot}$) [155]. It is still an open question as to whether this gap is real and the result of formation mechanisms, or simply due to observational biases [156]. Whether or not ground-based detectors will be able to distinguish between these regions in mass space is of great interest. Furthermore, from the distributions of black hole spin magnitudes and tilts (orientation of the spin relative to the orbital angular momentum), more can be learned about supernovae kicks and compact binary formation environments. Stellar-mass black hole spin measurements are currently done by modelling either the

accretion disk’s thermal continuum X-ray spectrum, or the profile of the broadened Fe K α line [157]. Both methods are fundamentally based on assumptions about the location of the inner edge of the accretion disk, and also depend sensitively on very complicated physical models of the disk and its emission. Gravitational waves will provide an entirely new method of measuring black hole spin which does not require the complicated modeling of accretion disk physics.

Of the stellar mass black hole angular momenta (spin) measurements made to date, half are found to have a magnitude $a \gtrsim 0.8$ [157]. With BBH observations, aLIGO and AdV will be able to provide independent measurements of the black hole spin magnitudes. Therefore, it will be interesting to evaluate how well aLIGO and AdV will be able to constrain the magnitude of the black holes’ component spins. The direction of the compact objects’ angular momenta is also of interest, with particular implications for formation mechanisms [158]. Measuring systems with component spins misaligned with the orbital angular momentum is outside of the scope of this project. However, this study does include systems with component spins that are both aligned and anti-aligned with the orbital angular momenta, and we will evaluate the ability of aLIGO and AdV to distinguish such systems from one another.

The standard technique for observing BBH mergers involves matched-filtering data taken from gravitational-wave observatories against “template” waveforms that should closely match potential astrophysical signals [54, 159, 55]. The observable BBH waveform includes the signal from the inspiral of the two black holes, as well as their merger and the resulting black hole’s ringdown. Search templates must include all of these features [160, 123]. As an alternative to matched-filter searches, a number of algorithms exist to perform searches for *unmodelled* gravitational-wave signals [161, 162, 163]. These algorithms do not require accurate knowledge of the waveforms to make observations, but are not as sensitive as matched-filter searches in cases where the waveform models are well understood.

Theoretical models of the inspiral, merger and ringdown of BBH systems are necessary to produce template banks for matched-filter searches and to use as model signals to test both matched-filter and unmodelled searches. The inspiral portion of the waveform can be modeled by analytic post-Newtonian (PN) calculations [164, 160], while numerical solutions of the General Relativity field equations are required to accurately model the final orbits and merger. Prior to breakthroughs in numerical

relativity (NR) in 2005 [?, 35, 105], template banks and search pipeline tests used only inspiral waveforms. Since 2007, NR waveforms have been used to calibrate analytical waveform models [94, 165, 166, 113, 167, 39, 168, 169]. Some of the analytical waveforms have been already employed in search pipelines [19]. However, there exists another useful and valuable avenue of communication between numerical relativists and gravitational-wave astronomers. As NR pushes into new regions of parameter space the waveforms can be used directly to test searches employing previously-calibrated templates, and the degree to which these searches prove to be insufficient can motivate both new template models and additional simulations.

The Numerical INjection Analysis (NINJA) project was created in 2008. The project uses recent advances in numerical relativity ([170] and references therein) to test analysis pipelines by adding numerically-modelled, physically-realistic signals to detector noise and attempting to recover these signals with search pipelines. The first NINJA project (NINJA-1) [122] utilized a total of 23 numerical waveforms, which were injected into Gaussian noise colored with the frequency sensitivity of initial LIGO and Virgo. These data were analyzed by nine data-analysis groups using both search and parameter-estimation algorithms [122].

However, there were four limitations to the NINJA-1 analysis. First, due to the computational cost of NR simulations, most waveforms included only ~ 10 orbits before merger. Therefore the waveforms were too short to inject over an astrophysically interesting mass range without introducing artifacts into the data. The lowest mass binary considered in NINJA-1 had a total mass of $35M_{\odot}$, whereas the mass of black holes could extend below $5M_{\odot}$ [156, 56]. Second, the waveforms were only inspected for obvious, pathological errors and no cross-checks were performed between the submitted waveforms. It was therefore difficult to assess the physical fidelity of the results. Third, the NINJA-1 data set contained stationary noise with the simulated signals already injected into the data. Since the data set lacked the non-Gaussian noise transients present in real detector data, it was not possible to fully explore the response of the algorithms in a real search scenario. Finally, the data set contained only 126 simulated signals, this precluded detailed statistical studies of the effectiveness of search and parameter estimation algorithms. Despite these limitations, the NINJA-1 project lead to a framework within which to perform injection studies using waveforms as calculated by the full nonlinear general theory of relativity, established

guidelines for such studies (in particular a well-defined format for the exchange of NR waveforms [171]), and clarified where further work was needed.

This lead to the initiation of the second NINJA project (NINJA-2), whose goals were to build and improve upon NINJA-1 and perform a systematic test of the efficiency of data-analysis pipelines in preparation for the Advanced detector era. A set of 60 NR waveforms were submitted by 8 numerical relativity groups for the NINJA-2 project [172]. These waveforms conform to a set of length and accuracy requirements, and are attached to PN inspiral signals to produce hybrid PN-NR waveforms that can be injected over the full range of physically relevant total binary masses. The construction and verification of these waveforms is described in a previous paper [172], and summarized here in section 5.2. In the Numerical-Relativity and Analytical-Relativity collaboration, a project complementary to the NINJA collaboration, 22 new NR waveforms were produced and rigorously analysed. These newly produced NR waveforms were compared to the most recent calibrated analytical models, finding that the loss of event rates due to modeling is below 3% [173]. In this paper we study the ability of the search algorithms used in the last of the Initial LIGO and Virgo science runs to observe numerically-modelled BBH waveforms from the set of 60 waveforms submitted to the NINJA-2 project. This is done using data taken during LIGO’s sixth and Virgo’s second science runs and recoloring that data to the sensitivities expected from early observation runs of aLIGO and AdV.

There are a wide range of search and parameter estimation algorithms available within the GW astronomy community: those that were used in past analyses of detector data, old algorithms that have been updated and re-tuned following the experience gained in those analyses, plus many new algorithms under development. For both practical reasons, and to mark a clear point in the development and refinement of these methods, this work employed *only* search and parameter estimation algorithms that were approved and used in the last initial-LIGO and Virgo science runs, *without any additional tuning or modifications* [161, 142, 19, 12]. By doing this we aim to provide a benchmark against which future algorithms can be compared.

A set of 7 numerical relativity waveforms, with masses ranging from $14.4M_{\odot}$ to $124M_{\odot}$ were added into the recolored data as an unbiased test of the process through which candidate events are identified for BBH waveforms. This data was distributed

to analysts who knew that such “blind injections” were present but had no information about the number, parameters or temporal location of these waveforms. This was similar to blind injection tests conducted by the LIGO and Virgo collaborations in their latest science runs [174]. Using a search for unmodelled gravitational wave transients we found that one of these signals was recovered, with an estimated false alarm rate of 1 every 47 years. The remaining 6 signals were consistent with background. Using a matched-filtered algorithm with a bank of BBH IMR waveforms, which were *not* calibrated against the NR signals used in NINJA-2, 6 of the signals were recovered with more significance than all background events. This allowed upper limits on the false alarm rate ranging between 1 every 5000 years and 1 every 40000 years to be placed on each blind injection. The remaining signal was not recovered due to having a low network signal-to-noise ratio and possessing a large anti-aligned spin, which was not modelled in the bank of waveforms used in the search.

Parameter estimation algorithms have come a long way since the first NINJA project. Previously these analyses were unable to estimate the parameters of high mass systems accurately due to the use of inspiral-only models on data with little measurable inspiral. We show that these tools are now capable of reliably providing parameter estimates for both low and high mass systems. For all but one injection the masses and spins of the black holes were recovered within the estimated 95% credible regions. The remaining injection suffered small systematic biases due to non-Gaussian features present in the noise, the modeling of which is an ongoing endeavor. We find that strong intrinsic degeneracies between the masses and black hole spins [175, 176] make it difficult to constrain the masses well, for 3 of the signals the presence of a neutron star cannot be ruled out. We also investigate the ability to constrain the sky localization of the various signals and demonstrate how even low power non-Gaussian noise transients in the data can effect the recovery of the intrinsic parameters of BBH systems.

We use large sets of known waveforms to assess the efficiency of the matched-filter BBH search algorithm as a function of the mass and angular momenta of the component black holes. These are the first such studies that have been done using real data recolored to second-generation noise curves, which include the non-Gaussian features that will be present in the data taken with Advanced LIGO and Advanced Virgo. As these results were obtained using the search pipelines and techniques that

were deployed in the final observing runs of Initial LIGO and Initial Virgo, they can therefore provide a benchmark against which improvements to the search techniques can be compared and assessed.

In our large-scale simulation studies we find evidence that incorporating search waveforms including the effects of spin will increase the efficiency of searches. The results shown here can be used, in the future, to compare with results of search pipelines including the effects of component spins that are aligned with the orbital angular momentum, which are currently under development [177, 127, 178]. We also assess the efficiency of the matched-filter BBH search algorithm to recover waveforms generated by different groups with the same parameters. We find that the efficiency of the matched-filter BBH search algorithm to recover different waveforms, generated by different groups, but with identical physical parameters, is indistinguishable up to statistical errors.

In this work we have not scaled observed masses and distances to account for cosmological effects, which will be important especially for high-mass binary black hole collisions. Therefore any masses and distances quoted should be interpreted as *observed* masses and *luminosity* distances.

The paper contains 4 analyses: First, an unmodelled search for the 7 blind injections. Second, a modelled search for the 7 blind injections using frequency domain TaylorF2 waveforms for templates with total mass $< 25M_{\odot}$ and effective-one-body inspiral-merger-ringdown waveforms for total mass $> 25M_{\odot}$. Third, a followup parameter estimation analysis on the 7 blind injections using three inspiral-merger-ringdown templates—PhenomB with no spin, EOBNRv2 and PhenomB with aligned spins. Finally, a large-scale simulation campaign, adding a large number of the submitted NR waveforms to the NINJA-2 recolored data and attempting to recover these with the modelled search pipeline.

The paper is organized as follows: In section 5.2 we briefly summarize the waveform catalogue described more fully in [172]. Section 5.3 describes the LIGO/Virgo data used and the processing that was done to make it resemble anticipated advanced-detector noise. Section 5.4 describes how the parameters for the blind injections were chosen, reports the values that were selected and describes the distribution of parameters used for the large-scale simulation campaign. Section 5.5 describes the various

detection and parameter estimation algorithms that were used in our analyses. Section 5.6 reports the results of the unmodelled and modelled searches for the 7 blind injections. Section 5.7 describes the results of the parameter estimation analysis on the blind injections. Section 5.8 describes the results of the large scale simulation campaign, which aims to quantify the sensitivity of the detection searches to the different hybrid waveforms. We conclude in section 5.9 with a discussion of how well the various algorithms performed, and implications for the Advanced detector era.

5.2 PN-NR Hybrid Waveforms

The NINJA-2 waveform catalog contains 60 PN-NR hybrid waveforms that were contributed by eight numerical relativity groups. This catalog and the procedures used to validate it are described in detail in [172]. We briefly summarize the NINJA-2 catalog here.

Each waveform in the NINJA-2 waveform catalog consists of a PN portion modelling the early inspiral, stitched to a numerical portion modelling the late inspiral, merger and ringdown. This ensures accurate modelling of the late portions of the waveform while simultaneously ensuring that waveforms are long enough to be scaled to masses as low as $10M_{\odot}$ without starting abruptly within the sensitive frequency band of the detectors. We require that for the NR portion of the waveform the amplitude be accurate to within 5% and the phase (as a function of gravitational-wave frequency) have an accumulated uncertainty over the inspiral, merger and ringdown of no more than 0.5 rad. Since we do not have access to exact waveforms we define “accuracy” by convergence of the numerical waveforms as resolution and waveform-extraction radius are increased. We also require at least five orbits of numerical data in order to ensure robust blending with the PN portion. No requirements were placed on the hybridization itself, although it is known that hybridization can introduce significant errors [115, 39, 179]. It was decided to limit NINJA-2 to systems without eccentricity, and with black-hole spins parallel or anti-parallel to the orbital angular momentum. This last condition avoids precession, which we do for two reasons; (i) precession greatly complicates waveform phenomenology and we prefer to first tackle a simpler subset which still maintains the main features of binary evolution and merger; and (ii) at the start of NINJA-2 the precessing-binary parameter space had




Figure 20 : Mass ratio q and dimensionless spins χ_i of the NINJA-2 hybrid waveform submissions. Reproduced from [172].

been sampled by only a handful of numerical simulations. Waveforms were submitted in the format described in [171], and data was provided as strain decomposed into spherical harmonics of weight -2 . Groups were encouraged to submit modes beyond $(l, m) = (2, \pm 2)$ and many did so. However the techniques to validate these higher modes are a current research topic. In order not to delay the NINJA-2 project it was decided to validate only the $(2, \pm 2)$ modes in [172] and employ only these modes for the first NINJA-2 analysis. Different groups employed different codes, as well as different methods for solving initial conditions, dealing with singularities, evolving Einstein’s equations, and extracting gravitational-wave information. In addition different PN approximants and different hybridization methods were used by different groups in constructing the full hybrid waveforms. It was found that the dominant source of disagreement between submissions was in the PN portion, and in particular overlaps between submissions were greater than 0.97 over the range of masses, including regions sensitive to differences in hybridization techniques. See [172] for details.

The parameter space for aligned-spin BBH systems is four dimensional; the masses and spin magnitudes of each of the two holes. However, in the absence of matter Einstein’s equations possess a mass invariance, and a solution obtained by numerical relativity or other method may be trivially rescaled to any total mass. We therefore eliminate total mass from the parameter space of submissions leaving the ratio of the two masses, denoted q , and the dimensionless spins denoted $\chi_{1,2}$ which must lie between $-1 < \chi_{1,2} < 1$.

Tables 3 and 4 give a summary of the submissions for systems where the masses of the two black holes are equal and unequal, respectively. The first column of Tables 3 and 4 gives a label for each waveform, to ease referring to them in later sections. These labels of the form “G2+20+20_T4” are constructed as follows: The first letter represents the group submitting the numerical simulation:

- F:** The numerical relativity group at Florida Atlantic University, also using the BAM code [180, 181, 182, 183].
- G:** The Georgia Tech group using MayaKracn [184, 185, 186, 187, 188, 189, 190]
- J:** The BAM (Jena) code, as used by the Cardiff-Jena-Palma-Vienna collaboration [191, 192, 167, 193, 181, 194]

- L:** The Lean Code, developed by Ulrich Sperhake [195, 196].
- Ll:** The Llama code, used by the AEI group and the Palma-Caltech groups [197, 198, 199]
- R:** The group from Rochester Institute of Technology, using the LazEv code [35, 200, 201, 202].
- S:** The SXS collaboration using the SpEC code [203, 204, 205, 206, 207, 130, 208, 209, 106, 210].
- U:** The group from The University of Illinois [211].

Immediately after this letter follows the mass-ratio $q = m_1/m_2$, where the black holes are labeled such that $q \geq 1$. Subsequently are the components of the initial dimensionless spin along the orbital angular momentum, multiplied by 100 (e.g. ‘+20’ corresponds to $\hat{L} \cdot \vec{S}_1/m_1^2 = 0.2$) of the more massive and the less massive black hole. The label closes with the Taylor-approximant being used for the PN portion of the waveform, with “T1” and “T4” representing TaylorT1 and TaylorT4, respectively. The Georgia Tech group submitted four pairs of simulations where each pair simulates systems with identical physical parameters, stitched to the same PN approximant. These waveforms are not identical however as each simulation within a pair has a different number of NR cycles and was generated at a different resolution. These are distinguished by appending “_1” and “_2” to the label.

Each NR group verified that their waveforms met the minimum NINJA-2 requirements as described above. The minimum-five-orbits requirement was easily verified by inspection, and the amplitude and phase uncertainties were estimated by convergence tests with respect to numerical resolution and waveform-extraction radius. The full catalog was then verified by the NINJA-2 collaboration. Submissions were inspected in the time and frequency domains to identify any obvious problems caused by hybridization or integration from the Newman-Penrose curvature scalar ψ_4 to strain. Where multiple simulations were available for the same physical parameters these simulations were compared using the matched-filter *overlap*. The inner product between two real waveforms $s_1(t)$ and $s_2(t)$ is defined as

$$(s_1 | s_2) = 4 \Re \int_0^\infty df \frac{\tilde{s}_1(f) \tilde{s}_2^*(f)}{S_n(f)} \quad (5.1)$$

where \tilde{x} denotes the Fourier transform of x and $S_n(f)$ is the power spectral density, which was taken to be the target sensitivity for the first advanced-detector runs, referred to as the “early aLIGO” PSD. This is described in more detail in section 5.3.

The overlap is then obtained by normalization and maximization over relative time and phase shifts, Δt and $\Delta\phi$.

$$\langle s_1 | s_2 \rangle := \max_{\Delta t, \Delta\phi} \frac{(s_1 | s_2)}{\sqrt{(s_1 | s_1)(s_2 | s_2)}}. \quad (5.2)$$

The investigations in [172] demonstrated that the submitted waveforms met the requirements as outlined above and in addition were consistent with each other to the extent expected. We therefore conclude that these submissions model real gravitational waves with sufficient accuracy to quantitatively determine how data-analysis pipelines will respond to signals in next-generation gravitational-wave observatories.

The NINJA-2 waveforms cover the 3-dimensional aligned-spin parameter space rather unevenly, as indicated in figure 20. The configurations available fall predominantly into two 1-dimensional subspaces: (i) Binaries of varying mass-ratio, but with non-spinning black holes. (ii) Binaries of black holes with equal-mass and equal-spin, and with varying spin-magnitude. Future studies, with additional waveforms covering the gaps that are clearly evident in figure 20 and waveforms including precession [212, 37, 173], would be useful to more fully understand the response of search codes across the parameter space, and would help to better tune analytical waveform models including inspiral, merger and ringdown phases.

5.3 Modified Detector Noise

In this section we describe the techniques used in this work to emulate data that will be taken by second generation gravitational wave observatories. This was accomplished by recoloring data taken from the initial LIGO and Virgo instruments to predicted 2015 – 2016 sensitivities. Recoloring initial LIGO and Virgo data allows the non-Gaussianity and non-stationarity of that data to be maintained.

The predicted sensitivity curves of the advanced detectors as a function of time can be found in the living document [27]. For this work we are interested in the sensitivity of the advanced detectors in 2015 – 2016 and used a previous prediction of

Table 3 : Summary of the contributions to the NINJA-2 waveform catalog with $m_1 = m_2$. Given are an identifying label, described in section 5.2, mass-ratio $q = m_1/m_2$ which is always 1 for these simulations, magnitude of the dimensionless spins $\chi_i = S_i/m_i^2$, orbital eccentricity e , frequency range of hybridization in $M\omega$, the number of numerical cycles from the middle of the hybridization region through the peak amplitude, and the post-Newtonian Taylor-approximant(s) used for hybridization.

Label	q	χ_1	χ_2	1000 e	100 $M\omega$ hyb.range	# NR cycles	pN Approx
S1-95-95_T1	1.0	-0.95	-0.95	1.00	3.3 – 4.1	18.42	T1
J1-85-85_T1	1.0	-0.85	-0.85	2.50	4.1 – 4.7	12.09	T1
J1-85-85_T4							T4
J1-75-75_T1	1.0	-0.75	-0.75	1.60	4.1 – 4.7	13.42	T1
J1-75-75_T4							T4
J1-50-50_T1	1.0	-0.50	-0.50	2.90	4.3 – 4.7	15.12	T1
J1-50-50_T4							T4
S1-44-44_T4	1.0	-0.44	-0.44	0.04	4.3 – 5.3	13.47	T4
Ll1-40-40_T1	1.0	-0.40	-0.40		6.1 – 8.0	6.42	T1
Ll1-40-40_T4							T4
J1-25-25_T1	1.0	-0.25	-0.25	2.50	4.5 – 5.0	15.15	T1
J1-25-25_T4							T4
Ll1-20-20_T1	1.0	-0.20	-0.20		5.7 – 7.8	8.16	T1
Ll1-20-20_T4							T4
J1+00+00_T1	1.0	0.00	0.00	1.80	4.6 – 5.1	15.72	T1
J1+00+00_T4							T4
G1+00+00_T4				3.00	5.5 – 7.5	9.77	T4
Ll1+00+00_F2					5.7 – 9.4	8.30	F2
S1+00+00_T4				0.05	3.6 – 4.5	22.98	T4
G1+20+20_T4_1	1.0	0.20	0.20	10.00	6.0 – 7.5	6.77	T4
G1+20+20_T4_2				6.00	5.5 – 7.5	10.96	T4
J1+25+25_T1	1.0	0.25	0.25	6.10	4.6 – 5.0	18.00	T1
J1+25+25_T4							T4
G1+40+40_T4_1	1.0	0.40	0.40	10.00	5.9 – 7.5	7.70	T4
G1+40+40_T4_2				6.00	5.5 – 7.5	12.02	T4
Ll1+40+40_T1					7.8 – 8.6	6.54	T1
Ll1+40+40_T4							T4
S1+44+44_T4	1.0	0.44	0.44	0.02	4.1 – 5.0	22.39	T4
J1+50+50_T1	1.0	0.50	0.50	6.10	5.2 – 5.9	15.71	T1
J1+50+50_T4							T4
G1+60+60_T4_1	1.0	0.60	0.60	12.00	6.0 – 7.5	8.56	T4
G1+60+60_T4_2				5.00	5.5 – 7.5	13.21	T4
J1+75+75_T1	1.0	0.75	0.75	6.00	6.0 – 7.0	14.03	T1
J1+75+75_T4							T4
G1+80+00_T4	1.0	0.80	0.00	13.00	5.5 – 7.5	12.26	T4
G1+80+80_T4_1	1.0	0.80	0.80	14.00	5.9 – 7.5	9.57	T4
G1+80+80_T4_2				6.70	5.5 – 7.5	14.25	T4
J1+85+85_T1	1.0	0.85	0.85	5.00	5.9 – 6.9	15.36	T1
J1+85+85_T4							T4
U1+85+85_T1				20.00	5.9 – 7.0	15.02	T1
G1+90+90_T4	1.0	0.90	0.90	3.00	5.8 – 7.5	15.05	T4
S1+97+97_T4	1.0	0.97	0.97	0.60	3.2 – 4.3	38.40	T4

Table 4 : Summary of the contributions to the NINJA-2 waveform catalog with $m_1 > m_2$. Given are an identifying label, described in section 5.2, mass-ratio $q = m_1/m_2$ magnitude of the dimensionless spins $\chi_i = S_i/m_i^2$, orbital eccentricity e , frequency range of hybridization in $M\omega$, the number of numerical cycles from the middle of the hybridization region through the peak amplitude, and the post-Newtonian Taylor-approximant(s) used for hybridization.

Label	q	χ_1	χ_2	$1000e$	$100 M\omega$ hyb.range	# NR cycles	pN Approx
J2+00+00_T1	2.0	0.00	0.00	2.30	6.3 – 7.8	8.31	T1
J2+00+00_T4							T4
G2+00+00_T4				2.50	5.5 – 7.5	10.42	T4
Ll2+00+00_F2					6.3 – 9.4	7.47	F2
S2+00+00_T2				0.03	3.8 – 4.7	22.34	T2
G2+20+20_T4	2.0	0.20	0.20	10.00	5.6 – 7.5	11.50	T4
J2+25+00_T1	2.0	0.25	0.00	2.00	5.0 – 5.6	15.93	T1
J2+25+00_T4							T4
J3+00+00_T1	3.0	0.00	0.00	1.60	6.0 – 7.1	10.61	T1
J3+00+00_T4							T4
S3+00+00_T2				0.02	4.1 – 5.2	21.80	T2
F3+60+40_T4	3.0	0.60	0.40	1.00	5.0 – 5.6	18.89	T4
J4+00+00_T1	4.0	0.00	0.00	2.60	5.9 – 6.8	12.38	T1
J4+00+00_T4							T4
L4+00+00_T1				5.00	5.1 – 5.5	17.33	T1
S4+00+00_T2				0.03	4.4 – 5.5	21.67	T2
S6+00+00_T1	6.0	0.00	0.00	0.04	4.1 – 4.6	33.77	T1
R10+00+00_T4	10.0	0.00	0.00	0.40	7.3 – 7.4	14.44	T4




Figure 21 : Left: predicted sensitivity curves for aLIGO and AdV. Shown are both the design curves and predicted 2015 – 2016 *early* sensitivity curves. Also shown is the early AdV noise curve rescaled such that the horizon distance for a $(10 M_{\odot}, 10 M_{\odot})$ binary system is equal to that obtained with the early aLIGO noise curve. Right: Horizon distance as a function of observed total mass for the early aLIGO and rescaled early AdV sensitivity curves. This plot is made considering only equal mass, non-spinning systems and calculated using the EOBNRv2 [166] waveform approximant. Results in this paper are generated from the early aLIGO noise curve and the rescaled early AdV curve.

the sensitivity curves for this time period as given in [213] and shown in the left panel of figure 21. These curves were used as the updated predictions given in [27] were not available when we began this study. We refer to the 2015 – 2016 predicted noise curves as the *early* sensitivity curves. It is clear from the figure that the predicted sensitivity of early AdV is significantly greater than that of the early aLIGO curve, when using the predictions given in [213]. In the right panel of figure 21 we show the distance at which optimally oriented, optimally located, non-spinning, equal mass binaries would be detected with a signal-to-noise ratio (SNR) of 8 using both noise curves. This is commonly referred to as the *horizon distance*. The early AdV noise curve was rescaled by a factor of 1.61 so that the sensitive distance for a $(10 M_{\odot}, 10 M_{\odot})$ binary merger would be equal to the early aLIGO noise curve. This rescaling was found to better reflect the updated predicted sensitivities presented in [27]. The results in this paper are generated using the early aLIGO and rescaled early AdV sensitivity curves.

As with the initial science runs, we expect data taken from these detectors, in the absence of gravitational-wave signals, to be neither Gaussian nor stationary. It is important that search pipelines demonstrate an ability to deal with these features.

To simulate data with advanced detector sensitivities and with realistic non-Gaussian and non-stationary features, we chose to use data recorded by initial LIGO and Virgo and recolor that data to the predicted early sensitivity curves of aLIGO and Adv. The data we chose to recolor was data taken during LIGO’s sixth science run and Virgo’s second science run.

The procedure for producing such *recolored data* was accomplished in the following steps, which were conducted separately for the two LIGO detectors and Virgo.

- Identify a two-month duration of initial detector data to be recolored
- Measure the power spectral density (PSD) for each distinct section of science mode data using the PSD estimation routines in the `lal` software package [85].
- Calculate an *average PSD* over the two month period by taking, for every discrete value of frequency recorded in the PSDs, the median value over each of the PSDs in the set.
- Remove any line features from the resulting PSD and from the predicted early noise curves. This is done because it is difficult to remove or introduce line features from the data without introducing unwanted artifacts. Therefore it is simpler to remove line features in the PSDs, which will have the effect of preserving the line features of the original data into the recolored data.
- For each frequency bin, record the median value of the PSD over each section of science mode
- Take the ratio of the median PSD and the predicted early advanced detector noise curve. This is the reweighting to be used when recoloring.
- Using the time domain filtering abilities of the `gstlal` software package [214], recolor the data using this reweighting factor.

In figure 22 we show some examples of the PSDs obtained from recoloring the data and compare with the predicted sensitivity curves. As there are some small stretches of data in the original science runs where the sensitivity was significantly different from the average, we show the 10 % and 90 % quantiles as well as the maximum and minimum values for the PSD of the recolored data. We notice that the sensitivity




Figure 22 : Sensitivity curves of the recolored data for the LIGO Hanford detector (left) and the Virgo detector (right). In both cases the black dashed line shows the predicted 2015 – 2016 sensitivity curve (with the scaling factor added for Virgo). The dark colored region indicates the range between the 10 % and 90 % quantiles of the PSD over time. The lighter region shows the range between the minimum and maximum of the PSD over time.

of the detector still varies with time, as in the initial data, and that the lines in the initial spectra are still present.

Non-Gaussian features present in the original data will still be present in the recolored data, albeit distorted by the recoloring process. An example of this is shown in figure 23 where we show the SNR time-series around a known glitch in both the original and recolored data. While the recoloring does have some effect on the glitch, the two SNR time series are very comparable. As in searches on the original data, we attempt to mitigate the effect of such features. A set of *data quality flags* were created for the initial detector data [215, 216]. These attempt to flag times where a known instrumental or environmental factor, which is known to produce non-Gaussian artifacts in the resulting strain data, was present. To simulate these data quality flags in our recolored data we simply used the same flags that were present in the original data and apply them to the recolored data.

5.4 Injection Parameters

As an unbiased test of the process through which candidate events are identified for BBH waveforms, 7 BBH waveforms were added to the recolored data. The analysts




Figure 23 : SNR time series in a 20 s window around a known glitch in the original data (left) and in the recolored data (right). While the SNR time series clearly change, the primary features of the glitch are preserved across the recoloring procedure. These SNR time series were obtained by matched-filtering a short stretch of recolored and original data against a (23.7,1.3) M_{\odot} template.

were aware that “blind injections” had been added, however the number and parameters of these simulated signals were not disclosed until the analyses were completed. This was similar to blind injection tests conducted by the LIGO and Virgo collaborations in their latest science runs [174]. These injections are self-blinded to ensure that no bias from knowing the parameters of the signal, or indeed whether a candidate event is a signal or a noise artifact, affects the analysis process.

The 7 waveforms added to the data were taken from the numerical relativity simulations discussed in section 5.2. The parameters of the blind injections are given in Table 5. The distribution of physical parameters used in these blind injections was not intended to represent any physical distribution. Instead, the injections were chosen to test the ability to recover BBH systems across a wide range of parameter space. We describe the results of searches for these blind injections in section 5.6 and of parameter estimation studies on these signals in section 5.7.

As well as these blind injections, a large number of (non-blind) simulated signals were subsequently analyzed to obtain sufficient statistics to adequately evaluate the sensitive distances at which the NR waveforms could be detected in the early aLIGO and early AdV simulated data sets. As mentioned in section 5.2, each of the NR waveforms is valid only at the values of mass ratio and dimensionless spins for which it was generated. However, it is possible to change the total mass, which just rescales the waveform, and the various orientation and location parameters, which will only

Table 5 : The details of the blind injections that were added to the NINJA-2 datasets prior to analysis. In this table the Event ID will be used throughout the paper to refer to specific injections. The network SNR of each injection is denoted by ρ_N . This is the sum of the overlaps of the injection with itself in each detector, using 30 Hz as the starting frequency in the overlap integrals. M denotes the total mass and q the mass ratio. χ denotes the spin on *each* black hole, in all 7 cases both black holes in the binary had the same spin. RA and dec give the right ascension and declination of the signals respectively. Dist. denotes the distance to the source. Detectors online lists the detectors for which data is present at the time of signal. Hybridization range gives the range of frequencies in which the signal is hybridized between the post-Newtonian and numerical components. Waveform label indicates which numerical waveform was used, as shown in Tables 3 and 4.

Event ID	Waveform label	ρ_N	q	M (M_\odot)	χ	RA (rad)	Dec. (rad)	Dist. (Mpc)	Detectors Online	Hybrid Range (Hz)
1	J4+00+00_T4	23.9	4	124	0.00	1.26	-0.76	569	HLV	15 – 18
2	Ll1-20-20_T4	14.1	1	35.5	-0.20	1.70	-0.03	244	HLV	52 – 71
3	Ll1+40+40_T4	16.2	1	14.4	0.40	4.18	0.07	170	HLV	175 – 193
4	G2+20+20_T4	15.1	2	26.8	0.20	2.19	-0.36	247	LV	68 – 90
5	L4+00+00_T1	19.2	4	19.1	0.00	1.68	0.14	83	HV	86 – 93
6	J1+25+25_T4	16.9	1	75.7	0.25	4.68	0.49	854	HV	20 – 21
7	J1-75-75_T1	9.8	1	19.3	-0.75	0.81	-0.07	292	HLV	69 – 79

affect the overall phase and amplitude of the signal. For each of the 60 NR waveforms given in Table 3 and used in results in section 5.8, a set of ~ 42000 simulated signals was generated, necessarily with mass ratios and spins that have values corresponding to the given NR waveform. The total mass was chosen from a uniform distribution between 10 and 100 M_\odot . The simulations were distributed uniformly in distance, however they were not injected beyond a distance where they could not possibly be detected. The mass-dependent maximum distance that we chose to use is given by

$$D_{\max} = \left(\frac{\mathcal{M}}{1.219 M_\odot} \right)^{5/6} 175 \text{ Mpc.} \quad (5.3)$$

Here $1.219 M_\odot$ is the chirp mass of a $(1.4+1.4) M_\odot$ binary system. The factor of $\mathcal{M}^{5/6}$ describes, to leading order, how the SNR of the inspiral-only portion of a compact-binary merger at fixed distance will scale with mass when the inspiral is bandwidth-limited [217, 55]. 175 Mpc is chosen because it is larger than the distance at which it would be possible to detect a $(1.4+1.4) M_\odot$ binary merger with the early noise curves. However, to include a large margin for safety ~ 7000 of the signals were generated with chirp-weighted distances between 175 and 350 Mpc. The orbital orientations, polarization angles and sky directions are all chosen from isotropic distributions. The

signal coalescence times are drawn from a uniform distribution within our analysis window. Coalescence times were limited to times where at least two observatories were operating and no data-quality flags were active. The results of analyses on the non-blind simulated signals are given in section 5.8.

5.5 Search Pipelines

The goal of this work was to evaluate the detection sensitivity to binary black hole systems, modelled from the latest numerical simulations, using the search pipelines that were used to search for gravitational-wave transient signals in data taken during the final initial LIGO and Virgo joint observing run. The two pipelines that were used to do this were the dedicated compact binary coalescence (CBC) search pipeline “`ihope`” [17, 16, 18, 174, 19, 142] and the unmodelled burst pipeline “Coherent WaveBurst” (cWB) [218, 11, 12, 219]. The `ihope` pipeline was developed as a search pipeline for detecting compact binary mergers. It employs a matched-filtering algorithm against a bank of template waveforms [142]. The `ihope` pipeline was used to search for CBC systems (not just binary black holes) with component masses $\in [1, 99] M_{\odot}$. As a complement to template-based specialized searches, cWB was developed as an all-purpose un-modeled search pipeline, hence, it does not require *a priori* knowledge of the signal waveforms. It is better suited for burst signals spanning a small time-frequency volume. Moreover, due to the lack of model constraints, cWB is more adversely affected by background noise than matched-filter searches. Past simulation studies with initial LIGO sensitivity curves have shown that cWB was sensitive to CBC mergers with total masses $\in [25, 500] M_{\odot}$ over wide regions of the binary parameter space [220].

In addition to the `ihope` and cWB detection pipelines we also use parameter estimation algorithms to provide estimates of the parameters of compact binary systems observed with the detection algorithms. In the following section we provide a brief overview of the detection and parameter estimation pipelines. The results of running these searches on the data containing the NINJA-2 blind injections are presented in section 5.6 and parameter estimation results given in section 5.7.

5.5.1 Coherent WaveBurst

Coherent WaveBurst is a multi-resolution algorithm for coherent detection and reconstruction of gravitational wave bursts [161]. The cWB algorithm has been used in various LIGO-Virgo burst searches [218, 11, 12] and more recently in the search for intermediate mass black hole binaries [219]. Within the framework of the constrained maximum likelihood analysis [161], cWB identifies GW signals in data from multiple detectors and provides estimates of the signal parameters, e.g. sky location and waveforms. Along with the reconstruction of un-modeled burst signals, which imply random polarization, cWB can perform loosely modeled likelihood analyses assuming different polarization states, i.e. elliptical, linear or circular.

The NINJA2 cWB analysis uses the elliptical polarization constraint [220, 219] and searches for signals in the frequency band from 32 Hz to 1024 Hz. The analysis is performed in several steps: first, the data streams from all GW detectors are processed with the Meyer's wavelet transformations with 6 different time-frequency resolutions of $4 \times 1/8$, $8 \times 1/16$, $16 \times 1/32$, $32 \times 1/64$, $64 \times 1/128$, $128 \times 1/256$ [Hz × s]. Then the data are conditioned with a linear predictor error filter to remove power lines, violin modes and other predictable data components. Triggers are reconstructed as the coherent sets of samples (pixels) identified in the time-frequency data. For each trigger the coherent statistics are then computed. These include the network correlation coefficient, cc and the network energy disbalance, Λ , which are used to enable the signal consistency selection cuts. The cWB detection statistic is the coherent network amplitude, η , which is used to rank the events and thereby establish the significance against a sample of background events obtained with the time-shift analysis [161, 219, 220]. This shifting procedure is typically performed thousands of times in order to accumulate sufficient statistics.

5.5.2 `ihope`

The `ihope` pipeline is designed to search for gravitational waves emitted by coalescing compact binaries [142]. It has been optimized for and used in LIGO and Virgo GW searches over the past decade [221, 17, 16, 15, 174, 19], and also in the mock Laser Interferometer Space Antenna (LISA) data challenges [222]. The NINJA-2 `ihope` analysis uses the same pipeline-tuning that was used in the searches performed during

the final initial LIGO and Virgo joint observing run [174].

The pipeline matched-filters the detector data against a bank of analytically modelled compact binary merger waveforms [55, 142]. Only nonspinning compact binary merger signals are used as filters and the bank is created so as to densely sample the range of possible binary masses [223]. For each detector, the filtering stage produces a sequence of *triggers* which are plausible events with a high signal to noise ratio $\text{SNR } \rho$. The algorithm proposed in [224] is used to keep only those that are found coincident in more than one detector across the network, which helps remove triggers due to noise. Knowledge of the instrument and its environment is used to further exclude triggers that are likely due to non-Gaussian noise transients, or *glitches*. Periods of heightened glitch rate are removed (*vetoed*) from the analysis. The time periods where the rate of glitches is elevated are divided into 3 *veto categories*. Periods of time flagged by category 1 and 2 vetoes are not included in the analysis as known couplings exist between instrumental problems and the gravitational-wave channel during these periods. Periods of time vetoed at category 3 are *likely* to have instrumental problems. A strong gravitational-wave signal can still be detected during category 3 times, but including these periods in the background estimate can compromise our ability to detect weaker signals in less glitchy periods of time. For this reason the search is performed both before and after category 3 vetoes are applied. The significance of events that survived category 1-3 vetoes were calculated using the background that also survived categories 1-3. The significance of events that survived category 2 but were vetoed at category 3 were calculated using background that survived categories 1-2.

Signal based consistency measures further help distinguish real signals from background noise triggers in those that are not vetoed and pass the coincidence test. The χ^2 statistic proposed in [225] quantifies the disagreement in the frequency evolution of the trigger and the waveform template that accumulated the highest SNR for it, cf. Eq. (4.14) of [225]. We weight the SNR with this statistic to obtain the *reweighted* SNRs for all coincident triggers. The exact weighting depends on the mass range the search is focused on, cf. Eq. (17,18) of [142]. The reweighted SNR is used as the ranking statistic to evaluate the significance, and thus the false alarm rate (FAR), of all triggers.

Following the division of the mass-parameter space used in [174, 19], we performed

both *low mass* and *high mass* `ihope` searches on the NINJA-2 data. The low-mass search focused on binaries with $2M_{\odot} \leq m_1 + m_2 < 25M_{\odot}$, and used frequency domain 3.5PN waveforms as templates [88, 226, 89]. The high-mass search instead focused on the mass-range $25M_{\odot} \leq m_1 + m_2 < 100M_{\odot}$, and used the effective-one-body inspiral-merger-ringdown model calibrated to numerical relativity, as described in [94]. The exact χ^2 -weighting used to define the re-weighted SNR varied between the two analyses [142]. The significance of the triggers found by both was estimated as follows. All coincident triggers are divided into 4 categories, i.e. HL, LV, HV and HLV, based on the detector combination they are found to be coincident in [174]. They are further divided into 3 *mass*-categories based on their chirp mass $\mathcal{M}_c = (m_1 m_2)^{3/5} (m_1 + m_2)^{-1/5}$ for the low-mass search, and 2 categories based on their template duration for the high-mass search [174]. The rate of background noise triggers, or *false alarms*, has been found to be significantly higher for shorter signals from more massive binaries, and also to be different depending on the detector combination, and these categorizations help segregate these effects for estimation of the background [174, 142].

For all the triggers the combined re-weighted SNR $\hat{\rho}$ is computed, which is the quadrature sum of re-weighted SNRs across the network of detectors. All triggers are then ranked according to their $\hat{\rho}$ in each of the mass/duration and coincidence sub-categories independently, allowing us to estimate the trigger false alarm rate (FAR) at a given threshold $\hat{\rho} = \hat{\rho}_0$. This is described by

$$\text{FAR } (\hat{\rho}_0) \simeq \frac{N(\hat{\rho} \geq \hat{\rho}_0)}{T_c}, \quad (5.4)$$

i.e. the number of background noise triggers in a given coincidence sub-category at least as loud as the threshold, $N(\hat{\rho} \geq \hat{\rho}_0)$, divided by the total time analyzed for that sub-category, T_c . The limiting precision on this quantity is of order $1/T_c$; thus, in the limit where *no* background events are louder than $\hat{\rho}$, we quote a FAR of less than $1/T_c$. The calculation of trigger FARs is described in more detail in [227, 17]. As the smallest FAR we can estimate is $1/T_c$, to get a more precise estimate for our detection candidates we simulate additional background time. We shift the time-stamps on the time-series of single detector triggers by Δt relative to the other detector(s), and treat the shifted time-series as independent coincident background time. All coincident triggers found in the shifted times would be purely due to background noise. We

repeat this process setting $\Delta t = \pm 5s, \pm 10s, \pm 15s, \dots$, recording all the time-shifted coincidences, until Δt is larger than the duration of the dataset itself. With the additional coincident background time T_c accumulated in this way, we can get a more precise estimate of the low FARs we expect for detection candidates.

The FARs computed this way are further multiplied by a trials factor that accounts for the fact that we rank events in their own template-mass and coincidence sub-categories independently, while each of the sub-categories corresponds to an (independent) analysis of the same stretch of interferometric data. This factor is discussed in detail in section IV of [174]. Taking the trials factor into account, the final combined FAR (cFAR) is reported in Table 7, and the results are described in detail in section 5.6.2.

5.5.3 Parameter estimation

The detection methods described above produce times of interest where a gravitational wave may be present in the data (i.e. triggers), along with point estimates of the compact object masses from the signal, independently in each detector. These triggers are followed up with the goal of estimating the posterior probability density function of the parameters that describe the signal and to evaluate the evidence of different waveform models. In order to do so, we use Bayesian methods, in which the data from all detectors are analysed coherently.

The Bayesian parameter estimation algorithms used in this work provide estimates of the posterior probability distribution function. The probability of a set of parameters $\vec{\theta}$ under a model M given the observed data d can be written as

$$p(\vec{\theta}|d, M) = \frac{p(\vec{\theta}|M)p(d|\vec{\theta}, M)}{p(d|M)}. \quad (5.5)$$

Here $p(d|\vec{\theta}, M)$ is the likelihood of observing the measured data given the set of parameters $\vec{\theta}$, and $p(d|M)$ is the marginal distribution of the data under model M , commonly referred to as the evidence. When only concerned with parameter estimation the evidence is a normalization constant that can be ignored. It becomes relevant however, when comparing how well two models do in describing the observed data.

By marginalizing over all model parameters the evidence is obtained

$$p(d|M) = \int p(\vec{\theta}|M)p(d|\vec{\theta}, M)d\vec{\theta}. \quad (5.6)$$

Given the evidence of two competing models, M_1 and M_2 , the support for M_1 over M_2 can be quantified via the Bayes factor

$$\mathcal{B}_{12} = \frac{p(d|M_1)}{p(d|M_2)}. \quad (5.7)$$

These techniques require the generation of $\sim 10^6 - 10^7$ model waveforms to probe the 9 (for non-spinning black holes) to 15 (for fully spinning black holes) dimensional parameter space of compact binary systems in circular orbit, making it infeasible at present to use numerical relativity simulations. Instead approximate models (*e.g.* post-Newtonian, effective one-body) that are computationally cheaper to produce are used to estimate the parameters of a measured signal. Numerous studies have assessed the statistical uncertainty in compact binary parameter estimates [228, 229, 230, 231], which use the same approximate model for injection and analysis. Few studies have been done to quantify the systematic uncertainty in parameter estimates due to the use of these approximate models [232, 233]. Numerical relativity simulations provide us with the most accurate waveforms currently available, making them ideal for quantifying the systematic uncertainties inherent with using approximate models. This mock data challenge is the first time such a study has been conducted using models that account for the component angular momenta of the compact objects.

The Markov Chain Monte Carlo sampler `lalinference_mcmc` [234, 235], and two nested sampling implementations `lalinference_nest` [230] and `MultiNest` [236] from the *LALInference* package of the LSC Algorithm Library [85] were used to follow up GW candidates from the detection search pipelines. Due to the computational burden, we carried out the analysis with `lalinference_mcmc`, and as a consistency check for selected candidates and waveforms posterior estimates were also obtained with `lalinference_nest` and `MultiNest`. For model comparisons we have calculated the evidence by marginalizing each posterior estimate using thermodynamic integration.

Each candidate was analyzed using two distinct waveform models: `PhenomB` and

`EOBNRv2`. Both models describe the IMR phases of the GW from a compact binary merger. `EOBNRv2` models non-spinning binaries using the effective-one-body (EOB) that re-sums the PN dynamics and energy flux, and describes the merger-ringdown signal as a superposition of quasi-normal modes [166]. `PhenomB` is a phenomenological model with a PN description of the inspiral phase building up on test-mass terms to 2PN order, fit to a set of spinning and non-spinning PN-NR hybrid waveforms [167]. Waveforms are generated in the frequency domain and model binaries with component spins aligned with the orbital angular momentum through a single spin parameter $\chi \equiv (1 + \delta)\chi_1/2 + (1 - \delta)\chi_2/2$. Here $\delta \equiv (m_1 - m_2)/M$ and $\chi_i \equiv S_i/m_i^2$, where S_i denotes the angular momentum of the i th component of the binary, and M the total mass of the system.

The mass-ratio dependent and higher order terms used in `PhenomB` are calibrated to PN-NR hybrids that cover the late-inspiral, merger and ringdown. Therefore the accuracy of the model is expected to decline with decreasing total mass, as the inspiral phase of the waveform becomes a larger fraction of the total SNR of the signal, especially for comparable mass binaries. At the time of the analysis however, it was the only IMR waveform, including spin effects, that was computationally feasible for use, making it the most physically relevant waveform for the analysis. `EOBNRv2` is more computationally expensive, but has been shown to be accurate enough for uncertainties in parameter estimates to be dominated by statistical error, rather than systematic [237]. It only models binaries with non-spinning components however, so the model is only relevant for non-spinning injections. `SEOBNRv1` is the successor to `EOBNRv2` that accounts for (aligned) spin [113], however it is currently too computationally expensive to be used for parameter estimation.

Due to the lack of astrophysical constraints on compact binary systems, it is difficult to physically motivate any particular choice for the prior distribution of the intrinsic parameters (i.e. masses and spins). For this study we have chosen to use distributions that are uniform in component masses and component spin magnitudes over the range of parameter values being injected. The prior distribution was also flat in coalescence time across a 200 ms window centered on the trigger time, isotropic in orientation angles (e.g. inclination), and volumetric, giving equal prior probability to all spatial locations.

5.6 Blind Injection Challenge Results

In this section we present the results of using the detection pipelines described in section 5.5 to search for the blind injections listed in Table 5.

5.6.1 Coherent WaveBurst

For the NINJA2 cWB analysis it was decided *a priori* to search for GW bursts in the entire available times during which all three detectors were operating (17.9 days) and to discard the remaining times. First the search was performed on a total of 12,000 time-lagged observation times, accumulating 563.7 years of effective background live time. The background events that survived the data quality and analysis selection cuts (i.e. $cc > 0.7$ and $\Lambda < 0.4$) were used for calculation of the significance of candidate events. This background sample contains all reconstructed events, most of which do not resemble expected compact coalescence waveforms, as we do not enforce any waveform model. As a result, our background distribution is populated by relatively high signal-to-noise ratio events.

After completing the background analysis, the zero lag live time was analysed. The search detected an on-source event showing a chirping waveform compatible with a compact binary coalescence at a SNR ~ 22.1 and $\eta = 7.1$. The FAR of the candidate was estimated at $\sim 1/47\ yr^{-1}$ from comparison with the burst reference background, yielding a false alarm probability (FAP) of ~ 0.001 . After the parameters of the blind injections were disclosed, this event was revealed to be the first blind injection of Table 5. As a follow-up analysis, we investigated *a posteriori* all the times of the blind injections, as well as those on 2-fold exclusive live time. We found that the rest of the injected signals are either reconstructed with extremely low η or missed, see Table 6. For massive systems, such as events 1 and 6, the cWB algorithm recovers a large fraction of the injected signal-to-noise ratio. For lighter binaries, as expected, the algorithm is largely sub-optimal ¹.

Table 6 : The cWB search and follow-up results. The Event Labels correspond to those of each blind injection given in Table 5. This association is based on the time of the candidates relative to the time of the injections. M denotes the total mass. The false alarm probability, FAR, and false alarm probability, FAP, of each event are estimated by comparison with the empirically-calculated background distribution of the corresponding network of detectors. All but the first event are well within the bulk of the corresponding FAR distributions. η is the network correlated amplitude, which is the main cWB detection statistic. The injected network SNR (ρ_{inj}^{net}) is the square root of the quadratic sum of the optimal SNR in each detector. The recovered network SNR (ρ_{rec}^{net}) is the cWB estimate of the injected network SNR. Events 5 and 7 were completely missed due to a low reconstructed SNR and/or because of nearby noise glitches.

Event ID	Event Label	M	1/FAR (yr)	FAP	Network	η	ρ_{rec}^{net}	ρ_{inj}^{net}
1	J4+00+00_T4	124	47	0.001	HLV	7.1	22.1	22.8
2	Ll1-20-20_T4	35.5	-	-	HLV	2.8	9.1	13.9
3	Ll1+40+40_T4	14.4	-	-	HLV	2.7	9.2	15.7
4	G2+20+20_T4	26.8	-	-	LV	1.6	7.4	14.1
5	L4+00+00_T1	19.1	-	-	HV	-	-	18.5
6	J1+25+25_T4	76.7	-	-	HV	2.0	13.8	15.9
7	J1-75-75_T1	19.3	-	-	HLV	-	-	9.5

5.6.2 `ihope`

The results of the low-mass and high-mass `ihope` searches are presented in Table 7. The Event IDs correspond to the Event IDs of the blind injections in Table 5. The mapping between the `ihope` candidates and the blind injections is based on the event times of each. All injections except for injection 7 were found with high significance in one or both searches. Event 7 was missed because the injection’s SNR was too small to be detected by the pipeline. The optimal SNR of this injection — obtained by finding the overlap of the injection with itself — was 5.7 in H, 6.0 in L, and 5.3 in V, giving a network SNR of 9.8. However, the injected SNR in Virgo was below the SNR threshold used by the `ihope` pipeline ($= 5.5$). This means that, at best, the event could only surpass threshold in H and L, giving a maximum recoverable network SNR of 8.2; the false alarm rate at this network SNR is order 10^3 per year.

For this analysis we used the same vetoes as were used in [174] and [19] applied to the corresponding times in the recolored data. After veto categories 1-3 were applied, the total analyzed time consisted of 0.6 days of coincident HL data, 5.4 days

¹Lately, a lot of work has been devoted to extend the sensitivity of the algorithm to lower total masses, which is part of the on-going upgrades of cWB.

Table 7 : The `ihope` search results. The Event IDs correspond to the Event ID of each blind injection given in Table 5; this association is based on the time of the candidates relative to the time of the injections. The cFARs are calculated from all possible 5 s time shifts over the entire two-month dataset. M and q give the total mass and mass ratio respectively that were recovered in each detector. The recovered SNR (ρ_{rec}) and re-weighted SNR ($\hat{\rho}$) are reported separately for each detector. To calculate the cFARs, the quadrature sum of $\hat{\rho}$ was used. Unless noted, the cFARs were calculated after category 1-3 vetoes were applied.

Event ID	1/cFAR (yr)	Detectors	M	q	ρ_{rec}	$\hat{\rho}$	Search
1	≥ 6200	H	99.8	24.7	18.6	12.3	High mass
		L	94.7	13.8	14.5	13.2	
		H	37.1	3.22	6.6	4.9	
2	$\geq 10000^*$	L	42.7	3.99	9.6	9.6	High mass
		V	40.3	2.47	9.2	9.2	
		L	13.8	1.15	12.4	11.6	
3	≥ 23000	V	14.2	1.41	5.9	5.2	Low mass Cat. 3
		H	13.7	1.15	7.9	7.5	
		L	13.8	1.15	12.4	11.6	
4	≥ 31000	V	14.2	1.41	5.9	5.2	Low mass Cat. 2
		L	24.8	1.14	9.0	8.7	
		V	25.5	1.55	12.4	12.4	
5	≥ 23000	L	25.0	1.80	8.5	8.5	Low mass
		V	25.0	1.43	10.9	9.2	
		H	19.5	4.27	16.2	15.6	
6	≥ 21000	V	22.2	6.24	8.8	8.1	Low mass
		H	72.4	5.19	10.6	10.6	
		V	65.8	1.19	14.7	12.0	
7	<i>Not found</i>						

* Only used LV triggers for computing significance of this event; see section 5.6.2.

† Only used HL triggers for computing significance of this event; see section 5.6.2.

of coincident LV data, 6.5 days of coincident HV data and 8.9 days of coincident HLV data. FARs were calculated in each bin using the time-shift method described in section 5.5.2, then combined over all bins.

Table 7 also gives the total masses and mass ratios that were recovered by the `ihope` pipeline in each detector for each candidates. We see that the values reported by `ihope` can vary substantially from the injected parameters. This is not surprising: many of the injections had spin, and one injection (Event 1) was outside of the mass range covered by the template bank. We also see that the high-mass search deviates from the actual mass parameters more than the low-mass search. This, too, is expected since the template bank in the high-mass search is more sparsely populated. In general, templates are placed in `ihope` so as to maximize detection probability across the parameter space while minimizing computational cost. `ihope` therefore only provides a rough estimate of candidate parameters. For more precise estimates we use the parameter estimation techniques described in section 5.5.3, the results of which are presented in section 5.7. For compact binary systems where most of the SNR is obtained from the inspiral, `ihope` is expected to give a good estimation of the chirp mass of the system [67, 90, 176]. For the lowest mass systems in this study, `ihope` is not recovering the chirp mass to within 5% accuracy. For the higher mass systems the error on the chirp mass recovered by `ihope` grows to over 100% for event 1.

The greatest concern for a detection pipeline like `ihope` is whether the mismatch between templates and signals is small enough so as not to lose a substantial amount of re-weighted SNR. The templates used in this search were able to recover enough SNR of the blind injections to make them stand significantly above background. One might think that the majority of the recovered SNR comes from templates matching the PN part of the injected waveforms. However, figure 24 shows the SNR recovered by `ihope` in the PN and NR parts of the injections as a fraction of the total available SNR. We see that most of the available NR SNR is recovered in every event even though template waveforms did not have merger and ringdown (in the case of events recovered by the low-mass search), or were not calibrated to these particular numerical waveforms (in the case of events recovered by the high-mass search). To more rigorously determine what effect mismatch between templates and signals may have on detection sensitivity, we perform a large scale injection campaign with the

NINJA-2 waveforms in section 5.8.

Initially we used 100 time shifts to identify candidate events. All of the coincident events associated with the blind injections were louder than all background in the 100 time shifts. These were the only events to be louder than all background. Using 100 time shifts we could only bound the cFAR of the events to $\lesssim 10 \text{ yr}^{-1}$, which is not small enough to claim a detection. To improve our estimate, we performed as many 5 s time shifts as possible in the NINJA-2 dataset. This is the same method that was used for the blind injection described in [174].

Two blind injections were found in all three detectors: Event 2 in the high-mass search and Event 3 in the low-mass search (before category 3 vetoes were applied). Estimating background using the extended slide method with three detectors adds computational complexity, and has not previously been performed (the blind injection in [174] was only coincident in two detectors). However, in both Events 2 and 3 one of the three detectors had significantly less $\hat{\rho}$ than the other two (H in Event 2 and V in Event 3). We therefore did not include the detector with the smallest $\hat{\rho}$ when estimating the extended background for these two events.

Event 6 was vetoed at category 3. We therefore calculated its significance only after the first two veto categories were applied. All of the other events survived category 3 vetoes. Naïvely, we expect these events to have lower FARs if their significance is calculated after categories 1-3 have been applied. However, for Event 3 the trigger in the H detector was vetoed at category 3, leaving only L and V. Since the H trigger contributed a substantial amount of the combined re-weighted SNR, we might expect the resulting FAR to be *higher* for this event after category 3. A method to deal with partially vetoed events like this has not been proposed. We therefore simply report both results here.

Event 4 was found with high significance by both the high-mass and low-mass searches. This is not surprising as the injected total mass was 26.8 M_\odot , which is close to the boundary between the two searches. Currently no method has been established on how to combine the results from the low-mass and high-mass searches. We therefore give both results here.




Figure 24 : SNR recovered by `ihope` as a fraction of total available SNR in each detector for each found injection. Hatched bars give the percentage of available SNR in the PN part of the injection; solid bars give the percentage of available SNR in the NR part. Color bars indicate the amount of SNR recovered by `ihope`. “LM” indicates events recovered by the low-mass search — which used 3.5PN waveforms for templates — “HM” indicates events recovered by the high-mass search — which used EOBNRv1 waveforms for templates. The PN SNR is determined by terminating the matched filter at the frequency half-way between the hybridization range; the NR part is given by filtering from that frequency and up. Remarkably, both the low-mass and high-mass searches recovered most of the NR SNR despite the templates not having merger and ringdown (low mass) or not calibrated to the numerical waveforms used for the injections (high mass).

5.7 Parameter Estimation Results

For each of the blind injections described in section 5.4 and 5.6, parameter distributions were estimated using both non-spinning and spin-aligned models, as functions of eleven and nine parameters, respectively. `PhenomB` was used both as an aligned-spin model, as well as a non-spinning model by fixing the effective spin χ to 0, which we will refer to as `PhenomB $\chi=0$` . In addition, the `EOBNRv2` model was used as an additional non-spinning waveform (see section 5.5.3 for model descriptions). From these posterior estimates, one can determine the marginalized distributions in parameters of interest, as well as the evidence for the given model (see eq. (5.6)).

Figure 25 shows the 95% credible region of the marginalized posterior in component mass space for the non-spinning and spinning models. The credible region is systematically larger for the spinning model due to the strong degeneracy between the mass ratio and spin magnitude for aligned-spin models [175, 176], illustrated in figure 26.

In a real detection scenario, we can never be certain that a source contains only non-spinning components. Therefore to evaluate the ability to distinguish between ‘typical’ stellar-mass black holes and those occupying the mass gap ($\sim 3 - 5M_\odot$), we must restrict our attention to the spinning `PhenomB` model. Event 3 is a spinning system where both components have a mass of $7.18M_\odot$. Despite being well above the mass gap, the degeneracy of the spinning model results in constraining the lower-mass component to be outside of the mass gap with only 46% confidence. Event 5, on the other hand, contains a lower-mass component well inside the mass gap, with a mass of $3.83M_\odot$. Knowing a priori that this is a non-spinning system, we would be able to constrain the mass to be within the gap with 100% certainty. However, when including spin, the `PhenomB` model only constrains it to be within the gap with 21% certainty. The estimates of the masses and spin are summarized in Table 8. These results highlight the need to take spin into account when making any statements about compact object mass from GW data. We note that spin-aligned systems are the most extreme case of this degeneracy; if spins are mis-aligned with respect to the binary angular momentum the binary precesses, which causes phase and amplitude modulations in the observed waveform. This effect provides additional information

to break the degeneracy between masses and spins that may result in tighter constraints on the component masses. However, we expect the exact details to depend on the actual parameters of the observed systems (masses, spins and orientation of the angular momenta), and studies are ongoing to address this issue.

If the waveform and noise models exactly describe the data, then these Bayesian credible intervals would be equivalent to frequentist confidence intervals, meaning that the true parameters would fall within the 95% credible interval, for example, for 95% of injections. Any errors in the models however, will introduce systematic biases that break this equivalence. In these analyses such biases could be introduced both by the use of model waveforms that only approximate those that were injected, and by our assumption that the noise is purely Gaussian. Since the noise used for this study is real (recolored) noise recorded by the initial LIGO and Virgo detectors, the non-Gaussianities inherent with real noise can introduce bias to the parameter estimates [238]. Such biases are most apparent in the first event. This non-spinning injection was loud enough that systematic uncertainties between the numerical relativity and EOBNRv2 waveforms should be less than the statistical uncertainty [237]. Omega scan [239] spectrograms show there to be a significant glitch in the Hanford detector at the time of the injection (see Fig. 27(a)). An additional MCMC analysis was performed on the same injection made into noiseless data, using the same PSD as estimated for the event 1 analysis. Figure 27(b) shows the 95% credible region of this analysis to indeed constrain the injected values, leading to the conclusion that the non-Gaussian features of the detector noise led to significant systematic biases. Such biases, to varying degrees, are likely for any signal in noise that is not properly modeled. It will be crucial for better noise models to be implemented before the first detections in order to avoid these biases in parameter estimates.

The top of figure 28 shows the evidence for each model for the events analyzed. From the non-spinning ($\text{PhenomB}_{\chi=0}$) and spinning (PhenomB) evidences, the support for the presence of spin in each signal can be quantified using the Bayes factor, defined in equation (5.7). The bottom of figure 28 shows the Bayes factor and associated error estimates.

These model comparisons show strong support for the presence of spin for events 3 ($\chi_{\text{inj}} = 0.4$) and 4 ($\chi_{\text{inj}} = 0.20$). Event 7 had the greatest spin magnitude at

$\chi_{\text{inj}} = -0.75$, however analyses showed little evidence of it. This can likely be attributed to the low network SNR of event 7 (see table 5), which was partly due to the faster phase evolution of systems with spins counter-aligned to the orbital angular momentum [240].

There has been much work done to quantify the ability of ground-based detectors to localize GW sources on the sky [241, 27, 242, 243]. The accuracy of such localizations will be important for triggering electromagnetic (EM) followup of GW detections. Though most of this sky-localization work has focused on binary neutron star mergers due to their likely association with short gamma ray bursts [244, 245, 246] and numerous other proposed emission mechanisms (e.g. r-process [247], etc.), there is still much to be learned from accurate binary black hole merger localization. Since such mergers have never been observed, the possibility of an unexpected EM emission mechanism warrants the EM followup of the first detections.

Figure 29 shows the two dimensional marginalized distribution for the sky position of each event. Estimates of each sky position and the areas of the 95% credible regions are given in Table 9. The constraints on sky position are remarkably consistent across the models used. This is of great importance for low latency sky localization efforts, where time is of the essence. These results suggest that the more physically accurate, and computationally expensive, waveform models do not provide significantly more precise or accurate estimates of source sky positions.

5.8 Sensitivity evaluation

A large set of simulated signals, distributed as described in section 5.4, was used to assess the sensitivity of the pipelines to observe numerical relativity signals buried in data taken from Initial LIGO and Initial Virgo and recoloured to predicted early advanced detector observing runs, as described in section 5.3. Here, we use the CBC search pipelines (low-mass and high-mass `ihope`) to assess search efficiency in the chosen mass region: total mass between 10 and 100 solar masses.

Each signal in these large simulation sets was added at a random time to the 2-month period of recolored data. Coalescence times were limited to times where at least two observatories were operating and no data-quality flags were active. We then search for each signal using the `ihope` pipeline, as described in section 5.5. In

Table 8 : Summary of the mass and spin estimates for events 1-7 for each of the models used. Median values are quoted along with the upper and lower bounds of the 95% credible intervals. The injected values for each event are included for reference.

Event ID	Model	$m_1 (\text{M}_\odot)$		$m_2 (\text{M}_\odot)$		χ	
		Median	True	Median	True	Median	True
1	PhenomB	$79.5^{68.6}_{93.6}$	99.2	$58.2^{36.7}_{71.6}$	24.8	$-0.184^{+0.417}_{-0.128}$	0
	PhenomB $_{\chi=0}$	$81.3^{72.6}_{96.1}$		$63.1^{40.2}_{72.7}$		-	
	EOBNRv2	$83.2^{74.1}_{96}$		$64.3^{37.7}_{74.8}$		-	
2	PhenomB	$22.1^{17.4}_{37.8}$	17.8	$13.6^{8.43}_{17.3}$	17.8	$-0.18^{+0.37}_{-0.0999}$	-0.2
	PhenomB $_{\chi=0}$	$32.2^{20.1}_{36.8}$		$9.82^{8.15}_{16.8}$		-	
	EOBNRv2	$27.9^{19.9}_{33.2}$		$11.8^{9.32}_{17.5}$		-	
3	PhenomB	$11^{7.32}_{26.2}$	7.18	$4.82^{2.55}_{7.04}$	7.18	$0.497^{+0.415}_{-0.923}$	0.4
	PhenomB $_{\chi=0}$	$7.64^{6.83}_{9.16}$		$6.04^{5.04}_{6.75}$		-	
	EOBNRv2	$11.9^{11.5}_{12.6}$		$3.91^{3.7}_{4.03}$		-	
4	PhenomB	$19.8^{12.9}_{82.6}$	17.9	$8.26^{2.78}_{12.4}$	8.95	$0.294^{+0.161}_{-0.862}$	0.2
	PhenomB $_{\chi=0}$	$13.4^{12.3}_{15.8}$		$11.2^{9.46}_{12.2}$		-	
	EOBNRv2	$13.3^{12.4}_{15.4}$		$11.5^{9.89}_{12.3}$		-	
5	PhenomB	$9.12^{7.38}_{16.9}$	15.3	$5.9^{3.57}_{7.22}$	3.83	$-0.36^{+0.434}_{-0.1}$	0
	PhenomB $_{\chi=0}$	$15.1^{14.3}_{15.6}$		$3.89^{3.75}_{4.09}$		-	
	EOBNRv2	$15^{14.5}_{15.4}$		$3.92^{3.81}_{4.05}$		-	
6	PhenomB	$42.4^{35.9}_{54.4}$	37.9	$32.5^{24.4}_{38.9}$	37.9	$0.277^{+0.0582}_{-0.468}$	0.25
	PhenomB $_{\chi=0}$	$39.2^{33.1}_{49.7}$		$27.5^{21}_{32.7}$		-	
	EOBNRv2	$40.1^{33.9}_{51}$		$27.7^{19.5}_{33.7}$		-	
7	PhenomB	$16.3^{10.1}_{39.8}$	9.67	$6.27^{3.18}_{9.61}$	9.67	$-0.332^{+0.673}_{-0.286}$	-0.75
	PhenomB $_{\chi=0}$	$24.8^{18.1}_{29.7}$		$4.45^{3.61}_{7.17}$		-	
	EOBNRv2	$23.1^{13.3}_{25.7}$		$4.8^{4.22}_{9.62}$		-	

Table 9 : Summary of the sky location estimates for events 1-7 for each of the models used. Median values are quoted along with the upper and lower bounds of the 95% credible intervals, as well as the area of the 95% credible regions in the sky. The injected values for each event are included for reference.

Event ID	Model	α		δ		Sky Area (sq. deg.)
		Median	True	Median	True	
1	PhenomB	$1.28_{1.33}^{1.23}$	1.26	$-0.767_{-0.684}^{0.807}$	-0.757	32.4
	$\text{PhenomB}_{\chi=0}$	$1.27_{1.32}^{1.22}$		$-0.764_{-0.712}^{0.803}$		26.6
	EOBNRv2	$1.26_{1.32}^{1.21}$		$-0.763_{-0.722}^{0.801}$		22.5
2	PhenomB	$1.67_{1.74}^{1.6}$	1.7	$0.0402_{-0.163}^{0.0877}$	-0.0276	103
	$\text{PhenomB}_{\chi=0}$	$1.67_{1.74}^{1.6}$		$0.0453_{-0.163}^{0.0827}$		100
	EOBNRv2	$1.67_{1.74}^{1.6}$		$0.0406_{-0.164}^{0.0921}$		100
3	PhenomB	$4.21_{4.26}^{4.17}$	4.18	$0.00761_{-0.086}^{0.0587}$	0.0684	41.7
	$\text{PhenomB}_{\chi=0}$	$4.24_{4.29}^{4.18}$		$-0.0377_{-0.0776}^{0.11}$		65.7
	EOBNRv2	$4.24_{4.3}^{4.17}$		$-0.0388_{-0.0761}^{0.126}$		80.0
4	PhenomB	$2.19_{2.41}^{2.01}$	2.19	$-0.333_{-0.132}^{0.441}$	-0.36	430
	$\text{PhenomB}_{\chi=0}$	$2.3_{2.45}^{2.08}$		$-0.36_{-0.197}^{0.459}$		301
	EOBNRv2	$2.3_{2.45}^{2.06}$		$-0.344_{-0.14}^{0.451}$		470
5	PhenomB	$1.84_{2.6}^{1.72}$	1.68	$0.376_{-0.44}^{0.193}$	0.144	215
	$\text{PhenomB}_{\chi=0}$	$1.85_{2.62}^{1.74}$		$0.377_{-0.45}^{0.209}$		199
	EOBNRv2	$1.86_{2.63}^{1.74}$		$0.389_{-0.455}^{0.223}$		150
6	PhenomB	$4.5_{5.78}^{4.44}$	4.68	$0.141_{-0.521}^{0.247}$	0.49	207
	$\text{PhenomB}_{\chi=0}$	$4.52_{5.84}^{4.44}$		$0.233_{-0.542}^{0.241}$		294
	EOBNRv2	$4.56_{5.85}^{4.44}$		$0.309_{-0.551}^{0.192}$		342
7	PhenomB	$0.83_{1.67}^{0.684}$	0.806	$-0.121_{-0.159}^{0.532}$	-0.0736	551
	$\text{PhenomB}_{\chi=0}$	$0.836_{1.84}^{0.662}$		$-0.117_{-0.17}^{0.595}$		716
	EOBNRv2	$0.836_{1.02}^{0.653}$		$-0.135_{-0.0922}^{0.495}$		627

the plots that follow we treat a simulated signal as “detected” if there was no louder background event in the 100 time-slide trials that were performed to estimate the search background. As two `iHope` searches are performed, “low-mass” and “high-mass”, we treat simulated signals as detected if *either* search recovers the signal with more significance than its corresponding 100 background trials.

The results of the injection campaign are summarized in figure 30, which shows the dependence of the search sensitive distance on the chosen NR waveform and the total mass value. The sensitive distance is defined as the volume-weighted average distance up to which a CBC signal with a given waveform and total mass can be detected, where we average over the source extrinsic parameters and over the varying detector noise levels in the data set via Monte Carlo integration. In order to obtain a clear comparison between different NR waveforms, we used *the same set of random parameters* (including total masses, coalescence times and orientation angles) for each set of ~ 24000 injected signals, thus statistical errors will tend to be correlated between different waveforms.

In figure 30, top left, we plot the sensitive distance as a function of total mass for a number of equal mass NR waveforms with varying component spin magnitudes. The two waveforms with anti-aligned spins have smaller sensitive distances than the other waveforms (see also [19]). This is expected; systems with anti-aligned spin emit less power in gravitational waves than non-spinning systems [84, 167] and also may not match well with the template bank of non-spinning waveforms that was used [127]. At the largest masses in this study we can detect the two aligned-spin waveforms at distance $\sim 10 - 15\%$ larger than the non-spinning waveform. However, at the smallest masses used, the non-spinning waveform can be seen to comparable distances. Again this is a combination of two factors. Aligned-spin waveforms emit more power in gravitational waves than non-spinning systems, but the bank of non-spinning waveforms will not capture all of that power. This emphasizes that investigating the use of template waveforms incorporating spin effects is a necessity for the advanced detector era. To illustrate this further the dotted lines in this figure represent a prediction of the sensitivity that would have been obtained if a bank of aligned-spin template waveforms had been used. This prediction was obtained by taking the sensitive distances found for non-spinning NR waveforms recovered in the current CBC searches, which use non-spinning inspiral-only or IMR templates, and multiplying them by the ratio

of optimal [matched filter] SNRs of (anti)aligned-spin NR waveforms to non-spinning NR waveforms, as a function of total mass. This is described by

$$D_{\text{sens}}^i(M) = \frac{\sigma^i(M)}{\sigma^{\text{NS}}(M)} D_{\text{sens}}^{\text{NS}}(M). \quad (5.8)$$

Here i is used to denote the aligned-spin waveform for which the sensitive distance, D_{sens}^i , is to be predicted. NS is used to denote the non-spinning waveform. The optimal matched filter SNR is proportional to the signal power σ , given by the inner product

$$\sigma^i = (h^i | h^i)^{1/2}, \quad (5.9)$$

where h^i is the spinning waveform labelled by i (or h^{NS} for the non-spinning waveform) for an optimally oriented and located binary at fixed distance.

We can see from the plot that the distance sensitivity to spinning waveforms using a non-spinning bank is not as large as the predicted values from using (anti)aligned-spin template banks. This is especially true for the highly aligned-spin waveform where the predicted sensitive distance is $\sim 15\%$ larger than the obtained distance for systems with total mass $> 40M_\odot$. We note that the actual sensitivity improvement of a search using (anti)aligned-spin templates may not be as much as predicted here because such a search would require a larger number of templates and therefore provide more chances to obtain large SNR when matched-filtered against the underlying noise. Therefore the detection threshold when performing an aligned-spin search will increase with respect to a search using a non-spinning template bank. However, even a factor of 10 increase in the number of *independent* templates will only increase the expected SNR of the loudest background event by less than 5%, if Gaussian noise is assumed [178]. Work is ongoing to assess how much larger template banks of aligned-spin BBH waveforms are when compared to banks restricted to only non-spinning waveforms and to accurately compare the performance of such searches.

In the top right panel of figure 30 we plot the horizon distance as a function of total mass for a number of non-spinning NR waveforms with varying mass ratio. The sensitivity to these systems, for the same total mass, increases as the mass ratio (which we take to be greater than or equal to 1) decreases. This is because, to leading order, the gravitational-wave power emitted during the inspiral phase is dependent on the chirp mass [217]; for the same total mass, a system with higher mass ratio will

have a smaller chirp mass.

The bottom left panel of figure 30 shows the horizon distance as a function of total mass for the 5 non-spinning, equal-mass waveforms that were submitted. As expected, there is no significant discrepancy between these waveforms. We remind the reader that the same set of source parameters was used for each of these 5 sets of injections. Therefore statistical errors are strongly correlated between the results for different waveforms. The dashed and dotted black lines on this plot represent a prediction of the sensitive distance to non-spinning signals for the early aLIGO and early AdV noise curves respectively. This prediction was obtained by calculating the horizon distance, defined in section 5.3, for the $G1 + 00 + 00_T4$ waveform for both the early aLIGO and early AdV sensitivity curves as a function of mass. This measurement is then rescaled by a factor of 2.26 to account for the fact that the observatories do not have equal power to all orientations and sky locations [93]. We note that the obtained results agree well with this prediction, and fall between the early aLIGO and early AdV predictions when the two diverge. This is expected, as detection in `iHope` is dominated by the sensitivity of the second most sensitive detector operating at the time [142]. For times when at least two observatories were operating, the NINJA-2 dataset approximately consists of 50% of time when only one of the LIGO detectors and Virgo were operating and 50% of time when both LIGO detectors were operating, including time when Virgo is operating and when it is not. As the early aLIGO sensitivity does not drop below the early AdV sensitivity for the mass range considered, we expect the obtained sensitivity curve to lie roughly in the middle of the two predictions, and this is what we observe. It is worth pointing out that the `iHope` search is able to achieve this Gaussian-noise-predicted sensitivity, even though this analysis is run on real data, which includes non-stationarity and non-Gaussian transients.

Finally, in the bottom right panel of figure 30 we show sensitive distances for a number of waveforms with unequal spins, or unequal masses and non-zero spins.

5.9 Conclusion

This paper presents the first systematic study to assess the ability to detect numerically modelled binary black hole data in real data taken from Initial LIGO and Virgo

and recolored to predicted sensitivity curves of Advanced LIGO and Advanced Virgo in early observing runs. Building upon the work of the first NINJA project, this work, the culmination of the second NINJA project, studies the ability to do gravitational wave astronomy on a set of 60 binary black hole hybrid waveforms submitted by 8 numerical relativity groups.

In this work, a set of 7 numerically modelled binary black hole waveforms were added into the recolored data. This data was distributed to analysts with no knowledge of the parameters of the systems. The unmodelled gravitational waveform search pipeline, cWB, was able to recover one of these signals with an estimated false alarm rate of 1 every 47 years. The matched-filtered compact binary merger search pipeline, `ihope`, using a bank of BBH IMR waveforms, which were not calibrated against the NR signals used in NINJA-2, was able to recover 6 of the waveforms with false alarm rate upper limits ranging between 1 every 300 years and 1 every 2500 years.

A range of parameter estimation codes were run on the 7 blind injections that were added to the data used in this work. Though only results from the MCMC sampler were shown, these results proved to be statistically equivalent to estimates produce by the nested sampling and multinest samplers. These results demonstrate that it will be difficult to produce precise estimate of black hole component masses and spins because of intrinsic degeneracies between these parameters in the emitted waveforms. For some of the BBH blind injections we find that a neutron-star–black-hole coalescence cannot be ruled out. We also demonstrate the sensitivity of current parameter estimation algorithms to non-Gaussian features in the data and explore the ability to perform sky-localization of BBH observations.

A large-scale monte-carlo study was conducted to assess the efficiency of the `ihope` search pipeline as a function of the mass and angular momenta of the component black holes. We find that for non-spinning equal mass waveforms the sensitivity of the `ihope` search pipeline in real noise, including non-Gaussian artifacts, agrees well with predictions obtained using a Gaussian-noise assumption. We have found evidence that adding waveform models that include the effects of spin into the search pipeline will increase the efficiency of binary black hole observation. We have also demonstrated that the ability to recover numerical relativity waveforms, with identical parameters, but submitted by different groups, is indistinguishable up to statistical errors.

These results represent the next step for the NINJA collaboration; they address

shortcomings in NINJA-1 while paving the way for future work. In a sense this paper represents a baseline, as it measures the ability of current gravitational-wave analyses to detect and recover the parameters of an important subset of possible BBH signals in non-Gaussian noise in the advanced detector era. From this baseline there are multiple directions in which NINJA can expand. On the NR front, groups are continuing to fill in the parameter space. As shown in figure 20, even within the subspace of systems with (anti-)aligned spins there are large regions left to explore. Although NINJA-2 chose not to consider precessing signals many groups already have or are working on such simulations [212, 248, 249, 250, 251, 112, 252, 253, 173, 169]. Similarly, although the analyses used only the $\ell = m = 2$ mode in this work, it is expected that higher modes will be important for detection and parameter recovery [75, 254, 255, 123, 166]. Additional modes have been provided for many of the waveforms in the NR catalog, although they have not yet been validated. In all cases, as additional waveforms and modes become available they can be injected into the noise allowing for systematic tests of both detection and parameter estimation analyses.

In parallel the detection and parameter estimation analyses continue to evolve and improve. There is much development work ongoing to improve the analytical waveform models that are used in analysis pipelines, particularly for inspiral-merger-ringdown waveforms. It seems likely that before the first aLIGO and AdV observation runs generic fast IMR precessing analytic models will be available [39, 113, 256, 257, 169]. Improvements in how detection pipelines deal with non-Gaussianities are being explored to attempt to achieve the maximum possible sensitivity to BBH signals across the parameter space. A number of efforts are ongoing to implement aligned-spin waveform models into search algorithms. As we have demonstrated here, this will increase sensitivity to BBH systems with aLIGO and AdV [177, 127, 178]. Work is also underway to develop more realistic models of detector noise for parameter estimation pipelines, which account for the non-stationarity and non-Gaussianity present in real noise [258]. Accounting for such features is expected to greatly reduce systematic biases in the recovered masses and spins, such as those seen in event 1. The results presented here can provide a measure against which these next-generation analyses can be compared, in a way that measures not only their response to signals but also to realistic noise.




Figure 25 : The 95% credible regions for the EOBNRv2 (blue), PhenomB $_{\chi=0}$ (red), and PhenomB (green) models. Injected values are indicated by the “x” and the neutron star and mass gap regions are indicated where relevant. A strong degeneracy between spin and mass ratio results in systematic biases and artificially strong constraints on mass estimates when spin is ignored (i.e. EOBNRv2, PhenomB $_{\chi=0}$). By accounting for spin the PhenomB model produces estimates consistent with the injected sources.




Figure 26 : The 90% (dashed), 95% (solid), and 99% (dotted) credible regions of the two-dimensional marginalized posterior for the mass ratio and effective spin χ using the PhenomB model. The strong correlation between mass ratio and spin is responsible for the systematically weaker constraints placed on component masses when analyzing signals with an aligned-spin model.




Figure 27 : (a) shows a spectrogram of the noise in Hanford centered on the end time of the injection (before the blind injection is added). A non-Gaussian feature is present with peak energy slightly after the end time of the injection. (b) shows the 95% credible regions for the EOBNRv2 analysis of the event 1 injection in real noise (solid) and noiseless data (dashed). Injected values are indicated by the “x”. The fact that the injected values are well outside of the 95% credible region when real noise is present, and not in noiseless data, leads to the conclusion the non-Gaussian feature in the noise led to significant systematic biases in parameter estimates.




Figure 28 : (Top) Evidences for the non-spinning (EOBNRv2 and $\text{PhenomB}_{\chi=0}$) and spin-aligned (PhenomB) models. (Bottom) Bayes factors showing the support for the spin-aligned model over the non-spinning $\text{PhenomB}_{\chi=0}$ model. Events 3 and 4 are found to be spinning with high certainty. Event 7, with the largest spin magnitude, has little support for spin, though this is likely due to its low SNR.




Figure 29 : 95% credible regions for the sky position of all seven events, using the EOBNRv2 (blue), $\text{PhenomB}_{\chi=0}$ (red), and PhenomB (green) models. Despite the substantial differences between these models, their sky localization ability is very consistent.




Figure 30 : Volume-weighted average sensitive distance as a function of total mass, for a variety of NR waveforms. The sensitive distance is defined as the volume-weighted average distance up to which a CBC signal with a given waveform and total mass can be detected, where we average over the source extrinsic parameters and over the varying detector noise levels in the data set via Monte Carlo integration. *Top left*—Waveforms with equal component masses and various spins. Here the dotted lines represent predicted performances if an aligned-spin search had been performed. Details of this prediction are given in section 5.8. *Top right*—Waveforms with zero component spins and various mass ratios. *Bottom left*—Waveforms with equal-mass, non-spinning components. The dashed black line represents the predicted sensitivity curve of non-spinning waveforms with the aLIGO sensitivity curve, as described in 5.8. The dotted black line represents the predicted sensitivity with the AdV sensitivity curve. *Bottom right*—Waveforms with unequal spins, or unequal component masses and non-zero spins. The waveform designations are given in Table 3. These plots were generated using the data described in section 5.3 and the distribution of signals described in section 5.4.

Chapter 6

Modeling of intermediate mass-ratio inspirals: Exploring the form of the self-force in the intermediate mass-ratio regime

6.1 Introduction

Testing Einstein’s theory of general relativity in the strong-field regime by directly detecting the gravitational waves (GWs) emitted by black hole (BH) binaries of extreme mass-ratio has revived interest in a fundamental problem in general relativity: that of the gravitational self-force acting on a mass particle that moves in the background of a more massive BH. The gravitational self-force arises as a result of the back reaction between the small compact object and its own gravitational field, which, at linear order in mass-ratio, corresponds to a linear perturbation of the central BH geometry. Theoretical work in this field has been making steady progress since the seminal contributions of Dirac on the electromagnetic self-force in flat spacetime [259], and the extension of this analysis to curved spacetime by DeWitt and Brehme [260]. The generalization of these early studies to the case of the gravitational self-force was accomplished independently by Quinn and Wald [261] and Mino, Sasaki and Tanaka [262]. More recent developments have introduced mathematical rigor in the theoretical derivation of the gravitational self-force, and have relaxed previous assumptions

with regard to the internal structure of the small compact object, i.e, it can now be a small Kerr BH or a small compact object made up of ordinary matter [31, 32].

Likewise, the actual computation of the gravitational self-force has evolved from simplified scalar-field models [263], to more sophisticated solutions that involve electromagnetic and gravitational problems in the context of Schwarzschild circular orbits. At present, the self-force program has succeeded in developing numerical codes to compute the gravitational self-force along generic orbits around a Schwarzschild BH, and actually implementing these computations to develop an accurate waveform model that describes the inspiral evolution of non-spinning stellar mass BHs onto supermassive non-spinning BHs [264]. The development of numerical algorithms to compute the self-force on a scalar charge moving along an eccentric-equatorial orbit of a Kerr BH has also been accomplished [265]. The extension of this algorithm to compute the gravitational self-force for Kerr inspirals is under development [265, 266].

The fact that the self-force program has focused on the computation of the self-force for spinless particles that inspiral into more massive BHs has a physical rationale. It is not just that such a problem would be more difficult to solve. It has also been shown that in the context of extreme-mass ratio inspirals (EMRIs), with typical mass-ratios $1:10^6$, the inclusion of small-body spin corrections in search templates will not allow us to measure the small body spin parameter with good accuracy [267]. Hence, from a data analysis perspective, the inclusion of small body spin effects is not necessary. Additionally, before attempting to compute the self-force for spinning particles, one may need to address a more pressing modeling issue for spinless particles: it has been shown that including conservative self-force corrections in the orbital phase of EMRIs may not be necessary for source detection, but they may still be necessary for accurate parameter reconstruction. Additionally, second order radiative corrections may contribute to the phase evolution at the same level as first-order conservative corrections [?, ?]. This then suggests that a waveform template that aims to provide an accurate description of the inspiral evolution of EMRIs may have to include both first-order conservative corrections, and second-order radiative corrections, as pointed out in [264].

In sharp contrast, modeling BH binaries with intermediate mass ratio, i.e., $1:10\text{--}1:1000$, (IMRIs) presents new challenges that can be neglected in the EMRI arena. In

the absence of fully general relativistic gravitational self-force corrections in this mass-ratio regime, some studies have assessed the importance of including post-Newtonian self-force corrections in search templates for spinning BHs of intermediate-mass that inspiral into supermassive Kerr BHs. This work has shown that the implementation of first-order post-Newtonian self-force corrections for spin-spin and spin-orbit couplings is essential to ensure the reliability of parameter estimation results [268]. These studies suggest that the computation and implementation of gravitational self-force corrections for spinning binaries is a pressing, important problem both from a theoretical and data analysis perspective.

In this article we shed light on the importance of including gravitational self-force corrections in search templates for non-spinning stellar mass BHs that inspiral into Schwarzschild BHs of intermediate-mass. As, at present, we have no access to accurate self-force calculations in this mass-ratio regime, it is important to shed light on the regime in which current self-force corrections do not render an accurate dynamical evolution of GW sources, and explore the range of applicability of the information we have at hand to develop accurate waveform templates in that mass-ratio regime. This study is particularly important in view of the ongoing upgrade of the LIGO detector [269]. Once advanced LIGO (aLIGO) begins observations, it will be possible to target the inspirals of neutron stars (NSs) and stellar-mass BHs into intermediate-mass BHs with masses $\sim 50M_{\odot} - 350M_{\odot}$ [270] —events which may take place in core-collapsed globular clusters [271, ?]. To perform this study we will make use of the recently developed effective-one-body (EOB) model that has been calibrated using numerical relativity (NR) simulations for non-spinning BH binaries of mass-ratio $q = m_1/m_2 = 1, 2, 3, 4$ and 6 [38]. It is worth pointing out that the actual calibration of this EOBNRv2 (EOBNR version two) model reproduces with great accuracy the features of true inspirals, and hence we will use this model as a benchmark to explore the form of the self-force in the intermediate-mass ratio regime. By construction, the EOBNRv2 model reproduces results in the test-mass particle limit, and also encodes self-force corrections that have been derived for small mass-ratios. In our analysis we will explicitly show these important modeling ingredients when we compare the predictions made by the EOBNRv2 and those obtained through black hole perturbation theory (BHPT).

Finally, we will make use of the EOBNRv2 model and perturbative results to

present a new prescription for the orbital frequency shift at the innermost stable circular orbit (ISCO), originally derived in [272] in the context of EMRIs. Our prescription reproduces exactly the self-force prediction for extreme-mass ratios and provides an accurate prediction in the intermediate and comparable-mass ratio regimes.

This paper is organized as follows. In Section 6.2 we present a succinct description of the EOBNRv2 model. In Section 6.3 we derive an IMRI waveform model that accurately captures the features of true inspirals, as compared with the EOBNRv2 model. The derivation of this model will enable us to explore what the form of the self-force should be in the intermediate-mass-ratio regime so as to accurately reproduce the orbital dynamics obtained through NR simulations. In Section 6.4 we derive a new prescription for the gravitational self-force correction to the orbital frequency at the ISCO that encodes results from the extreme, intermediate and comparable-mass ratio regimes. Finally, we summarize our results in Section 6.5.

6.2 Effective One Body model

The Effective One Body (EOB) model was recently calibrated for non-spinning BH binaries of mass ratios $q = m_1/m_2 = 1, 2, 3, 4$ and 6 by comparison to NR simulations [38]. In this Section we briefly describe this model.

6.2.1 EOB dynamics

The EOB is a scheme that maps the dynamics of the two body problem in general relativity to that of one object moving in the background of an effective metric. In the non-spinning limit this metric takes the form

$$ds_{\text{eff}}^2 = -A(r) dt^2 + \frac{D(r)}{A(r)} dr^2 + r^2 \left(d\Theta^2 + \sin^2 \Theta d\Phi^2 \right), \quad (6.1)$$

where (r, Φ) stand for the dimensionless radial and polar coordinates, respectively. The conjugate momenta of these quantities is given by (p_r, p_Φ) . Since p_r diverges near the horizon, it is convenient to replace it by the momentum conjugate to the EOB *tortoise* radial coordinate r_* , i.e.,

$$\frac{dr_*}{dr} = \frac{\sqrt{D(r)}}{A(r)}. \quad (6.2)$$

Using this coordinate transformation, the effective EOB Hamiltonian can be written as [38]

$$H^{\text{eff}}(r, p_{r_*}, p_\Phi) \equiv \mu \hat{H}^{\text{eff}}(r, p_{r_*}, p_\Phi) = \mu \sqrt{p_{r_*}^2 + A(r) \left[1 + \frac{p_\Phi^2}{r^2} + 2(4 - 3\eta) \eta \frac{p_{r_*}^4}{r^2} \right]}, \quad (6.3)$$

where $\mu = m_1 m_2 / (m_1 + m_2)$, $M = m_1 + m_2$ and $\eta = \mu/M$ stand for the reduced and total mass of the system, and the symmetric-mass ratio, respectively. Additionally, the real EOB Hamiltonian is given by [38]

$$H^{\text{real}}(r, p_{r_*}, p_\Phi) \equiv \mu \hat{H}^{\text{real}}(r, p_{r_*}, p_\Phi) = M \sqrt{1 + 2\eta \left(\frac{H^{\text{eff}} - \mu}{\mu} \right)} - M. \quad (6.4)$$

Furthermore, to ensure the existence and η -continuity of a last stable orbit (ISCO) as well as the existence and η -continuity of an η -deformed analog of the light-ring (the last stable orbit of a massless particle), these metric coefficients must be Padé resummed. At present these coefficients are available at (pseudo) 5PN and 3PN order for $A(r)$ and $D(r)$, respectively,

$$D(r) = \frac{r^3}{(52\eta - 6\eta^2) + 6\eta r + r^3}, \quad (6.5)$$

and

$$A(r) = \frac{\text{Num}(A)}{\text{Den}(A)}, \quad (6.6)$$

with

$$\begin{aligned} \text{Num}(A) &= r^4 [-64 + 12a_4 + 4a_5 + a_6 + 64\eta - 4\eta^2] \\ &+ r^5 [32 - 4a_4 - a_5 - 24\eta], \end{aligned} \quad (6.7)$$

and

$$\begin{aligned}
\text{Den}(A) = & 4 a_4^2 + 4 a_4 a_5 + a_5^2 - a_4 a_6 + 16 a_6 + (32 a_4 \\
& + 16 a_5 - 8 a_6) \eta + 4 a_4 \eta^2 + 32 \eta^3 + r [4 a_4^2 + a_4 a_5 \\
& + 16 a_5 + 8 a_6 + (32 a_4 - 2 a_6) \eta + 32 \eta^2 + 8 \eta^3] \\
& + r^2 [16 a_4 + 8 a_5 + 4 a_6 + (8 a_4 + 2 a_5) \eta + 32 \eta^2] \\
& + r^3 [8 a_4 + 4 a_5 + 2 a_6 + 32 \eta - 8 \eta^2] \\
& + r^4 [4 a_4 + 2 a_5 + a_6 + 16 \eta - 4 \eta^2] \\
& + r^5 [32 - 4 a_4 - a_5 - 24 \eta] ,
\end{aligned} \tag{6.8}$$

where $a_4 = [94/3 - (41/32) \pi^2] \eta$, and a_5, a_6 are adjustable parameters which were determined by minimizing the inspiral phase difference between the NR and EOB (2,2) modes in [38].

The EOB equations of motion that describe the orbital dynamics of the BH binary are given by [273]

$$\frac{dr}{d\hat{t}} = \frac{A(r)}{\sqrt{D(r)}} \frac{\partial \hat{H}^{\text{real}}}{\partial p_{r_*}}(r, p_{r_*}, p_\Phi), \tag{6.9a}$$

$$\frac{d\Phi}{d\hat{t}} = \frac{\partial \hat{H}^{\text{real}}}{\partial p_\Phi}(r, p_{r_*}, p_\Phi), \tag{6.9b}$$

$$\frac{dp_{r_*}}{d\hat{t}} = -\frac{A(r)}{\sqrt{D(r)}} \frac{\partial \hat{H}^{\text{real}}}{\partial r}(r, p_{r_*}, p_\Phi) + {}^{\text{nK}}\hat{\mathcal{F}}_\Phi \frac{p_{r_*}}{p_\Phi}, \tag{6.9c}$$

$$\frac{dp_\Phi}{d\hat{t}} = {}^{\text{nK}}\hat{\mathcal{F}}_\Phi, \tag{6.9d}$$

where $\hat{t} \equiv t/M$, $\hat{\Omega} \equiv d\Phi/d\hat{t} \equiv M\Omega$ and the radiation-reaction force ${}^{\text{nK}}\hat{\mathcal{F}}_\Phi$ is given by [38]

$${}^{\text{nK}}\hat{\mathcal{F}}_\Phi = -\frac{1}{\eta v_\Omega^3} \frac{dE}{dt}, \tag{6.10}$$

with $v_\Omega \equiv \hat{\Omega}^{1/3}$. An important improvement in the EOB formalism over models that used Padé resummation of Taylor approximants to the energy flux is the implementation of a resummed energy flux of the form

$$\frac{dE}{dt} = \frac{v_\Omega^6}{8\pi} \sum_{\ell=2}^8 \sum_{m=1}^{\ell} m^2 \left| \frac{\mathcal{R}}{M} h_{\ell m} \right|^2, \quad (6.11)$$

where \mathcal{R} is the distance to the source and h_{lm} 's represent the multipoles of the waveform, defined through the following relation

$$h_+ - i h_\times = \frac{M}{\mathcal{R}} \sum_{l=2}^{\infty} \sum_{m=-l}^{m=l} Y_{-2}^{lm} h_{lm}, \quad (6.12)$$

where Y_{-2}^{lm} represent the spin weighted -2 spherical harmonics and h_+ and h_\times stand for the two gravitational wave polarizations. Since the numerical relativity simulations used to calibrate the EOB model were found to satisfy the condition $h_{\ell m} = (-1)^\ell h_{\ell -m}^*$ with great accuracy, where $*$ denotes complex conjugate, one can also assume that the analytical modes that enter the sum in Eq. 6.11 also satisfy this property. Furthermore, because $|h_{\ell,m}| = |h_{\ell,-m}|$, the sum in Eq. 6.11 extends only over positive m modes.

6.2.2 Modelling of the inspiral and plunge evolution

In the EOB scheme, the inspiral and plunge phases are described by the product of several factors, namely,

$$h_{\ell m}^{\text{insp-plunge}} = h_{\ell m}^F N_{\ell m}, \quad (6.13)$$

where the function $N_{\ell m}$ is introduced to ensure that the EOB model reproduces: a) the shape of the NR amplitudes $|h_{\ell m}|$ near their maxima; and b) the timelag between the maxima of the $|h_{\ell m}|$ and the maxima of $|h_{22}|$, obtained from NR data. On the other hand, the factorized resummed modes $h_{\ell m}^F$ are given by

$$h_{\ell m}^F = h_{\ell m}^{(N,\epsilon)} \hat{S}_{\text{eff}}^{(\epsilon)} T_{\ell m} e^{i\delta_{\ell m}} (\rho_{\ell m})^\ell, \quad (6.14)$$

where ϵ stands for the parity of $h_{\ell m}$, i.e., $\epsilon = 1$ if $\ell + m$ is even, and $\epsilon = 0$ for odd $\ell + m$.

The factor $h_{\ell m}^{(N,\epsilon)}$ stands for the Newtonian contribution, defined in Eqs.(15)-(18) of [38]. The remaining terms $\hat{h}_{\ell m}^{(\epsilon)} = \hat{S}_{\text{eff}}^{(\epsilon)} T_{\ell m} e^{i\delta_{\ell m}} (\rho_{\ell m})^\ell$ represent a resummed version of all PN corrections, which have the structure $\hat{h}_{\ell m}^{(\epsilon)} = 1 + \mathcal{O}(x)$, where x is the gauge-invariant object $x = \hat{\Omega}^{2/3}$.

Regarding the structure of the $\hat{S}_{\text{eff}}^{(\epsilon)}$ factor, we note that in the even parity case, which corresponds to mass moments, the leading order source of GW radiation is given by the energy density. Therefore, the source factor can be defined as $\hat{S}_{\text{eff}}^{(\epsilon=0)} = \hat{H}^{\text{eff}}(r, p_{r*}, p_\Phi)$ [274]. On the other hand, in the odd-parity case, which is associated to current modes, the angular momentum \hat{L}_{eff} turns out to be a factor in the Rege-Wheeler-Zerilli odd-parity multipoles in the limit of small mass-ratio η [274]. Hence, one can define $\hat{S}_{\text{eff}}^{(\epsilon=1)} = \hat{L}_{\text{eff}} = p_\Phi v_\Omega$ [38].

Furthermore, considering a Schwarzschild background of mass $M_{\text{ADM}} = H^{\text{real}}$, the tail term $T_{\ell m}$ is a resummed version of an infinite number of logarithmic terms that enter the transfer function between the near-zone and far-zone waveforms. Since this complex object only resums the leading logarithms of tail effects, one needs to introduce an additional dephasing factor $\delta_{\ell m}$, which is related to subleading logarithms. The final building block is given by $(\rho_{\ell m})^\ell$, which was introduced to enhance the agreement of the EOB model with NR in the strong-field regime. The explicit expressions for these various quantities can be found in Eqs. (19)-(21) and Appendix

B of [38].

Having described the building blocks of the EOB model, we will now describe how to go about in the actual construction of the EOB model using NR simulations. The first step in the calibration of the EOBNR model consists of aligning the waveforms at low frequency, following the procedure outlined in [38]. This approach is used to minimize the phase difference between the NR and EOB (ℓ, m) modes using the prescription

$$\mathcal{T}(\Delta t, \Delta\phi) = \int_{t_1}^{t_2} (\phi^{\text{EOB}}(t + \Delta t) + \Delta\phi - \phi^{\text{NR}}(t))^2 dt, \quad (6.15)$$

where $\Delta t/\Delta\phi$ are time/phase shifts, respectively, over which the minimization is performed. The time window (t_1, t_2) is chosen so as to maximize the length of the NR waveform, but making sure that junk radiation does not contaminate the numerical data.

The numerical h_{22} is usually called the leading multipolar waveform because, compared to all the other multipoles, it provides the leading contribution to the amplitude of the full waveform $h(t)$. Additionally, once the NR and EOB $(\ell, m) = (2, 2)$ modes are aligned using the prescription given by Eq. (6.15), the peak of the numerical h_{22} takes place at the same time the orbital frequency $\hat{\Omega}$ reaches its peak. Put in different words, the time at which the numerical h_{22} reaches its maximum and the EOB light-ring time (innermost circular orbit for a massless particle) are coincident.

To calibrate the EOB dynamics, determined by Eqs. (6.9a)-(6.9d), one minimizes the phase difference between the leading NR and EOB $(\ell, m) = (2, 2)$ modes during the inspiral phase. This phase minimization procedure enables us to constrain the value of the doublet (a_5, a_6) . As pointed out in [38], the calibration of the adjustable parameters (a_5, a_6) is not unique. For instance, [38] and [275] provide different values for the doublet (a_5, a_6) which reproduce with great accuracy NR simulations for equal and comparable-mass non-spinning BH binaries. Even if the calibration of these parameters is degenerate, one can instill some physics in the way (a_5, a_6) are determined. In [38], these parameters are modeled as smooth functions of η , such that they reproduce the self-force prediction of the orbital frequency shift at the ISCO in the test-mass particle limit $\eta \rightarrow 0$, i.e., for $a_6(\eta \rightarrow 0)/\eta$ and $a_5(\eta \rightarrow 0)/\eta$ the EOB

model reproduces the result [272]

$$M\Omega_{\text{ISCO}} = \frac{1}{6\sqrt{6}} (1 + 1.2512\eta + \mathcal{O}(\eta^2)). \quad (6.16)$$

It is worth pointing out that in contrast with the model developed in [276], the development of the EOBNRv2 model did not require the computation of higher-order PN corrections in the adjustable parameters ρ_{22} and δ_{22} . The actual expression for these two parameters had already been determined analytically in previous studies [275].

Having determined the EOB dynamics, the N_{22} coefficients are computed and implemented in the energy flux resummed prescription given by Eq. (6.11), one can determine the rest of the EOB adjustable parameters, namely, higher-order PN corrections in $\rho_{\ell m}/\delta_{\ell m}$ for $(\ell, m) \neq (2, 2)$, by minimizing the amplitude/phase difference between the remaining numerical and EOB multipolar waveforms used in the calibration. These higher-order corrections for $(\ell, m) \neq (2, 2)$ are included in the inspiral waveform prescription, Eq. (6.13), but not in the energy flux, Eq. (6.11).

6.2.3 Merger and ring-down calibration of the EOB model

The merger of two non spinning BHs generates a distorted Kerr BH whose gravitational radiation can be modeled using a superposition of quasi normal modes (QNMs), which are described by the indices (ℓ, m, n) , where (ℓ, m) denote the mode and n specifies the tone. Each of these modes has a complex frequency $\sigma_{\ell mn}$ given by

$$\sigma_{\ell mn} = \omega_{\ell mn} - i/\tau_{\ell mn}, \quad (6.17)$$

where the real/imaginary part $\omega_{\ell mn}/\tau_{\ell mn}^{-1}$ corresponds to the frequency/inverse damping time of each QNM. These two observables are uniquely determined by the mass and spin of the Kerr BH formed after merger [?]. The prescription used to compute these quantities is given by [38]

$$\frac{M_f}{M} = 1 + \left(\sqrt{\frac{8}{9}} - 1 \right) \eta - 0.4333\eta^2 - 0.4392\eta^3, \quad (6.18a)$$

$$\frac{a_f}{M_f} = \sqrt{12}\eta - 3.871\eta^2 + 4.028\eta^3. \quad (6.18b)$$

It is worth pointing out that a mode (ℓ, m) always consists of a superposition of two different frequencies/damping times. These ‘twin modes’ are given by $\omega'_{\ell mn} = -\omega_{\ell-mn}$ and $\tau'_{\ell mn} = \tau_{\ell-mn}$. However, when considering two initially non-spinning BHs, the mirror solutions are degenerate in the modulus of the frequency and damping time, and hence one has that $\omega_{\ell mn} > 0$ and $\tau_{\ell mn} > 0$.

Following [38], the merger-ringdown waveform may be written as follows

$$h_{\ell m}^{\text{merger-RD}}(t) = \sum_{n=0}^{N-1} A_{\ell mn} e^{-i\sigma_{\ell mn}(t-t_{\text{match}}^{\ell m})}, \quad (6.19)$$

where N is the number of overtones included in the model, i.e., $N = 8$, and $A_{\ell mn}$ are complex amplitudes which will be determined by smoothly matching the inspiral-plunge waveform (Eq. (6.13)) with its merger ring-down counterpart (Eq. (6.19)).

To determine the complex amplitudes $A_{\ell mn}$, one defines $t_{\text{match}}^{\ell m}$ as the time at the amplitude maximum of the $h_{\ell m}^{\text{EOB}}$ mode, namely, $t_{\text{match}}^{\ell m} = t_{\max}^{\Omega} + \Delta t_{\max}^{\ell m}$, and demand continuity of the waveform at $N - 2$ points sampled in the time range $[t_{\text{match}}^{\ell m} - \Delta t_{\text{match}}^{\ell m}, t_{\text{match}}^{\ell m}]$, and ensure the continuity and differentiability of the waveforms at $t_{\text{match}}^{\ell m} - \Delta t_{\text{match}}^{\ell m}$ and $t_{\text{match}}^{\ell m}$ (see Eqs. (36a)-(36c) in [38]).

Finally, the full waveform can be written as

$$h_{\ell m} = h_{\ell m}^{\text{insp-plunge}} \mathcal{H}(t_{\text{match}}^{\ell m} - t) + h_{\ell m}^{\text{merger-RD}} \mathcal{H}(t - t_{\text{match}}^{\ell m}). \quad (6.20)$$

where $\mathcal{H}(t)$ is the Heaviside step function.

This is the model we shall use in the following Section to develop an IMRI waveform model to explore the form of the self-force in the intermediate-mass ratio regime. We will consider three different systems with mass ratios 17:100, 10:100 and 1:100. We have chosen these systems because: a) the EOBNRv2 model was calibrated by comparison to NR simulations for events with mass-ratio 1:1-1:6, and hence systems with component masses 17:100 should be accurately modeled using EOBNRv2; b) the actual construction of the EOB model is such that it reproduces the expected dynamics of systems with small mass-ratio, e.g., $\sim 1:100$. This modeling statement will not be taken for granted in our subsequent analysis. We will show in the following Section that this is indeed the case by comparing results between EOBNRv2 and those obtained using Teukolsky data; c) we will explore an additional case, 1:10, for

which the EOBNRv2 is the best model currently available to shed light on the form that self-force corrections should have so as to reproduce the dynamical evolution of these type of GW sources. Studying these type of sources is also important to try to bridge the gap in the parameter space covered by current waveform models.

6.3 Self-force corrections in the intermediate-mass-ratio regime

In the previous Section, we described the model we shall now use to build a waveform model for intermediate-mass-ratio inspirals (IMRIs). Using this IMRI model we will explore whether the inclusion of available self-force corrections, which have been computed in the extreme-mass-ratio regime [277, 278, 272], are able to reproduce the inspiral evolution of intermediate-mass-ratio binary black holes.

6.3.1 IMRI self-force model

In this Section we introduce the key elements we need to build our IMRI model which is simple and flexible enough to explore the form of the self-force in the intermediate-mass-ratio regime, and that, at the same time is able to reproduce with great accuracy the binary’s dynamical evolution as predicted by EOBNRv2. In the following we assume that EOBNRv2 dynamics provides a good description of the actual binary black hole dynamical evolution. This is a reasonable assumption because, at present, EOBNRv2 is the best interface to translate NR simulations into a waveform model that uses coordinates which can be related to physical units [279]. This desirable modeling approach is obtained by comparing EOB gravitational waveforms to NR waveforms as seen by an observer at infinity [131]. The fact that this model has been calibrated to NR simulations of mass-ratios 1:1-1:6 also means that such a model provides the adequate arena for the studies we want to carry out. Having said that, we now describe the approach we shall follow to build our IMRI model:

1. We start with an ansatz for the orbital frequency evolution of our IMRI model which is inspired by recent studies on the inclusion of linear order self-force corrections in the EOB approach [280]. We develop this prescription so as to faithfully reproduce the EOBNRv2 orbital frequency evolution (See Figure 31). An accurate modeling of the orbital frequency is necessary to build the gauge-invariant object $x = \hat{\Omega}^{2/3}$, which is a crucial element in our analysis.

2. We propose an ansatz to model the IMRI gravitational wave angular momentum flux which we calibrate so as to faithfully reproduce its EOBNRv2 counterpart (see Figures 32 and 33). Note that the EOBNRv2 angular momentum flux used for this calibration is constructed by summing over 35 leading and subleading waveform multipoles $h_{\ell m}$, with $2 \leq \ell \leq 8$, $1 \leq m \leq \ell$. We include all these modes so as to model the radiative part of the self-force in our IMRI model as accurately as possible.
3. Having derived an accurate prescription for the orbital frequency and the angular momentum flux, we make use of a prescription for the angular momentum that goes beyond the test-mass particle limit and which includes conservative self-force corrections. This expression for $\hat{L}_z(x)$ encodes conservative self-force corrections in the redshift observable z_{SF} (see Eq. (6.29)).
4. Because the prescriptions for the orbital frequency and the flux of angular momentum of our IMRI model reproduce their EOBNRv2 counterparts with great accuracy (see Figures 31 and 32), we can now use these objects to constrain the form of the gauge-invariant expression of the angular momentum $\hat{L}_z(x)$ in Eq. (6.27) below by demanding internal consistency in our model, i.e., by reproducing faithfully the dynamical evolution of the binary black hole as predicted by EOBNRv2. This is equivalent to exploring the form of the red-shift observable z_{SF} which reproduces the binary's dynamical evolution as predicted by EOBNRv2. This calibration procedure enable us to reproduce the expected inspiral evolution point to point to better than one part in a thousand, as shown in Figures 34 and 35.
5. Having determined the form of z_{SF} that reproduces the dynamical evolution of the systems considered in this analysis, we compare the gauge-invariant expression of $\hat{L}_z(x)$ using our results and self-force corrections obtained in the context of EMRIs. We show that available self-force corrections do not reproduce the late inspiral evolution for systems with mass-ratio 1:6 and 1:10, but that they provide a fair description for systems with mass-ratio 1:100. This result is to be expected because these available self-force corrections were derived in the small mass-ratio limit.

Having described the approach to be followed, we start off by describing the ansatz we use to model the orbital frequency. To do this, we follow [275] and [281], and use the following prescription

$$\Omega = \frac{A}{\hat{H}} \frac{p_\phi}{r^2}. \quad (6.21)$$

Furthermore, assuming circular orbits, we can simplify Eq. (6.21) using the following relations for the conservative Hamiltonian \hat{H}

$$\hat{H} = \sqrt{A(u)(1 + j_0^2 u^2)}, \quad \text{with } p_\phi = j_0, \quad \text{and } j_0^2 = -\frac{A'(u)}{(u^2 A(u))'}, \quad (6.22)$$

where $u = 1/r$ and $'$ denotes d/du . Using a prescription similar to that introduced in [280], we shall use the following ansatz for the potential $A(u)$

$$A_{\text{ansatz}}(u) = 1 - 2u + \eta \left(\sqrt{1 - 3u} k_{\text{fit}} - u \left(1 + \frac{1 - 4u}{\sqrt{1 - 3u}} \right) \right), \quad \text{with } k_{\text{fit}} = 2u \frac{1 + a_1 u + a_2 u^2}{1 + a_3 u + a_4 u^2 + a_5 u^3}. \quad (6.23)$$

Under these assumptions, the prescription for the orbital frequency takes the simple form

$$\Omega(u)_{\text{ansatz}} = u^{3/2} \sqrt{-\frac{A'_{\text{ansatz}}(u)}{2}}. \quad (6.24)$$

The values of the a_i parameters in the function k_{fit} are given in Table 10. It is worth pointing out that this prescription captures accurately the orbital phase evolution of the sources described above all the way down to ISCO and slightly beyond with much less computational complexity than EOBNRv2. The actual comparison between this scheme and the EONBRv2 model is shown in Figure 31. Notice that our IMRI model does pretty well from large r to the fast-motion strong-field regime in all three cases shown. Notice that we are modeling the potential $A_{\text{ansatz}}(u)$ using the coordinate u and not M/x as done in [280]. The rationale for doing this is to use a simple, yet accurate, prescription for the orbital frequency that captures faithfully the EOBNRv2 orbital frequency evolution. We could have also used a different prescription for this object, i.e., a PN series [?], a Padé resummed expression, etc. The purpose this object

serves at this stage in the analysis is to reproduce accurately the orbital evolution predicted by EOBNRv2 with less computational complexity.

η	a_1	a_2	a_3	a_4	a_5
$\frac{1700}{13689}$	-7.458	15.0179	-7.208	12.152	6.103
$\frac{10}{121}$	-8.037	17.059	-7.826	14.589	5.393
$\frac{100}{10201}$	-9.535	22.984	-9.456	21.862	3.085

Table 10 : Ω_{ansatz} fit coefficients

Having derived a prescription for the orbital evolution, we require a prescription to generate the inspiralling trajectory of the stellar mass compact object. The first step to achieve this consists of deriving a consistent model for the flux of angular momentum. The model for the flux of angular momentum that we introduce in the following Section sums up the effect of including all the dominant and subdominant modes currently available in the literature, i.e., $2 \leq \ell \leq 8$, $1 \leq m \leq \ell$, with the advantage that the inclusion of extra modes does not impact the computational cost of generating the IMRI waveform.

6.3.2 Radial evolution prescription

An important component of the IMRI waveform model is the prescription used for the fluxes of energy and angular momentum. Deriving an accurate expression for these quantities is essential to capture the main features of true inspirals, as shown in [282] in the context of EMRIs.

As discussed in Section 6.2, in the EOBNRv2 model the flux of energy is constructed using the prescription given by Eq. (6.11), which is obtained by summing over the modes ($h_{\ell m}$) with $2 \leq \ell \leq 8$, $1 \leq m \leq \ell$. Using this prescription as input data and the fact that in the EOB formalism the following relation is fulfilled [280]

$$\dot{E} = \Omega \dot{L}_z, \quad (6.25)$$

we derive a prescription for the gravitational-wave angular momentum flux which is valid from early inspiral all the way to the ISCO and slightly beyond. This prescription, which encapsulates the contribution from 35 dominant and subdominant modes,

is inspired by the modeling of accurate EMRI waveform models introduced in [282], i.e.,

$$\left(\dot{L}_z\right)_{\text{fit}} = -\frac{32}{5} \frac{\mu^2}{M} \frac{1}{r^{7/2}} \left[1 - \frac{1247}{336} \frac{1}{r} + 4\pi \frac{1}{r^{3/2}} - \frac{44711}{9072} \frac{1}{r^2} + \frac{1}{r^{5/2}} \left(c_{2.5}^1 + c_3^2 \frac{1}{r^{1/2}} + c_{3.5}^3 \frac{1}{r} \right) \right], \quad (6.26)$$

and the coefficients in Eq. (6.26) are given in Table 11.

η	$c_{2.5}^1$	c_3^2	$c_{3.5}^3$
$\frac{1700}{13689}$	-121.903	551.141	-694.699
$\frac{10}{121}$	-110.657	360.261	-156.016
$\frac{100}{10201}$	-75.255	333.449	-363.505

Table 11 : \dot{L}_z fit coefficients

Figures 32 and 33 present the comparison between the EOBNRv2 flux, which includes all currently known dominant and subdominant modes, and the calibrated angular momentum flux (\dot{L}_z) . Notice that this computationally inexpensive scheme does pretty well from large r all the way to the ISCO and slightly beyond. Hence, this approach will enable us to capture the main features of the inspiral evolution of the systems under consideration in the regime of interest with good accuracy. Modelling the prescription of the flux of angular momentum in our IMRI waveform using all the dominant and subdominant modes in the EOBNRv2 model is equivalent to modeling the radiative part of the self-force with the best information currently available in the literature.

In Figure 33 we present a comparison between the flux of angular momentum predicted by the EOBNRv2 model and the fit to Teukolsky data proposed in [282] for extreme-mass-ratio inspirals. Note the remarkable agreement between both formalisms all the way down to the ISCO. This comparison also confirms that the construction of the EOBNRv2 captures the main features of inspirals with small mass-ratios which are modeled using black hole perturbation theory. This comparison also shows that the EOBNRv2 model encodes fairly well the radiative part of the self-force for systems with small mass-ratio.

Having found a prescription for the angular momentum that captures the dynamics of IMRIs, we can now use it to generate the inspiral trajectory of the stellar mass compact object that inspirals into an IMBH using the relation

$$\frac{dr}{dt} = \frac{dL_z}{dt} \frac{dr}{dL_z}. \quad (6.27)$$

The first term on the right hand side of Eq. (6.27) can be obtained from Eq. (6.26). With regard to the derivative of the angular momentum with respect to the radial coordinate, we need to use a prescription for the angular momentum that goes beyond the test-mass particle limit, as conservative self-force corrections play a more significant role in this regime. Hence, we use the relation derived by Barausse et al in [280] which includes conservative self-force corrections, i.e.,

$$\hat{L}_z(x) = \frac{L_z}{\mu M} = \frac{1}{\sqrt{x(1-3x)}} + \eta \left(-\frac{1}{3\sqrt{x}} z'_{\text{SF}}(x) + \frac{1}{6\sqrt{x}} \frac{4-15x}{(1-3x)^{3/2}} \right) + \mathcal{O}(\eta^2), \quad (6.28)$$

where $x = (M\Omega)^{2/3}$ and $'$ stands for d/dx . Furthermore,

$$z_{\text{SF}}(x) = 2x \frac{1 + b_1 x + b_2 x^2}{1 + b_3 x + b_4 x^2 + b_5 x^3}, \quad (6.29)$$

and the various coefficients b_i were derived in [280] using available self-force data. In this section we derive the coefficients in Eq. (6.29) that reproduce the actual inspiral evolution all the way down to the ISCO. Note that in Eq. (6.27) we have in place a prescription for the angular momentum flux which faithfully reproduces the expected loss of angular momentum even beyond the ISCO, as compared to the EOBNRv2 model. Hence, the only ingredient that needs to be tuned to reproduce the expected inspiral evolution is contained in Eq. (6.28). We have followed this approach to explore the form that $\hat{L}_z(x)$ should have. Put in different words, the form that the self-force redshift observable $z_{\text{SF}}(x)$ should have to reproduce an inspiral trajectory consistent with EOBNRv2 all the way down to the ISCO. The various coefficients of Eq. (6.29) that generate such an inspiral orbit are given in Table 12.

In Figure 34 we show that this prescription captures with great accuracy the features of EOBNRv2 inspirals. To recap, the IMRI model incorporates both first-order conservative corrections –through the construction of the gauge invariant expression of

η	b_1	b_2	b_3	b_4	b_5
$\frac{1700}{13689}$	-3.062	0.760	-3.261	0.550	-6.000
$\frac{10}{121}$	-3.260	0.892	-3.475	0.600	-6.829
$\frac{100}{10201}$	-3.370	1.570	-3.730	0.970	-7.280

Table 12 : $z_{\text{SF}}(x)$ fit coefficients

the angular momentum in Eq. (6.28)— and first-order radiative-corrections through the construction of the flux of angular momentum in Eq. (6.26). Notice that the latter object encodes the best information currently available in the literature, since it includes the contribution from 35 dominant and subdominant (ℓ, m) modes of the multipolar waveform to the angular momentum flux. Using this IMRI model, we have been able to constrain the form of the angular momentum $L_z(x)$ that reproduces the inspiral evolution predicted by the EOBNRv2 model.

In order to show the importance of increasing our knowledge of the self-force in the intermediate-mass-ratio regime, we present in Figure 36 three different curves which describe the time evolution of three binary systems during late inspiral. These plots show that if we use a waveform model (unfitted) that includes: a) an accurate prescription for the orbital frequency $\hat{\Omega}$; b) a prescription for the flux of angular momentum that incorporates the contribution from all dominant and subdominant (ℓ, m) modes; c) an invariant expression for the angular momentum using the object $x = \hat{\Omega}^{2/3}$ (see Eq. (6.28)), and; d) available self-force corrections to constrain the coefficients b_i [280], then such a model would generate an inspiral trajectory that deviates from the expected orbital evolution, in particular near the ISCO.

Figure 36 also shows that our model actually predicts the expected orbital evolution all the way down to the ISCO, in agreement with the EOBNRv2 model, and requires a different set of coefficients b_i in the redshift observable $z_{\text{SF}}(x)$, as compared with results in the extreme-mass-ratio limit quoted in [280]. Notice that this is not only a modeling issue. It is an indication that implementing available self-force data into IMRI waveform models will not render the correct inspiral evolution, at the very least for the cases we have considered. This is an important result of this paper. In order to substantiate this statement, in the following Section we will compute the value of the gauge invariant angular momentum $L_z(x)$ at the ISCO within both the self-force formalism and our IMRI model. We will also show that the evolution of

$L_z(x)$ is consistent between both formalisms during early inspiral, but that the form of this object differs as we near the ISCO.

One may also expect that for binaries with small mass-ratios, available self-force corrections may provide a fairly good description of the inspiral evolution. This is what we actually see in the bottom panel of Figure 36. We will also show in the following Section that the evolution of $L_z(x)$ for binaries with mass-ratio 1:100 is pretty consistent from early inspiral to the ISCO with EOBNRv2. This may not be surprising, since current self-force data have been obtained in the context of EMRIs and the EOBNRv2 has been calibrated so as to reproduce the dynamical evolution of binary black holes of small mass-ratio [274, 283, 284, 285]. This exercise then suggests that it may be necessary to go beyond first-order conservative corrections to reproduce accurately the expected dynamical evolution of intermediate-mass ratio systems with $\eta \sim 10^{-2} - 10^{-1}$.

To carry out the analysis described above in the following Section, we will start by using the EOBNRv2 model to compute the orbital frequency ISCO shift for binaries of mass ratio 1:1, 1:2, 1:3, 1:4, 1:5, 1:6, 1:10 and 1:100, along with perturbative results in the context of extreme-mass-ratio inspirals. We will use this expression for the orbital frequency ISCO shift to evaluate the gauge-invariant object x and then compute the value of the angular momentum at the ISCO using Eq. (6.28). We should also acknowledge the fact that EOB has not yet been calibrated using NR simulations of systems with mass-ratio 1:100. It is expected that EOBNRv2 provides a fair description of the dynamical evolution of these types of sources, and we show in the following section that both EOBNRv2 and perturbative calculations provide a very consistent modeling of the gauge-invariant angular momentum \hat{L}_z from early inspiral to the ISCO for binaries with mass-ratio 1:100. However, this study should be compared to accurate NR simulations of mass-ratio 1:100 [286] once these are generated with several gravitational waveform cycles before merger.




Figure 31 : The panels show the orbital frequency, as a function of the radial coordinate r , obtained using the fit described in the text (solid red line), and the orbital frequency predicted by the EOB model (solid black line). The systems shown in the panels correspond to binaries of component masses $17M_{\odot} + 100M_{\odot}$ (top-left panel), $10M_{\odot} + 100M_{\odot}$ (top-right) and $1M_{\odot} + 100M_{\odot}$ (bottom-left panel). The bottom-right panel demonstrates the accuracy with which our IMRI model reproduces the orbital frequency predicted by the EOBNRv2 for the systems 17:100 (dashed blue), 10:100 (dashed-dot red) and 1:100 (solid black). Note that the spikes are due to artifacts in the interpolation function used to plot the EOBNRv2 orbital frequency and the numerical fit used to reproduce it. Also notice that the discrepancy between the data and the fit is always smaller than one part in a thousand.




Figure 32 : We compare the calibrated flux of angular momentum described in Eq. (6.26) against the prediction of the EOBNRv2 model for binaries of component masses $17M_\odot + 100M_\odot$ (top left panel), $10M_\odot + 100M_\odot$ (top right panel), and $1M_\odot + 100M_\odot$ (bottom-left panel). The bottom-right panel shows the accuracy with which the IMRI model reproduces the EOBNRv2 angular momentum flux, $\dot{L} = dL_z/dt$, for the systems: 17:100 (dashed blue), 10:100 (dashed-dot red) and 1:100 (solid black). The spikes in the bottom-right panel are due to numerical artifacts of the interpolating function used to plot the EOBNRv2 angular momentum flux and the numerical fit to reproduce it. The fit is such that its discrepancy to the EOBNRv2 is always smaller than one part in a thousand.




Figure 33 : The left panel shows the angular momentum flux for binaries of mass ratio 1:100 computed using the EOBNRv2 model and the fit to Teukolsky data introduced in [282]. Both formalisms present a remarkable agreement all the way down to the ISCO. Note that the angular momentum flux fit to Teukolsky data is valid only from early inspiral until the ISCO. The right panel shows the relative difference between the two prescriptions for the angular momentum flux $\dot{L} = dL_z/dt$.




Figure 34 : The panels show the radial and azimuthal evolution for a $17M_\odot + 100M_\odot$ system (top panels), $10M_\odot + 100M_\odot$ system (middle panels), and $1M_\odot + 100M_\odot$ system (bottom panels), obtained using the IMRI model compared against the EOBNRv2 model.




Figure 35 : The panels show the accuracy with which the IMRI model proposed in the paper reproduces the radial (right panel) and azimuthal (left-panel) time evolution predicted by the EOBNRv2 model for the systems $17M_{\odot} + 100M_{\odot}$ (dashed blue), $10M_{\odot} + 100M_{\odot}$ (dashed-dot red), and $1M_{\odot} + 100M_{\odot}$ (solid black). Notice that our model reproduces the EOBNRv2 orbital and azimuthal evolution point to point with an accuracy better than one part in a thousand.




Figure 36 : The panels show the radial evolution using the EOBNRv2 model, the IMRI model described in the text, and a model (Unfitted) which is the same as the IMRI model described in text except for the fact that the coefficients used in Eq. (6.29) for the function $z_{\text{SF}}(x)$ were derived using available self-force data from extreme-mass ratio calculations [280]. The plots correspond to binaries of component masses $17M_{\odot} + 100M_{\odot}$ (top-left panel), $10M_{\odot} + 100M_{\odot}$ (top-right panel) and $1M_{\odot} + 100M_{\odot}$ (bottom-left panel). The bottom-right panel shows that the orbital evolution predicted by a model that incorporates self-force corrections from extreme-mass-ratio inspirals (EMRIs) deviates from the orbital evolution predicted by the EOBNRv2 model at late inspiral. This discrepancy is more noticeable for systems with mass-ratios 17:100 (dashed blue) and 10:100 (dashed-dot red). Furthermore, for smaller mass-ratios, 1:100 (solid black), the discrepancy becomes comparatively smaller, as expected.

6.4 ISCO shift: connecting the extreme, intermediate and comparable-mass ratio regimes

In the previous Sections we have mentioned that during the inspiral of a stellar mass compact object of mass m_2 into a supermassive BH of mass m_1 , the radiative part of the self-force drives the inspiral evolution of the small object, whereas its conservative part has a cumulative effect on the orbital phase evolution [287]. These two effects have been considered in the development of EMRI waveform templates [288, 289, 290, 291, 282, 292, 293].

We shall now consider a novel effect that was explored by Barack & Sago [272]. They have shown that the self-force also introduces shifts in the innermost stable circular orbit radius and frequency. For a test-mass particle, these two quantities are given by

$$r_{\text{ISCO}} = 6m_1, \quad m_1\Omega_{\text{ISCO}} = \frac{1}{6\sqrt{6}}, \quad (6.30)$$

whereas, for finite η , in the Lorenz gauge, these two quantities take the form [272]

$$\Delta r_{\text{ISCO}} = -3.269(\pm 3 \times 10^{-3})m_2, \quad \frac{\Delta\Omega_{\text{ISCO}}}{\Omega_{\text{ISCO}}} = 0.4870(\pm 6 \times 10^{-4})\frac{m_2}{m_1}. \quad (6.31)$$

Since Barack & Sago carried out these calculations in Lorenz gauge, it was necessary to translate these results into coordinates that are commonly used for GW observations, i.e., asymptotically flat coordinates. This exercise has been done for the orbital frequency, which is a gauge invariant object, and hence can be compared to results obtained in alternative formalisms, such as PN theory or the EOB approach. In [277] and [279], the authors derive the ‘renormalization’ factor that translates results of Lorenz-gauge calculations into physical units. Applying this renormalization technique, one finds that the ISCO frequency is given by

$$M\Omega_{\text{ISCO}} = \frac{1}{6\sqrt{6}}(1 + 1.2512\eta + \mathcal{O}(\eta^2)), \quad (6.32)$$

with $M = m_1 + m_2$. This prediction can be compared with PN-based ISCO calculations at different orders of accuracy [294]

$$\begin{aligned} M\Omega_{\text{ISCO}}^{\text{2PN}} &= \frac{1}{6\sqrt{6}} \left(1 + \frac{7}{12}\eta + \mathcal{O}(\eta^2) \right), \\ M\Omega_{\text{ISCO}}^{\text{3PN}} &= \frac{1}{6\sqrt{6}} \left(1 + \left(\frac{565}{288} - \frac{41}{768}\pi^2 \right)\eta + \mathcal{O}(\eta^2) \right). \end{aligned} \quad (6.33)$$

The EOB approach has also been used to describe the orbital frequency shift at ISCO. Damour suggested in [279] that a fit for the ISCO orbital frequency shift that incorporates results from the gravitational self-force and NR simulations may be a quadratic polynomial in η of the form [279]

$$M\Omega_{\text{ISCO}}^{\text{Damour}} = \frac{1}{6\sqrt{6}} (1 + 1.25\eta + 1.87\eta^2). \quad (6.34)$$

In this paper we build up on this analysis and update this estimate using results from the self-force program, and making use of the EOBNRv2 model, which has been calibrated to NR simulations [38]. The prescription for the orbital frequency ISCO shift that we propose below reproduces accurately the results predicted by the self-force program for EMRIs, and also reproduces the best data currently available for intermediate and comparable-mass systems. To compute the ISCO orbital frequency shift within the EOB formalism, we use the equation derived in [279], namely,

$$2A(u)A'(u) + 4u(A'(u))^2 - 2uA(u)A''(u) = 0, \quad (6.35)$$

where $u = 1/r$ and $'$ stands for d/du . This ISCO condition can be rewritten in terms of the radial coordinate as [294]

$$rA(r)A''(r) - 2r(A'(r))^2 + 3A(r)A'(r) = 0, \quad (6.36)$$

where $' = d/dr$. We shall use the metric coefficient $A(r)$ quoted in [38], which includes the Padé expression for $A(r)$ at 5PN order. Having obtained the value for r_{isco} using Eq. (6.36), we evaluate the angular orbital frequency $M\Omega_{\text{ISCO}}$ at this fiducial value using Eq. (10b) of [38] with $p_r = 0$.

Using a variety of events, including extreme, $\eta \sim 10^{-5}$, and intermediate, $\eta \sim 10^{-2} - 10^{-1}$, mass ratio inspirals, we derive a quadratic polynomial fit in η for the

ISCO orbital frequency shift for these types of events, namely

$$M\Omega_{\text{ISCO}}^{\text{fit}} = \frac{1}{6\sqrt{6}} (1 + 1.05786\eta + 2.12991\eta^2). \quad (6.37)$$

We found that at second order in η , this numerical fit does not reproduce accurately the orbital frequency ISCO shift for small-mass ratios [272]. We can fix these problems using a prescription of the form

$$M\Omega_{\text{ISCO}}^{\text{fit}} = \frac{1}{6\sqrt{6}} (1 + 1.2512\eta - 0.0553751\eta^2 + 5.78557\eta^3). \quad (6.38)$$

At this level of accuracy we exactly reproduce the prediction for EMRIs for small η , as well as the most-up-to-date results for binaries modeled using the EOBNRv2 scheme. We compare the range of applicability of this numerical expression, along the other various approximations mentioned above, in Figure 37.




Figure 37 : ISCO shift using various approximations as described in the text. The ‘Numerical Data’ has been obtained using calculations in the extreme-mass ratio regime [272] and the EOBNRv2. The prescription that encapsulates results from the extreme, intermediate and comparable mass-ratio regime is labeled as ‘Numerical Fit’ and is given by Eq. (6.38) in the main text.

With our prescription for $M\Omega_{\text{ISCO}}$, we can evaluate the value of the gauge-invariant angular momentum flux at the ISCO. To do so, we compute the value of the angular momentum, see Eq. (6.28), using available self-force data (SF Fit), i.e., we use Eq. (6.32) to compute the shift in the orbital frequency at the ISCO, and then use Eq. (6.28) in conjunction with the values for the b_i coefficients of Eq. (6.29) quoted

in [280], i.e., self-force corrections derived in the extreme-mass-ratio limit. We also present results for an ‘incomplete model’, in which we use Eq. (6.38) to evaluate the value of the orbital frequency at the ISCO, and then use Eq. (6.28) with the set of b_i coefficients quoted in [280]. Finally, the IMRI model encodes all the results derived in this paper, namely, we use the prescription for the shift of the orbital frequency at the ISCO in Eq. (6.38), and the prescription for the angular momentum (Eq. (6.28)) using the corrections quoted in Table 12. Table 13 shows that, in accord with Figure 36, for binaries with symmetric mass-ratio $\eta \sim 0.01$, the value of the angular momentum evaluated at ISCO is fairly consistent between the two models. However, the predicted value of the angular momentum at ISCO becomes more discrepant for binaries with $\eta \sim 0.1$. This then suggests that the evolution of intermediate-mass ratio inspirals cannot be fully captured by using self-force calculations from extreme-mass ratio inspirals. This can be better visualized in Figure 38 where we show the angular momentum during inspiral all the way down to the ISCO using two formalism, namely, the IMRI prescription and a model that includes available self-force corrections, which is labeled as ‘Self Force’.

η	$L_z(x_{\text{ISCO}})$		
	SF Fit	Incomplete	IMRI
$\frac{1700}{13689}$	3.3389	3.3522	3.2918
$\frac{10}{121}$	3.3876	3.3878	3.3254
$\frac{100}{10201}$	3.4561	3.4561	3.4429

Table 13 : The Table shows the value of the angular momentum L_z as a function of the gauge invariant object $x = (M\Omega)^{2/3}$ evaluated at the ISCO radius (see Eq. (6.28)). The SF (self-force) values are computed using the prescription given in Eq. (6.32) to compute the orbital frequency shift, and Eq. (6.28) with the coefficients b_i quoted in [280], i.e., evaluated in the context of extreme-mass-ratio inspirals. Incomplete stands for a prescription in which we use the prescription for the orbital frequency given by Eq. (6.38), and the angular momentum prescription given by Eq. (6.28) with the coefficients b_i quoted in [280]. The b_i values of the IMRI model are obtained using a model that reproduces the dynamical evolution of intermediate-mass-ratio inspirals, as compared with the EOBNRv2 model, and for which we have derived a new prescription for the red-shift observable $z_{\text{SF}}(x)$ which actually reproduces the features of true inspirals.

Table 13 and Figures 36, 38 show that as the mass-ratio increases, the discrepancy between a model that incorporates available self-force corrections [278] and one that has been calibrated to NR simulations becomes more pronounced. For events with

mass-ratio 1:10 this difference looks slightly larger than for those of mass-ratio 17:100. Part of the reason for this behavior may be the fact that the model we have used to perform this analysis, EOBNRv2, provides the best prescription currently available for the inspiral evolution of sources with mass-ratios 1:1-1:6. It is expected that the model provides a good description of the inspiral evolution of sources with mass-ratio 1:10, and hence we have extended its realm of applicability to shed light on the form that the self-force is expected to have so as to reproduce the inspiral evolution of these type of sources. Thus, a conservative conclusion we can draw at this stage is that, using the best waveform model currently available in the literature, we have shown that one may not be able to accurately model the dynamics of sources with mass-ratio 1:10 using available self-force calculations. This trend is also present in the case of events with mass-ratio 17:100 and the results in this case are more conclusive. We have shown that a waveform model that includes available self-force corrections will not be able to capture faithfully the inspiral evolution of these GW sources. These results suggest that one may need to include higher-order corrections in the self-force to capture the behavior of true inspirals in the intermediate-mass-ratio regime. Finally, our work is also a consistency check on the internal structure of the EOB model, since we have shown that the EOBNRv2 renders a good prescription for the inspiral evolution via the flux of angular momentum and that the angular momentum prescription is consistent with results obtained from perturbative calculations.




Figure 38 : The panels show the invariant angular momentum $L_z(x)$, with $x = (M\Omega)^{2/3}$, from early inspiral to the ISCO using two different prescriptions. The ‘IMRI’ prescription reproduces accurately the expected inspiral evolution, as compared with results from the EOBNRv2 model—which has been calibrated to NR simulations. The ‘Self-Force’ prescription encapsulates self-force corrections derived in the context of EMRIs [278, 280]. Note that for the three binary systems studied, $17M_\odot + 100M_\odot$ (top-left panel), $10M_\odot + 100M_\odot$ (top-right panel), and $1M_\odot + 100M_\odot$ (bottom-left panel), the ‘Self Force’ prescription for the angular momentum presents a deviation from its expected value that becomes more noticeable near the ISCO. The bottom-right panel shows the fractional accuracy between the two different prescriptions for the angular momentum for the systems 17:100 (dashed blue), 10:100 (dashed-dot red) and 1:100 (solid black). Notice also that, as expected, for binaries with small mass-ratio the ‘Self Force’ and ‘IMRI’ prescriptions are fairly consistent all the way down to the ISCO. As in the previous plots, the spikes in the bottom-right panel are due to numerical artifacts.

6.5 Conclusions

In this paper we have developed a code to reproduce the analysis presented in [38]. Using this code we generated the inspiral evolution of three different systems to perform an exploratory study of the importance of including accurate self-force corrections in search templates that aim to detect non-spinning intermediate-mass ratio inspirals. The choice of the systems to perform this study reflects the knowledge we have at present on the dynamical evolution of non-spinning BH binaries ($17M_{\odot} + 100M_{\odot}$), what we want to know ($10M_{\odot} + 100M_{\odot}$), and a special system that provides reassurance that the dynamics of intermediate-mass-ratio inspirals with typical mass-ratios $\eta \sim 0.01$ can be captured using perturbative theory, as shown in [286, 295].

The EOBNRv2 model we have used as a benchmark to carry out this study has the advantage of encoding the best information currently available of non-spinning BH binaries, is capable of reproducing the expected dynamics in the test-mass particle limit, and also includes corrections taken from self-force corrections in the extreme-mass-ratio limit. We have confirmed these statements for systems with $\eta \sim 0.01$ by showing that the EOBNRv2 does predict the expected form of the angular momentum flux, as compared to Teukolsky data, as well as the evolution of the gauge-invariant angular momentum, as compared to self-force data.

In order to explore the form of the self-force in the intermediate-mass-ratio regime, we developed an IMRI model that reproduces the inspiral evolution predicted by the EOBNRv2 model, and which enable us to explore the form that the self-force should have in this mass-ratio regime so as to reproduce the binary's dynamical evolution as predicted by the best available interface to NR simulations. We have found that for systems with component masses $17M_{\odot} + 100M_{\odot}$, available self-force corrections do not accurately reproduce the inspiral evolution. We have shown that there is a clear deviation from the true inspiral trajectory near the ISCO. We have also explored this issue in greater detail by showing that the gauge-invariant angular momentum does deviate from the current self-force prediction near the ISCO. When we extend the realm of applicability of the EOBNRv2 to binaries of mass-ratio 1:10, we observe a similar behaviour. This exploratory study then suggests that it may be necessary to extend conservative corrections beyond the linear order to accurately capture the true inspiral evolution for sources with intermediate-mass-ratio. Furthermore, once

NR simulations of systems with mass-ratios 1:10, 1:15 and 1:100 have reached enough resolution near merger, we will be in a good position to calibrate the EOB model so as to reproduce the true dynamical evolution for these type of sources [286, 296, 297]. Such a model will enable us to further probe the parameter space to develop IMRI models, as the one proposed in this paper, that capture the features of true inspirals at a very inexpensive computational cost.

We have also used our EOBNRv2 code to derive a prescription for the orbital frequency ISCO shift that encapsulates results from the extreme, intermediate and comparable-mass ratio regimes. Our prescription is the first in the literature that reproduces the self-force result in the appropriate limit. We have made use of this new prescription to estimate the value of the angular momentum at the ISCO using our IMRI prescription and available self-force data. We have found a clear discrepancy for systems with mass-ratios $\eta \sim 0.1$, but have confirmed that for systems with small mass ratios $\eta \sim 0.01$ both predictions are fairly consistent.

This study has also shed light on a pressing issue that needs to be addressed before aLIGO begins observations, namely, at present we use templates in searches for GW sources whose actual dynamics are currently unknown. For instance, searches for BH binaries with total mass $25M_{\odot} - 200M_{\odot}$, and individual masses from $3M_{\odot}$ to $99M_{\odot}$ have been carried out, but not a single model has been calibrated using high-resolution NR simulations for mass-ratios smaller than 1:6. Hence, it is important that numerical relativists and search template developers interact more closely so as to run NR simulations which cover regions in parameter space where future GW detectors may detect GW sources. This meaningful interaction will be important from a theoretical perspective, as we will be able to further our knowledge of the self-force, and from a data analysis perspective, as we will be in a stronger position to develop accurate search templates for use in data analysis.

The approach we have outlined in this paper is the initial step to construct inspiral-merger-ringdown IMRI waveforms. Having derived a consistent prescription for the inspiral evolution, we can use a similar approach to that described in [298, ?] to include merger and ring-down in a physically consistent way. Developing complete IMRI waveforms may be useful for aLIGO data analysis, as they will provide the accuracy of more complex waveform models at an inexpensive computational cost.

6.6 Appendix

In this Section we show that the approach outlined in the main body of the paper can still be used to model equal-mass (EM) binaries with great accuracy at an inexpensive computational cost. To do so we consider a system with total mass $20M_{\odot}$. Following the method described in Section 6.3, we start by deriving accurate prescriptions for the orbital frequency Ω_{ansatz} and angular momentum flux \dot{L}_z . Thereafter we derive the corrections that should be implemented in the gauge-invariant expression for the angular momentum (see Eq. (6.28)) to accurately reproduce the inspiral evolution predicted by the EOBNRv2 model.

The prescriptions for the orbital frequency, angular momentum flux and angular momentum which reproduce the inspiral evolution for the $10M_{\odot}+10M_{\odot}$ binary system shown in Figure 39 require the following set of coefficients (compare Tables 10, 11, 12)

$$\begin{aligned}\Omega(u)_{\text{ansatz}} &: a_1 = -5.644, \quad a_2 = 11.003, \quad a_3 = -5.390, \quad a_4 = 8.794, \quad a_5 = 0.45839 \\ \dot{L}_z &: c_{2.5}^1 = -90.566, \quad c_3^2 = 304.941, \quad c_{3.5}^3 = -339.500 \\ z_{\text{SF}} &: b_1 = -5.803, \quad b_2 = 14.171, \quad b_3 = -7.074, \quad b_4 = 22.810, \quad b_5 = -25.580\end{aligned}$$

Having shown that the approach outlined in the paper is still effective to model EM binaries, we consider worthwhile developing a model that encapsulates the physics of binaries of comparable and intermediate-mass-ratio. In order to extend our IMRI model into the comparable mass-ratio regime, we require to implement some improvements in the model. First and foremost, we require expressions for the orbital frequency evolution (see Eq. (6.24)) and angular momentum (see Eq. (6.28)) which incorporate conservative corrections at second-order in mass ratio η . We can draw this conclusion by comparing the bottom-right panel of Figure 38 with the bottom panel of Figure 39. The former plot shows that for IMRIs the prescription for the angular momentum deviates from the self-force prediction near the ISCO. However, the latter plot shows that in the EM case the prescription for the angular momentum deviates considerably from the self-force prediction even during the inspiral evolution, far away

from the ISCO. Because the prescription we have used for the angular momentum in both cases includes corrections at liner order in mass-ratio, this suggest that to derive a model that unifies both regimes we require an expression for the angular momentum which includes second-order conservative corrections. We can follow a similar strategy to encapsulate in a single prescription the radiative piece of the self-force for comparable and intermediate mass-ratio systems by deriving second-order radiative corrections in the prescription for the angular momentum flux.

Having derived expressions for Ω_{ansatz} , \dot{L}_z and $L_z(x)$ which include corrections at second-order in mass-ratio, we will be able to tune the coefficients of these objects and derive numerical fits for the coefficients in terms of the mass-ratio η of the system. This approach will provide a unified description for the inspiral evolution of IMRIs and comparable-mass systems. We will pursue these studies in the future.




Figure 39 : The panels show the accuracy with which our model can reproduce the dynamics of an equal mass (EM) binary system with masses ($10M_{\odot} + 10M_{\odot}$), as compared with the EOBNRv2 model. The top panels show that our model can reproduce the orbital angular evolution and the flux of angular momentum predicted by the EOBNRv2 model point to point with an accuracy better than one part in a thousand. The middle panels show that our model can reproduce with a similar accuracy the radial and azimuthal evolution of a $10M_{\odot} + 10M_{\odot}$ binary system. The bottom panel shows that the expression for the gauge-invariant angular momentum $L_z(x)$ for an EM mass deviates clearly from the EMRI prescription even before nearing the ISCO.

Chapter 7

Self-forced evolutions of an implicit rotating source: a natural framework to model comparable and intermediate mass-ratio systems from inspiral through ringdown

7.1 Introduction

The black hole (BH) mass function in the local Universe is a strongly bi-modal distribution that identifies two main families: stellar-mass BHs with typical masses $\sim 10M_{\odot}$ observed in Galactic X-ray binaries [299] and, more recently, in globular clusters [300], and supermassive BHs with masses $\gtrsim 10^5 M_{\odot}$ observed to be present in most galactic nuclei [301, 302]. However, a population of X-ray sources with luminosities in excess of $10^{39} \text{ erg s}^{-1}$ has recently been observed, and *Chandra* and *XMM-Newton* spectral observations of these ultra-luminous X-ray sources (ULXs) revealed cool disc signatures that were consistent with the presence of intermediate mass BHs (IMBHs) with masses $10^{2-4} M_{\odot}$ [303, 304, 305]. Subsequent observations have shown that these ULXs have spectral and temporal signatures that are not

consistent with the sub-Eddington accretion regime that is expected for IMBHs at typical ULX luminosities. Rather, these later studies suggest that many ULXs are powered by super-Eddington accretion onto $\lesssim 100M_{\odot}$ BH remnants. Nevertheless, recent work by Swartz et al. [306] has demonstrated that as well as the high mass X-ray binaries that characterise most ULXs, there is a subpopulation of ULXs that seem to be powered by a separate physical mechanism. These objects have typical luminosities $L \gtrsim 10^{41} \text{ erg s}^{-1}$, which cannot be explained by close to maximal radiation from super-Eddington accretion onto massive BHs formed in low metallicity regions [307, 308, 309]. Several hyper-luminous X-ray sources, including M82 X-1, ESO 243-49 HLX-1, Cartwheel N10 and CXO J122518.6+144545, present the best indirect evidence for the existence of IMBHs [310, 311, 312, 313]. In particular, the colocation of M82 X-1 with a massive, young stellar cluster, the features of its power spectrum, and some reported transitions between a hard state and a thermal dominant state, make this object a strong IMBH candidate [314, 315, 316, 317]. Recent searches of archival *Chandra* and *XMM-Newton* data sets have also uncovered two new hyper-luminous X-ray sources with luminosities in excess of $10^{39} \text{ erg s}^{-1}$. These sources are the most promising IMBH candidates currently known, although the highest possible super-Eddington accretion rate onto the largest permitted BH remnant cannot yet be ruled out [318]. This increasing body of observational evidence [319, 320], and the fact that the existence of IMBHs provides a compelling explanation for the initial seeding of supermassive BHs present in most galactic nuclei [321, 322, 323] has revived the quest for these elusive objects.

Since hyper-luminous X-ray sources are rare, and our knowledge about their astrophysical properties is still limited, we may have to use a different means to search for IMBHs in order to improve our knowledge about the channels that lead to the formation of these objects, and to shed light on their astrophysical properties, such as mass and spin distributions [324]. In this chapter we explore the use of observations in the gravitational wave (GW) spectrum to gain insight into the properties of IMBHs.

The current upgrade of the Laser Interferometer Gravitational Wave Observatory (LIGO) and its international partners Virgo and Kagra [269, 325, 326], will enable the detection of GWs from coalescences involving IMBHs with masses $50M_{\odot} \lesssim M \lesssim 500M_{\odot}$, if these instruments achieve their target sensitivity down to the low-frequency

cutoff at 10Hz (See Figure 40) [327]. Advanced LIGO (aLIGO) and Advanced Virgo are expected to have greatest sensitivity in the 60Hz - 500Hz range, with a peak at ~ 60 Hz (see Figure 40). Proposed third generation detectors, such as the Einstein Telescope [328], which aim to extend the frequency range of ground-based detectors down to 1Hz, while also maintaining high frequency sensitivity up to 10kHz, will enhance our ability to search for GWs emitted by sources that involve BHs with masses between $10^{2-4} M_{\odot}$ [329, 330, 331].

A promising channel for detection of IMBHs is through the emission of gravitational radiation during the coalescence of stellar-mass compact remnants — neutron stars (NSs) or BHs — with IMBHs in core-collapsed globular clusters. This expectation is backed up by numerical simulations of globular clusters [332, 333, 334, 335, 336, 337, 338, 339] which suggest that IMBHs could undergo several collisions with stellar-mass compact remnants during the lifetime of the cluster through a variety of mechanisms, including gravitational radiation, Kozai resonances and binary exchange processes. As discussed in [340], the most likely mechanism for the formation of binaries involving a stellar-mass compact remnant and an IMBH is hardening via three body interactions, with an expected detection rate of $\sim 1 - 10 \text{ yr}^{-1}$ with ground-based observatories [340, 341].




Figure 40 : The panel shows the expected sensitivity for two configurations of the Einstein Telescope (ET), namely, ETD (black), ETB (blue) and LIGO’s Zero Detuned High Power (ZDHP) configuration (red). The vertical axis measures $F_{\text{normalized}} = (f/f_{\text{max}})^{-7/6} \sqrt{S_n(f_{\text{max}})/S_n(f)}$, where f_{max} is the maximum of the corresponding power spectral density, $S_n(f)$.

The frequency of the dominant quadrupolar harmonic in the GWs emitted at the

innermost stable circular orbit (ISCO) for a binary of non-spinning objects is

$$f_{\text{ISCO}} = 4.4 \text{kHz} \left(\frac{M_\odot}{M} \right), \quad (7.1)$$

so for a typical intermediate mass–ratio coalescence (IMRC) with total mass $M \gtrsim 100M_\odot$, advanced detectors will observe the late inspiral, merger and ringdown. However, the heaviest IMRCs, with masses $\gtrsim 500M_\odot$, will coalesce out of band or at the low frequency limit of the bandwidth. Hence, merger and ringdown — which intrinsically generate a signal-to-noise ratio (SNR) suppressed by a factor of symmetric mass-ratio — see Table 14 below — relative to the SNR generated during the inspiral phase— will significantly contribute to the SNR of IMRCs over a considerable portion of the detectable mass-range [342]. In order to get the most information from GW observations of IMRCs, it is therefore necessary to develop waveform models that incorporate the inspiral, merger and ringdown in a physically consistent way. In order to make progress in this direction, we previously developed a waveform model that combined results from Black Hole Perturbation Theory (BHPT) and post-Newtonian (PN) theory to explore the information that could be obtained from observations of IMRCs with the EinsteinTelescope [330, 331]. Although this model provided an important step in exploring the science that could be done with IMRC observations, the model was limited.

In [343, 344, 345, 346] we explored using the self-force formalism [287, 347] to develop a waveform model with a robust description of the dynamical evolution of IMRCs during the inspiral phase. These were found to be effective when used to carry out matched-filter based searches for inspiral-only IMRCs [342], but searches for IMRCs in the advanced detector era will require waveform models that include not only the inspiral but also the merger and ringdown [342]. The model described in [330] included merger and ring down but without the self-force driven inspiral. It would be ideal to develop a consistent inspiral, merger, ringdown waveform model by comparison to numerical relativity simulations, as described in [?, 348]. However, numerical relativity simulations for systems with mass-ratios $q = m_1/m_2 \lesssim 1/10$ are very computationally expensive at present [?]. We can circumvent this problem by making use of recent breakthroughs in the self-force program that have shown that the conservative part of the self-force can reproduce, with good accuracy, results

from numerical relativity simulations of comparable-mass binary systems [349]. Furthermore, the recent computation of the self-force inside the ISCO equips us to now develop models that better reproduce the true dynamics of black hole binaries in the strong field regime [350].

In addition to IMRCs, which may be detected by second generation detectors, and with more likelihood by third generation detectors, we have explored the suitability of using self-forced evolutions to model the mergers of systems involving NSs and stellar mass BHs [351]. Since NSBH mergers are promising GW sources for second generation detectors, we need to develop accurate and computationally inexpensive templates appropriate for these events. Current efforts to model NSBH systems have generally used PN approximants, evaluated to the highest post-Newtonian order available, but these approximates are not sufficiently reliable to model these events [352, 353]. In Figure 41, we show the phase difference between the PN approximant TaylorT4 [354] and the EOB model introduced in [?, 348]. This exhibits a substantial discrepancy near the ISCO. These considerations have impelled us to develop a different approach to model events with the typical mass- ratios expected for NSBH binaries.

In this chapter we combine recent developments in the self-force program, in PN theory and in numerical relativity to develop a model that describes the inspiral, merger and ringdown of IMRCs and comparable mass-ratio systems that could be detected by second and third generation ground-based GW detectors. In Section 7.2 we discuss the modelling of the conservative part of the self-force. We show that using the linear in mass-ratio self force results for binaries with mass-ratios $q \gtrsim 1/6$ gives a system without an ISCO. We then discuss the implications of this result for the modelling of comparable and intermediate mass-ratio binaries. Thereafter, we describe the approach we use to model the radiative part of the self-force for the inspiral evolution. Having constructed the inspiral part of the self-forced waveform model, we extend the transition scheme of Ori and Thorne [298] by including finite mass-ratio corrections, and modelling the orbital phase evolution using the implicit rotating source (IRS) model. We adopt this description for the late-time radiation in order to provide a smooth progression from late inspiral to ringdown. We show that this approach provides the correct orbital frequency evolution in the vicinity of the light-ring. Finally, we construct the ringdown waveform using both a sum of quasinormal modes and the late-time radiation waveform evolution predicted by




Figure 41 : The phase discrepancy in radians between the PN approximant TaylorT4, and the Effective One Body model, shown as a function of time from $r = 30M$ to the point when the TaylorT4 model reaches the ISCO. The systems have mass-ratio, q , total mass, M_{tot} , and final phase discrepancy, $\Delta\Phi$: $(q, M_{\text{tot}}, \Delta\Phi) = (1/6, 7M_{\odot}, 21.5 \text{ rads})$ (top-left), $(1/8, 9M_{\odot}, 30.2 \text{ rads})$ (top-right), $(1/10, 11M_{\odot}, 70.1 \text{ rads})$ (bottom-left) and $(1/15, 16M_{\odot}, 83.2 \text{ rads})$ (bottom-right) respectively.

the IRS model, and show their equivalence. Section 7.3 presents a summary of our findings and future directions of work.

7.2 Modeling

7.2.1 Nomenclature

Throughout this chapter we will use units with $G = c = 1$, unless otherwise stated. We will consider BH binaries on circular orbits with component masses m_1, m_2 , such that $m_1 < m_2$. We assume that the binary components are non spinning. We will use several combinations of the masses $m_{1,2}$ in the following Sections, which are summarized in Table 14.

Binary masses	
m_1	mass of inspiralling compact object
m_2	mass of central compact object
$M = m_1 + m_2$	total mass of binary system
$q = \frac{m_1}{m_2}$	mass–ratio
$\mu = \frac{m_1 m_2}{m_1 + m_2}$	reduced mass
$\eta = \frac{\mu}{M}$	symmetric mass–ratio

Table 14 : The table summarizes the nomenclature we will use throughout our analysis.

Having defined the variables to be used in the subsequent sections, we shall now describe the construction of the self-forced waveform model. The model consists of four building blocks — the inspiral, the transition, the plunge and the ringdown phases. The next section describes the inspiral evolution.

7.2.2 Inspiral evolution

We model the inspiral phase evolution in the context of the Effective One Body (EOB) formalism [355], i.e., we consider the scenario in which the dynamics of a binary system is mapped onto the motion of a test particle in a time-independent and spherically symmetric Schwarzschild space-time with total mass M :

$$ds_{\text{EOB}}^2 = -A(r)dt^2 + B(r)dr^2 + r^2d\Omega^2, \quad (7.2)$$

where the potentials A , B are known to 3PN order [356, 357]. In the test-mass particle limit $\eta \rightarrow 0$, these potentials recover the Schwarzschild results, namely:

$$A(u, \eta \rightarrow 0) = B^{-1}(u, \eta \rightarrow 0) = 1 - 2u, \quad \text{with} \quad u = \frac{M}{r}. \quad (7.3)$$

In the EOB formalism, the orbital frequency evolution can be derived from a Hamiltonian, H_{EOB} [355], given by:

$$H_{\text{EOB}} = M\sqrt{1 + 2\eta(H_{\text{eff}} - 1)}, \quad (7.4)$$

using the general Hamiltonian equation:

$$\frac{d\phi}{dt} = M\Omega = \frac{\partial H_{\text{EOB}}}{\partial L} = \frac{u^2 L(x) A(u)}{H(u) H_{\text{eff}}(u)}, \quad (7.5)$$

where

$$\begin{aligned} H_{\text{eff}}(u) &= \frac{A(u)}{\sqrt{\tilde{A}(u)}}, \quad \tilde{A}(u) = A(u) + \frac{1}{2} u A'(u), \\ \text{and} \quad H(u) &= \sqrt{1 + 2\eta(H_{\text{eff}} - 1)}. \end{aligned} \quad (7.6)$$

Recent work in the self-force formalism has enabled the derivation of gravitational self-force corrections to the EOB potential $A(u) \rightarrow 1 - 2u + \eta a(u) + \mathcal{O}(\eta^2)$ [280]. Deriving this gravitational self-force contribution, $a(u)$, is equivalent to including all PN corrections to the EOB potential $A(u)$ at linear order in η . We shall now briefly describe the construction of the gravitational self-force contribution $a(u)$, emphasizing the fact that this contribution encodes information about the strong-field regime of the gravitational field.

As shown by Detweiler and Whiting [358], the gravitational self-force corrected worldline can be interpreted as a geodesic in a smooth perturbed spacetime with metric

$$g_{\alpha\beta} = g_{\alpha\beta}^0(m_2) + h_{\alpha\beta}^R, \quad (7.7)$$

where the regularized R field is a smooth perturbation associated with m_1 . Detweiler proposed a gauge invariant relation to handle the conservative effect of the gravitational self-force in circular motion [359, 360]:

$$z_1(\Omega) = \sqrt{1 - 3x} \left(1 - \frac{1}{2} h_{uu}^{R,F} + q \frac{x}{1 - 3x} \right), \quad (7.8)$$

where x is the gauge-invariant dimensionless frequency parameter given by $x = (M\Omega)^{2/3}$, $h_{uu}^{R,G}$ is a double contraction of the regularised metric perturbation with the four-velocity, u^μ , $h_{uu}^{R,G} = h_{\mu\nu}^{R,G} u^\mu u^\nu$, the label G indicates the gauge used to evaluate the metric perturbation and the label F indicates that this expression is valid within the class of asymptotically flat gauges. In [350], $z_1(\Omega)$ was calculated in Lorenz

gauge and the following gauge transformation can be used to link the asymptotically flat $h_{uu}^{R,F}$ metric perturbation to its Lorenz-gauge counterpart $h_{uu}^{R,L}$:

$$h_{uu}^{R,F} = h_{uu}^{R,L} + 2q \frac{x(1-2x)}{(1-3x)^{3/2}}. \quad (7.9)$$

Hence, inserting Eq. (7.9) into Eq. (7.8) leads to

$$z_1(\Omega) = \sqrt{1-3x} \left(1 - \frac{1}{2} h_{uu}^{R,L} - 2q \frac{x(1-2x)}{(1-3x)^{3/2}} + q \frac{x}{1-3x} \right). \quad (7.10)$$

The numerical data obtained in [350] for $h_{uu}^{R,L}$ from $x > 1/5$ was new and the numerical accuracy for $x < 1/5$ was also much improved compared to previous results [359, 277]. The EoB potential $a(x)$ can be constructed from $h_{uu}^{R,L}$ via

$$a(x) = -\frac{1}{2} (1-3x) \tilde{h}_{uu}^{R,L} - 2x\sqrt{1-3x}, \quad (7.11)$$

with $\tilde{h}_{uu}^{R,L} = q^{-1} h_{uu}^{R,L}$. In [350], a global fit formula for $a(x)$ was given a dense sample of numerical values over the entire range $0 < x < \frac{1}{3}$. The numerical fit for $a(x)$ that we use in this study is taken from Eq.(54) of [350]. This global analytic fit for $a(x)$ can be recast using the relation

$$a(x) = 2x^3 \frac{(1-2x)}{\sqrt{1-3x}} a_E(x). \quad (7.12)$$

Using the above dictionary, the model for $a(x)$ in this chapter reproduces the numerical data points for the function $a_E(x)$ to within a maximal absolute difference of 1.2×10^{-5} over the domain $0 < x < \frac{1}{3}$. The corresponding self-force corrected energy and angular momentum are given by [350, 280]

$$\begin{aligned} E(u(x)) &= E_0(x) + \eta \left(-\frac{1}{3} \frac{x}{\sqrt{1-3x}} a'(x) + \frac{1}{2} \frac{1-4x}{(1-3x)^{3/2}} a(x) - E_0(x) \left(\frac{1}{2} E_0(x) + \frac{x}{3} \frac{1-6x}{(1-3x)^{3/2}} \right) \right), \\ L_z(u(x)) &= L_0(x) + \eta \left(-\frac{1}{3} \frac{x}{\sqrt{x(1-3x)}} a'(x) - \frac{1}{2} \frac{1}{\sqrt{x}(1-3x)^{3/2}} a(x) - \frac{1}{3} \frac{1-6x}{\sqrt{x}(1-3x)^{3/2}} (E_0(x) - E(u(x))) \right), \end{aligned}$$

with $u(x) = x \left(1 + \eta \left[\frac{1}{6} a'(x) + \frac{2}{3} \left(\frac{1-2x}{\sqrt{1-3x}} - 1 \right) \right] \right),$

where $E_0(x)$ and $L_0(x)$ are given by

$$E_0(x) = \frac{1 - 2x}{\sqrt{1 - 3x}} - 1, \quad (7.16)$$

$$L_0(x) = \frac{1}{\sqrt{x(1 - 3x)}}. \quad (7.17)$$




Figure 42 : The panels show the energy and angular momentum given by Eqs. (7.13)-(7.14), respectively. We show the functional form of these parameters for binary systems with mass-ratio values, from top to bottom, $q \in [0, 1/100, 1/20, 1/10, 1/6, 1/5, 1]$.

In Figure 42, we show the effect of these conservative corrections on the orbital parameters.

As discussed in [280], if one makes use of the self-force expression for the energy given by Eq. (7.13), and minimize it with respect to the orbital frequency, then one finds that binary systems with mass-ratios $q \in [1, 1/2, 1/3]$ do not have an ISCO in this model. It was argued in [280] that deriving self-force results in the strong field regime may alleviate this problem. We have explored this issue, and have found that using linear self-force corrections that are valid all the way to the light ring (last unstable circular orbit for massless particles) does not fix this problem for comparable mass-ratio systems. In Figure 43, we show that the existence of an ISCO is guaranteed for BH binaries with symmetric mass-ratio $\eta \lesssim 6/49$ (or $q \lesssim 1/6$), and its location may be approximated by

$$x_{\text{ISCO}} = \frac{1}{6} (1 + 0.83401\eta + 4.59483\eta^2). \quad (7.18)$$

It remains to be seen whether the inclusion of second-order conservative corrections

gives an ISCO for binaries with mass-ratios $q \gtrsim 1/6$.




Figure 43 : The location of the innermost stable circular orbit is determined by the condition $dE/dx = 0$. The panel shows dE/dx as a function of the gauge invariant quantity $x = (M\Omega)^{2/3}$. The various curves represent binary systems with mass-ratios, from top to bottom, $q \in [0, 1/100, 1/20, 1/10, 1/6, 1/5, 1]$. Note that binaries with mass-ratios $q \gtrsim 1/6$ do not have an ISCO in this model.

In summary, the building blocks we use to construct the conservative dynamics are

- The orbital frequency evolution is computed using Eq. (7.5) with the gravitational self-force contribution included in the potential $A(u) = 1 - 2u + \eta a(u)$.
- Eq. (7.5) is evaluated using the self-force-corrected expression for the angular momentum, $L(x)$, given by Eq. (7.14). The self-force-corrected expression for the energy, given in Eq. (7.13) is only used to determine the point at which the inspiral ends and the transition region begins.

Eq. (7.5) provides a very accurate modeling of the orbital frequency from early inspiral through the ISCO. However, the post-ISCO time evolution of this prescription does not render an accurate representation of the orbital frequency as compared to numerical relativity simulations. This is a problem that has been addressed in the EOB formalism by introducing a phenomenological approach —the so-called non-quasi-circular coefficients— that enabled them to reproduce the orbital evolution extracted from numerical simulations [?]. The approach we will follow to circumvent this problem is described in detail in Section 7.2.4, and consists of embedding the self-force formalism in the context of the implicit rotating source model [361] after the ISCO.

This completes the description of the conservative part of the self-force. We now describe how to couple this with the radiative part of the self-force to model the inspiral evolution.

7.2.3 Dissipative dynamics

A consistent self-force evolution model that incorporates first-order in mass-ratio conservative corrections should also include second-order radiative corrections. The model we described in the previous section was constructed including first order conservative corrections. However, second-order self-force radiative corrections are not known at present. Several studies have demonstrated the importance of including the missing second order corrections to the radiative part of the self-force, both for source detection and for parameter estimation [362, 363, 346, 344, 343].

We use a new prescription for the energy flux that uses the first-order in mass ratio terms derived in [364], including PN corrections up to 22nd PN order

$$\begin{aligned} (\dot{E})_{\text{PN}} = & -\frac{32}{5} \frac{\mu^2}{M} x^{7/2} \left[1 - \frac{1247}{336} x + 4\pi x^{3/2} - \frac{44711}{9072} x^2 - \frac{8191}{672} \pi x^{5/2} \right. \\ & + x^3 \left\{ \frac{6643739519}{69854400} + \frac{16}{3} \pi^2 - \frac{1712}{105} \gamma_E - \frac{856}{105} \ln(16x) \right\} - \frac{16285}{504} \pi x^{7/2} \\ & + x^4 \left\{ -\frac{323105549467}{3178375200} + \frac{232597}{4410} \gamma_E - \frac{1369}{126} \pi + \frac{39931}{294} \ln(2) - \frac{47385}{1568} \ln(3) + \frac{232597}{4410} \ln(x) \right\} \\ & + x^{9/2} \left\{ \frac{265978667519}{745113600} \pi - \frac{6848}{105} \gamma_E \pi - \frac{13696}{105} \pi \ln(2) - \frac{6848}{105} \pi \ln(x) \right\} \\ & \left. + \text{higher order corrections up to 22PN order} \right]. \end{aligned} \quad (7.19)$$

We include higher-order in mass-ratio terms using the exponential resummation approach described in [362]. In this approach, the energy flux is

$$\left(\frac{dE}{dt} \right)_{\text{hybrid}} = \mathcal{L}_0 \exp(\mathcal{L}_\eta), \quad (7.20)$$

where \mathcal{L}_0 denotes the leading-order in mass-ratio PN energy flux given in Eq. (7.19), and \mathcal{L}_η incorporates mass-ratio corrections to the highest PN order available [365, 366, 362], and additional corrections characterised by a set of unknown coefficients,

b_i

$$\begin{aligned}\mathcal{L}_\eta = & \left[x \left[-\frac{35}{12} \eta + b_1 \eta^2 \right] + 4\pi x^{3/2} \left[b_2 \eta + b_3 \eta^2 \right] + x^2 \left[\frac{9271}{504} \eta + \frac{65}{18} \eta^2 \right] + \pi x^{5/2} \left[-\frac{583}{24} \eta + b_4 \eta^2 \right] \right. \\ & \left. + x^3 \left[\eta \left(-\frac{134543}{7776} + \frac{41}{48} \pi^2 \right) - \frac{94403}{3024} \eta^2 - \frac{775}{324} \eta^3 \right] + \pi x^{7/2} \left[\frac{214745}{1728} \eta + \frac{193385}{3024} \eta^2 \right] \right].\end{aligned}$$

The coefficients b_i were taken to be constant in [362], but we found that a better match to the EOB phase evolution could be obtained by allowing an additional dependence on mass-ratio in these terms (see Eqs. (7.23)-(7.25) below). We constrain the b_i coefficients by ensuring that the phase evolution of this model reproduces the phase evolution predicted by the EOB model introduced in [?, 348], which was calibrated to the phase evolution of compact binaries observed in numerical relativity simulations. To do so, we implemented the EOB model [?] and performed a Monte Carlo simulation to optimize the values of the b_i coefficients (see Figure 44). The optimization was done in two stages. We started by considering the three coefficients b_1 , b_2 and b_4 , sampling a wide range of parameter space, namely $b_i \in [-200, 200]$. We constrained the duration of the waveform from early inspiral to the light-ring to be similar to its EOB counterpart. Waveforms that differed from their EOB counterparts by more than 10^{-4} seconds were discarded. Once the region under consideration had been sparsely sampled, we focused on regions of parameter space where the orbital phase evolution was closest to the EOB evolution, and finely sampled these to obtain the optimal values for the coefficients. We found that this approach enabled us to reproduce the EOB phase evolution with a phase discrepancy of the order ~ 1 rad. After constraining b_1 , b_2 and b_4 , we explored whether including additional corrections could further improve the phase evolution, by adding η corrections beyond 3PN order. Such corrections were found to have a negligible impact on the actual phase evolution. This is not difficult to understand, since such corrections are of order $(\mathcal{O}(\eta^4), \mathcal{O}(\eta^3))$, at (3PN, 3.5PN) respectively. We found a similar behavior when we added leading order mass-ratios corrections beyond 4PN order. Thus, we took a different approach: having derived the optimal value for b_1 , b_2 and b_4 , we took these results as initial seeds for an additional MC simulation in which b_3 was also included in Eq. (7.21), and repeated the optimization procedure. The results of these simulations are shown

in Figure 44.

We carried out several different Monte Carlo runs to find the ‘optimal’ optimization interval, meaning the range of radial separations over which we tried to best match the phase evolution relative to the EOB model. We found that starting the optimization at $r = 30M$ gave results that performed moderately well at early inspiral, but that underperformed at late inspiral, leading to phase discrepancies of order ~ 3 rads. Starting the optimization at $r = 20M$ instead decreased the phase discrepancy with respect to the former case by a factor of 10 during early inspiral, and enabled us to reproduce the phase evolution in the EOB model for all the mass-ratios considered to within the accuracy of the numerical waveforms used to calibrate the EOB model in [?, 348]. Implementing these numerically optimized higher-order η corrections in Eq. (7.21) leads to:

$$\begin{aligned} \mathcal{L}_\eta &= \left[x \left[-\frac{35}{12}\eta + B_1 \right] + 4\pi x^{3/2} B_2 + x^2 \left[\frac{9271}{504}\eta + \frac{65}{18}\eta^2 \right] + \pi x^{5/2} \left[-\frac{583}{24}\eta + B_3 \right] \right. \\ &\quad \left. + x^3 \left[\eta \left(-\frac{134543}{7776} + \frac{41}{48}\pi^2 \right) - \frac{94403}{3024}\eta^2 - \frac{775}{324}\eta^3 \right] + \pi x^{7/2} \left[\frac{214745}{1728}\eta + \frac{193385}{3024}\eta^2 \right] \right]. \end{aligned} \quad (7.22)$$

where:

$$B_1 = \frac{1583.650 - 11760.507\eta}{1 + 142.389\eta - 981.723\eta^2} \eta^2, \quad (7.23)$$

$$B_2 = \frac{-12.081 + 35.482\eta}{1 - 4.678\eta + 13.280\eta^2} \eta + \frac{19.045 - 240.031\eta}{1 - 18.461\eta + 74.142\eta^2} \eta^2, \quad (7.24)$$

$$B_3 = \frac{51.814 - 980.100\eta}{1 - 13.912\eta + 88.797\eta^2} \eta^2. \quad (7.25)$$

This improved prescription for the energy flux, which incorporates second-order mass-ratio corrections to the PN expansion up to 3.5PN order, is sufficient to generate a model whose phase evolution reproduces with excellent accuracy the phase evolution predicted by EOB throughout inspiral and merger (see Figure 45).

Given the energy flux defined by Eqs (7.19)–(7.21), we generate the inspiral trajectory using the simple prescription

$$\frac{dx}{dt} = \frac{dE}{dt} \frac{dx}{dE}, \quad (7.26)$$

where we have used the mass-ratio corrected energy —Eq. (7.13)— to compute dE/dx . Figure 45 shows that for binaries with mass-ratio $q = 1/6$, the phase discrepancy between our self-force model and EOB is $\lesssim 0.5$ rads at the light-ring, which is within the numerical accuracy of the simulations used to calibrate EOB. It has been shown recently that EOB remains accurate for mass-ratios up to $q = 1/8$ [367]. In that regime the phase discrepancy between this model and EOB at the light-ring is less than 1 rad, as shown in Figure 45. For binaries with $q = 1/10$, the phase discrepancy at the light-ring is $\lesssim 1.2$ rads, which is still within the numerical accuracy of available simulations [296, 297].

It must be emphasized that even if we only use the inspiral evolution to model binaries with mass-ratios that typically describe NSBH binaries, our self-force evolution model performs better than TaylorT4, since we can reduce the phase discrepancy between TaylorT4 and EOB at the last stable circular orbit by a factor of (~ 40 , ~ 70) for binaries with $q = (1/6, 1/10)$ and total mass $M_{\text{tot}} \in (7M_{\odot}, 11M_{\odot})$ (see Figure 41).

Figure 45 conveys an important message — deriving the second order corrections to the radiative part of the self-force may well provide a robust framework to describe in a single model not only events that are naturally described by BHPT, such as the mergers of stellar mass compact objects with supermassive BHs in galactic nuclei [346, 344, 264, 289, 290, 287, 368], but also events that are more naturally described by PN or numerical methods, in particular the coalescences of compact object binaries with comparable or intermediate mass-ratios [346, 268, 330, 331, 267].

To finish this Section, we describe the approach followed to construct the gravitational waveform from the inspiral trajectory. At leading post-Newtonian order, a general inspiral waveform can be written as

$$h(t) = -(h_+ - i h_{\times}) = \sum_{\ell=2}^{\infty} \sum_{m=-\ell}^{\ell} h^{\ell m} {}_{-2}Y_{\ell m}(\iota, \Phi). \quad (7.27)$$

If only the leading-order modes $(\ell, m) = (2, \pm 2)$ are included, the inspiral waveform




Figure 44 : The (top, bottom) panels show the results of the optimization runs that were used to constrain the values of the b_i coefficients given in Eq. (7.21). The panels show the results for binaries of mass–ratio $q \in [1/6, 1/8]$, and total mass $M_{\text{tot}} = [7M_\odot, 9M_\odot]$. The ‘optimal’ value for the coefficients has been chosen by ensuring that the flux prescription minimizes the phase discrepancy between the EOB model and our self-force model. The color bar shows the phase difference squared between both models, which is integrated from $r = 20M$ all the way to the light-ring.

components are given by

$$h_+(t) = \frac{4\mu r^2 \dot{\phi}^2}{D} \left(\frac{1 + \cos^2 \iota}{2} \right) \cos [2(\phi(t) + \Phi)], \quad (7.28)$$

$$h_\times(t) = \frac{4\mu r^2 \dot{\phi}^2}{D} \cos \iota \sin [2(\phi(t) + \Phi)], \quad (7.29)$$

where D is the distance to the source. Since the orbital evolution will deviate from a circular trajectory during late inspiral ($\dot{r} \neq 0$), we must consider more general orbits in which both \dot{r} and $\dot{r}\dot{\phi}$ are non-negligible. For such orbits, the Newtonian GW




Figure 45 : The panels show the orbital phase evolution of a self-force model making use of optimized PN energy flux given by Eq. (7.20) and the phase evolution as predicted by the EOB model. The [top/bottom] panels exhibit this evolution for a compact binary with mass-ratio $q = [(1/6, 1/8), (1/10, 1/15)]$, and total mass $M_{\text{tot}} = [(7M_{\odot}, 9M_{\odot}), (11M_{\odot}, 16M_{\odot})]$, respectively.

polarizations are given by [369]:

$$\begin{aligned} h_+(t) &= \frac{2\mu}{D} \left\{ (1 + \cos^2 \iota) \left[\cos [2(\phi(t) + \Phi)] \left(-\dot{r}^2 + r^2 \dot{\phi}^2 + \frac{1}{r} \right) \right. \right. \\ &\quad \left. \left. + 2r \dot{r} \dot{\phi} \sin [2(\phi(t) + \Phi)] \right] + \left(-\dot{r}^2 - r^2 \dot{\phi}^2 + \frac{1}{r} \right) \sin^2 \iota \right\}, \end{aligned} \quad (7.30)$$

$$\begin{aligned} h_\times(t) &= \frac{4\mu}{D} \cos \iota \left\{ \sin [2(\phi(t) + \Phi)] \left(-\dot{r}^2 + r^2 \dot{\phi}^2 + \frac{1}{r} \right) \right. \\ &\quad \left. - 2r \dot{r} \dot{\phi} \cos [2(\phi(t) + \Phi)] \right\}, \end{aligned} \quad (7.31)$$

where \dot{r} can be computed using

$$\frac{dr}{dt} = -\frac{1}{u^2} \frac{du}{dx} \frac{dx}{dt}.$$

Having described the construction of the inspiral evolution, we shall now describe the approach followed to smoothly connect the late inspiral evolution onto the plunge phase. The adiabatic prescription given by Eq. (7.26) breaks down when $dE/dx \rightarrow 0$. Hence, we need a scheme that enables us to match the late inspiral phase onto the plunge phase. We will do this by modifying the “transition” phase developed by Ori and Thorne [288] by including finite mass–ratio corrections.

7.2.4 Transition and plunge phases

In this Section we describe an extension of the transition phase model introduced by Ori and Thorne [288]. The basic idea behind this approach can be understood by studying the motion of an inspiralling object in terms of the effective potential, $V(r, L)$, which takes the following simple form for a Schwarzschild BH [370]:

$$V(r, L) = \left(1 - \frac{2}{r}\right) \left(1 + \frac{L^2}{r^2}\right). \quad (7.32)$$

Throughout the inspiral, the body moves along a nearly circular orbit, and hence the ratio of the energy flux to the angular momentum flux is given by:

$$\frac{dE}{d\tau} = \Omega \frac{dL}{d\tau}. \quad (7.33)$$

Hence, near the ISCO, the energy and angular momentum of the body satisfy the following relations:

$$E \rightarrow E_{\text{ISCO}} + \Omega_{\text{ISCO}} \xi, \quad (7.34)$$

$$L \rightarrow L_{\text{ISCO}} + \xi. \quad (7.35)$$

Re-writing the effective potential, Eq. (7.32), in terms of $\xi = L - L_{\text{ISCO}}$, one notices that during early inspiral, $\xi \gg 0$, the motion of the object is adiabatic, and the object sits at the minimum of the potential—as shown in the left panel of Figure 46. However, as the object nears the ISCO, the minimum of the potential moves inward

due to radiation reaction. At some point, the body's inertia prevents the body from staying at the minimum of the potential, and adiabatic inspiral breaks down [288] — illustrated in the right-hand panel of Figure 46.




Figure 46 : Left panel: The object sits at the minimum of the effective potential, Eq. (7.32), which corresponds to the case $\xi = L - L_{\text{ISCO}} \gg 0$. Right panel. Blue (top) curve: radial geodesic motion, which corresponds to $\xi = L - L_{\text{ISCO}} \gg 0$; Red (middle) curve: the object nears the ISCO and the orbit shrinks due to radiation reaction. Note that the minimum of the potential has moved inwards ($\xi = 0.35$). Yellow (bottom) curve: body's inertia prevents it from staying at the minimum of the potential, and adiabatic inspiral breaks down ($\xi = 0$). At this point the transition regime takes over the late inspiral evolution [288]. Note: this plot is based on Figure 1 of [288].

The equation that governs the radial motion during the transition regime is found by linearising the equation

$$\left(\frac{dr}{d\tau} \right)^2 = E(r)^2 - V(r, L), \quad (7.36)$$

using Eqs. (7.34), (7.35), and

$$\frac{d\xi}{d\tau} = \kappa \eta, \quad \text{with} \quad \kappa = \left[\frac{32}{5} \Omega^{7/3} \frac{\dot{\mathcal{E}}}{\sqrt{1-3u}} \right]_{\text{ISCO}}, \quad (7.37)$$

where $\dot{\mathcal{E}}$ is the general relativistic correction to the Newtonian, quadrupole-moment formula [298]. We now extend these Eqs. by including finite mass-ratio corrections. Eq. (7.36) can be replaced by

$$\frac{dx}{dt} = \frac{u^2(1-2u)}{E(x)} \left(\frac{du}{dx} \right)^{-1} \left[E(x)^2 - V(u(x), L(x)) \right]^{1/2}, \quad (7.38)$$

where we have used

$$\frac{d\tau}{dt} = \frac{1 - 2u(x)}{E(x)}, \quad (7.39)$$

and the expressions for the energy and angular momentum are given by Eqs. (7.13), (7.14). In order to linearize Eq. (7.38) we replace $E(x)$ and $L(x)$ by Eqs (7.34) and (7.35) respectively.




Figure 47 : The left panel shows the effective potential for a Schwarzschild BH without including finite mass-ratio corrections. Note that the minimum of the potential takes place at the ISCO, which can be determined using Eq. (7.18). The right panel exhibits the influence of finite mass-ratio corrections on the effective potential used to modify Ori and Thorne transition regime [288]. The curves represent binaries, from top to bottom, with mass-ratios $q \in [0, 1/100, 1/20, 1/10, 1/6, 1/5, 1]$.

As discussed in [288], since these equations use the η -corrected values for $E(x_{\text{ISCO}})$, $L(x_{\text{ISCO}})$ and Ω_{ISCO} , then they remain valid even for finite mass-ratio η [288]. In Figure 47 we show the effect that these finite mass-ratio η corrections have on the effective potential $V(x, L(x))$. We determine the point at which the transition regime starts by carrying out a stability analysis near the ISCO using dE/dx . As shown in Figure 43, the ISCO is determined by the relation $dE/dx = 0$. We have found that the relation

$$\left(\frac{dE}{dx} \right) \Big|_{\text{transition}} = -0.054 + \frac{1.757 \times 10^{-4}}{\eta}, \quad (7.40)$$

provides a robust criterion to mark the start of the transition regime for binaries with mass-ratios $1/100 < q < 1/6$.

In [298], the authors only kept terms linear in ξ , but we have explored which higher order terms had a noticeable impact on the evolution by examining their impact on the length and phasing of the waveform. We found that terms $\propto \xi$ and $\propto (u - u_{\text{ISCO}})\xi$

were important, but corrections at order $\mathcal{O}(\xi^2)$ could be ignored even for comparable mass-ratio systems.

We model the evolution of the orbital frequency during the transition regime and thereafter in a different manner to that proposed by Ori and Thorne [288]. In order to ensure that the late-time evolution of the orbital frequency of our self-force model is as close as possible to the orbital evolution extracted from numerical relativity simulations, we incorporate the late-time frequency evolution that was derived by Baker et al [361] in their implicit rotating source (IRS) model, namely:

$$\frac{d\phi}{dt} = \Omega_i + (\Omega_f - \Omega_i) \left(\frac{1 + \tanh(\ln \sqrt{\varkappa} + (t - t_0)/b)}{2} \right)^{\varkappa}, \quad (7.41)$$

where Ω_i is the value of the orbital frequency when the transition regime begins, and Ω_f is the value of the frequency at the light ring, which corresponds to $\omega_{\ell mn}/m$, where $\omega_{\ell mn}$ is the fundamental quasi-normal ringing frequency ($n = 0$) for the fundamental mode $(\ell, m) = (2, 2)$ of the post-merger black hole (see Eq. (7.45) below). The constant mass-dependent coefficient t_0 is computed by ensuring that $d\Omega/dt$ peaks at a time $t = t_0$. The parameter \varkappa is computed by enforcing continuity between the first order time derivative of the orbital frequency as predicted by the self-force evolution —Eq. (7.5)— and that given by the first order time derivative of Eq. (7.41).

At the end of the plunge phase, we match the plunge waveforms, which are generated using Eqs. (7.30), (7.31), onto the $l = m = 2, n = 0$ ringdown mode since this dominates the ringdown radiation. In the following Section we will describe in detail the procedure followed to attach the ringdown waveform.

After the transition regime, the equations of motion we use to model the plunge phase are: the second order time derivative of Eq. (7.38) which gives the radial evolution, and Eq. (7.41) which describes the orbital frequency evolution. We determine the point at which to attach the plunge phase by integrating Eq. (7.38) backwards in time, and finding the point at which the transition and plunge equations of motion smoothly match.

In Figures 48 and 49 we show the evolution obtained by combining Ori and Thorne's [288] transition approach for the radial motion with the frequency evolution proposed by Baker et al [361]. In all the cases shown in Figures 48 and 49, the orbital frequency peaks and saturates at a value given by $\omega_{\ell mn}/m$. This can be understood




Figure 48 : (Top, bottom) panels: the left panel shows the inspiral, transition and plunge radial evolution for a BH binary of mass-ratio $q = (1/6, 1/8)$ — and total mass $M_{\text{tot}} \in (7M_{\odot}, 9M_{\odot})$ — using the coordinate transformation given by Eq. (7.15). The right panel shows the orbital frequency $M\Omega$ from late inspiral all the way to the light ring. The evolution starts from an initial radial value $r = 30M$.

if we analyze the asymptotic behavior of Eq. (7.41) near the light-ring, i.e.,

$$\frac{d\phi}{dt} \approx \Omega_f - (\Omega_f - \Omega_i) e^{-2(t-t_0)/b}. \quad (7.42)$$

Recasting Eq. (7.41) in this form, enables us to identify the constant coefficient b with the e-folding rate for the decay of the fundamental quasinormal mode (QNM). As discussed previously, Eq. (7.41) predicts the expected orbital evolution during late inspiral and onward. To provide a unified description from late inspiral through to ringdown, we have decided to adopt the IRS approach, since this framework allows us to smoothly transition from late inspiral onto the plunge phase, and finally describe the ringdown waveform as a natural consequence of the IRS strain-rate amplitude




Figure 49 : As in Figure 48, but with the (top, bottom) panels showing the radial and orbital frequency evolution for binaries with mass-ratios $q = (1/10, 1/15)$, and total mass $M_{\text{tot}} \in (11M_{\odot}, 16M_{\odot})$, respectively. As before, the evolution starts from an initial radial value $r = 30M$.

decay relation $A^2(t) \propto \dot{\Omega} \bar{\Omega}$ [361]. In other words, since Eq. (7.42) has the correct behavior near the light-ring as predicted by BHPT, the IRS model provides a natural framework to attach the ringdown waveform at the end of the plunge phase. We will discuss this feature in further detail in the following Section.

7.2.5 Ringdown Waveform

Numerical relativity simulations have shown that coalescing binary BHs in general relativity lead to the formation of a distorted rotating remnant, which radiates GWs while it settles down into a stationary Kerr BH [371, 372]. The GWs emitted during this intermediate phase resemble a ringing bell. Hence, this type of radiation is commonly known in the literature as ringdown radiation, and consists of a superposition of QNMs — first discovered in numerical studies of the scattering of GWs in

the Schwarzschild spacetime by Vishveshwara [373]. QNMs are damped oscillations whose frequencies and damping times are uniquely determined by the mass and spin of the post-merger Kerr BH. The frequency $\hat{\omega}$ of each QNM has two components: the real part represents the oscillation frequency, and the imaginary part corresponds to the inverse of the damping time:

$$\hat{\omega} = \omega_{\ell mn} - i/\tau_{\ell mn}. \quad (7.43)$$

As discussed above, the observables $\omega_{\ell mn}, \tau_{\ell mn}$ are uniquely determined by the final mass, M_f , and final spin, q_f , of the post-merger Kerr BH. To determine M_f , we use the analytic phenomenological expression for the final mass of the BH remnant that results from the merger of generic binary BHs on circular quasi-orbits introduced in [374], namely,

$$\frac{M_f}{M} = 1 - \left(1 - \frac{2\sqrt{2}}{3}\right) \eta - 0.543763 \eta^2. \quad (7.44)$$

This expression reproduces the expected result in the test mass particle limit, and also reproduces results from currently available numerical relativity simulations [374, 375]. We determine the final spin of the BH remnant q_f using the fit proposed in [376]:

$$q_f = \sqrt{12} \eta + s_1 \eta^2 + s_2 \eta^3, \quad (7.45)$$

with:

$$s_1 = -3.454 \pm 0.132, \quad s_2 = 2.353 \pm 0.548. \quad (7.46)$$

This prescription is consistent with the numerical relativity simulations described in [376, 375], and reproduces test mass limit predictions. This compact formula is also consistent with the prescriptions introduced in [377, 378]. The largest discrepancy between Eq. (7.45) and those derived in [377, 378] is $\lesssim 2.5\%$ for binaries with $q \lesssim 1/6$. The ringdown waveform is given by [372, 361]

$$h(t) = -(h_+ - i h_\times) = \frac{M_f}{D} \sum_{\ell mn} \mathcal{A}_{\ell mn} e^{-i(\omega_{\ell mn} t + \phi_{\ell mn})} e^{-t/\tau_{\ell mn}}, \quad (7.47)$$

where $\mathcal{A}_{\ell mn}$ and $\phi_{\ell mn}$ are constants to be determined by smoothly matching the plunge waveform onto the subsequent ringdown. The ringdown portion of the self-force waveform model constructed in this chapter includes the mode $\ell = m = 2$ and the tones $n = 0, 1, 2$. The approach we follow to attach the leading mode and overtones is the following:

- In order to ensure continuity between the plunge and ringdown waveforms, we use the end of the plunge waveform — Eqs. (7.30), (7.31) — to construct an interpolation function $F(t)$. The interpolation method used to construct $F(t)$ is a cubic spline.
- We match the plunge waveform onto the leading mode $\ell = m = 2, n = 0$ of the ringdown waveform, Eq. (7.47), at the point where the amplitude of the plunge waveform peaks, t_{\max} . Attaching the mode requires $F(t = t_{\max})$ and $F'(t = t_{\max})$ which are computed from the interpolation function. These conditions fix two constants per polarisation.
- To attach the first overtone, $\ell = m = 2, n = 1$, we insert into Eq. (7.47) the constants determined by attaching the leading mode as seeds to compute the amplitude and phase coefficients for the first overtone by enforcing continuity at $t_{\max} + dt$.
- Finally, we insert into Eq. (7.47) the value of the amplitude and phase coefficients previously determined for the leading mode and first overtone, and determine the four remaining constants by enforcing continuity at $t_{\max} + 2 dt$.

Having described the methodology followed to construct complete waveforms for comparable and intermediate mass-ratio systems, we finish this Section by putting together all these various pieces to construct sample waveforms for a few systems with mass-ratio $q \in [1/6, 1/8, 1/10, 1/15]$, and total mass $M_{\text{tot}} \in (7M_{\odot}, 9M_{\odot}, 11M_{\odot}, 16M_{\odot})$ in Figure 50.

7.2.6 Ringdown waveform construction in the context of an Implicit Rotating Source

Having described the ringdown waveform construction as a sum of quasinormal modes, we finish this Section by exhibiting the power of the IRS model to describe the




Figure 50 : The panels show sample waveforms from inspiral to ringdown for systems with mass-ratios $q \in [1/6, 1/8]$ —and total mass $M_{\text{tot}} \in (7M_{\odot}, 9M_{\odot})$ — (top panels—from left to right) and $q \in [1/10, 1/15]$ —and total mass $M_{\text{tot}} \in (11M_{\odot}, 16M_{\odot})$ — (bottom panels —from left to right). The inspiral evolution for the [top, bottom] panels starts from $r = [30M, 25M]$.

ringdown evolution. In the IRS model, the strain-rate amplitude decay is given by [361]:

$$A^2(t) = 16\pi \dot{E} \approx 16\pi \xi \Omega \dot{\Omega}. \quad (7.48)$$

Using Eq. (7.42), in the limit $\Omega \rightarrow \Omega_f$, the amplitude decay is given by

$$A_0^2 e^{-2t/\tau} \approx \frac{32\pi \xi \Omega_f}{b} (\Omega_f - \Omega_i) e^{-2(t-t_0)/\tau}. \quad (7.49)$$

Hence, the late-time amplitude in the IRS model is given by

$$A_{\ell m}^2 \approx 16\pi \xi_{\ell m} \omega_{\ell m} \dot{\omega}_{\ell m}, \quad (7.50)$$

where the value of $\xi_{\ell m}$ is set by ensuring amplitude continuity at the light-ring. In Figure 51 we explicitly show the equivalence of the ringdown waveform construction both in the IRS approach and using the sum of QNMs utilized in the previous Section. This detailed analysis shows that the IRS is a powerful tool to model the late time portions of the waveforms in a unified way.




Figure 51 : The panels show sample the late-time evolution of waveforms whose ringdown phase is modeled using the implicit rotating source (IRS) model and a sum of quasinormal modes (QNMs). The systems shown correspond to binaries with mass-ratios $q \in [1/6, 1/8]$ —and total mass $M_{\text{tot}} \in (7M_{\odot}, 9M_{\odot})$ — (top panels—from left to right) and $q \in [1/10, 1/15]$ —and total mass $M_{\text{tot}} \in (11M_{\odot}, 16M_{\odot})$ — (bottom panels —from left to right). The inspiral evolution for the [top, bottom] panels starts from $r = [30M, 25M]$.

7.2.7 Summary

In this Section we briefly summarize the key ingredients that were used to develop the waveform model described in this chapter:

- Inspiral evolution

- The building blocks of the inspiral evolution are the expressions for the energy, E , and angular momentum, L , — Eqs. (7.13) and (7.14) — that include gravitational self-force corrections and are valid over the domain $0 < x < \frac{1}{3}$ [350].
- The orbital frequency is modeled using Eq. (7.5). This prescription encapsulates gravitational self-force corrections that render a better phase evolution when calibrated against EOB.
- The inspiral trajectory is modeled using the simple prescription (7.26). This scheme is no longer valid near ISCO, where $dE/dx = 0$ for binaries with $q \leq 6$.
- We construct the inspiral waveform using Eqs. (7.30), (7.31).

When the inspiralling object nears the ISCO, we need to invoke the ‘transition scheme’ introduced by Ori and Thorne [288], which enables us to smoothly attach the late inspiral evolution onto the plunge phase.

- Transition phase

- The transition regime starts at a point when dE/dx satisfies Eq. (7.40).
- The equations of motion that govern the transition phase are (7.34), (7.35). These relations are valid, since the motion near the ISCO is nearly-circular.
- Using Eqs. (7.34), (7.35), we linearize the second order time derivative of Eq. (7.38).
- In order to reproduce the orbital phase evolution predicted by numerical simulations from the ISCO to the light-ring, we modify the original transition phase by smoothly matching the inspiral orbital phase evolution, Eq. (7.5), onto the IRS model, Eq. (7.41) at the start of the transition phase.

- Plunge phase

- The equations of motion that govern the plunge phase are given by the second order time derivative of Eq. (7.38), and Eq. (7.41).
- We integrate these relations backwards in time to find the point at which both the transition and plunge equations of motion smoothly match. The transition phase ends at this point.
- Near the light-ring Eq. (7.41) has the behavior predicted by BHPT, which enables us to smoothly match the plunge phase onto the ringdown.
- Both the transition and plunge waveforms are constructed using Eqs. (7.30) and (7.31).

Having constructed the waveform from inspiral to ringdown, we derived second-order radiative corrections to improve the radiative evolution of the waveform model. This was necessary in order to construct a waveform model that is internally consistent, i.e., since the orbital elements include first-order conservative corrections, then radiative corrections should enter the fluxes at second order. We have derived these corrections by enforcing a close agreement between our self-force model and EOB, and have shown that our model can reproduce the orbital phase evolution predicted by EOB within the numerical error of the simulations used to calibrate this model for a variety of mass-ratios.

- Ringdown phase

- The ringdown waveform is constructed using Eq. (7.47).
- We use the plunge waveform to construct an interpolation function $F(t)$, and then use this function to attach the leading mode $\ell = m = 2$, $n = 0$ at the point where the amplitude of the plunge waveform peaks, t_{\max} . We enforce continuity by ensuring that $F(t_{\max}) = h_{\text{RD}}^{n=0}$ and $F'(t_{\max}) = h'_{\text{RD}}^{n=0}$.
- We include the first and second overtone $n = 1$, $n = 2$ in the ringdown waveform.
- Using the IRS model, we have shown that having knowledge of the time evolution of the orbital frequency provides sufficient information to construct the amplitude decay during ringdown. Hence, we can construct

an alternative ringdown waveform using only Eq. (7.50), and ensuring smooth continuity with the plunge waveform.

- Finally, we have explicitly shown that the implicit rotating source approach provides a natural transition from late-time radiation to ringdown that is equivalent to ringdown waveform modeling based on a sum of QNMs.

Throughout the chapter we have emphasized the fact that our model provides a more reliable framework to model binaries whose components are non-spinning, and with mass-ratios $q \leq 1/6$, as compared to available PN approximants. It is worth emphasizing that our model is also computationally inexpensive. All the waveforms we generated to constrain the higher-order η corrections in the energy flux—Eq. (7.22)—can be generated in fractions of a second. A direct comparison between our code and EOB shows that, averaged over 500 realizations, our code is $\sim 20\%$ faster than the optimized version of the EOB code currently available in the LIGO Scientific Collaboration LAL library. It should be emphasized, though, that our code at present has not been optimized, and hence, compared to EOB our model is expected to significantly reduce the cost of waveform generation, making it relatively more viable for parameter estimation efforts. This is a key feature in our model that enabled us to sample a wide region of parameter space to constrain the B_i coefficients in Eq. (7.22). Furthermore, these self-forced waveforms do not need to be highly sampled near the light-ring, hence decreasing the speed with which they can be generated, because the prescription we have used to model the late-time orbital frequency evolution provides the correct evolution near the light-ring. This particular feature also enables us to match the plunge waveform onto its ringdown counterpart without having to interpolate the orbital frequency evolution using a phenomenological approach. The model is internally consistent and the only phenomenology invoked during its construction is related to currently unknown physics, i.e., higher order radiative corrections to the energy flux. Once these corrections are formally derived in the near future, the flexibility of our model will enable us to replace the radiative corrections that we have currently computed by numerical optimization. At that stage, we will be able to describe in a single unified model the dynamical evolution of binaries whose mass-ratios range from the extreme to the comparable regime.

7.3 Conclusions

In this chapter we have developed a self-force waveform model that is capable of reproducing with good accuracy the inspiral, merger and ringdown of binaries with mass-ratios $q \leq 1/6$. This work suggests that a model that incorporates first order conservative self-force corrections in the orbital elements, and second-order radiative corrections in the dissipative piece of the self-force may suffice to describe in a unified manner the coalescence of binaries with mass-ratios that range from the comparable to the extreme. Our model includes conservative self-force corrections that have been derived in the strong-field regime [350]. Using these results, we find that binaries with mass-ratios $q \gtrsim 1/6$ do not have an ISCO. For systems with $q \leq 1/6$, we have derived a simple relation that provides the location of the ISCO in terms of the symmetric mass-ratio (see Eq. (7.18)). To describe the inspiral evolution, we have derived second-order corrections to the energy flux by minimizing the phase discrepancy between our self-force model and the EOB model [?, 348] for a variety of mass-ratios. We have shown that our model reproduces the phase evolution of the EOB model within the accuracy of available numerical simulations for a variety of mass-ratios.

This chapter also presents an extension of the “transition regime” developed by Ori and Thorne [288] to smoothly match the late inspiral evolution onto the plunge phase. We have found that using the inspiral phase expression for the orbital frequency during the plunge phase does not render an accurate description of the actual orbital evolution as predicted by numerical simulations. Hence, we have embedded the self-force framework in the IRS model proposed by Baker et al [361] to ensure that our model is as close as possible to the orbital evolution predicted by numerical relativity simulations. The implementation of this prescription ensures that the orbital frequency saturates near the light-ring, which facilitates matching onto the ringdown phase. We have shown that the IRS model provides a natural transition onto the ringdown phase that is equivalent to a ringdown waveform construction based on a sum of QNMs.

The motivation for constructing this model is two-fold: to exhibit the versatility of the self-force formalism to accurately describe the evolution of binaries beyond the extreme mass-ratio limit; and to provide a tool that can be used to explore the

information that could be extracted by GW detectors that target binaries with comparable and intermediate mass-ratios. Current studies have only explored the use of PN approximants to model the merger of NSBH binaries, despite their inadequacy to capture the evolution of these systems [379, 352, 353] (see Figure 41). Comparing Figures 41 and 45, we conclude that even if second generation GW detectors were only capable of capturing the inspiral evolution of NSBH mergers, our self-force model would be better equipped to describe these events. The construction of this IRS self-force model constitutes an important step towards the construction of more reliable waveforms to describe IMRCs. In citeSmith:2013, it was shown that Huerta–Gair (HG) waveforms — which are closely related to the model developed in this chapter, and which were also based on a consistent combination of BHPT and self-force corrections — work in the relevant mass range for advanced detectors. The new model introduced in this article, improves upon these waveforms and should therefore not only work in the regime of interest to advanced detectors but over a wider parameter space that could be applicable to third generation detectors or later observations.

Having developed a strong foundation to model binaries on circular orbits whose components are not rotating, it is necessary to incorporate more ingredients to the waveform model to capture GW signals from binaries whose components have significant spin [380, ?, 381, 382, 267, 383, 384, 354], or systems that form in core-collapsed globular clusters, and hence are expected to have non-negligible eccentricity at merger [385, 386]. In order to do so, we require input both from the self-force program — which is making substantial progress towards the computation of the self-force in a Kerr background [387, 388, 389]— and from numerical simulations [?], in particular from binaries with typical mass-ratios $1/20 < q < 1/10$ to explore the performance of our self-force model in this currently unconstrained regime. Undoubtedly, the development and implementation of improved numerical algorithms [390] to carry out these simulations will facilitate the realization of these studies in the near future.

Recent observational discoveries [300] suggest that NSBH mergers may also occur in globular clusters. Hence, in these dense stellar environments we may expect that multiple n -body interactions and binary exchange processes may lead to the formation of binaries on eccentric orbits. Detectors such as the Einstein Telescope, operating at low frequencies $\sim 1\text{Hz}$, may be capable of detecting the signature of eccentricity

during the early inspiral. In order to assess these effects, we aim to extend the model introduced in this chapter by including eccentricity, making use of self-force corrections for generic orbits in a Schwarzschild background [264].

Chapter 8

Conclusions

Bibliography

- [1] R. A. Hulse and J. H. Taylor. *Discovery of a pulsar in a binary system.* *ApJL*, 195:L51–L53, January 1975.
- [2] J.M. Weisberg, J.H. Taylor, and L.A. Fowler. GRAVITATIONAL WAVES FROM AN ORBITING PULSAR. *Sci.Am.*, 245:66–74, 1981.
- [3] J. H. Taylor and J. M. Weisberg. Further experimental tests of relativistic gravity using the binary pulsar PSR 1913 + 16. *Astrophys. J.*, 345:434–450, October 1989.
- [4] Marta Burgay, N. D’Amico, A. Possenti, R.N. Manchester, A.G. Lyne, et al. An Increased estimate of the merger rate of double neutron stars from observations of a highly relativistic system. *Nature*, 426:531–533, 2003.
- [5] F. A. E. Pirani. *Acta Physica Polonica*, 15:389, 1956.
- [6] G. F. Moss, L. R. Miller, and Forward R. L. Photon-noise-limited laser transducer for gravitational antenna. *Applied Optics*, 10:2495–2498, 1971.
- [7] R Weiss. Electromagnetically coupled broadband gravitational wave antenna. *Quarterly Progress Report of the Research Laboratory of Electronics of the Massachusetts Institutue of Technology*, 105:54, April 1972.
- [8] R. P. Drever et al. A gravity-wave detector using optical cavity sensing. In E. Schmutzler, editor, *Proceedings of the Ninth International Conference on General Relativity and Gravitation*, pages 265–267. Cambridge University Press, Cambridge, UK, 1983.
- [9] Peter R. Saulson. Fundamentals of interferometric gravitational wave detectors. 1995.

- [10] Eirini Messaritaki. Report on the first binary black hole inspiral search in LIGO data. *Class.Quant.Grav.*, 22:S1119–S1128, 2005.
- [11] J. Abadie et al. All-sky search for gravitational-wave bursts in the first joint LIGO-GEO-Virgo run. *Phys.Rev.*, D81:102001, 2010.
- [12] J. Abadie et al. All-sky search for gravitational-wave bursts in the second joint LIGO-Virgo run. *Phys.Rev.*, D85:122007, 2012.
- [13] B.P. Abbott et al. Search for gravitational wave ringdowns from perturbed black holes in LIGO S4 data. *Phys.Rev.*, D80:062001, 2009.
- [14] J. Abadie et al. Search for Gravitational Waves from Low Mass Compact Binary Coalescence in LIGO’s Sixth Science Run and Virgo’s Science Runs 2 and 3. *Phys.Rev.*, D85:082002, 2012.
- [15] J. Abadie et al. Search for Gravitational Waves from Compact Binary Coalescence in LIGO and Virgo Data from S5 and VSR1. *Phys.Rev.*, D82:102001, 2010.
- [16] B.P. Abbott et al. Search for Gravitational Waves from Low Mass Compact Binary Coalescence in 186 Days of LIGO’s fifth Science Run. *Phys.Rev.*, D80:047101, 2009.
- [17] B.P. Abbott et al. Search for Gravitational Waves from Low Mass Binary Coalescences in the First Year of LIGO’s S5 Data. *Phys.Rev.*, D79:122001, 2009.
- [18] J. Abadie et al. Search for gravitational waves from binary black hole inspiral, merger and ringdown. *Phys.Rev.*, D83:122005, 2011.
- [19] J. Aasi et al. Search for Gravitational Waves from Binary Black Hole Inspiral, Merger and Ringdown in LIGO-Virgo Data from 2009-2010. *Phys.Rev.*, D87:022002, 2013.
- [20] B. Abbott et al. Setting upper limits on the strength of periodic gravitational waves using the first science data from the GEO 600 and LIGO detectors. *Phys.Rev.*, D69:082004, 2004.

- [21] B. Abbott et al. First all-sky upper limits from LIGO on the strength of periodic gravitational waves using the Hough transform. *Phys.Rev.*, D72:102004, 2005.
- [22] Alicia M. Sintes. Recent results on the search for continuous sources with LIGO and GEO 600. *J.Phys.Conf.Ser.*, 39:36–38, 2006.
- [23] J. Abadie et al. Beating the spin-down limit on gravitational wave emission from the Vela pulsar. *Astrophys.J.*, 737:93, 2011.
- [24] C. Palomba. Searches for continuous gravitational wave signals and stochastic backgrounds in LIGO and Virgo data. 2012.
- [25] Gregory M. Harry et al. Advanced LIGO: The next generation of gravitational wave detectors. *Class.Quant.Grav.*, 27:084006, 2010.
- [26] F. Acernese et al. Advanced Virgo Baseline Design, 2009. [Virgo Technical Document VIR-0027A-09].
- [27] J. Aasi et al. Prospects for Localization of Gravitational Wave Transients by the Advanced LIGO and Advanced Virgo Observatories. 2013.
- [28] J. Abadie et al. Predictions for the Rates of Compact Binary Coalescences Observable by Ground-based Gravitational-wave Detectors. *Class.Quant.Grav.*, 27:173001, 2010.
- [29] Luc Blanchet. Gravitational radiation from post-Newtonian sources and inspiralling compact binaries. *Living Rev.Rel.*, 9:4, 2006.
- [30] A. Buonanno and T. Damour. Effective one-body approach to general relativistic two-body dynamics. *Phys.Rev.*, D59:084006, 1999.
- [31] S. E. Gralla and R. M. Wald. A rigorous derivation of gravitational self-force. *Classical and Quantum Gravity*, 25(20):205009, October 2008.
- [32] S. E. Gralla and R. M. Wald. Derivation of Gravitational Self-Force. *ArXiv e-prints*, July 2009.
- [33] Frans Pretorius. Evolution of binary black hole spacetimes. *Phys.Rev.Lett.*, 95:121101, 2005.

- [34] Frans Pretorius. Simulation of binary black hole spacetimes with a harmonic evolution scheme. *Class.Quant.Grav.*, 23:S529–S552, 2006.
- [35] Manuela Campanelli, C.O. Lousto, P. Marronetti, and Y. Zlochower. Accurate evolutions of orbiting black-hole binaries without excision. *Phys.Rev.Lett.*, 96:111101, 2006.
- [36] Simulating eXtreme Spacetimes collaboration.
- [37] Abdul H. Mroue et al. A catalog of 174 binary black-hole simulations for gravitational-wave astronomy. *Phys.Rev.Lett.*, 111:241104, 2013.
- [38] Yi Pan, Alessandra Buonanno, Michael Boyle, Luisa T. Buchman, Lawrence E. Kidder, et al. Inspiral-merger-ringdown multipolar waveforms of nonspinning black-hole binaries using the effective-one-body formalism. *Phys.Rev.*, D84:124052, 2011. 26 pages, 25 figures.
- [39] L. Santamaria, F. Ohme, P. Ajith, B. Bruegmann, N. Dorband, et al. Matching post-Newtonian and numerical relativity waveforms: systematic errors and a new phenomenological model for non-precessing black hole binaries. *Phys.Rev.*, D82:064016, 2010.
- [40] M. Brett Deaton, Matthew D. Duez, Francois Foucart, Evan O'Connor, Christian D. Ott, et al. Black Hole-Neutron Star Mergers with a Hot Nuclear Equation of State: Outflow and Neutrino-Cooled Disk for a Low-Mass, High-Spin Case. *Astrophys.J.*, 776:47, 2013.
- [41] Philipp Msta, Sherwood Richers, Christian D. Ott, Roland Haas, Anthony L. Piro, et al. Magnetorotational Core-Collapse Supernovae in Three Dimensions. *Astrophys.J.*, 785:L29, 2014.
- [42] Kentaro Somiya. Detector configuration of KAGRA: The Japanese cryogenic gravitational-wave detector. *Class.Quant.Grav.*, 29:124007, 2012.
- [43] S.J. Waldman. The Advanced LIGO Gravitational Wave Detector. 2011.
- [44] S. W. Hawking and W. Israel. *Three Hundred Years of Gravitation*. Cambridge University Press, 1987.

- [45] Thibault Damour, Alessandro Nagar, Mark Hannam, Sascha Husa, and Bernd Bruegmann. Accurate Effective-One-Body waveforms of inspiralling and coalescing black-hole binaries. *Phys.Rev.*, D78:044039, 2008.
- [46] Alessandra Buonanno, Yi Pan, Harald P. Pfeiffer, Mark A. Scheel, Luisa T. Buchman, et al. Effective-one-body waveforms calibrated to numerical relativity simulations: Coalescence of non-spinning, equal-mass black holes. *Phys.Rev.*, D79:124028, 2009.
- [47] Thibault Damour and Alessandro Nagar. An Improved analytical description of inspiralling and coalescing black-hole binaries. *Phys.Rev.*, D79:081503, 2009.
- [48] Thibault Damour and Alessandro Nagar. Comparing Effective-One-Body gravitational waveforms to accurate numerical data. *Phys.Rev.*, D77:024043, 2008.
- [49] Thibault Damour, Alessandro Nagar, Ernst Nils Dorband, Denis Pollney, and Luciano Rezzolla. Faithful Effective-One-Body waveforms of equal-mass coalescing black-hole binaries. *Phys.Rev.*, D77:084017, 2008.
- [50] Alessandra Buonanno and Thibault Damour. Transition from inspiral to plunge in binary black hole coalescences. *Phys.Rev.*, D62:064015, 2000. 52 pages, 21 figures, ReVTex, epsfig; few misprints corrected.
- [51] Thibault Damour, Bala R. Iyer, Piotr Jaranowski, and B.S. Sathyaprakash. Gravitational waves from black hole binary inspiral and merger: The Span of third postNewtonian effective one-body templates. *Phys.Rev.*, D67:064028, 2003.
- [52] Emanuele Berti, Vitor Cardoso, and Clifford M. Will. On gravitational-wave spectroscopy of massive black holes with the space interferometer LISA. *Phys.Rev.*, D73:064030, 2006.
- [53] Yasushi Mino, Misao Sasaki, Masaru Shibata, Hideyuki Tagoshi, and Takahiro Tanaka. Black hole perturbation: Chapter 1. *Prog.Theor.Phys.Suppl.*, 128:1–121, 1997.
- [54] L A Wainstein and V D Zubakov. *Extraction of signals from noise*. Prentice-Hall, Englewood Cliffs, NJ, 1962.

- [55] Bruce Allen, Warren G. Anderson, Patrick R. Brady, Duncan A. Brown, and Jolien D.E. Creighton. FINDCHIRP: An Algorithm for detection of gravitational waves from inspiraling compact binaries. *Phys.Rev.*, D85:122006, 2012.
- [56] Feryal Ozel, Dimitrios Psaltis, Ramesh Narayan, and Jeffrey E. McClintock. The Black Hole Mass Distribution in the Galaxy. *Astrophys.J.*, 725:1918–1927, 2010.
- [57] B.S. Sathyaprakash and S.V. Dhurandhar. Choice of filters for the detection of gravitational waves from coalescing binaries. *Phys.Rev.*, D44:3819–3834, 1991.
- [58] R. Balasubramanian, B.S. Sathyaprakash, and S.V. Dhurandhar. Gravitational waves from coalescing binaries: Detection strategies and Monte Carlo estimation of parameters. *Phys.Rev.*, D53:3033–3055, 1996.
- [59] T.A. Apostolatos. Search templates for gravitational waves from precessing, inspiraling binaries. *Phys.Rev.*, D52:605–620, 1995.
- [60] Kipp Cannon et al. Singular value decomposition applied to compact binary coalescence gravitational-wave signals. *Phys.Rev.*, D82:044025, 2010.
- [61] Benjamin J. Owen. Search templates for gravitational waves from inspiraling binaries: Choice of template spacing. *Phys.Rev.*, D53:6749–6761, 1996.
- [62] B.S. Sathyaprakash. Filtering postNewtonian gravitational waves from coalescing binaries. *Phys.Rev.*, D50:7111–7115, 1994.
- [63] S. Babak, R. Balasubramanian, D. Churches, T. Cokelaer, and B.S. Sathyaprakash. A Template bank to search for gravitational waves from inspiralling compact binaries. I. Physical models. *Class.Quant.Grav.*, 23:5477–5504, 2006.
- [64] Benjamin J. Owen and B.S. Sathyaprakash. Matched filtering of gravitational waves from inspiraling compact binaries: Computational cost and template placement. *Phys.Rev.*, D60:022002, 1999.
- [65] Thomas Cokelaer. Gravitational waves from inspiralling compact binaries: Hexagonal template placement and its efficiency in detecting physical signals. *Phys.Rev.*, D76:102004, 2007.

- [66] LIGO (David Shoemaker). Advanced LIGO anticipated sensitivity curves. Technical report, LIGO Document T0900288-v3, 2009.
- [67] Curt Cutler and Eanna E. Flanagan. Gravitational waves from merging compact binaries: How accurately can one extract the binary's parameters from the inspiral wave form? *Phys.Rev.*, D49:2658–2697, 1994.
- [68] Serge Droz, Daniel J. Knapp, Eric Poisson, and Benjamin J. Owen. Gravitational waves from inspiraling compact binaries: Validity of the stationary phase approximation to the Fourier transform. *Phys.Rev.*, D59:124016, 1999.
- [69] Piotr Jaranowski and Gerhard Schaefer. The Binary black hole dynamics at the third postNewtonian order in the orbital motion. *Annalen Phys.*, 9:378–383, 2000.
- [70] Piotr Jaranowski and Gerhard Schaefer. The Binary black hole problem at the third postNewtonian approximation in the orbital motion: Static part. *Phys.Rev.*, D60:124003, 1999.
- [71] Thibault Damour, Piotr Jaranowski, and Gerhard Schaefer. Dimensional regularization of the gravitational interaction of point masses. *Phys.Lett.*, B513:147–155, 2001.
- [72] Luc Blanchet, Guillaume Faye, Bala R. Iyer, and Siddhartha Sinha. The Third post-Newtonian gravitational wave polarisations and associated spherical harmonic modes for inspiralling compact binaries in quasi-circular orbits. *Class.Quant.Grav.*, 25:165003, 2008. 57 pages, no figures.
- [73] Alessandra Buonanno, Bala Iyer, Evan Ochsner, Yi Pan, and B.S. Sathyaprakash. Comparison of post-Newtonian templates for compact binary inspiral signals in gravitational-wave detectors. *Phys.Rev.*, D80:084043, 2009.
- [74] D.J.A. McKechnie. On the use of higher order waveforms in the search for gravitational waves emitted by compact binary coalescences. 2011. Ph.D. thesis.
- [75] Larne Pekowsky, James Healy, Deirdre Shoemaker, and Pablo Laguna. Impact of Higher-order Modes on the Detection of Binary Black Hole Coalescences. 2012.

- [76] Jose A. Gonzalez, Ulrich Sperhake, and Bernd Bruegmann. Black-hole binary simulations: The Mass ratio 10:1. *Phys.Rev.*, D79:124006, 2009.
- [77] Denis Pollney, Christian Reisswig, Erik Schnetter, Nils Dorband, and Peter Diener. High accuracy binary black hole simulations with an extended wave zone. *Phys.Rev.*, D83:044045, 2011.
- [78] Carlos O. Lousto, Hiroyuki Nakano, Yosef Zlochower, and Manuela Campanelli. Intermediate Mass Ratio Black Hole Binaries: Numerical Relativity meets Perturbation Theory. *Phys.Rev.Lett.*, 104:211101, 2010.
- [79] Luisa T. Buchman, Harald P. Pfeiffer, Mark A. Scheel, and Bela Szilagyi. Simulations of non-equal mass black hole binaries with spectral methods. *Phys.Rev.*, D86:084033, 2012.
- [80] P. Ajith, Michael Boyle, Duncan A. Brown, Bernd Brugmann, Luisa T. Buchman, et al. The NINJA-2 catalog of hybrid post-Newtonian/numerical-relativity waveforms for non-precessing black-hole binaries. *Class.Quant.Grav.*, 29:124001, 2012.
- [81] Thibault Damour, Piotr Jaranowski, and Gerhard Schaefer. On the determination of the last stable orbit for circular general relativistic binaries at the third postNewtonian approximation. *Phys.Rev.*, D62:084011, 2000.
- [82] Thibault Damour, Bala R. Iyer, and Alessandro Nagar. Improved resummation of post-Newtonian multipolar waveforms from circularized compact binaries. *Phys.Rev.*, D79:064004, 2009.
- [83] A. Buonanno and T. Damour. Effective one-body approach to general relativistic two-body dynamics. *Phys.Rev.*, D59:084006, 1999.
- [84] Alessandra Buonanno, Yanbei Chen, and Thibault Damour. Transition from inspiral to plunge in precessing binaries of spinning black holes. *Phys.Rev.*, D74:104005, 2006.
- [85] LSC Algorithm Library. <https://www.lsc-group.phys.uwm.edu/daswg/projects/lalsuite.html>.

- [86] Jon Mathews and Robert L. Walker. *Mathematical Methods of Physics*. W.A. Benjamin; 2nd edition, 1970.
- [87] Luc Blanchet, Thibault Damour, Bala R. Iyer, Clifford M. Will, and Alan.G. Wiseman. Gravitational radiation damping of compact binary systems to second postNewtonian order. *Phys.Rev.Lett.*, 74:3515–3518, 1995.
- [88] Luc Blanchet, Guillaume Faye, Bala R. Iyer, and Benoit Joguet. Gravitational wave inspiral of compact binary systems to 7/2 postNewtonian order. *Phys.Rev.*, D65:061501, 2002.
- [89] Luc Blanchet, Thibault Damour, Gilles Esposito-Farese, and Bala R. Iyer. Gravitational radiation from inspiralling compact binaries completed at the third post-Newtonian order. *Phys.Rev.Lett.*, 93:091101, 2004.
- [90] Eric Poisson and Clifford M. Will. Gravitational waves from inspiraling compact binaries: Parameter estimation using second postNewtonian wave forms. *Phys.Rev.*, D52:848–855, 1995.
- [91] Lee Lindblom, Benjamin J. Owen, and Duncan A. Brown. Model Waveform Accuracy Standards for Gravitational Wave Data Analysis. *Phys.Rev.*, D78:124020, 2008.
- [92] Lee Lindblom, John G. Baker, and Benjamin J. Owen. Improved Time-Domain Accuracy Standards for Model Gravitational Waveforms. *Phys.Rev.*, D82:084020, 2010.
- [93] Lee Samuel Finn and David F. Chernoff. Observing binary inspiral in gravitational radiation: One interferometer. *Phys.Rev.*, D47:2198–2219, 1993.
- [94] Alessandra Buonanno, Yi Pan, John G. Baker, Joan Centrella, Bernard J. Kelly, et al. Toward faithful templates for non-spinning binary black holes using the effective-one-body approach. *Phys.Rev.*, D76:104049, 2007.
- [95] B.S. Sathyaprakash and B.F. Schutz. Physics, Astrophysics and Cosmology with Gravitational Waves. *Living Rev.Rel.*, 12:2, 2009.

- [96] A. Tutukov and L. Yungelson. Evolution of close binary with relativistic component. *Nauchnye Informatsii*, 27:86, 1973.
- [97] A. Tutukov and L. Yungelson. Evolution of massive close binaries. *Nauchnye Informatsii*, 27:70, 1973.
- [98] R.F. Webbink. Double white dwarfs as progenitors of R Coronae Borealis stars and Type I supernovae. *Astrophys.J.*, 277:355–360, 1984.
- [99] I. Kowalska, T. Regimbau, T. Bulik, M. Dominik, and K. Belczynski. Effect of metallicity on the gravitational-wave signal from the cosmological population of compact binary coalescences. 2012.
- [100] Chris L. Fryer, Krzysztof Belczynski, Grzegorz Wiktorowicz, Michal Dominik, Vicky Kalogera, et al. Compact Remnant Mass Function: Dependence on the Explosion Mechanism and Metallicity. *Astrophys.J.*, 749:91, 2012.
- [101] Michal Dominik, Krzysztof Belczynski, Christopher Fryer, Daniel Holz, Emanuele Berti, et al. Double Compact Objects I: The Significance of the Common Envelope on Merger Rates. *Astrophys.J.*, 759:52, 2012.
- [102] Krzysztof Belczynski and Michal Dominik. A test of black hole natal kick mechanism by the first gravitational radiation detections. 2012.
- [103] J. Abadie et al. Search for Gravitational Wave Bursts from Six Magnetars. *Astrophys.J.*, 734:L35, 2011.
- [104] G.L. Turin. An introduction to matched filters. *Information Theory, IRE Transactions on*, 6(3):311–329, 1960.
- [105] John G. Baker, Joan Centrella, Dae-Il Choi, Michael Koppitz, and James van Meter. Gravitational wave extraction from an inspiraling configuration of merging black holes. *Phys.Rev.Lett.*, 96:111102, 2006.
- [106] Lee Lindblom, Mark A. Scheel, Lawrence E. Kidder, Robert Owen, and Oliver Rinne. A New generalized harmonic evolution system. *Class.Quant.Grav.*, 23:S447–S462, 2006.

- [107] Frans Pretorius. Binary Black Hole Coalescence. 2007.
- [108] Mark Hannam. Status of black-hole-binary simulations for gravitational-wave detection. *Class.Quant.Grav.*, 26:114001, 2009.
- [109] Ian Hinder. The Current Status of Binary Black Hole Simulations in Numerical Relativity. *Class.Quant.Grav.*, 27:114004, 2010.
- [110] Harald P. Pfeiffer. Numerical simulations of compact object binaries. *Class.Quant.Grav.*, 29:124004, 2012.
- [111] <http://www.black-holes.org/SpEC.html>.
- [112] Abdul H. Mroue and Harald P. Pfeiffer. Precessing Binary Black Holes Simulations: Quasicircular Initial Data. 2012.
- [113] Andrea Taracchini, Yi Pan, Alessandra Buonanno, Enrico Barausse, Michael Boyle, et al. Prototype effective-one-body model for nonprecessing spinning inspiral-merger-ringdown waveforms. *Phys.Rev.*, D86:024011, 2012.
- [114] Michael Boyle. Uncertainty in hybrid gravitational waveforms: Optimizing initial orbital frequencies for binary black-hole simulations. *Phys.Rev.*, D84:064013, 2011.
- [115] Ilana MacDonald, Samaya Nissanke, Harald P. Pfeiffer, and Harald P. Pfeiffer. Suitability of post-Newtonian/numerical-relativity hybrid waveforms for gravitational wave detectors. *Class.Quant.Grav.*, 28:134002, 2011.
- [116] Ilana MacDonald, Abdul H. Mroue, Harald P. Pfeiffer, Michael Boyle, Lawrence E. Kidder, et al. Suitability of hybrid gravitational waveforms for unequal-mass binaries. *Phys.Rev.*, D87:024009, 2013.
- [117] Frank Ohme, Mark Hannam, and Sascha Husa. Reliability of complete gravitational waveform models for compact binary coalescences. *Phys.Rev.*, D84:064029, 2011.
- [118] Mark Hannam, Sascha Husa, Frank Ohme, and P. Ajith. Length requirements for numerical-relativity waveforms. *Phys.Rev.*, D82:124052, 2010.

- [119] Parameswaran Ajith, S. Babak, Y. Chen, M. Hewitson, B. Krishnan, et al. Phenomenological template family for black-hole coalescence waveforms. *Class.Quant.Grav.*, 24:S689–S700, 2007.
- [120] Benjamin Aylott, John G. Baker, William D. Boggs, Michael Boyle, Patrick R. Brady, et al. Status of NINJA: The Numerical INjection Analysis project. *Class.Quant.Grav.*, 26:114008, 2009.
- [121] Lucia Santamaria, Badri Krishnan, and John T. Whelan. Searching for numerically-simulated signals of black hole binaries with a phenomenological template family. *Class.Quant.Grav.*, 26:114010, 2009.
- [122] Benjamin Aylott, John G. Baker, William D. Boggs, Michael Boyle, Patrick R. Brady, et al. Testing gravitational-wave searches with numerical relativity waveforms: Results from the first Numerical INjection Analysis (NINJA) project. *Class.Quant.Grav.*, 26:165008, 2009.
- [123] Duncan A. Brown, Prayush Kumar, and Alexander H. Nitz. Template banks to search for low-mass binary black holes in advanced gravitational-wave detectors. *Phys. Rev. D*, 87:082004, Apr 2013.
- [124] Bela Szilagyi. Stretching the limits of NR (NRDA 2013). http://www.grg.uib.es/NRDA13/slides/Szilagyi_StretchingTheLimitsOfNR.pdf.
- [125] J. Abadie et al. Results from the ninja-2 mock data challenge. In Prep.
- [126] Ian W. Harry, Bruce Allen, and B.S. Sathyaprakash. A Stochastic template placement algorithm for gravitational wave data analysis. *Phys.Rev.*, D80:104014, 2009.
- [127] P. Ajith, N. Fotopoulos, S. Privitera, A. Neunzert, and A.J. Weinstein. An effectual template bank for the detection of gravitational waves from inspiralling compact binaries with generic spins. 2012.
- [128] Gian Mario Manca and Michele Vallisneri. Cover art: Issues in the metric-guided and metric-less placement of random and stochastic template banks. *Phys.Rev.*, D81:024004, 2010.

- [129] B.S. Sathyaprakash. Mother templates for gravitational wave chirps. *Class.Quant.Grav.*, 17:L157–162, 2000.
- [130] Mark A. Scheel, Michael Boyle, Tony Chu, Lawrence E. Kidder, Keith D. Matthews, et al. High-accuracy waveforms for binary black hole inspiral, merger, and ringdown. *Phys.Rev.*, D79:024003, 2009.
- [131] Michael Boyle, Duncan A. Brown, Lawrence E. Kidder, Abdul H. Mroue, Harald P. Pfeiffer, et al. High-accuracy comparison of numerical relativity simulations with post-Newtonian expansions. *Phys.Rev.*, D76:124038, 2007.
- [132] Piotr Jaranowski and Gerhard Schaefer. Third postNewtonian higher order ADM Hamilton dynamics for two-body point mass systems. *Phys.Rev.*, D57:7274–7291, 1998.
- [133] Luc Blanchet, Thibault Damour, and Gilles Esposito-Farese. Dimensional regularization of the third postNewtonian dynamics of point particles in harmonic coordinates. *Phys.Rev.*, D69:124007, 2004.
- [134] Thibault Damour, Piotr Jaranowski, and Gerhard Schaefer. Equivalence between the ADM-Hamiltonian and the harmonic coordinates approaches to the third postNewtonian dynamics of compact binaries. *Phys.Rev.*, D63:044021, 2001.
- [135] Luc Blanchet and Bala R. Iyer. Third postNewtonian dynamics of compact binaries: Equations of motion in the center-of-mass frame. *Class.Quant.Grav.*, 20:755, 2003.
- [136] Luc Blanchet, Guillaume Faye, Bala R. Iyer, and Benoit Joguet. Gravitational wave inspiral of compact binary systems to 7/2 postNewtonian order. *Phys.Rev.*, D65:061501, 2002.
- [137] Luc Blanchet, Thibault Damour, Gilles Esposito-Farese, and Bala R. Iyer. Dimensional regularization of the third post-Newtonian gravitational wave generation from two point masses. *Phys.Rev.*, D71:124004, 2005.

- [138] Luc Blanchet and Bala R. Iyer. Hadamard regularization of the third post-Newtonian gravitational wave generation of two point masses. *Phys.Rev.*, D71:024004, 2005.
- [139] Jolien D.E. Creighton and Warren G. Anderson. *Gravitational-Wave Physics and Astronomy: An Introduction to Theory, Experiment and Data Analysis*. Wiley-VCH, 2011.
- [140] Kipp Cannon, J.D. Emberson, Chad Hanna, Drew Keppel, and Harald Pfeiffer. Interpolation in waveform space: enhancing the accuracy of gravitational waveform families using numerical relativity. *Phys.Rev.*, D87(4):044008, 2013.
- [141] Mark A. Scheel, Bela Szilagyi, and Harald P. Pfeiffer. "Private communication."
- [142] S. Babak, R. Biswas, P.R. Brady, D.A. Brown, K. Cannon, et al. Searching for gravitational waves from binary coalescence. *Phys.Rev.*, D87:024033, 2013.
- [143] Numerical Relativity - Analytical Relativity Collaboration (NRAR).
- [144] Eliu A. Huerta, Prayush Kumar, and Duncan A. Brown. Accurate modeling of intermediate-mass-ratio inspirals: exploring the form of the self-force in the intermediate-mass-ratio regime. *Phys.Rev.*, D86:024024, 2012.
- [145] Michael Boyle. Angular velocity of gravitational radiation from precessing binaries and the corotating frame. 2013.
- [146] Patricia Schmidt, Mark Hannam, and Sascha Husa. Towards models of gravitational waveforms from generic binaries: A simple approximate mapping between precessing and non-precessing inspiral signals. *Phys.Rev.*, D86:104063, 2012.
- [147] C. Capano, Y. Pan, and A. Buonanno. Impact of higher harmonics in searching for gravitational waves from non-spinning binary black hole. In Prep.
- [148] T. Accadia et al. Status of the Virgo project. *Class.Quant.Grav.*, 28:114002, 2011.
- [149] K. Kuroda. Status of LCGT. *Class.Quant.Grav.*, 27:084004, 2010.

- [150] Bala Iyer, Tarun Souradeep, C S Unnikrishnan, Sanjeev Dhurandhar, Sendhil Raja, and Sengupta A., *LIGO-India: Proposal of the Consortium for Indian Initiative in Gravitational-wave Observations (IndIGO)*, LIGO Tech. Rep. LIGO-M1100296-v2. <https://dcc.ligo.org/LIGO-M1100296-v2/public>, 2013.
- [151] C.S. Unnikrishnan. IndIGO and LIGO-India: Scope and plans for gravitational wave research and precision metrology in India. *Int.J.Mod.Phys.*, D22:1341010, 2013.
- [152] K S Thorne. Gravitational radiation. In S. W. Hawking and W. Israel, editors, *Three hundred years of gravitation*, chapter 9, pages 330–458. Cambridge University Press, Cambridge, 1987.
- [153] J. Abadie et al. Predictions for the Rates of Compact Binary Coalescences Observable by Ground-based Gravitational-wave Detectors. *Class.Quant.Grav.*, 27:173001, 2010.
- [154] Vassiliki Kalogera and Gordon Baym. The maximum mass of a neutron star. *Astrophys.J.*, 470:L61–L64, 1996.
- [155] Charles D. Bailyn, Raj K. Jain, Paolo Coppi, and Jerome A. Orosz. The Mass distribution of stellar black holes. *Astrophys.J.*, 499:367, May 1997.
- [156] Will M. Farr et al. The Mass Distribution of Stellar-Mass Black Holes. *Astrophys.J.*, 741:103, 2011.
- [157] Jeffrey E. McClintock et al. Measuring the Spins of Accreting Black Holes. *Class.Quant.Grav.*, 28:114009, 2011.
- [158] Krzysztof Belczynski, Ronald E. Taam, Emmanouela Rantsiou, and Marc van der Sluys. Black Hole Spin Evolution: Implications for Short-hard Gamma Ray Bursts and Gravitational Wave Detection. *Astrophys.J.*, 2007.
- [159] L A Wainstein and V D Zubakov. *Statistical Theory of Signal Detection*. Pergamon, London, UK, 1968.

- [160] Alessandra Buonanno, Bala Iyer, Evan Ochsner, Yi Pan, and B.S. Sathyaprakash. Comparison of post-Newtonian templates for compact binary inspiral signals in gravitational-wave detectors. *Phys.Rev.*, D80:084043, 2009.
- [161] S. Klimenko, I. Yakushin, A. Mercer, and Guenakh Mitselmakher. Coherent method for detection of gravitational wave bursts. *Class.Quant.Grav.*, 25:114029, 2008.
- [162] Antony C Searle, Patrick J Sutton, Massimo Tinto, and Graham Woan. Robust Bayesian detection of unmodelled bursts. *Class.Quant.Grav.*, 25:114038, 2008.
- [163] Patrick J. Sutton et al. X-Pipeline: An Analysis package for autonomous gravitational-wave burst searches. *New J.Phys.*, 12:053034, 2010.
- [164] Luc Blanchet. Gravitational radiation from post-Newtonian sources and inspiralling compact binaries. *Living Rev.Rel.*, 9:4, 2006.
- [165] Thibault Damour and Alessandro Nagar. An Improved analytical description of inspiralling and coalescing black-hole binaries. *Phys.Rev.*, D79:081503, 2009.
- [166] Yi Pan et al. Inspiral-merger-ringdown multipolar waveforms of nonspinning black-hole binaries using the effective-one-body formalism. *Phys.Rev.*, D84:124052, 2011.
- [167] P. Ajith et al. Inspiral-merger-ringdown waveforms for black-hole binaries with non-precessing spins. *Phys.Rev.Lett.*, 106:241101, 2011.
- [168] Thibault Damour, Alessandro Nagar, and Sebastiano Bernuzzi. Improved effective-one-body description of coalescing nonspinning black-hole binaries and its numerical-relativity completion. *Phys.Rev.*, D87:084035, 2013.
- [169] Andrea Taracchini et al. Effective-one-body model for black-hole binaries with generic mass ratios and spins. 2013.
- [170] Joan Centrella, John G. Baker, Bernard J. Kelly, and James R. van Meter. Black-hole binaries, gravitational waves, and numerical relativity. *Rev.Mod.Phys.*, 82:3069, 2010.

- [171] D. Brown et al. Data formats for numerical relativity waves. 2007.
- [172] P. Ajith et al. The NINJA-2 catalog of hybrid post-Newtonian/numerical-relativity waveforms for non-precessing black-hole binaries. *Class.Quant.Grav.*, 29:124001, 2012.
- [173] Ian Hinder et al. Error-analysis and comparison to analytical models of numerical waveforms produced by the NRAR Collaboration. *Class.Quant.Grav.*, 31:025012, 2013.
- [174] J. Abadie et al. Search for Gravitational Waves from Low Mass Compact Binary Coalescence in LIGO’s Sixth Science Run and Virgo’s Science Runs 2 and 3. *Phys.Rev.*, D85:082002, 2012.
- [175] Emily Baird, Stephen Fairhurst, Mark Hannam, and Patricia Murphy. Degeneracy between mass and spin in black-hole-binary waveforms. *Phys.Rev.*, D87:024035, 2013.
- [176] Mark Hannam, Duncan A. Brown, Stephen Fairhurst, Chris L. Fryer, and Ian W. Harry. When can gravitational-wave observations distinguish between black holes and neutron stars? *Astrophys.J.*, 766:L14, 2013.
- [177] Duncan A. Brown, Ian Harry, Andrew Lundgren, and Alexander H. Nitz. Detecting binary neutron star systems with spin in advanced gravitational-wave detectors. *Phys.Rev.*, D86:084017, 2012.
- [178] I.W. Harry et al. Investigating the effect of precession on searches for neutron-star-black-hole binaries with Advanced LIGO. 2013.
- [179] Frank Ohme. Analytical meets numerical relativity - status of complete gravitational waveform models for binary black holes. *Class.Quant.Grav.*, 29:124002, 2012.
- [180] Wolfgang Tichy and Pedro Marronetti. A Simple method to set up low eccentricity initial data for moving puncture simulations. *Phys.Rev.*, D83:024012, 2011.

- [181] Bernd Bruegmann et al. Calibration of Moving Puncture Simulations. *Phys.Rev.*, D77:024027, 2008.
- [182] Pedro Marronetti et al. Binary black holes on a budget: Simulations using workstations. *Class.Quant.Grav.*, 24:S43–S58, 2007.
- [183] Bernd Bruegmann, Wolfgang Tichy, and Nina Jansen. Numerical simulation of orbiting black holes. *Phys.Rev.Lett.*, 92:211101, 2004.
- [184] James Healy, Frank Herrmann, Ian Hinder, Deirdre M. Shoemaker, Pablo Laguna, et al. Superkicks in Hyperbolic Encounters of Binary Black Holes. *Phys.Rev.Lett.*, 102:041101, 2009.
- [185] James Healy, Pablo Laguna, Richard A. Matzner, and Deirdre M. Shoemaker. Final Mass and Spin of Merged Black Holes and the Golden Black Hole. *Phys.Rev.*, D81:081501, 2010.
- [186] Tanja Bode, Roland Haas, Tamara Bogdanovic, Pablo Laguna, and Deirdre Shoemaker. Relativistic Mergers of Supermassive Black Holes and their Electromagnetic Signatures. *Astrophys.J.*, 715:1117–1131, 2010.
- [187] Frank Herrmann, Ian Hinder, Deirdre M. Shoemaker, Pablo Laguna, and Richard A. Matzner. Binary Black Holes: Spin Dynamics and Gravitational Recoil. *Phys.Rev.*, D76:084032, 2007.
- [188] James Healy, Janna Levin, and Deirdre Shoemaker. Zoom-Whirl Orbits in Black Hole Binaries. *Phys.Rev.Lett.*, 103:131101, 2009.
- [189] Tanja Bode et al. Mergers of Supermassive Black Holes in Astrophysical Environments. *Astrophys.J.*, 744:45, 2012.
- [190] Ian Hinder, Birjoo Vaishnav, Frank Herrmann, Deirdre Shoemaker, and Pablo Laguna. Universality and final spin in eccentric binary black hole inspirals. *Phys.Rev.*, D77:081502, 2008.
- [191] Sascha Husa, Jose A. Gonzalez, Mark Hannam, Bernd Bruegmann, and Ulrich Sperhake. Reducing phase error in long numerical binary black hole evolutions with sixth order finite differencing. *Class.Quant.Grav.*, 25:105006, 2008.

- [192] Mark Hannam, Sascha Husa, Bernd Bruegmann, and Achamveedu Gopakumar. Comparison between numerical-relativity and post-Newtonian waveforms from spinning binaries: The Orbital hang-up case. *Phys.Rev.*, D78:104007, 2008.
- [193] Mark Hannam, Sascha Husa, Frank Ohme, Doreen Muller, and Bernd Bruegmann. Simulations of black-hole binaries with unequal masses or nonprecessing spins: Accuracy, physical properties, and comparison with post-Newtonian results. *Phys.Rev.*, D82:124008, 2010.
- [194] Mark Hannam, Sascha Husa, Ulrich Sperhake, Bernd Bruegmann, and Jose A. Gonzalez. Where post-Newtonian and numerical-relativity waveforms meet. *Phys.Rev.*, D77:044020, 2008.
- [195] Ulrich Sperhake. Binary black-hole evolutions of excision and puncture data. *Phys.Rev.*, D76:104015, 2007.
- [196] Ulrich Sperhake et al. Eccentric binary black-hole mergers: The Transition from inspiral to plunge in general relativity. *Phys.Rev.*, D78:064069, 2008.
- [197] Denis Pollney and Christian Reisswig. Gravitational memory in binary black hole mergers. *Astrophys.J.*, 732:L13, 2011.
- [198] C. Reisswig, N.T. Bishop, D. Pollney, and B. Szilagyi. Characteristic extraction in numerical relativity: binary black hole merger waveforms at null infinity. *Class.Quant.Grav.*, 27:075014, 2010.
- [199] Denis Pollney, Christian Reisswig, Erik Schnetter, Nils Dorband, and Peter Diener. High accuracy binary black hole simulations with an extended wave zone. *Phys.Rev.*, D83:044045, 2011.
- [200] Carlos O. Lousto, Hiroyuki Nakano, Yosef Zlochower, and Manuela Campanelli. Intermediate Mass Ratio Black Hole Binaries: Numerical Relativity meets Perturbation Theory. *Phys.Rev.Lett.*, 104:211101, 2010.
- [201] Carlos O. Lousto, Hiroyuki Nakano, Yosef Zlochower, and Manuela Campanelli. Intermediate-mass-ratio black hole binaries: Intertwining numerical and perturbative techniques. *Phys.Rev.*, D82:104057, 2010.

- [202] Hiroyuki Nakano, Yosef Zlochower, Carlos O. Lousto, and Manuela Campanelli. Intermediate-mass-ratio black hole binaries II: Modeling Trajectories and Gravitational Waveforms. *Phys.Rev.*, D84:124006, 2011.
- [203] Harald P. Pfeiffer, Lawrence E. Kidder, Mark A. Scheel, and Saul A. Teukolsky. A Multidomain spectral method for solving elliptic equations. *Comput.Phys.Commun.*, 152:253–273, 2003.
- [204] Mark A. Scheel et al. Solving Einstein’s equations with dual coordinate frames. *Phys.Rev.*, D74:104006, 2006.
- [205] Geoffrey Lovelace, Michael Boyle, Mark A. Scheel, and Bela Szilagyi. Accurate gravitational waveforms for binary-black-hole mergers with nearly extremal spins. *Class.Quant.Grav.*, 29:045003, 2012.
- [206] Bela Szilagyi, Lee Lindblom, and Mark A. Scheel. Simulations of Binary Black Hole Mergers Using Spectral Methods. *Phys.Rev.*, D80:124010, 2009.
- [207] Geoffrey Lovelace, Mark.A. Scheel, and Bela Szilagyi. Simulating merging binary black holes with nearly extremal spins. *Phys.Rev.*, D83:024010, 2011.
- [208] Spectral Einstein Code.
- [209] Michael Boyle et al. High-accuracy comparison of numerical relativity simulations with post-Newtonian expansions. *Phys.Rev.*, D76:124038, 2007.
- [210] Michael Boyle and Abdul H. Mroue. Extrapolating gravitational-wave data from numerical simulations. *Phys.Rev.*, D80:124045, 2009.
- [211] Zachariah B. Etienne, Yuk Tung Liu, Stuart L. Shapiro, and Thomas W. Baumgarte. General relativistic simulations of black-hole-neutron-star mergers: Effects of black-hole spin. *Phys.Rev.*, D79:044024, 2009.
- [212] Manuela Campanelli, Carlos O. Lousto, Hiroyuki Nakano, and Yosef Zlochower. Comparison of Numerical and Post-Newtonian Waveforms for Generic Precessing Black-Hole Binaries. *Phys.Rev.*, D79:084010, 2009.
- [213] J Aasi et al. *Possible scenarios for Commissioning and Early observing with the Second Generation Detectors*, LIGO Tech. Rep. LIGO-G1000176, 2010.

- [214] LSC Algorithm Library for GStreamer.
- [215] J. Aasi et al. The characterization of Virgo data and its impact on gravitational-wave searches. *Class.Quant.Grav.*, 29:155002, 2012.
- [216] J. Aasi et al. *Characterizing the LIGO instruments in their sixth science run*, in preparation, 2014.
- [217] P.C. Peters and J. Mathews. Gravitational radiation from point masses in a Keplerian orbit. *Phys.Rev.*, 131:435–439, 1963.
- [218] B. Abbott et al. Search for gravitational-wave bursts in LIGO data from the fourth science run. *Class.Quant.Grav.*, 24:5343–5370, 2007.
- [219] J. Abadie et al. Search for Gravitational Waves from Intermediate Mass Binary Black Holes. *Phys.Rev.*, D85:102004, 2012.
- [220] C. Pankow et al. A Burst search for gravitational waves from binary black holes. *Class.Quant.Grav.*, 26:204004, 2009.
- [221] B. Abbott et al. Search for gravitational waves from binary inspirals in S3 and S4 LIGO data. *Phys.Rev.*, D77:062002, 2008.
- [222] Stanislav Babak et al. The Mock LISA Data Challenges: From Challenge 1B to Challenge 3. *Class.Quant.Grav.*, 25:184026, 2008.
- [223] S. Babak, R. Balasubramanian, D. Churches, T. Cokelaer, and B.S. Sathyaprakash. A Template bank to search for gravitational waves from inspiralling compact binaries. I. Physical models. *Class.Quant.Grav.*, 23:5477–5504, 2006.
- [224] C.A.K. Robinson, B.S. Sathyaprakash, and Anand S. Sengupta. A Geometric algorithm for efficient coincident detection of gravitational waves. *Phys.Rev.*, D78:062002, 2008.
- [225] Bruce Allen. A chi**2 time-frequency discriminator for gravitational wave detection. *Phys.Rev.*, D71:062001, 2005.

- [226] Luc Blanchet, Bala R. Iyer, and Benoit Joguet. Gravitational waves from inspiralling compact binaries: Energy flux to third postNewtonian order. *Phys.Rev.*, D65:064005, 2002.
- [227] Drew Keppel. *Signatures and dynamics of compact binary coalescences and a search in LIGO's S5 data*. PhD thesis, California Institute of Technology, 2009.
- [228] Marc van der Sluys et al. Parameter estimation for signals from compact binary inspirals injected into LIGO data. *Class.Quant.Grav.*, 26:204010, 2009.
- [229] V. Raymond et al. Degeneracies in Sky Localisation Determination from a Spinning Coalescing Binary through Gravitational Wave Observations: A Markov-Chain Monte-Carlo Analysis for two Detectors. *Class.Quant.Grav.*, 26:114007, 2009.
- [230] J. Veitch and A. Vecchio. Bayesian coherent analysis of in-spiral gravitational wave signals with a detector network. *Phys.Rev.*, D81:062003, 2010.
- [231] V. Raymond et al. The Effects of LIGO detector noise on a 15-dimensional Markov-chain Monte-Carlo analysis of gravitational-wave signals. *Class.Quant.Grav.*, 27:114009, 2010.
- [232] Philippe Canitrot. Systematic errors for matched filtering of gravitational waves from inspiraling compact binaries. *Phys.Rev.*, D63:082005, 2001.
- [233] Curt Cutler and Michele Vallisneri. LISA detections of massive black hole inspirals: Parameter extraction errors due to inaccurate template waveforms. *Phys.Rev.*, D76:104018, 2007.
- [234] M. V. van der Sluys et al. Gravitational-Wave Astronomy with Inspiral Signals of Spinning Compact-Object Binaries. *Astrophys. J.*, 688:L61–L65, December 2008.
- [235] Marc van der Sluys et al. Parameter estimation of spinning binary inspirals using Markov-chain Monte Carlo. *Class.Quant.Grav.*, 25:184011, 2008.
- [236] F. Feroz, M.P. Hobson, and M. Bridges. MultiNest: an efficient and robust Bayesian inference tool for cosmology and particle physics. *Mon.Not.Roy.Astron.Soc.*, 398:1601–1614, 2009.

- [237] Tyson B. Littenberg, John G. Baker, Alessandra Buonanno, and Bernard J. Kelly. Systematic biases in parameter estimation of binary black-hole mergers. *Phys.Rev.*, D87:104003, 2013.
- [238] J. Aasi et al. Parameter estimation for compact binary coalescence signals with the first generation gravitational-wave detector network. *Phys.Rev.*, D88:062001, 2013.
- [239] Tomoki Isogai. Used percentage veto for LIGO and virgo binary inspiral searches. *J.Phys.Conf.Ser.*, 243:012005, 2010.
- [240] Manuela Campanelli, C.O. Lousto, and Y. Zlochower. Spinning-black-hole binaries: The orbital hang up. *Phys.Rev.*, D74:041501, 2006.
- [241] J. Veitch et al. Estimating parameters of coalescing compact binaries with proposed advanced detector networks. *Phys.Rev.*, D85:104045, 2012.
- [242] Samaya Nissanke, Jonathan Sievers, Neal Dalal, and Daniel Holz. Localizing compact binary inspirals on the sky using ground-based gravitational wave interferometers. *Astrophys.J.*, 739:99, 2011.
- [243] Stephen Fairhurst. Source localization with an advanced gravitational wave detector network. *Class.Quant.Grav.*, 28:105021, 2011.
- [244] David Eichler, Mario Livio, Tsvi Piran, and David N. Schramm. Nucleosynthesis, Neutrino Bursts and Gamma-Rays from Coalescing Neutron Stars. *Nature*, 340:126–128, 1989.
- [245] Ramesh Narayan, Bohdan Paczynski, and Tsvi Piran. Gamma-ray bursts as the death throes of massive binary stars. *Astrophys.J.*, 395:L83–L86, 1992.
- [246] Wen-Fai Fong and Edo Berger. The Locations of Short Gamma-ray Bursts as Evidence for Compact Object Binary Progenitors. *Astrophys.J.*, 776:18, 2013.
- [247] S. Rosswog, C. Freiburghaus, and F.K. Thielemann. Nucleosynthesis calculations for the ejecta of neutron star coalescences. *Nuclear Physics A*, 688:344–348, May 2001.

- [248] Wolfgang Tichy and Pedro Marronetti. Binary black hole mergers: Large kicks for generic spin orientations. *Phys.Rev.*, D76:061502, 2007.
- [249] Wolfgang Tichy and Pedro Marronetti. The Final mass and spin of black hole mergers. *Phys.Rev.*, D78:081501, 2008.
- [250] Patricia Schmidt, Mark Hannam, Sascha Husa, and P. Ajith. Tracking the precession of compact binaries from their gravitational-wave signal. *Phys.Rev.*, D84:024046, 2011.
- [251] R. O’Shaughnessy, B. Vaishnav, J. Healy, Z. Meeks, and D. Shoemaker. Efficient asymptotic frame selection for binary black hole spacetimes using asymptotic radiation. *Phys.Rev.*, D84:124002, 2011.
- [252] R. O’Shaughnessy, J. Healy, L. London, Z. Meeks, and D. Shoemaker. Is J enough? Comparison of gravitational waves emitted along the total angular momentum direction with other preferred orientations. *Phys.Rev.*, D85(6):084003, 2012.
- [253] R. O’Shaughnessy, L. London, J. Healy, and D. Shoemaker. Precession during merger: Strong polarization changes are observationally accessible features of strong-field gravity during binary black hole merger. *Phys.Rev.*, D87(4):044038, 2013.
- [254] James Healy, Pablo Laguna, Larne Pekowsky, and Deirdre Shoemaker. Template Mode Hierarchies for Binary Black Hole Mergers. *Phys.Rev.*, D88:024034, 2013.
- [255] Sean T. McWilliams, Bernard J. Kelly, and John G. Baker. Observing mergers of non-spinning black-hole binaries. *Phys.Rev.*, D82:024014, 2010.
- [256] Yi Pan et al. Inspiral-merger-ringdown waveforms of spinning, precessing black-hole binaries in the effective-one-body formalism. 2013.
- [257] Mark Hannam et al. Twist and shout: A simple model of complete precessing black-hole-binary gravitational waveforms. 2013.

- [258] Tyson B. Littenberg, Michael Coughlin, Benjamin Farr, and Will M. Farr. Fortifying the characterization of binary mergers in LIGO data. 2013.
- [259] P. A. M. Dirac. Classical Theory of Radiating Electrons. *Royal Society of London Proceedings Series A*, 167:148–169, August 1938.
- [260] B. S. DeWitt and R. W. Brehme. Radiation damping in a gravitational field. *Annals of Physics*, 9:220–259, February 1960.
- [261] T. C. Quinn and R. M. Wald. Axiomatic approach to electromagnetic and gravitational radiation reaction of particles in curved spacetime. *Phys. Rev. D*, 56:3381–3394, September 1997.
- [262] Y. Mino, M. Sasaki, and T. Tanaka. *Gravitational radiation reaction to a particle motion*. *Phys. Rev. D*, 55:3457–3476, March 1997.
- [263] P. Canizares and C. F. Sopuerta. Tuning time-domain pseudospectral computations of the self-force on a charged scalar particle. *Classical and Quantum Gravity*, 28(13):134011, July 2011.
- [264] Niels Warburton, Sarp Akcay, Leor Barack, Jonathan R. Gair, and Norichika Sago. Evolution of inspiral orbits around a Schwarzschild black hole. *Phys. Rev.*, D85:061501, 2012.
- [265] N. Warburton and L. Barack. Self force on a scalar charge in Kerr spacetime: eccentric equatorial orbits. *ArXiv e-prints*, March 2011.
- [266] N. Warburton and L. Barack. Self-force on a scalar charge in Kerr spacetime: Circular equatorial orbits. *Phys. Rev. D*, 81(8):084039–+, April 2010.
- [267] E.A. Huerta and Jonathan R Gair. Importance of including small body spin effects in the modelling of extreme and intermediate mass-ratio inspirals. *Phys. Rev.*, D84:064023, 2011.
- [268] E.A. Huerta, Jonathan R. Gair, and Duncan A. Brown. Importance of including small body spin effects in the modelling of intermediate mass-ratio inspirals. II Accurate parameter extraction of strong sources using higher-order spin effects. *Phys. Rev.*, D85:064023, 2012.

- [269] G. M. Harry and LIGO Scientific Collaboration. Advanced LIGO: the next generation of gravitational wave detectors. *Class. Quantum Grav.*, 27(8):084006, April 2010.
- [270] D. A. Brown, J. Brink, H. Fang, J. R. Gair, C. Li, G. Lovelace, I. Mandel, and K. S. Thorne. Prospects for Detection of Gravitational Waves from Intermediate-Mass-Ratio Inspirals. *Physical Review Letters*, 99(20):201102–+, November 2007.
- [271] M. C. Miller and E. J. M. Colbert. Intermediate-Mass Black Holes. *International Journal of Modern Physics D*, 13:1–64, January 2004.
- [272] L. Barack and N. Sago. Gravitational Self-Force Correction to the Innermost Stable Circular Orbit of a Schwarzschild Black Hole. *Phys. Rev. Lett.*, 102(19):191101–+, May 2009.
- [273] Alessandra Buonanno and Thibault Damour. Transition from inspiral to plunge in binary black hole coalescences. *Phys. Rev. D*, 62:064015, Aug 2000.
- [274] T. Damour, B. R. Iyer, and A. Nagar. Improved resummation of post-Newtonian multipolar waveforms from circularized compact binaries. *Phys. Rev. D*, 79(6):064004–+, March 2009.
- [275] T. Damour and A. Nagar. The Effective One Body description of the Two-Body problem. *ArXiv e-prints*, June 2009.
- [276] A. Taracchini, Y. Pan, A. Buonanno, E. Barausse, M. Boyle, T. Chu, G. Lovelace, H. P. Pfeiffer, and M. A. Scheel. A prototype effective-one-body model for non-precessing spinning inspiral-merger-ringdown waveforms. *ArXiv e-prints*, February 2012.
- [277] N. Sago, L. Barack, and S. Detweiler. Two approaches for the gravitational self-force in black hole spacetime: Comparison of numerical results. *Phys. Rev. D*, 78(12):124024–+, December 2008.
- [278] L. Barack and N. Sago. *Gravitational self-force on a particle in circular orbit around a Schwarzschild black hole*. *Phys. Rev. D*, 75(6):064021–+, March 2007.

- [279] Thibault Damour. Gravitational Self Force in a Schwarzschild Background and the Effective One Body Formalism. *Phys.Rev.*, D81:024017, 2010.
- [280] Enrico Barausse, Alessandra Buonanno, and Alexandre Le Tiec. The complete non-spinning effective-one-body metric at linear order in the mass ratio. *Phys.Rev.*, D85:064010, 2012.
- [281] Sebastiano Bernuzzi and Alessandro Nagar. Binary black hole merger in the extreme-mass-ratio limit: a multipolar analysis. *Phys.Rev.*, D81:084056, 2010.
- [282] J. R. Gair and K. Glampedakis. *Improved approximate inspirals of test bodies into Kerr black holes*. *Phys. Rev. D*, 73(6):064037–+, March 2006.
- [283] Yi Pan, Alessandra Buonanno, Ryuichi Fujita, Etienne Racine, and Hideyuki Tagoshi. Post-Newtonian factorized multipolar waveforms for spinning, non-precessing black-hole binaries. *Phys.Rev.*, D83:064003, 2011.
- [284] N. Yunes, A. Buonanno, S. A. Hughes, Y. Pan, E. Barausse, M. C. Miller, and W. Throwe. Extreme Mass-Ratio Inspirals in the Effective-One-Body Approach: Quasi-Circular, Equatorial Orbits around a Spinning Black Hole. *ArXiv e-prints*, September 2010.
- [285] Nicolas Yunes, Alessandra Buonanno, Scott A. Hughes, Yi Pan, Enrico Barausse, et al. Extreme Mass-Ratio Inspirals in the Effective-One-Body Approach: Quasi-Circular, Equatorial Orbits around a Spinning Black Hole. *Phys.Rev.*, D83:044044, 2011.
- [286] C. O. Lousto and Y. Zlochower. Orbital Evolution of Extreme-Mass-Ratio Black-Hole Binaries with Numerical Relativity. *Physical Review Letters*, 106(4):041101, January 2011.
- [287] L. Barack. TOPICAL REVIEW: Gravitational self-force in extreme mass-ratio inspirals. *Class. Quantum Grav.*, 26(21):213001–+, November 2009.
- [288] A. Ori and K. S. Thorne. Transition from inspiral to plunge for a compact body in a circular equatorial orbit around a massive, spinning black hole. *Phys. Rev. D*, 62(12):124022–+, December 2000.

- [289] L. Barack and C. Cutler. *LISA capture sources: Approximate waveforms, signal-to-noise ratios, and parameter estimation accuracy.* *Phys. Rev. D*, 69(8):082005–+, April 2004.
- [290] J. R. Gair, L. Barack, T. Creighton, C. Cutler, S. L. Larson, E. S. Phinney, and M. Vallisneri. *Event rate estimates for LISA extreme mass ratio capture sources.* *Class. Quantum Grav.*, 21:1595–+, October 2004.
- [291] S. Babak, H. Fang, J. R. Gair, K. Glampedakis, and S. A. Hughes. ‘‘*Kludge*’’ gravitational waveforms for a test-body orbiting a Kerr black hole. *Phys. Rev. D*, 75(2):024005–+, January 2007.
- [292] L. Barack and C. Cutler. Confusion noise from LISA capture sources. *Phys. Rev. D*, 70(12):122002–+, December 2004.
- [293] P. Amaro-Seoane, J. R. Gair, M. Freitag, M. C. Miller, I. Mandel, C. J. Cutler, and S. Babak. *TOPICAL REVIEW: Intermediate and extreme mass-ratio inspirals—astrophysics, science applications and detection using LISA.* *Class. Quantum Grav.*, 24:113–+, September 2007.
- [294] M. Favata. Conservative self-force correction to the innermost stable circular orbit: Comparison with multiple post-Newtonian-based methods. *Phys. Rev. D*, 83(2):024027–+, January 2011.
- [295] Ulrich Sperhake, Vitor Cardoso, Christian D. Ott, Erik Schnetter, and Helvi Witek. Extreme black hole simulations: collisions of unequal mass black holes and the point particle limit. *Phys. Rev.*, D84:084038, 2011.
- [296] Carlos O. Lousto, Hiroyuki Nakano, Yosef Zlochower, and Manuela Campanelli. Intermediate-mass-ratio black hole binaries: Intertwining numerical and perturbative techniques. *Phys. Rev.*, D82:104057, 2010.
- [297] Hiroyuki Nakano, Yosef Zlochower, Carlos O. Lousto, and Manuela Campanelli. Intermediate-mass-ratio black hole binaries II: Modeling Trajectories and Gravitational Waveforms. *Phys. Rev.*, D84:124006, 2011.
- [298] D. Kennefick and A. Ori. *Radiation-reaction-induced evolution of circular orbits of particles around Kerr black holes.* *Phys. Rev. D*, 53:4319–4326, April 1996.

- [299] J. E. McClintock and R. A. Remillard. *Black hole binaries*, pages 157–213. April 2006.
- [300] M. Morscher, S. Umbreit, W. M. Farr, and F. A. Rasio. Retention of Stellar-mass Black Holes in Globular Clusters. *Astrophys. J. Lett.*, 763:L15, January 2013.
- [301] A. Merloni and S. Heinz. A synthesis model for AGN evolution: supermassive black holes growth and feedback modes. *MNRAS*, 388:1011–1030, August 2008.
- [302] M. Fukugita and P. J. E. Peebles. The Cosmic Energy Inventory. *Astrophys. J.*, 616:643–668, December 2004.
- [303] J. M. Miller, A. Zezas, G. Fabbiano, and F. Schweizer. XMM-Newton Spectroscopy of Four Bright Ultraluminous X-Ray Sources in the Antennae Galaxies (NGC 4038/4039). *Astrophys. J.*, 609:728–734, July 2004.
- [304] J. M. Miller, A. C. Fabian, and M. C. Miller. Revealing a Cool Accretion Disk in the Ultraluminous X-Ray Source M81 X-9 (Holmberg IX X-1): Evidence for an Intermediate-Mass Black Hole. *Astrophys. J.*, 607:931–938, June 2004.
- [305] J. M. Miller. *Present evidence for intermediate mass black holes in ULXs and future prospects*, pages 227–238. January 2005.
- [306] D. A. Swartz, R. Soria, A. F. Tennant, and M. Yukita. A Complete Sample of Ultraluminous X-ray Source Host Galaxies. *Astrophys. J.*, 741:49, November 2011.
- [307] L. Zampieri and T. P. Roberts. Low-metallicity natal environments and black hole masses in ultraluminous X-ray sources. *MNRAS*, 400:677–686, December 2009.
- [308] K. Belczynski, T. Bulik, C. L. Fryer, A. Ruiter, F. Valsecchi, J. S. Vink, and J. R. Hurley. On the Maximum Mass of Stellar Black Holes. *Astrophys. J.*, 714:1217–1226, May 2010.
- [309] K. Ohsuga and S. Mineshige. Global Structure of Three Distinct Accretion Flows and Outflows around Black Holes from Two-dimensional Radiation-magnetohydrodynamic Simulations. *Astrophys. J.*, 736:2, July 2011.

- [310] H. Matsumoto, T. G. Tsuru, K. Koyama, H. Awaki, C. R. Canizares, N. Kawai, S. Matsushita, and R. Kawabe. Discovery of a Luminous, Variable, Off-Center Source in the Nucleus of M82 with the Chandra High-Resolution Camera. *Astrophys. J. Lett.*, 547:L25–L28, January 2001.
- [311] Sean Farrell, Natalie Webb, Didier Barret, Olivier Godet, and Joana Rodrigues. An Intermediate-mass Black Hole of Over 500 Solar Masses in the Galaxy ESO 243-49. *Nature*, 460:73–75, 2009.
- [312] F. P. A. Wolter Ginevra Trinchieri. Chandra observations of the ULX N10 in the Cartwheel galaxy. *ArXiv e-prints*, March 2010.
- [313] P. G. Jonker, M. A. P. Torres, A. C. Fabian, M. Heida, G. Miniutti, and D. Pooley. A bright off-nuclear X-ray source: a type IIn supernova, a bright ULX or a recoiling supermassive black hole in CXOJ122518.6+144545. *MNRAS*, 407:645–650, September 2010.
- [314] Simon F. Portegies Zwart, Holger Baumgardt, Piet Hut, Junichiro Makino, and Stephen L.W. McMillan. The Formation of massive black holes through collision runaway in dense young star clusters. *Nature*, 428:724, 2004.
- [315] T. E. Strohmayer and R. F. Mushotzky. Discovery of X-Ray Quasi-periodic Oscillations from an Ultraluminous X-Ray Source in M82: Evidence against Beaming. *Astrophys. J. Lett.*, 586:L61–L64, March 2003.
- [316] P. Kaaret and H. Feng. Confirmation of the 62 Day X-Ray Periodicity from M82. *Astrophys. J.*, 669:106–108, November 2007.
- [317] H. Feng and P. Kaaret. Identification of the X-ray Thermal Dominant State in an Ultraluminous X-ray Source in M82. *Astrophys. J. Lett.*, 712:L169–L173, April 2010.
- [318] A. D. Sutton, T. P. Roberts, D. J. Walton, J. C. Gladstone, and A. E. Scott. The most extreme ultraluminous X-ray sources: evidence for intermediate-mass black holes? *MNRAS*, 423:1154–1177, June 2012.
- [319] M. Trenti. Dynamical evidence for intermediate mass black holes in old globular clusters. *ArXiv Astrophysics e-prints*, December 2006.

- [320] M. Coleman Miller and E. J. M. Colbert. Intermediate-Mass Black Holes. *International Journal of Modern Physics D*, 13:1–64, 2004.
- [321] M. Volonteri. Formation of supermassive black holes. *A&ARv*, 18:279–315, July 2010.
- [322] R. Schneider, A. Ferrara, P. Natarajan, and K. Omukai. First Stars, Very Massive Black Holes, and Metals. *Astrophys. J.*, 571:30–39, May 2002.
- [323] Q. Yu and S. Tremaine. Observational constraints on growth of massive black holes. *MNRAS*, 335:965–976, October 2002.
- [324] I. Mandel. Spin distribution following minor mergers and the effect of spin on the detection range for low-mass-ratio inspirals. *ArXiv e-prints*, July 2007.
- [325] F. Acernese et al. Virgo status. *Class. Quantum Grav.*, 25(18):184001–+, September 2008.
- [326] K. Somiya and the KAGRA Collaboration. Detector configuration of KAGRA—the Japanese cryogenic gravitational-wave detector. *Class. Quantum Grav.*, 29(12):124007, June 2012.
- [327] David Shoemaker. Advanced LIGO anticipated sensitivity curves, 2010. <https://dcc.ligo.org/cgi-bin/DocDB>ShowDocument?docid=2974>.
- [328] A. Freise, S. Chelkowski, S. Hild, W. Del Pozzo, A. Perreca, and A. Vecchio. Triple Michelson interferometer for a third-generation gravitational wave detector. *Class. Quantum Grav.*, 26(8):085012–+, April 2009.
- [329] J. R. Gair, I. Mandel, M. C. Miller, and M. Volonteri. Exploring intermediate and massive black-hole binaries with the Einstein Telescope. *General Relativity and Gravitation*, 43:485–518, February 2011.
- [330] E. A. Huerta and J. R. Gair. Intermediate-mass-ratio inspirals in the Einstein Telescope. I. Signal-to-noise ratio calculations. *Phys. Rev. D*, 83(4):044020–+, February 2011.

- [331] E. A. Huerta and J. R. Gair. Intermediate-mass-ratio inspirals in the Einstein Telescope. II. Parameter estimation errors. *Phys. Rev. D*, 83(4):044021–+, February 2011.
- [332] Y. Taniguchi, Y. Shioya, T. G. Tsuru, and S. Ikeuchi. Formation of Intermediate-Mass Black Holes in Circumnuclear Regions of Galaxies. *PASJ*, 52:533–537, June 2000.
- [333] M. C. Miller and D. P. Hamilton. Four-Body Effects in Globular Cluster Black Hole Coalescence. *Astrophys. J.*, 576:894–898, September 2002.
- [334] H. Mouri and Y. Taniguchi. Runaway Merging of Black Holes: Analytical Constraint on the Timescale. *Astrophys. J. Lett*, 566:L17–L20, February 2002.
- [335] H. Mouri and Y. Taniguchi. Mass Segregation in Star Clusters: Analytic Estimation of the Timescale. *Astrophys. J.*, 580:844–849, December 2002.
- [336] K. Gültekin, M. C. Miller, and D. P. Hamilton. Growth of Intermediate-Mass Black Holes in Globular Clusters. *Astrophys. J.*, 616:221–230, November 2004.
- [337] K. Gültekin, M. C. Miller, and D. P. Hamilton. Three-Body Dynamics with Gravitational Wave Emission. *Astrophys. J.*, 640:156–166, March 2006.
- [338] R. M. O’Leary, F. A. Rasio, J. M. Fregeau, N. Ivanova, and R. O’Shaughnessy. Binary Mergers and Growth of Black Holes in Dense Star Clusters. *Astrophys. J.*, 637:937–951, February 2006.
- [339] R. M. O’Leary, R. O’Shaughnessy, and F. A. Rasio. Dynamical interactions and the black-hole merger rate of the Universe. *Phys. Rev. D*, 76(6):061504, September 2007.
- [340] I. Mandel, D. A. Brown, J. R. Gair, and M. C. Miller. Rates and Characteristics of Intermediate Mass Ratio Inspirals Detectable by Advanced LIGO. *Astrophys. J.*, 681:1431–1447, July 2008.
- [341] J. Abadie, B. P. Abbott, R. Abbott, M. Abernathy, T. Accadia, F. Acernese, C. Adams, R. Adhikari, P. Ajith, B. Allen, et al. TOPICAL REVIEW: Predictions for the rates of compact binary coalescences observable by ground-based

- gravitational-wave detectors. *Class. Quantum Grav.*, 27(17):173001, September 2010.
- [342] R. J. E. Smith, I. Mandel, and A. Vecchio. Studies of waveform requirements for intermediate mass-ratio coalescence searches with advanced gravitational-wave detectors. *Phys. Rev. D*, 88(4):044010, August 2013.
 - [343] E. A. Huerta and J. R. Gair. Influence of conservative corrections on parameter estimation for extreme-mass-ratio inspirals. *Phys. Rev. D*, 79(8):084021–+, April 2009.
 - [344] E. A. Huerta and J. R Gair. On the relevance of gravitational self-force corrections on parameter estimation errors for extreme-mass-ratio inspirals. *ArXiv e-prints*, September 2010.
 - [345] E. A. Huerta and J. R. Gair. Erratum: Influence of conservative corrections on parameter estimation for extreme-mass-ratio inspirals [Phys. Rev. D 79, 084021 (2009)]. *Phys. Rev. D*, 84(4):049903, August 2011.
 - [346] E. A. Huerta, P. Kumar, and D. A. Brown. Accurate modeling of intermediate-mass-ratio inspirals: Exploring the form of the self-force in the intermediate-mass-ratio regime. *Phys. Rev. D*, 86(2):024024, July 2012.
 - [347] E. Poisson, A. Pound, and I. Vega. The Motion of Point Particles in Curved Spacetime. *Living Reviews in Relativity*, 14:7, September 2011.
 - [348] T. Damour, A. Nagar, and S. Bernuzzi. Improved effective-one-body description of coalescing nonspinning black-hole binaries and its numerical-relativity completion. *Phys. Rev. D*, 87(8):084035, April 2013.
 - [349] A. Le Tiec, E. Barausse, and A. Buonanno. Gravitational Self-Force Correction to the Binding Energy of Compact Binary Systems. *Phys. Rev. Lett.*, 108(13):131103, March 2012.
 - [350] S. Akcay, L. Barack, T. Damour, and N. Sago. Gravitational self-force and the effective-one-body formalism between the innermost stable circular orbit and the light ring. *Phys. Rev. D*, 86(10):104041, November 2012.

- [351] LIGO Scientific Collaboration, Virgo Collaboration, J. Aasi, J. Abadie, B. P. Abbott, R. Abbott, T. D. Abbott, M. Abernathy, T. Accadia, F. Acernese, et al. Prospects for Localization of Gravitational Wave Transients by the Advanced LIGO and Advanced Virgo Observatories. *ArXiv e-prints*, April 2013.
- [352] A. Buonanno, B. R. Iyer, E. Ochsner, Y. Pan, and B. S. Sathyaprakash. Comparison of post-Newtonian templates for compact binary inspiral signals in gravitational-wave detectors. *Phys. Rev. D* , 80(8):084043, October 2009.
- [353] A. H. Nitz, A. Lundgren, D. A. Brown, E. Ochsner, et al. Accuracy of gravitational waveform models for observing neutron-star–black-hole binaries in Advanced LIGO. *ArXiv e-prints*, July 2013.
- [354] Alessandra Buonanno, Yan-bei Chen, and Michele Vallisneri. Detecting gravitational waves from precessing binaries of spinning compact objects: Adiabatic limit. *Phys.Rev.*, D67:104025, 2003.
- [355] Thibault Damour. Coalescence of two spinning black holes: An effective one-body approach. *Phys. Rev. D* , 64(12):124013, Nov 2001.
- [356] A. Buonanno and T. Damour. Effective one-body approach to general relativistic two-body dynamics. *Phys. Rev. D* , 59(8):084006, April 1999.
- [357] T. Damour, P. Jaranowski, and G. Schäfer. Determination of the last stable orbit for circular general relativistic binaries at the third post-Newtonian approximation. *Phys. Rev. D* , 62(8):084011, October 2000.
- [358] S. Detweiler and B. F. Whiting. Self-force via a Green’s function decomposition. *Phys. Rev. D* , 67(2):024025, January 2003.
- [359] S. Detweiler. Consequence of the gravitational self-force for circular orbits of the Schwarzschild geometry. *Phys. Rev. D* , 77(12):124026, June 2008.
- [360] S. Detweiler. Elementary development of the gravitational self-force. *ArXiv e-prints*, August 2009.
- [361] J. G. Baker, W. D. Boggs, J. Centrella, B. J. Kelly, S. T. McWilliams, and J. R. van Meter. Mergers of nonspinning black-hole binaries: Gravitational radiation characteristics. *Phys. Rev. D* , 78(4):044046, August 2008.

- [362] S. Isoyama, R. Fujita, N. Sago, H. Tagoshi, and T. Tanaka. Impact of the second-order self-forces on the dephasing of the gravitational waves from quasi-circular extreme mass-ratio inspirals. *Phys. Rev. D*, 87(2):024010, January 2013.
- [363] L. M. Burko and G. Khanna. Self-force gravitational waveforms for extreme and intermediate mass ratio inspirals. II. Importance of the second-order dissipative effect. *Phys. Rev. D*, 88(2):024002, July 2013.
- [364] R. Fujita. Gravitational Waves from a Particle in Circular Orbits around a Schwarzschild Black Hole to the 22nd Post-Newtonian Order. *Progress of Theoretical Physics*, 128:971–992, November 2012.
- [365] L. Blanchet, G. Faye, B. R. Iyer, and B. Joguet. Gravitational-wave inspiral of compact binary systems to $7/2$ post-Newtonian order. *Phys. Rev. D*, 65(6):061501, March 2002.
- [366] L. Blanchet, A. Buonanno, and G. Faye. Tail-induced spin-orbit effect in the gravitational radiation of compact binaries. *Phys. Rev. D*, 84(6):064041, September 2011.
- [367] Y. Pan, A. Buonanno, A. Taracchini, M. Boyle, L. E. Kidder, A. H. Mroue, H. P. Pfeiffer, M. A. Scheel, B. Szilagyi, and A. Zenginoglu. Stability of non-spinning effective-one-body model in approximating two-body dynamics and gravitational-wave emission. *ArXiv e-prints*, November 2013.
- [368] J. R. Gair, M. Vallisneri, S. L. Larson, and J. G. Baker. Testing General Relativity with Low-Frequency, Space-Based Gravitational-Wave Detectors. *Living Reviews in Relativity*, 16:7, September 2013.
- [369] A. Gopakumar and B. R. Iyer. Second post-Newtonian gravitational wave polarizations for compact binaries in elliptical orbits. *Phys. Rev. D*, 65(8):084011, April 2002.
- [370] C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. San Francisco: W.H. Freeman and Co., 1973, 1973.

- [371] E. Berti, V. Cardoso, and M. Casals. Eigenvalues and eigenfunctions of spin-weighted spheroidal harmonics in four and higher dimensions. *Phys. Rev. D*, 73(2):024013, January 2006.
- [372] E. Berti, V. Cardoso, and C. M. Will. Gravitational-wave spectroscopy of massive black holes with the space interferometer LISA. *Phys. Rev. D*, 73(6):064030–+, March 2006.
- [373] C.V. Vishveshwara. Scattering of Gravitational Radiation by a Schwarzschild Black-hole. *Nature*, 227:936–938, 1970.
- [374] E. Barausse, V. Morozova, and L. Rezzolla. On the Mass Radiated by Coalescing Black Hole Binaries. *Astrophys. J.*, 758:63, October 2012.
- [375] L. Rezzolla, P. Diener, E. N. Dorband, D. Pollney, C. Reisswig, E. Schnetter, and J. Seiler. The Final Spin from the Coalescence of Aligned-Spin Black Hole Binaries. *Astrophys. J. Lett*, 674:L29–L32, February 2008.
- [376] A. Buonanno, Y. Pan, J. G. Baker, J. Centrella, B. J. Kelly, S. T. McWilliams, and J. R. van Meter. Approaching faithful templates for nonspinning binary black holes using the effective-one-body approach. *Phys. Rev. D*, 76(10):104049, November 2007.
- [377] E. Barausse and L. Rezzolla. Predicting the Direction of the Final Spin from the Coalescence of Two Black Holes. *Astrophys. J. Lett*, 704:L40–L44, October 2009.
- [378] L. Rezzolla, E. Barausse, E. N. Dorband, D. Pollney, C. Reisswig, J. Seiler, and S. Husa. Final spin from the coalescence of two black holes. *Phys. Rev. D*, 78(4):044002, August 2008.
- [379] D. A. Brown, P. Kumar, and A. H. Nitz. Template banks to search for low-mass binary black holes in advanced gravitational-wave detectors. *Phys. Rev. D*, 87(8):082004, April 2013.

- [380] F. Foucart, M. B. Deaton, M. D. Duez, L. E. Kidder, I. MacDonald, C. D. Ott, H. P. Pfeiffer, M. A. Scheel, B. Szilagyi, and S. A. Teukolsky. Black-hole-neutron-star mergers at realistic mass ratios: Equation of state and spin orientation effects. *Phys. Rev. D*, 87(8):084006, April 2013.
- [381] M. Saijo, K.-I. Maeda, M. Shibata, and Y. Mino. Gravitational waves from a spinning particle plunging into a Kerr black hole. *Phys. Rev. D*, 58(6):064005–+, September 1998.
- [382] L. M. Burko. Orbital evolution of a test particle around a black hole. II. Comparison of contributions of spin-orbit coupling and the self-force. *Phys. Rev. D*, 69(4):044011–+, February 2004.
- [383] Luc Blanchet, Alessandra Buonanno, and Guillaume Faye. Erratum: Higher-order spin effects in the dynamics of compact binaries. ii. radiation field [phys. rev. d 74, 104034 (2006)]. *Phys. Rev. D*, 75:049903, Feb 2007.
- [384] L. Blanchet, A. Buonanno, and G. Faye. Higher-order spin effects in the dynamics of compact binaries. II. Radiation field. *Phys. Rev. D*, 74(10):104034–+, November 2006.
- [385] R. M. O’Leary, B. Kocsis, and A. Loeb. Gravitational waves from scattering of stellar-mass black holes in galactic nuclei. *MNRAS*, 395:2127–2146, June 2009.
- [386] E. A. Huerta and D. A. Brown. Effect of eccentricity on binary neutron star searches in advanced LIGO. *Phys. Rev. D*, 87(12):127501, June 2013.
- [387] H. Yang, F. Zhang, A. Zimmerman, and Y. Chen. The scalar Green function of the Kerr spacetime. *ArXiv e-prints*, November 2013.
- [388] S. R. Dolan, L. Barack, and B. Wardell. Self-force via m-mode regularization and 2+1D evolution. II. Scalar-field implementation on Kerr spacetime. *Phys. Rev. D*, 84(8):084001, October 2011.
- [389] A. Pound, C. Merlin, and L. Barack. Gravitational self-force from radiation-gauge metric perturbations. *ArXiv e-prints*, October 2013.

- [390] F. Zhang and B. Szilágyi. Joint approach for reducing eccentricity and spurious gravitational radiation in binary black hole initial data construction. *Phys. Rev. D*, 88(8):084033, October 2013.