

Multiple Constraints and Non-regular Solution in Deep Declarative Network

Suikei Wang

A thesis submitted for the degree of
Master of Machine Learning and Computer Vision
The Australian National University

October 2020

© Suikei Wang 2020

Except where otherwise indicated, this thesis is my own original work.

Suikei Wang
30 October 2020

to my parents, for their unconditional love

Acknowledgments

The past two years at the Australian National University have been an invaluable experience for me. When I started my Master of Machine Learning and Computer Vision at the beginning of 2019, I could barely understand lectures, knew little about the country, and had never heard of the term "convex optimization". It is unbelievable that I have been doing a research project on this topic for a whole year. ANU has the top tier research group in this realm and how honored I am to be a postgraduate student here.

First and foremost, I would like to express my deep and sincere gratitude to my supervisor, Stephen Gould. I knew Stephen before the admission of the program and how privileged I am to work and learn with one of the most brilliant minds in our field. He always has an insightful and high-level view on this topic. More importantly, Stephen is extremely kind and patient as he is always willing to discuss and share ideas with us during the weekly meetings. Although sometimes I am stuck in some problems he used to give me constructive suggestions.

I would like to thank Dylan Campbell and Miaomiao Liu – another two giants of the Australian Centre for Robotic Vision – for offering my supervision and being my second examiner on my thesis. Dylan is always enthusiasm and knowledgeable, and his ideas and works inspire me a lot in my research. Although I did not work with Miaomiao before, her works in the computer vision area are very impressive, which make me work hardly to complete this research. I want to thank her for her time examining my thesis seminar.

During my research, I have very fun collaboration and discussion with collaborators in our research group: Jiteng Ma, Jingyi Zhang, Yuchong Yao, Ruikai Cui, and Wenbo Du. Although we almost never meet in person due to the COVID-19, we still share many ideas related to our research through the social network. I greatly enjoyed the teamwork with them.

Outside the research group, I am very lucky to be surrounded by many great friends. Yao Liu, Ce Wang and Nuocheng Tian – We almost study together every single day during the academic year, even in the public holiday. I am always impressed by their insightful ideas and I really appreciate the brainstorm with them. Mingrui Zhao – As my closest partner, a charming Engineering & Mathematics double-degree undergraduate student and my math "teacher", he always encourages me to pursue my academic ambition and I learned a lot from him.

I thank my parents: Shiwu Wang and Hongyin Shen. As traditional Asian parents, they never restrict my life planning, and always encourage me to fulfill my dream. I hope that they will a little proud of me.

Abstract

Put your abstract here.

Contents

Acknowledgments	vii
Abstract	ix
1 Introduction	1
1.1 Motivation	1
1.2 Thesis Outline	1
1.3 Contribution	2
I Deep Declarative Network: Multiple Constrained Declarative Nodes	3
2 An Overview of Numerical Optimization	5
2.1 Theory of Optimization	5
2.1.1 Existence of Optimizers	5
2.1.2 Optimality Conditions for Unconstrained Problems	7
2.1.3 Optimality Conditions for Constrained Problems	9
2.2 Solution of Unconstrained and Constrained Optimization Problems . .	13
2.2.1 Unconstrained Optimization	13
2.2.2 Equality Constrained Optimization	15
2.2.3 Inequality Constrained Optimization	17
2.3 Differentiable Neural Network	18
2.4 Summary	19
3 Deep Declarative Network	21
3.1 An Overview of Deep Declarative Network	21
3.1.1 Declarative Node	21
3.1.2 Learning	22
3.2 Back-propagation Through Declarative Nodes	23
3.2.1 Unconstrained	23
3.2.2 Equality Constrained	24
3.2.3 Inequality Constrained	26
3.3 Examples of Declarative Nodes	27
3.3.1 Implementation Details	27
3.3.2 Equality Constrained	28
3.3.3 Inequality Constrained	29
3.4 Future Work of the Deep Declarative Network	30

3.5	Summary	31
-----	-------------------	----

II	Deep Declarative Nodes: Non-regular Solution	33
4	An Overview of Regular and Non-regular Solution	35
5	Solutions of Non-regular Point	37
5.1	Overdetermined System	37
5.1.1	Least-Squared Method	37
5.1.2	Conjugate Gradient and Preconditioning	38
5.2	Rank Deficiency	38
5.2.1	Orthogonal Matching Pursuit Algorithm	38
5.2.2	Strategies in Applications	38
5.3	Non-convex Problems	38
5.3.1	Augumented Lagrange Method	38
5.3.2	Non-linear Lagrangian Duality	38
5.4	Future Work of the Solution for Non-regular Points	38
5.5	Summary	38
6	Conclusion	39
	Appendix	45
A	An Overview of Numerical Optimization	45
A.1	Theory of Optimization	45
A.1.1	Proof of Theorem 2.3	45
A.1.2	Proof of Theorem 2.4	45
A.1.3	Proof of Theorem 2.6	46
A.1.4	Proof of Theorem 2.8	46
A.1.5	Proof of Lemma 2.10	47
A.2	Solution of Unconstrained and Constrained Optimization Problems	48
A.2.1	Proof of Equation 2.3	48
A.2.2	Proof of Equation 2.4 [Gould et al., 2019]	48

List of Figures

2.1	Contour Graph of $f(x, y) = x^4 - 4xy + y^4$	8
2.2	Function $f(x, y)$ at $x = 0$, $x = -1$ and $x = 1$	9
2.3	Feasible set of constraints c_1 , c_2 and c_3	11
2.4	Contour of a constrained problem	12
2.5	Constrained function $c(x) = \ x\ _2^2 - 1$	16
3.1	End-to-end learnable declarative node	22
3.2	Different scenarios for the solution to inequality constrained nodes	27
3.3	Plots of the function y (top) and the gradient (bottom) sweeping the first component of the input x_1 while holding the other elements of x constant	29
3.4	Plots of the function y (top) and the gradient (bottom) sweeping the first component of the input x_1 while holding the other elements of x constant	30
3.5	Plots of the function y (top) and the gradient (bottom) sweeping the first component of the input x_1 while holding the other elements of x constant	31

Introduction

1.1 Motivation

1.2 Thesis Outline

Following the two central themes we mentioned above, this thesis consists of two parts – PART I Deep Declarative Network: Multiple Constrained Declarative Nodes and PART II Deep Declarative Nodes: Non-regular Solution.

PART I focus on the regular points of multiple constrained declarative nodes in the deep declarative network, with some basic examples of different constraints nodes.

In Chapter 2, we first give an overview of the theory of optimization, with the discussion of the optimality. Next, we formally define the unconstrained, equality constrained and inequality constrained problems. We then discuss several related works on the solutions to these problems. Finally, the differentiable neural network, an application of numerical optimization in machine learning is briefly discussed with various modern deep learning algorithms based on it.

In Chapter 3, we present the details of the deep declarative network. We begin with the overview structure and how it works through the specific declarative nodes. We then introduce the learning progress of the deep declarative network. We analyze the backpropagation through the declarative nodes in different constrained problems. Finally, we give multiple examples of the implementation of the deep declarative nodes in both equality constrained and inequality constrained optimization problems. Also, we point out the limitation of current deep declarative nodes and address some ideas for future improvements in the optimization process. We also look to the practical application of the deep declarative network in computer vision tasks, which will be foreseeable. This chapter is based on works of Gould et al. [2019].

PART II is an extension of the deep declarative nodes in non-regular solution problems, which cannot be solved through the traditional numerical optimization methods. Detailedly,

In Chapter 4, we give an overview of the non-regular solution problems. We list the general non-regular solution cases: Overdetermined system, rank deficiency, and non-convex feasible set. We also introduce various related works for solving the problems when the gradient result is not a regular point.

In Chapter 5, we demonstrate various possible solutions for each non-regular point case. Since for non-regular solution problems, there is no exact solution, we can only approximate the closed result. Thus, we also compare and discuss the results of each method on minimizing the final loss. Finally, we point out some future possible extension works for solving non-regular point problems.

We will finally present the conclusion in Chapter 6. Proofs for the important theorems and definitions are given in the Appendix.

1.3 Contribution

Part I

Deep Declarative Network: Multiple Constrained Declarative Nodes

An Overview of Numerical Optimization

In this chapter, we aim to provide readers with an overview of numerical optimization. We begin with the theory of optimization (Section 2.1), from the existence of optimizers, to the optimality conditions for both unconstrained and constrained problems with duality. As the theoretical background of optimization, this field provides a solid solution for the algorithm.

We then formally define the optimization of unconstrained and constrained problems in Section 2.2 and describe the general regular solution for these problems based on the gradient calculation.

Next, we discuss briefly the differentiable optimization in the neural network, which is a novel end-to-end network structure involving optimization problems in each layer with and without constraints. (Section 2.3). Finally, we give a summary of the numerical optimization for solving unconstrained and constrained problems in Section 2.4.

2.1 Theory of Optimization

2.1.1 Existence of Optimizers

In optimization, a basic question is to determine the existence of a global minimizer for a given function f . There are several sufficient conditions on f to guarantee the existence, and the optimizer falls in the feasible set of solutions. For a feasible set, some related definitions are following:

Definition 2.1. A subset $\Omega \in \mathbb{R}^n$ is called

- *bounded* if there is a constant $R > 0$ such that $\|x\| \leq R$ for all $x \in \Omega$
- *closed* if the limit point of any convergent sequence in Ω always lies in Ω
- *compact* if any sequence $\{x_k\}$ in Ω contains a subsequence that converges to a point in Ω

The following result gives a characterization of compact sets in \mathbb{R} . When we find the minimum or maximum solution for the problem, there exists a lower bound or upper bound but not necessarily an optimal solution. Therefore, we have some additional requirements.

Firstly, we give the definition of compact sets in Lemma 2.2. [Oman, 2017] gives a brief proof.

Lemma 2.2 (Bolzano-Weierstrass theorem). A subset Ω in \mathbb{R}^n is *compact* if and only if it is bounded and closed.

We also assume that the function f is continuous and " $+\infty$ at infinity". More precisely, $f(x) \rightarrow +\infty$ if $|x| \rightarrow +\infty$. Such a function is called *inf-compact* or *coercive*. [Nocedal and Wright, 2006] Then the problem can be restricted to a bounded set and existence of a global minimum x^* is guaranteed: a continuous function has a minimum on a compact set. This theorem is defined as follows and the proof is given in Appendix A.1.1.

Theorem 2.3. [Nocedal and Wright, 2006] If f is a continuous function defined on a compact set Ω in \mathbb{R} , then f has a global minimizer x^* on Ω i.e. there exists $x^* \in \Omega$ such that $f(x^*) \leq f(x)$ for all $x \in \Omega$.

More general, based on the definition of coercive function f , we can give following theorem. Proof is given in Appendix A.1.2.

Theorem 2.4. [Nocedal and Wright, 2006] If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a continuous coercive function, then f has at least one global minimizer.

Theorem 2.4 requires the continuity of f which is somewhat restrictive for applications. However, we can replace it by the lower semi-continuity of f which is a rather weaker condition.

Definition 2.5. Let $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\pm\infty\}$. Then f is called *lower semi-continuous* at a point $x_0 \in \mathbb{R}^n$ if for any sequence $f(x_k)$ converging to x_0 here holds $f(x_0) \leq \lim_{k \rightarrow \infty} f(x_k)$. f is called *lower semi-continuous* if f is lower semi-continuous at every point.

Recall our assumptions on function f , it is a continuous function, which is always lower semi-continuous. However, lower semi-continuous functions are not necessarily continuous. For instance, a binary function equals to 0 when $x \leq 0$ and equals to 1 when $x > 0$ is not continuous at $x_0 = 0$. However, since it is greater than 0 for all x and $f(0) = 0$, we have $f(0) = 0 \leq \liminf_{x \rightarrow 0} f(x)$ and it is lower semi-continuous at $x_0 = 0$.

The theorem of the existence of the optimizer of lower semi-continuous function is given as follows and the proof is given in Appendix A.1.3

Theorem 2.6. [Nocedal and Wright, 2006] Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a lower semi-continuous function. If f has a nonempty, compact sublevel set $D := \{x \in \mathbb{R}^n : f(x) \leq \alpha\}$, then f achieves a global minimizer on \mathbb{R} .

Also, we introduce the definition of convex function and convex set which are important in regular optimization problems.

Definition 2.7. A function f is convex when

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y) \quad \text{for all } x, y, \text{ and } \alpha \in]0, 1[$$

A set $C \subset \mathbb{R}^n$ is convex when

$$\alpha x + (1 - \alpha)y \in C \quad \text{for all } x, y \text{ in } C, \text{ and } \alpha \in]0, 1[$$

The problem we are going to discuss in this part is convex and regular, which means its gradient can be computed and the solution exists. However, although the existence of the optimizer is sufficient, for different problems, the optimality conditions are different. In the next two sections, we will give necessary and sufficient conditions for both unconstrained and constrained problems.

2.1.2 Optimality Conditions for Unconstrained Problems

Firstly, we consider the unconstrained minimization problem

$$\min_{x \in \mathbb{R}^n} f(x), \tag{2.1}$$

where f is given function on \mathbb{R}^n .

In order to determine the minimizer, it is important to understand what can happen at a minimizer, and at what condition a point must be a minimizer. Now we have to recognize the optimum point. There are two necessary conditions and one sufficient condition given below [Nocedal and Wright, 2006]. The proof is given in Appendix A.1.4.

Theorem 2.8. *Necessary and Sufficient Conditions. Let $f : \Omega \rightarrow \mathbb{R}$ be a function defined on a set $\Omega \subset \mathbb{R}^n$ and let x^* be an interior point of Ω that is a local minimizer of f .*

Necessary conditions:

- (NC1) *If f is differentiable at x^* , then x^* is a critical point of f , i.e. $\nabla f(x^*) = 0$.*
- (NC2) *If f is twice continuous differentiable on Ω , then the Hessian $\nabla^2 f(x^*)$ is positive semidefinite.*

Sufficient condition (SC1): if x^ is such that $\nabla f(x^*) = 0$ and $\nabla^2 f(x^*)$ is positive definite, then x^* is a local minimum. (i.e. $f(x) \geq f(x^*)$ for x close to x^*)*

Any point satisfying (NC1) as the minimizer of f is called a *critical* or *stationary* point of f . In the objective function f is convex, (NC1) is also the sufficient condition for the global minimum of the solution.

Let us see an example of unconstrained minimization problem. Supposed we have to determine the minimization of function

$$f(x, y) = x^4 - 4xy + y^4$$




Figure 2.1: Contour Graph of $f(x, y) = x^4 - 4xy + y^4$

From the definition of function f , it is clear that f is continuous. Then we can expand f by writing

$$f(x, y) = (x^4 + y^4) \left(1 - \frac{4xy}{x^4 + y^4} \right)$$

we can see f is coercive. Also, we give the contour graph of function f in Figure 2.1. Therefore f has global minimizers which are critical points. Now according to (NC1), we can find the global minimizer through solving the derivative of f equaling to zero:

$$0 = \nabla f(x, y) = \begin{pmatrix} 4x^3 - 4y \\ -4x + 4y^3 \end{pmatrix}$$

Thus, $y = x^3$ and $x = y^3$. Consequently $y = y^9$, i.e.


$$0 = y - y^9 = y(1 - y^8) = y(1 - y^4)(1 + y^4) = y(1 - y)(1 + y)(1 + y^2)(1 + y^4)$$

This implies $y = 0, 1, -1$. Thus f has three critical points $(0, 0), (1, 1), (-1, -1)$. Then we can evaluate f as these points since they may be local minimizer:

$$f(0, 0) = 0, \quad f(1, 1) = -2, \quad f(-1, -1) = -2$$

It achieves the same global minimum value on $(1, 1)$ and $(-1, -1)$. Therefore, they are both global minimizers of f . Figure 2.2 shows the function $f(x, y)$ at these two optimal points.

From this example, we verify that through (NC1), we can find the global minimizer. However, not all continuous functions with critical points have any maximizer

Figure 2.2: Function $f(x, y)$ at $x = 0$, $x = -1$ and $x = 1$

or minimizer. If the function goes to infinity along its axes or a line, it does not have any maximizer or minimizer although it has a critical point. The condition of the minimizer as the critical point is that the function f should be a convex function with continuous first partial derivatives.

Let us move to the sufficient condition (SC1). The result obtained under this theorem is best possible for general functions. Specifically, for a convex function f is defined on a convex set $\Omega \subset \mathbb{R}^n$, any local minimizer of f is also a global minimizer. Moreover, if a function f is strictly convex, it has at most one global minimizer.

2.1.3 Optimality Conditions for Constrained Problems

A general formulation for constrained optimization problems is as follows:

$$\begin{aligned} & \text{minimize } f(x) \\ & \text{subject to } \begin{cases} c_i(x) = 0 & \text{for } i = 1, \dots, m_e, \\ c_i(x) \leq 0 & \text{for } i = m_e + 1, \dots, m \end{cases} \end{aligned} \quad (2.2)$$

where f and c_i are smooth real-valued functions on \mathbb{R}^n , and m_e and m are nonnegative integers with $m_e < m$. We set

$$\mathcal{E} := \{1, \dots, m_e\} \quad \text{and} \quad \mathcal{I} := \{m_e + 1, \dots, m\}$$

as index sets of equality constraints and inequality constraints, respectively.

Here, f is so-called the objective function, and $c_i, i \in \mathcal{E}$ and \mathcal{I} are equality constraints and inequality constraints respectively.

To solve the optimization problem (2.2), we define the feasible set of it to be

$$\mathcal{F} := \{x \in \mathbb{R}^n : c_i(x) = 0 \text{ for } i \in \mathcal{E} \text{ and } c_i(x) \leq 0 \text{ for } i \in \mathcal{I}\}$$

Any point $x \in \mathcal{F}$ is called a feasible point of (2.2) and we call (2.2) infeasible if $\mathcal{F} = \emptyset$. Also, in this feasible set, a feasible point $x^* \in \mathcal{F}$ is called a local minimizer of (2.2) if it is the minimum solution in a neighborhood (strict local minimizer if it is the only one minimum solution). The definition of the global minimizer and strict global minimizer is similar, whose neighborhood is the whole feasible set.

Let us move to the constraints in this problem. For equality constraints, they are strictly equivalent. However, for inequality constraints, there are some exceptions. Let x^* be a local minimizer of (2.2). If there is an index $i \in \mathcal{I}$ such that $c_i(x^*) < 0$, then, x^* is still the local minimizer of the problem obtained by deleting i -th constraint. In this situation, we say that the i -th constraint is inactive at x^* since it does not have any effect on the solution. A general definition of active and inactive inequality constraints is as follows:

Definition 2.9. At a feasible point $x \in \mathcal{F}$, the index $i \in \mathcal{I}$ is said to be *active* if $c_i(x) = 0$ and *inactive* if $c_i(x) < 0$.

In the next chapter, we will give different processes for different cases of active or inactive inequality constraints in the deep declarative nodes. In this chapter, we only focus on the necessary and sufficient conditions for a feasible point x to be a local minimizer of (2.2). These conditions will be derived by considering the change of f on the feasible set along with certain directions. We give the lemma for the condition of local minimizer $x^* \in \mathcal{F}$ as follows, which can be proved through Taylor's formula in Appendix A.1.5.

Lemma 2.10. If $x^* \in \mathcal{F}$ is a local minimizer of (2.2), then

$$d^T \nabla f(x^*) \geq 0 \quad \text{for all } d \in T_{x^*} \mathcal{F}$$

where $T_{x^*} \mathcal{F}$ is the set of all vectors tangent to \mathcal{F} .


However, we may not be able to extract useful results from this lemma, since $T_{x^*} \mathcal{F}$ depends only on the geometry of \mathcal{F} but not on the constraints functions c_i . Not all local minimum falls on the boundary of the constraint function, which is a part of $T_{x^*} \mathcal{F}$. Therefore, it is necessary to introduce linearized feasible directions to give a characterization of $T_{x^*} \mathcal{F}$ in terms of c_i .

Definition 2.11. Given $x \in \mathcal{F}$, we define

$$\text{LFD}(x) := \left\{ d \in \mathbb{R}^n : d^T \nabla c_i(x) = 0 \text{ for } i \in \mathcal{E}; d^T \nabla c_i(x) \leq 0 \text{ for } i \in \mathcal{I} \cap \mathcal{A}(x) \right\}$$

and call it the set of linearized feasible directions of \mathcal{F} at x .

Heuristically, for $i \in \mathcal{E}$ we should travel along directions d with $d^T \nabla c_i(x) = 0$ in order to stay on the curve $c_i(x) = 0$; for $i \in \mathcal{I}$ we should travel along directions with

Figure 2.3: Feasible set of constraints c_1 , c_2 and c_3

$d^T \nabla c_i(x) \leq 0$ in order to stay in the region $c_i(x) \leq 0$. Let us see an example of the linearized feasible directions and the tangent. Supposed we are considering a set \mathcal{F} with variables $(x, y) \in \mathbb{R}^2$ and three inequality constraints functions:

$$c_1(x, y) = x - 1 \leq 0$$

$$c_2(x, y) = -y \leq 0$$

$$c_3(x, y) = y^2 - x \leq 0$$

We can illustrate the feasible set of constraints c_1 , c_2 and c_3 in Fig 2.3. The active set of $0 = (0, 0)$ is $\{2, 3\}$, since $c_1(0) = -1 < 0$, which is inactive. And we can get the derivative of c_2 and c_3 at 0:

$$\nabla c_2(0) = (0, -1)^T \quad \text{and} \quad \nabla c_3(0) = (-1, 0)^T$$

Then we have the linearized feasible directions on $x = 0$:

$$\begin{aligned} \text{LFD}(0) &= \left\{ d \in \mathbb{R}^2 : d^T \nabla c_2(0) \leq 0 \text{ and } d^T \nabla c_3(0) \leq 0 \right\} \\ &= \{ d \in \mathbb{R}^2 : d \geq 0 \} \end{aligned}$$

which equals to the set of all vectors tangent to the feasible set $T_0 \mathcal{F}$.

Unlike the unconstrained optimization problem, the first order necessary condition of the existence of the optimizer is different since we should consider its linearized feasible directions and constraints feasibility. This is so-called the Karush-Kuhn-Tucker theorem:

Theorem 2.12 (Karush-Kuhn-Tucker Theorem). *Let $x^* \in \mathcal{F}$ be a local minimizer of*




Figure 2.4: Contour of a constrained problem

problem (2.2). If

$$T_{x^*}\mathcal{F} = \text{LFD}(x^*),$$

then there exists $\lambda^* = (\lambda_1^*, \dots, \lambda_m^*)^T \in \mathbb{R}^m$ such that

$$\nabla f(x^*) + \sum_{i \in \mathcal{E} \cup \mathcal{I}} \lambda_i^* \nabla c_i(x^*) = 0, \quad (\text{Lagrangian stationary})$$

$$\left. \begin{array}{l} c_i(x^*) = 0 \quad \text{for all } i \in \mathcal{E}, \\ c_i(x^*) \leq 0 \quad \text{for all } i \in \mathcal{I}, \end{array} \right\} \quad (\text{primal feasibility})$$

$$\lambda_i^* \geq 0 \quad \text{for all } i \in \mathcal{I}, \quad (\text{dual feasibility})$$

$$\lambda_i^* c_i(x^*) = 0 \quad \text{for all } i \in \mathcal{E} \cup \mathcal{I}. \quad (\text{complementary slackness})$$

This set of equations are Karush-Kuhn-Tucker (KKT) conditions and a point x^* is called a KKT point if there exists λ^* such that (x^*, λ^*) satisfies the KKT conditions.

For constrained optimization problem, the classic solution is using Lagrange multipliers [Bertsekas, 2014]. Figure 2.4 shows an example of constrained problem, where $f(x, y) = d_i, i = 1, 2, 3$ are the contours of different solution of the objective function and $c(x, y) = 0$ is the equality constraint of the problem.

For multiple constraints, this method introduces the function

$$\mathcal{L}(x, \lambda) := f(x) + \sum_{i \in \mathcal{E} \cup \mathcal{I}} \lambda_i c_i(x)$$

which is called the Lagrange function. x is the primal variables and $\lambda_i, i = 1, \dots, m$ are the Lagrange multipliers or the dual variables. According to the Lagrange multipliers method, we can solve this problem through the gradient of the Lagrange

function:

$$\nabla_x \mathcal{L}(x, \lambda) = \nabla f(x) + \sum_{i \in \mathcal{E} \cup \mathcal{I}} \lambda_i \nabla c_i(x)$$

Therefore, the first equation in KKT conditions can be written as

$$\nabla_x \mathcal{L}(x^*, \lambda^*) = 0$$

2.2 Solution of Unconstrained and Constrained Optimization Problems

According to the sufficient conditions for unconstrained optimization problems, we can easily compute the optimality through the first and second derivative of the objective function. For equality and inequality constrained problems, the introduction of Lagrangian \mathcal{L} is useful for their closed-form solution. Gould et al. [2016] collected both argmin and argmax bi-level optimization results with and without constraints, which also provide insightful examples of these cases. Amos and Kolter [2017] also present a solution for exact, constrained optimization within a neural network. In this thesis, we only focus on argmin problems, but the argmax problems have similar results.

In this section, we are going to provide some background for the solution of both unconstrained and constrained optimization problems, which is based on the gradient of the regular point.

2.2.1 Unconstrained Optimization

For unconstrained optimization problems, the solution is easy to obtain since we only need to focus on the optimality of the objective function. We consider an objective function $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$:

$$y(x) \in \operatorname{argmin} f(x, y)$$

The derivative of $y(x)$ with respect to x is

$$\frac{dy(x)}{dx} = -[\frac{\partial^2 f}{\partial y(x)^2}]^{-1} \frac{\partial^2 f}{\partial x \partial y(x)} \quad (2.3)$$

which can be proved through differentiating and chain rule. [A.2.1]

A very classic example of the unconstrained minimization problem based on a closed convex nonempty set is the L2 norm $\|\cdot\|_2$. Let $\Omega \in \mathbb{R}^n$ be a closed convex nonempty set. For any $x \in \mathbb{R}^n$, the minimization problem is defined as follows:

$$\min_{y \in \Omega} \|y - x\|_2^2$$

This problem has a unique minimizer, which can be denoted by $P_\Omega(x)$, the Euclidean projection of x onto Ω .

Proof. Let $m := \inf_{y \in \Omega} \|y - x\|_2^2$. Since $\Omega \neq \emptyset$, we have $0 \leq m < \infty$. Let $\{y_k\} \subset \Omega$ be a minimizing sequence such that $\|y_k - x\|_2^2 \rightarrow m$ as $k \rightarrow \infty$. Thus $\|y_k - x\|_2^2 \leq m + 1$ for large k which implies that $\|y_k\|_2 \leq \|x\|_2 + \sqrt{m+1}$ for large k . Therefore $\{y_k\}$ is a bounded sequence. Consequently $\{y_k\}$ has a convergent subsequence $\{y_{k_l}\}$ with limit y^* . Since Ω is closed, we have $y^* \in \Omega$. Thus

$$m = \lim_{l \rightarrow \infty} \|y_{k_l} - x\|_2^2 = \|y^* - x\|_2^2$$

which means that m is achieved at y^* , i.e. the given minimization problem has a solution.

Next we show that the given minimization problem has a unique solution by contradiction. If the solution is not unique, let y_0 and y_1 be two distinct solutions. Then for $0 < t < 1$ we set $y_t = ty_1 + (1-t)y_0$. Since Ω is convex, we have $y_t \in \Omega$. Thus

$$\begin{aligned} \|y_0 - x\|_2^2 &= \|y_1 - x\|_2^2 \leq \|y_t - x\|_2^2 = \|t(y_1 - x) + (1-t)(y_0 - x)\|_2^2 \\ &= t^2 \|y_1 - x\|_2^2 + (1-t)^2 \|y_0 - x\|_2^2 + 2t(1-t) \langle y_1 - x, y_0 - x \rangle \\ &= t \|y_1 - x\|_2^2 + (1-t) \|y_0 - x\|_2^2 - (t-t^2) \|y_1 - x\|_2^2 \\ &\quad - (1-t - (1-t)^2) \|y_0 - x\|_2^2 + 2t(1-t) \langle y_1 - x, y_0 - x \rangle \\ &= t \|y_1 - x\|_2^2 + (1-t) \|y_0 - x\|_2^2 \\ &\quad - t(1-t) \left(\|y_1 - x\|_2^2 + \|y_0 - x\|_2^2 - 2 \langle y_1 - x, y_0 - x \rangle \right) \\ &= \|y_0 - x\|_2^2 - t(1-t) \|y_1 - y_0\|_2^2 \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product on \mathbb{R}^n . Therefore $t(1-t) \|y_1 - y_0\|_2^2 \leq 0$ for $0 < t < 1$ and thus $\|y_1 - y_0\|_2^2 \leq 0$. So $y_1 = y_0$ which is a contradiction.

Overall, the minimization problem defined above has a unique minimizer. \square

There are many different methods to solve the unconstrained optimization problem since generally, we treat this kind of problem as basic one. There are two most classical methods, Newton method [Newton and Colson, 1736] and the Method of Steepest Descent [Debye, 1909]. The former one, Newton method starts from an initial guess x_0 and defines a sequence $\{x_k\}$ iteratively according to some rule. It uses the tangent line of the objective function f at x_k to replace f and uses the root of $L(x) = 0$, where $L(x)$ is the updated $f(x)$ as the next iterate x_{k+1} . Finally, the iteration is terminated as long as the difference between x_k and x_{k+1} less than a preassigned small number. The later one, steepest descent is a basic gradient method, which decreases the value of the objective function in a direction of most rapid change. The change rate of a function f at x in the direction u , a unit vector in \mathbb{R} is determined by the directional derivative. Therefore, at x the value of f decrease fastest in the direction $u = -\nabla f(x) / \|\nabla f(x)\|$, which leads to the gradient method: we update the x through the direction with the step length.

2.2.2 Equality Constrained Optimization

Constrained problems are usually more complicated since the solution is restricted on a boundary or in a feasible region. For equality constraints, the basic case is the linear equality constraints $A\mathbf{y} = \mathbf{b}$. Again, we consider an objective function $f : \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}$. Let $A \in \mathbb{R}^{p \times m}$ and $\mathbf{b} \in \mathbb{R}^p$. A is a set of p linear equations as constraints $A\mathbf{y} = \mathbf{b}$. The problem is defined as follows:

$$\begin{aligned} \mathbf{y}(x) \in \arg \min_{\mathbf{y} \in \mathbb{R}^m} & f(x, \mathbf{y}) \\ \text{subject to} & A\mathbf{y} = \mathbf{b} \end{aligned}$$

The derivative of $\mathbf{y}(x)$ with respect to x is

$$\frac{d\mathbf{y}(x)}{dx} = \left(H^{-1} A^T \left(A H^{-1} A^T \right)^{-1} A H^{-1} - H^{-1} \right) \mathbf{b} \quad (2.4)$$

where $H = \partial^2 f(x, \mathbf{y}) / \partial \mathbf{y}(x)^2$ and $\mathbf{b} = \partial^2 f(x, \mathbf{y}) / \partial x \partial \mathbf{y}(x)$.

The solution in 2.4 can be proved through the Lagrange multipliers [Bertsekas, 2014] in A.2.2. More generally, constraints can be non-linear. That means we cannot use A as a weight matrix for constrained parameters anymore. Therefore, we define the equality constraints problem using a set of m constraints functions $c(x, \mathbf{y})$:

$$\begin{aligned} \mathbf{y}(x) \in \arg \min_{\mathbf{y} \in \mathbb{R}^m} & f(x, \mathbf{y}) \\ \text{subject to} & c_i(x, \mathbf{y}) = 0, \quad i = 1, \dots, m \end{aligned}$$

Solution for general multiple non-linear equality constraints is discussed in the chapter of deep declarative network nodes. Here, we are giving a simple example of non-linear equality constrained optimization problem.

For any given nonzero vector $\mathbf{y} \in \mathbb{R}^n$, we define the minimization problem as follows:

$$\begin{aligned} \text{minimize} & -\mathbf{x}^T \mathbf{y} \\ \text{subject to} & \|\mathbf{x}\|_2^2 = 1 \end{aligned}$$

From the constraint defined above, we can write the constraint function as $c(x) = \|\mathbf{x}\|_2^2 - 1$ and illustrate it in Fig 2.5. Differentiating $c(x)$ with respect to x , we get $\nabla c(x) = \mathbf{x} \neq 0$. Therefore, it follows the definition of LFD 2.11 and the theorem of KKT 2.12, which means that every local minimizer of this problem is a KKT point. Now we can write the Lagrangian function:

$$\mathcal{L}(x, \lambda) = -\mathbf{x}^T \mathbf{y} + \lambda (\|\mathbf{x}\|_2^2 - 1)$$

and the KKT conditions are:

$$\nabla_x \mathcal{L} = -\mathbf{y} + 2\lambda \mathbf{x} = 0, \quad \|\mathbf{x}\|_2^2 = 1$$

From $-\mathbf{y} + 2\lambda \mathbf{x} = 0$ and $\mathbf{y} \neq 0$ defined in the question, we must have $\lambda \neq 0$ and




Figure 2.5: Constrained function $c(x) = \|x\|_2^2 - 1$

$x = y/2\lambda$. Combined with $\|x\|_2^2 = 1$, we get

$$4\lambda^2 = \|y\|_2^2 \Leftrightarrow \lambda = \pm \frac{\|y\|_2}{2}$$

Therefore, consequently, we have $x = \pm \frac{y}{\|y\|_2}$. For each x , we can compute its corresponding value of the objective function:

$$\begin{aligned} x &= \frac{y}{\|y\|_2}, -x^T y = -\|y\|_2 \\ x &= -\frac{y}{\|y\|_2}, -x^T y = \|y\|_2 \end{aligned}$$

Obviously, the minimum is achieved $-\|y\|_2$ at $x = \frac{y}{\|y\|_2}$.

Algorithms for solving constrained problems are various. For basic linear programming, which means that all functions involved are linear, we can transform it into standard form with matrix A , then solve the problem using Lagrangian function based on the KKT condition.

Penalty method[Yeniay, 2005], a function determining when a point x is feasible or not, is used to replace the constrained problem with an unconstrained one. For a minimization problem $f(x)$, the penalty function $P(x)$ associated with a penalty parameter are introduced to combine with $f(x)$ and now we are going to solve a series of unconstrained problems. These problems have converged solutions of the original constrained problem.

2.2.3 Inequality Constrained Optimization

Similar to equality constrained problems, inequality constrained problem usually defined the solution in a feasible set. In general, the standard form of inequality constrained problem is negative constraints:

$$\begin{aligned} y(x) \in \arg \min_{y \in \mathbb{R}^m} & f(x, y) \\ \text{subject to} & c_i(x, y) \leq 0, \quad i = 1, \dots, m \end{aligned}$$

Solutions for inequality constrained problems are various. According to the properties of inequality constraints, active and inactive constraints have different criteria. We aim to find the gradient of the optimal solution, $f'(x)$, based on the inequality constrained argmin function. Panier et al. [1988] proposed a globally convergent algorithm for solving the minimization of smooth objective function based on smooth inequality constraints. This algorithm is based on the Quasi-Newton [Dennis and Moré, 1977] iteration for the solution of the first order condition of the optimality in KKT. An updated version of this algorithm, a new QP-free method demonstrated by Qi and Qi [2000], emphasizes the feasibility of all iterates. It reformulates the KKT optimality condition Fischer–Burmeister function [Jiang, 1999] for nonlinear complementarity problems. The classical solution is still based on the Lagrange multipliers. Bertsekas [2014] proposed that for one-sided inequality constrained problems, it cannot be converted to equality constrained problem. Therefore, it introduced the method minimizing the augmented Lagrangian with respect to x for various value of the Lagrange parameters, which is presented by Powell [1969] and Hestenes [1969]:

$$\begin{aligned} \bar{L}_c(x, z, \lambda, \mu) = & f(x) + \lambda' h(x) + \frac{1}{2} c |h(x)|^2 \\ & + \sum_{j=1}^r \left\{ \mu_j [g_j(x) + z_j^2] + \frac{1}{2} c |g_j(x) + z_j^2|^2 \right\} \end{aligned}$$

The minimization of this augmented Lagrangian can be found through computing the first order derivative with respect to z explicitly for each fixed x .

More recently, Gould et al. [2016] introduced a method approximating the gradient of the inequality constrained problem based on ideas from interior-point methods [Boyd et al., 2004]. It gives a demonstration of log-barrier function, which transforms the original constrained problem into a unconstrained minimization problem.

$$\phi(x, y) = \sum_{i=1}^m \log(-c_i(x, y)) \quad (2.5)$$

$$\text{minimize}_y \quad t f(x, y) - \sum_{i=1}^m \log(-c_i(x, y)) \quad (2.6)$$

Equation 2.5 is the log-barrier function, which takes the sum of the logarithm for all constraints. Then subtracting it in the unconstrained minimization problem approximates the original inequality constrained problem. t in Equation 2.6 is a scaling

factor for duality gap control if the solution set is convex.

Similar to the solution for unconstrained and equality constrained problem, minimizing Equation 2.6 is based on the gradient and hessian of the log-barrier function. Therefore, we can compute the approximation of the inequality constrained objective function.

2.3 Differentiable Neural Network

If a problem is differentiable, that means the solution of this problem can be back-propagated. In neural networks, back-propagation [Goodfellow et al., 2016] is widely used to train the feedforward neural network, especially for supervised learning. Therefore, in deep neural networks, we can treat constrained optimization as an individual layer. Recently, there are several works on end-to-end differentiable convex optimization in the neural network, since this type of layer provides inductive bias for different problems, which is very practical.

OptNet [Amos and Kolter, 2017] is a very classical differentiable layer neural network. Each layer in the end-to-end deep neural network is intergraded into optimization problems, which can capture and encode complex dependencies and constraints between hidden variables. It specifically considers the quadratic programs, which are general convex optimization problems. Similarly, SATNet [Wang et al., 2019], a differentiable maximum satisfiability solver, is also intergraded into end-to-end deep learning systems. Besides, it combines the solver with the traditional convolutional network. Both OptNet and SATNet are applied to solving the Sudoku puzzles, which is a very basic constrained logical problem. To make it more general and efficient, Agrawal et al. [2019] demonstrate an approach based on disciplined convex programs, which is a subclass of the classical convex optimization problems. The affine map introduced in this paper represents the disciplined parametrized program.

A fashion application in the differentiable network is the Perspective-n-Points (PnP) solver. Chen et al. [2020] present BPnP based on PnP solver, performing geometric optimization in computer vision tasks. It back-propagates gradient through PnP accurately and effectively since there is a differentiable function in the optimizer block. Besides, for blind PnP problems in the 3D computer vision task, Campbell et al. [2020] propose an end-to-end network based on the differentiating optimization solutions, which is robust and outperforming.

Apart from the above, the differentiable neural network has many practical and powerful applications. Amos and Yarats [2019] introduce a differentiable variant cross-entropy method for non-convex optimization objective function. Again, due to the differentiable feature of the network, the output of the cross-entropy method is differentiable with respect to the parameters in the objective function, even it is non-convex. In 3D reconstruction tasks, some implicit shape and texture are difficult to represent. Hence, Niemeyer et al. [2020] introduce a differentiable rendering formulation, which makes the network learn them from input images directly since implicit differentiation can learn the depth gradients. Also, some research has been

done to simplify the differentiable neural network since the computational cost and complexity of the differential operators can be very high in different tasks. The architecture proposed by Chen and Duvenaud [2019] is cheap and efficient, which sets the Jacobian matrix into diagonal and hollow. It also changes the backward progress into automatic differentiation, which is more effective and lightweight.

2.4 Summary

In Chapter 2, firstly, we introduce the numerical optimization briefly with some necessary conditions and theorems. For the general convex optimization problem, the existence of the local or global optimizers can be determined by the feasible set. Next, the optimality of both unconstrained and constrained problems are discussed. For unconstrained problems, we only need to follow the necessary and sufficient conditions to find the global minimum of the solution. For constrained problems, it should also satisfy the KKT condition. Then we compare existing algorithms on the solution to these problems. Here, for constrained optimization problems, we have to consider the linearity of constraints. Specifically, the activity of inequality constraints can also be solved with different algorithms. Finally, since this thesis is based on the end-to-end differentiable network, some related works of the application are described since these works inspire the deep declarative network in the next chapter. In the next chapter, we are going to describe the deep declarative network in detail with examples.

Deep Declarative Network

In this chapter, we will cover the structure and nodes in the deep declarative network: from its learning process to the back-propagation.

Before delving into the details of the back-propagation in different constraints cases, we give an overview of the deep declarative network in Section 3.1. In particular, the basic structure of the network and the details of declarative nodes are described according to Gould et al. [2019]. The learning progress of the network is also given. We hope this will give readers a better sense of what is the deep declarative network and how it works.

In Section 3.2, we present the details of the back-propagation in different constrained problems. The gradient computation results are based on the implicit differentiation and different in constrained problems. We discuss this part based on the regular solution and compare it with the general solution in the previous chapter.

Next we present the examples of constrained optimization problems with both linear and non-linear, equality and inequality constraints in Section 3.3. We also provide more implementation details of the deep declarative nodes.

Finally, we summarize the deep declarative network and its solution in different constrained problems under the regular point.

3.1 An Overview of Deep Declarative Network

3.1.1 Declarative Node

In deep declarative network, it defines the solution of a constrained optimization problem with parameter $x \in \mathbb{R}^n$ as the output of each node $y \in \mathbb{R}^m$. The general optimization problem can be defined as

$$y \in \arg \min_{u \in C} f(x, u) \quad (3.1)$$

where f is the objective function $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$, and $C \in \mathbb{R}^m$ is the set of constraints parameterized by x .

Apart from the traditional forward processing mapping node, deep declarative node does not explicitly define the transforming function from the input to the out-




Figure 3.1: End-to-end learnable declarative node

put. It defines the input-output relationship implicitly by an objective and constraints optimization problem, where the solution of the problem is the output.

Figure 3.1 shows the forward and backward pass of the declarative node. In the forward evaluation pass, the output of the declarative y is computed as the solution of some minimization problem $f(x, u; \theta)$. We use D to denote the total derivative with respect to the independent variables. Therefore, in the backward pass, the gradient of the global objective function with respect to the output $DJ(y)$ is back-propagated. Its value is computed through the chain rule based on the gradients with respect to the input $DJ(x)$ and parameters $DJ(\theta)$.

Since the definition of deep declarative nodes is very general, it can be embedded within another network for solving subproblems such as robust fitting. However, we may not be able to find the gradient when the feasible set is discrete, or the declarative node is low efficiency to evaluate. As non-regular solution cases, the nonexistent gradient problem will be discussed in the next part. In the next subsection, the learning details of the deep declarative network are described.

3.1.2 Learning

Since in declarative nodes, there is no explicit forward function defined, we can directly compute the optimal solution y through some algorithms. Under this assumption, when we performing the back-propagation, we can compute the gradient of the output from each node with respect to the corresponding input through the implicit differentiation directly. This can be treated as a bi-level optimization problem[Bard, 1998] where the parameterized constraints as a lower-level problem blinds variables in the objective function, an upper-level problem. Combining the schematic illustration in Figure 3.1, the problem can be defined formally as

$$\begin{aligned} & \text{minimize} && J(x, y) \\ & \text{subject to} && y \in \arg \min_{u \in C} f(x, u) \end{aligned} \quad (3.2)$$

We may have additional layers to make the objective function $J(x, y)$ depend on y , which is a function of x . In general, it is the sum of loss terms and regularization terms. We can solve this minimization problem through the gradient descent as

follows:

$$D J(x, y) = D_X J(x, y) + D_Y J(x, y) D y(x) \quad (3.3)$$

where $D_X J(x, y)$ is the partial derivatives of $J(x, y)$ with respect to x and $D_Y J(x, y)$ is the partial derivatives of $J(x, y)$ with respect to y . We used to use D_X and D_Y to denote the partial derivatives. We decompose the total derivatives of $J(x, y)$ as the sum of the partial derivatives with the chain rule. In application, we can consider it as the sum of gradients for losses on training examples.

The lower-level objective function f can be simpler. If it is the only term involving y in the upper-level objective function J , that means $J(x, y) = g(x, f(x, y))$ and the lower-level problem is actually unconstrained with $u \in C = \mathbb{R}^m$. Under this condition, the calculation of the gradient can be expanded using chain rule through both $D_X J(x, y)$ and $D_Y J(x, y)$:

$$\begin{aligned} D J(x, y) &= D_X g(x, f) + D_F g(x, f) (D f + D_Y f D y) \\ &= D_X g(x, f) + D_F g(x, f) D f \end{aligned} \quad (3.4)$$

where $D_Y f(x, y) = 0$ since y is the minimum of $f(x, y)$ and $f(x, y)$ is an unconstrained problem, its partial derivative should be zero.

In addition, for the solution y , we should verify its regularity.

Definition 3.1 (Regular Point). [Gould et al., 2019] A feasible point u is said to be *regular* if the equality constraints gradient $D_U h_i$ and the active inequality constraints gradients $D_U g_i$ are linearly independent, or there are no equality constraints and the inequality constraints are all inactive at u .

Therefore, in unconstrained problems, we can consider the solution is regular by default. However in constrained problems, especially the inequality constrained problems, we should be aware of the feasible set is continuous or not.

3.2 Back-propagation Through Declarative Nodes

Let us focus back on the more general case with y involving in different terms. The backward pass is different in different sub-classes of declarative nodes. We consider three common cases based on Equation 3.1.

3.2.1 Unconstrained

Firstly, the most basic case is the unconstrained problem. Consider a function $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$, we have

$$y \in \arg \min_{u \in C} f(x, u) \quad (3.5)$$

We make the assumption that the solution of this problem, $y(x)$ exists, and in the neighborhood of the point $(x, y(x))$, f is second-order differentiable. Therefore, we

can compute the derivative of y with respect to x is

$$Dy(x) = -H^{-1}B$$

where $H = D_{YY}^2 f(x, y(x)) \in \mathbb{R}^{m \times m}$ is the second-order derivative of f with respect to y , and it is a non-singular matrix. $B = D_{XY}^2 f(x, y(x)) \in \mathbb{R}^{m \times n}$ is the second-order derivative of f with respect to y and x (the derivative of $D_Y f(x, y)$ with respect to x).

The proof of this solution is similar to the proof of Equation 2.3: setting the partial derivative of $f(x, y)$ with respect to the optimal y as 0, then transposing and differentiating both sides according to the implicit function theorem:

$$\begin{aligned} 0_{m \times n} &= D(D_Y f(x, y))^T \\ &= D_{XY}^2 f(x, y) + D_{YY}^2 f(x, y) Dy(x) \end{aligned} \quad (3.6)$$

After rearrangement, we get

$$Dy(x) = - (D_{YY}^2 f(x, y))^{-1} D_{XY}^2 f(x, y) \quad (3.7)$$

Since this is the unconstrained case, for any stationary point of $f(x, y)$, the result is valid. In the following constrained cases, the optimal solution is actually the stationary point of the Lagrangian.

3.2.2 Equality Constrained

Secondly, we consider the equality constrained problem, that means the feasible set is defined by p nonlinear equality constraints. Consider functions $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ and $h : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^p$, we have

$$\begin{aligned} y(x) &\in \arg \min_{u \in \mathbb{R}^m} f(x, u) \\ \text{subject to } &h_i(x, u) = 0, i = 1, \dots, p \end{aligned} \quad (3.8)$$

where $h = [h_1, \dots, h_p]^T$ are a set of constraints.

Again, we make the assumption that the solution of this problem, $y(x)$ exists and both f and all constraints in h are second-order differentiable. Also, we should consider the Jacobian matrix of h with respect to y is full rank since we need the optimal point is regular. Then we can calculate the derivative of y with respect to x is

$$Dy(x) = H^{-1}A^T \left(AH^{-1}A^T \right)^{-1} \left(AH^{-1}B - C \right) - H^{-1}B$$

where

$$\begin{aligned} A &= D_Y h(x, y) \in \mathbb{R}^{p \times m} \\ B &= D_{XY}^2 f(x, y) - \sum_{i=1}^p \lambda_i D_{XY}^2 h_i(x, y) \in \mathbb{R}^{m \times n} \\ C &= D_X h(x, y) \in \mathbb{R}^{p \times n} \\ H &= D_{YY}^2 f(x, y) - \sum_{i=1}^p \lambda_i D_{YY}^2 h_i(x, y) \in \mathbb{R}^{m \times m} \end{aligned}$$

and $\lambda \in \mathbb{R}^p$ can be solved through the system $\lambda^T A = D_Y f(x, y)$.

Similar to the solution of linear equality constrained problem in Equation 2.4, we can form the Lagrangian by the method of Lagrange multipliers[Bertsekas, 2014]:

$$\mathcal{L}(x, y, \lambda) = f(x, y) - \sum_{i=1}^p \lambda_i h_i(x, y) \quad (3.9)$$

We introduce the Lagrange multipliers λ to set the stationary point of this Lagrangian is (y, λ) . Then we differentiate \mathcal{L} with respect to y and λ separately, which are both resulting in 0 since y is the optimality. We have two cases at the optimal point y . The first one is that the partial derivatives of f with respect to y equals to zero: $D_Y f(x, y) = 0 \in \mathbb{R}^{1 \times m}$. That means we transfer this problem to an unconstrained problem since its solution satisfies the constraints directly. Under this case, λ can be set as 0 directly. The second one is that the partial derivatives of f with respect to y is non-zero vector, and it is orthogonal to the constraint surface, which is controlled by the set of equality constraints $h(x, y) = 0$. In this case, from the derivative of \mathcal{L} with respect to y equaling to zero, we have

$$D_Y f(x, y) = \sum_{i=1}^p \lambda_i D_Y h_i(x, y) = \lambda^T A \quad (3.10)$$

where A is the same as the defined above.

To solve this equation for λ , if we want to compute explicitly, it has a unique analytic solution $\lambda = (AA^T)^{-1} A(D_Y f)^T$.

Next, we compute the second-order derivative of \mathcal{L} with respect to x . It still equals to zero in both functions. Solving the equation through variable elimination [Boyd et al., 2004] we have

$$D\lambda(x) = \left(AH^{-1}A^T\right)^{-1} \left(AH^{-1}B - C\right) \quad (3.11)$$

$$Dy(x) = H^{-1}A^T \left(AH^{-1}A^T\right)^{-1} \left(AH^{-1}B - C\right) - H^{-1}B \quad (3.12)$$

which is our result.

The problem and solution defined in Section 2.2.2 is the simpler case of this general one. Also, if we only have one constraint, the matrix A and vector λ can be

simplified as vector and scalar separately. Moreover, for linear only constraints, the representation of matrix H and B are also simpler since $\lambda = 0$.

3.2.3 Inequality Constrained

Lastly, inequality constrained problems are more complicated since we have to consider the solution for active and inactive constraints. Also, previous works mentioned in Section 2.2.3 are approximation results or focusing on the convex problems. Here, the declarative nodes consider the the problem containing both equality and inequality constraints. Consider functions $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ and $h : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^p$, and $g : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^q$, we have

$$\begin{aligned} y(x) \in \arg \min_{u \in \mathbb{R}^m} f(x, u) \\ \text{subject to} \quad & h_i(x, u) = 0, i = 1, \dots, p \\ & g_i(x, u) \leq 0, i = 1, \dots, q \end{aligned} \quad (3.13)$$

where $h = [h_1, \dots, h_p]^T$ are still a set of equality constraints, and $g = [g_1, \dots, g_q]^T$ are a set of inequality constraints.

In declarative nodes, it present a more general result based on activity of all inequality constraints at the optimal point $y(x)$. Also, assumptions of the existence of the optimal point $y(x)$ and second-order differentiation of f , g , h remain true. We combine the equality constraints and inequality constraints together in $\tilde{h} = [h_1, \dots, h_p, g_1, \dots, g_q]$ and the derivative of \tilde{h} with respect to y is a full rank matrix to keep the regularity. We have

$$Dy(x) = H^{-1}A^T \left(AH^{-1}A^T \right)^{-1} \left(AH^{-1}B - C \right) - H^{-1}B \quad (3.14)$$

where

$$\begin{aligned} A &= D_Y \tilde{h}(x, y) \in \mathbb{R}^{(p+q) \times m} \\ B &= D_{XY}^2 f(x, y) - \sum_{i=1}^p \lambda_i D_{XY}^2 \tilde{h}_i(x, y) \in \mathbb{R}^{m \times n} \\ C &= D_X \tilde{h}(x, y) \in \mathbb{R}^{(p+q) \times n} \\ H &= D_{YY}^2 f(x, y) - \sum_{i=1}^{p+q} \lambda_i D_{YY}^2 \tilde{h}_i(x, y) \in \mathbb{R}^{m \times m} \end{aligned}$$

and $\lambda \in \mathbb{R}^{p+q}$ satisfies $\lambda^T A = D_Y f(x, y)$ with $\lambda_i \leq 0$ for $i = p+1, \dots, p+q$, which is almost the same as the solution of the equality constrained problem.

However, in inequality constraints, the gradient is discontinuous since the Lagrange multipliers λ for active inequality constraints are zero. We separate the solution of inequality constrained declarative nodes into 3 scenarios, which is illustrated in Figure 3.2. For any inequality constraints $g(x, u) \leq 0$, the constraint can be active or inactive at the optimal solution y . In the first scenario, the constraint is inactive at the solution y , which means $g(x, y) < 0$ and it is completely in the feasible set




Figure 3.2: Different scenarios for the solution to inequality constrained nodes

(y_1). Therefore we have $D_Y f(x, y) = 0$ and we can consider it as an unconstrained problem. In the second scenario, the constraint is active at y , but it is orthogonal to the constraint surface (y_2). Then we have $D_Y f(x, y) \neq 0$ and $\lambda \neq 0$. Then we can consider this inequality constraint as an equality one since the solution falls on the boundary of the feasible set, but the gradient of negative and pointing outside the set. The last scenario, the constraint is active and $D_Y f(x, y) = 0$ when the solution is on the boundary and it is a local minimum (y_3). For the backward propagation, in this case, we can choose both solutions of unconstrained or equality constrained gradient.

3.3 Examples of Declarative Nodes

3.3.1 Implementation Details

We give the implementation of both equality and inequality constraints with linear and nonlinear cases in raw Python under the gradient calculation package Auto-grad[Maclaurin et al., 2015]. In this subsection, we are going to show the implementation details of the basic deep declarative nodes.

Constraints definition. In the multiple equality constraints node and inequality constraints node, equality constraints and inequality constraints are defined in two functions separately. If there are more than one constraint, they are stored in a 1-d array. All constraints equal to zero or less than 0, which means all parameters in the equations are in the left-hand side and the right-hand side is always zero. There is a specific checking function defined to check this. In the linear equality constraints node, the set of linear constraints are defined as a single matrix A and its corresponding vector b in the initialization directly.

Gradient computation. To compute the gradient of the solution under the con-

straints, we need to calculate the Jacobian and Hessian matrix, which are the first-order and second-order derivative matrix of the constraints. For matrix H in the general solution of the constrained problem, it is costly to compute the inverse of it. Therefore, we transfer the problem to solving a linear system $Hx_1 = A^T$ and $Hx_2 = B$ where $x_1 = H^{-1}A^T$ and $x_2 = H^{-1}B$ to solve them. Cholesky decomposition, an algorithm for solving the linear system efficiently, is applied to this problem. It decomposes a Hermitian positive-definite matrix into the product of a lower triangular matrix with real and positive diagonal entries with its conjugate transpose, which is the unique decomposition. In our problems, since matrix H is a symmetric real positive-definite matrix, it can be decomposed through this algorithm effectively. The solution of the Lagrange multipliers λ similar, which is the solution of a linear system $\lambda^T A = D_Y f(x, y)$ through the least-square solution solver.

Optimality checking. For constrained optimization problems, according to the optimality conditions defined in Section 2.1.3, we should check if the first-order optimality condition is satisfied or not. We define a function in all cases for checking the optimality: the gradient of constraint is zero at optimal point ($D_Y h(x, y) = 0$), or the gradient of objective function equals to the product of the Lagrange multipliers and the gradient of constraints ($D_Y f(x, y) = \lambda D_Y h(x, y)$).

Exception handling. Since our assumptions are based on the regular solution, and not all problems have a regular optimal point, we define the exception cases for this problem. When we solve the linear system for the Lagrange multipliers λ , if the solution does not falls as a regular point, we may get a null value in λ due to the non-existence of the gradient on a non-regular solution. We are going to discuss the solution for this specific case in the next part. For here, we just throw the exception once there is a null value in the Lagrange multipliers λ .

3.3.2 Equality Constrained

Firstly, we consider a single linear equality constrained problem, minimizing the KL-divergence between the input x and output y subject to the output forming a valid probability vector, which can be formally defined as

$$\begin{aligned} y = & \underset{u}{\operatorname{argmin}} & -\sum_{i=1}^n x_i \log u_i \\ & \text{subject to} & \sum_{i=1}^n u_i = 1 \end{aligned} \quad (3.15)$$

where the positivity constraint on y is automatically satisfied by the domain of the log function.

A nice feature of this problem is that we can solve it in closed-form as

$$y = \frac{1}{\sum_{i=1}^n x_i} x.$$

Now we are going to solve this problem via an iterative method with derivative of deep declarative node. Set $n = 5$ and $m = 5$, which means the input $x \in \mathbb{R}^5$ and the output $y \in \mathbb{R}^5$. We begin from a random feasible solution, which is the




Figure 3.3: Plots of the function y (top) and the gradient (bottom) sweeping the first component of the input x_1 while holding the other elements of x constant

normalization of the value y , then perform the gradient update. Figure 3.3 shows the function and gradient sweeping the first component of the input x_1 from 0.1 to 10.0 while holding the other elements of x constant.

Now we consider a multiple non-linear equality constrained problem which is defined formally as

$$\begin{aligned} y = \operatorname{argmin}_u \quad & \sum_{i=1}^n x_i u_i^2 \\ \text{subject to} \quad & \sum_{i=1}^{n-1} u_i^2 = 1 \\ & \sum_{i=1}^n u_i = 0 \end{aligned} \quad (3.16)$$

Same as the previous example, we instantiate the multiple equality constraints nodes in deep declarative nodes with $n = 3$ and $m = 3$. We begin from a feasible solution, $\cos^2 x + \sin^2 x - \sin x - \cos x = 0$, performing the gradient descent until the convergence. Figure 3.4 shows the function y and its gradient changes.

3.3.3 Inequality Constrained

For inequality constrained problems, we consider a similar problem based on the multiple equality constrained problem above. The only difference is that we restrict




Figure 3.4: Plots of the function y (top) and the gradient (bottom) sweeping the first component of the input x_1 while holding the other elements of x constant

the first parameter u_1 is less than u_2 . Therefore, the problem is defined as

$$\begin{aligned}
 y = \operatorname{argmin}_u \quad & \sum_{i=1}^n x_i u_i^2 \\
 \text{subject to} \quad & \sum_{i=1}^{n-1} u_i^2 = 1 \\
 & \sum_{i=1}^n u_i = 0 \\
 & u_1 - u_2 < 0
 \end{aligned} \tag{3.17}$$

The deep declarative node is the instantiation of inequality constraints nodes with $n = 3$ and $m = 3$. Same as the feasible solution as the beginning, we set $x = \pi/6$, which satisfies the inequality constraints since $\sin(\pi/6) < \cos(\pi/6)$. Figure 3.5 shows the function y and its gradient changes. The gradient flattened to some extreme value at the beginning of the iteration then converge to almost zero.

3.4 Future Work of the Deep Declarative Network

There are various possible challenging extensions of the deep declarative network in both theory and application. As a differentiable network, it combines the exact functionality solution with the iterative gradient method. In this section, we give several future foreseeable extensions of the declarative network comparing with other state-of-the-art differentiable models in both theory optimization and applications in computer vision tasks.

We give the solution to the deep declarative nodes of constrained problems un-




Figure 3.5: Plots of the function y (top) and the gradient (bottom) sweeping the first component of the input x_1 while holding the other elements of x constant

der the assumption that the solution of the problem $y(x)$ exists, and it should be a regular point whose gradient can be computed. Also, both objective function and constraints are required to be first and second-order differentiable, which also means that they are continuous instead of discrete. Therefore, in some computer vision tasks such as binary constrained classification problem may not be able to solve using this approach. Some relevant extension methods are discussed in PART II. Also, the robustness of the declarative nodes can be improved by introducing attention to specific constraints.

In solving the real computer vision tasks, comparing to other differentiable models, the deep declarative network can be applied to the visual Sudoku problem. Also, many constrained optimization problems in the graph can be solved through this method. Future work in applications can focus on providing a more specific algorithm for different tasks.

3.5 Summary

In Chapter 3, we describe the deep declarative network from its structure to its implementation with examples of different constrained problems. As a differentiable neural network, the deep declarative network defines its nodes to process input implicitly through the optimization problem, which can find the optimal solution directly without the verification of the local or global minimum. Its learning progress is also based on the derivative of the solution and the gradient of its constraints. We

also introduce the back-propagation through deep declarative nodes in three subclasses: unconstrained, constrained, and inequality constrained problems. All representations of the gradient in declarative nodes have assumptions of the existence of the optimal point, its regularity, and its first and second-order derivative. In the next chapter, we are going to give an overview of the non-regular solution with related previous works.

Part II

Deep Declarative Nodes: Non-regular Solution

An Overview of Regular and Non-regular Solution

Solutions of Non-regular Point

In the last chapter, we discussed various previous works in solving the non-regular point problem. However, not all these approaches are suitable for deep declarative nodes and they may not be able to optimize each parameter in the node. In this chapter, we set out to tackle solving this problem efficiently in the extension of the deep declarative network.

For each scenario, we provide two practical solutions in Section 5.1 (overdetermined system), Section 5.2 (rank deficiency problems) and Section 5.3 (non-convex cases) separately. More specifically, we introduce the least-squares method and conjugate gradient preconditioning approach for the overdetermined system, which are practical and classical approximation algorithms. For rank deficient problems, we firstly propose a greedy strategy, the orthogonal matching pursuit algorithms to recover the matrix as a full rank one. We also present that in the application of the deep declarative network, how to avoid the non-regular point and minimize the approximation error. In the last scenario, we provide two theoretical methods based on the optimality conditions of the optimization problem, which are extensions of the traditional Lagrange multipliers approach. We finally prospect some future improvements for the non-regular solution in the optimization problem in Section 5.4. The summary of this chapter is given in the end.

5.1 Overdetermined System

5.1.1 Least-Squared Method

As a classical approach to approximate the solution of the overdetermined system in linear analysis, the least-squares method is powerful and empirical in many prediction problems by minimizing the sum of the squares of the residuals. [Datta, 2010]

We begin with the linear system of equations. Supposed we have such a linear system

$$\mathbf{Ax} = \mathbf{b} \tag{5.1}$$

where $\mathbf{A} \in \mathbb{R}^{m \times n}$ with $m > n$, $\mathbf{x} \in \mathbb{R}^n$ and $\mathbf{b} \in \mathbb{R}^m$. Therefore, there are more equations than unknowns, which is an overdetermined system and there is no solution making $\mathbf{Ax} = \mathbf{b}$ for all \mathbf{x} . \mathbf{b} is also not in the column subspace of \mathbf{A} , hence \mathbf{b} is

actually not a linear combination of the column vectors of \mathbf{A} . With the least-square method, we want to find an \mathbf{x} which makes the residual vector $\mathbf{r} \in \mathbb{R}^m$ approaching zero:

$$\mathbf{r} = \mathbf{b} - \mathbf{Ax}, \text{ for each element in } \mathbf{r} : r_i = b_i - \sum_{j=1}^n a_{ij}x_j, \quad i = 1, \dots, m \quad (5.2)$$

The solution \mathbf{x} in Equation 5.2 given by the least squares method minimizes $\|\mathbf{r}\|_2 = \|\mathbf{b} - \mathbf{Ax}\|_2$, which is also the sum of errors:

$$\|\mathbf{r}\|_2^2 = \mathbf{r}^T \mathbf{r} = (\mathbf{b} - \mathbf{Ax})^T (\mathbf{b} - \mathbf{Ax}) = \mathbf{b}^T \mathbf{b} - \mathbf{x}^T \mathbf{A}^T \mathbf{b} - \mathbf{b}^T \mathbf{Ax} + \mathbf{x}^T \mathbf{A}^T \mathbf{Ax} \quad (5.3)$$

5.1.2 Conjugate Gradient and Preconditioning

5.2 Rank Deficiency

5.2.1 Orthogonal Matching Pursuit Algorithm

In a linear system $Ax = b$, if A is rank deficient, there are two problems. The first problem is that there is not an exact solution at all. Since the range of A does not span the entire \mathbb{R}^n (if A has n columns) but only an $(N - 1)$ -dimensional subspace. In this case, we can solve exactly for x only if b is in this subspace. The second problem is that there are actually infinity solutions, since A has a non-empty null space, which means that there is a 1-dimensional subspace of vectors x that gives $Ax = 0$. If we use the least-squares solution, the pseudoinverses of A multiply by b , it provides a solution for both problems. For the first problem, it gives the closest x , which is also the one with the smallest $\|b - Ax\|$. At the same time, for the second problem, out of all solutions that share the smallest approximation error and it gives the one with the smallest norm, which is the minimum-norm solution. Therefore, applying the least-squares method to rank deficient may not be able to obtain the best solution for all columns in the matrix.

We consider the orthogonal matching pursuit algorithm (OMP), which is a greedy compressed approach widely used in signal recovery.

5.2.2 Strategies in Applications

5.3 Non-convex Problems

5.3.1 Augmented Lagrange Method

5.3.2 Non-linear Lagrangian Duality

5.4 Future Work of the Solution for Non-regular Points

5.5 Summary

Conclusion

Bibliography

- AGRAWAL, A.; AMOS, B.; BARRATT, S.; BOYD, S.; DIAMOND, S.; AND KOLTER, J. Z., 2019. Differentiable convex optimization layers. In *Advances in neural information processing systems*, 9562–9574. (cited on page 18)
- AMOS, B. AND KOLTER, J. Z., 2017. Optnet: Differentiable optimization as a layer in neural networks. *arXiv preprint arXiv:1703.00443*, (2017). (cited on pages 13 and 18)
- AMOS, B. AND YARATS, D., 2019. The differentiable cross-entropy method. *arXiv preprint arXiv:1909.12830*, (2019). (cited on page 18)
- BARD, J. F., 1998. *Practical Bilevel Optimization: Algorithms and Applications*, vol. 30. Springer Science & Business Media. (cited on page 22)
- BERTSEKAS, D. P., 2014. *Constrained optimization and Lagrange multiplier methods*. Academic press. (cited on pages 12, 15, 17, and 25)
- BOYD, S.; BOYD, S. P.; AND VANDENBERGHE, L., 2004. *Convex optimization*. Cambridge university press. (cited on pages 17 and 25)
- CAMPBELL, D.; LIU, L.; AND GOULD, S., 2020. Solving the blind perspective-n-point problem end-to-end with robust differentiable geometric optimization. *arXiv preprint arXiv:2007.14628*, (2020). (cited on page 18)
- CHEN, B.; PARRA, A.; CAO, J.; LI, N.; AND CHIN, T.-J., 2020. End-to-end learnable geometric vision by backpropagating pnp optimization. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, 8100–8109. (cited on page 18)
- CHEN, R. T. AND DUVENAUD, D. K., 2019. Neural networks with cheap differential operators. In *Advances in Neural Information Processing Systems*, 9961–9971. (cited on page 19)
- DATTA, B. N., 2010. *Numerical linear algebra and applications*, vol. 116. Siam. (cited on page 37)
- DEBYE, P., 1909. Näherungsformeln für die zylinderfunktionen für große werte des arguments und unbeschränkt veränderliche werte des index. *Mathematische Annalen*, 67, 4 (1909), 535–558. (cited on page 14)
- DENNIS, J. E., JR AND MORÉ, J. J., 1977. Quasi-newton methods, motivation and theory. *SIAM review*, 19, 1 (1977), 46–89. (cited on page 17)

-
- GOODFELLOW, I.; BENGIO, Y.; COURVILLE, A.; AND BENGIO, Y., 2016. *Deep learning*, vol. 1. MIT press Cambridge. (cited on page 18)
- GOULD, S.; FERNANDO, B.; CHERIAN, A.; ANDERSON, P.; CRUZ, R. S.; AND GUO, E., 2016. On differentiating parameterized argmin and argmax problems with application to bi-level optimization. *arXiv preprint arXiv:1607.05447*, (2016). (cited on pages 13 and 17)
- GOULD, S.; HARTLEY, R.; AND CAMPBELL, D., 2019. Deep declarative networks: A new hope. *arXiv preprint arXiv:1909.04866*, (2019). (cited on pages xii, 1, 21, 23, and 48)
- HESTENES, M. R., 1969. Multiplier and gradient methods. *Journal of optimization theory and applications*, 4, 5 (1969), 303–320. (cited on page 17)
- JIANG, H., 1999. Global convergence analysis of the generalized newton and gauss-newton methods of the fischer-burmeister equation for the complementarity problem. *Mathematics of Operations Research*, 24, 3 (1999), 529–543. (cited on page 17)
- MACLAURIN, D.; DUVENAUD, D.; AND ADAMS, R. P., 2015. Autograd: Effortless gradients in numpy. In *ICML 2015 AutoML Workshop*, vol. 238, 5. (cited on page 27)
- NEWTON, I. AND COLSON, J., 1736. *The Method of Fluxions and Infinite Series; with Its Application to the Geometry of Curve-lines... Translated from the Author's Latin Original Not Yet Made Publick. To which is Subjoin'd a Perpetual Comment Upon the Whole Work...* by J. Colson. (cited on page 14)
- NIEMEYER, M.; MESCHEDER, L.; OECHSLE, M.; AND GEIGER, A., 2020. Differentiable volumetric rendering: Learning implicit 3d representations without 3d supervision. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, 3504–3515. (cited on page 18)
- NOCEDAL, J. AND WRIGHT, S., 2006. *Numerical optimization*. Springer Science & Business Media. (cited on pages 6 and 7)
- OMAN, G., 2017. A short proof of the bolzano-weierstrass theorem. *The College Mathematics Journal*, (2017). (cited on page 6)
- PANIER, E. R.; TITS, A. L.; AND HERSKOVITS, J. N., 1988. A qp-free, globally convergent, locally superlinearly convergent algorithm for inequality constrained optimization. *SIAM Journal on Control and Optimization*, 26, 4 (1988), 788–811. (cited on page 17)
- POWELL, M. J., 1969. A method for nonlinear constraints in minimization problems. *Optimization*, (1969), 283–298. (cited on page 17)
- QI, H.-D. AND QI, L., 2000. A new qp-free, globally convergent, locally superlinearly convergent algorithm for inequality constrained optimization. *SIAM Journal on Optimization*, 11, 1 (2000), 113–132. (cited on page 17)

-
- WANG, P.-W.; DONTI, P. L.; WILDER, B.; AND KOLTER, Z., 2019. Satnet: Bridging deep learning and logical reasoning using a differentiable satisfiability solver. *arXiv preprint arXiv:1905.12149*, (2019). (cited on page 18)
- YENİAY, Ö., 2005. Penalty function methods for constrained optimization with genetic algorithms. *Mathematical and computational Applications*, 10, 1 (2005), 45–56. (cited on page 16)

Appendix

A An Overview of Numerical Optimization

A.1 Theory of Optimization

A.1.1 Proof of Theorem 2.3

Proof. Let

$$m := \inf\{f(x) : x \in \Omega\}$$

By the definition of m we may pick a sequence $\{x_k\} \subset \Omega$ with $f(x_k) \rightarrow m$ as $k \rightarrow \infty$. Because Ω is compact, we can extract a convergent subsequence $\{x_{k_j}\}$ from $\{x_k\}$. Let $x^* \in \Omega$ denote the limit point of $\{x_{k_j}\}$. Since f is continuous, $f(x^*) = \lim_{j \rightarrow \infty} f(x_{k_j}) = m$. Thus m is finite and x^* is a global minimizer of f on Ω .

When $\Omega = \mathbb{R}^n$, we need to impose conditions on f at infinity to guarantee the existence of a global minimizer. \square

A.1.2 Proof of Theorem 2.4

Proof. Let $m := \inf\{f(x) : x \in \mathbb{R}^n\}$, and take a sequence $\{x_k\}$ such that

$$f(x_k) \rightarrow m \quad \text{as } k \rightarrow \infty.$$

Since f is coercive, $\{x_k\}$ must be bounded; otherwise it has a subsequence $\{x_{k_j}\}$ with $\|x_{k_j}\| \rightarrow \infty$ as $j \rightarrow \infty$, and hence $m = \lim_{j \rightarrow \infty} f(x_{k_j}) = +\infty$, a contradiction.

Thus there is $r > 0$ such that

$$\{x_k\} \subset \{x \in \mathbb{R}^n : \|x\| \leq r\}.$$

Because $\{x \in \mathbb{R}^n : \|x\| \leq r\}$ is compact, $\{x_k\}$ has a convergent subsequence $\{x_{k_j}\}$

with $x_{k_j} \rightarrow x^*$ as $j \rightarrow \infty$. In view of the continuity of f , we have

$$f(x^*) = \lim_{j \rightarrow \infty} f(x_{k_j}) = m$$

Therefore m is finite and f achieves its minimum on \mathbb{R}^n at x^* □

A.1.3 Proof of Theorem 2.6

Proof. We may assume that $\alpha > f_* := \inf \{f(x) : x \in \mathbb{R}^n\}$. Let $\{x_k\}$ be a minimizing sequence for f , i.e.

$$f(x_k) \rightarrow f_* \quad \text{as } k \rightarrow \infty$$

Then there is an N such that $f(x_k) \leq \alpha$ for all $k \geq N$, that is, $x_k \in D$ for all $k \geq N$. Since D is compact, $\{x_k\}_{k=N}^\infty$ has a convergent subsequence $\{x_{k_j}\}$ with $x_{k_j} \rightarrow x_* \in D$ as $j \rightarrow \infty$. In view of the lower semi-continuity of f , we have

$$f(x_*) \leq \lim_{j \rightarrow \infty} f(x_{k_j}) = f_*$$

By the definition of f_* we must have $f(x_*) = f_*$. Therefore f achieves its minimum on \mathbb{R} at x_* . □

A.1.4 Proof of Theorem 2.8

Proof. (NC1): First, recall that for any $v \in \mathbb{R}^n$ there holds

$$v^T \nabla f(x^*) = D_v f(x^*) = \lim_{t \searrow 0} \frac{f(x^* + tv) - f(x^*)}{t}.$$

Since x^* is a local minimizer, we have

$$f(x^* + tv) - f(x^*) \geq 0 \quad \text{for small } |t|.$$

Therefore

$$v^T \nabla f(x^*) \geq 0 \quad \text{for all } v \in \mathbb{R}^n.$$

In particular this implies $(-v)^T \nabla f(x^*) \geq 0$ and thus

$$v^T \nabla f(x^*) \leq 0 \quad \text{for all } v \in \mathbb{R}^n.$$

Therefore $v^T \nabla f(x^*) = 0$ for all $v \in \mathbb{R}^n$. Taking $v = \nabla f(x^*)$ gives $\|\nabla f(x^*)\|^2 = 0$ which shows that $\nabla f(x^*) = 0$ □

Proof. (NC2): Recall that for any $v \in \mathbb{R}^n$ and small $t > 0$ there is $0 < s < 1$ such that

$$f(x^* + tv) = f(x^*) + tv^T \nabla f(x^*) + \frac{1}{2} t^2 v^T \nabla^2 f(x^* + stv) v.$$

Since x^* is a local minimizer of f , we have $f(x^* + tv) \geq f(x^*)$ and $\nabla f(x^*) = 0$ by (NC1). Therefore

$$\frac{1}{2}t^2 v^T \nabla^2 f(x^* + stv) v = f(x^* + tv) - f(x^*) \geq 0.$$

This implies that

$$v^T \nabla^2 f(x^* + stv) v \geq 0.$$

Taking $t \rightarrow 0$ gives

$$v^T \nabla^2 f(x^*) v \geq 0 \quad \text{for all } v \in \mathbb{R}^n$$

i.e. $\nabla^2 f(x^*)$ is semi-definite. □

Proof. (SC1): Since $\nabla^2 f(x)$ is continuous and $\nabla^2 f(x^*) \geq 0$, we can find $r > 0$ such that

$$B_r(x^*) \subset \Omega \quad \text{and} \quad \nabla^2 f(x) > 0 \text{ for all } x \in B_r(x^*).$$

By Taylor's formula we have

$$f(x) = f(x^*) + \nabla f(x^*) \cdot (x - x^*) + \frac{1}{2}(x - x^*)^T \nabla^2 f(\hat{x})(x - x^*)$$

where $\hat{x} := x^* + t(x - x^*)$ for some $0 < t < 1$.

It is clear that $\hat{x} \in B_r(x^*)$ and hence $\nabla^2 f(\hat{x}) > 0$ which implies that

$$(x - x^*)^T \nabla^2 f(\hat{x})(x - x^*) > 0 \quad \text{for } x \neq x^*$$

Consequently

$$f(x) > f(x^*) + \nabla f(x^*) \cdot (x - x^*)$$

for all $x \in B_r(x^*)$ with $x \neq x^*$.

Since $\nabla f(x^*) = 0$, we can obtain $f(x) > f(x^*)$ for all $x \in B_r(x^*)$ with $x \neq x^*$. □

A.1.5 Proof of Lemma 2.10

Proof. For $d \in T_{x^*}\mathcal{F}$, we have $z_k \subset \mathcal{F}$ and t_k such that

$$z_k \rightarrow x^*, \quad 0 < t_k \rightarrow 0 \quad \text{and} \quad \frac{z_k - x^*}{t_k} \rightarrow d$$

as $k \rightarrow \infty$. As $f(x^*) \leq f(z_k)$, by Taylor's formula we have

$$\begin{aligned} f(x^*) &\leq f(z_k) = f(x^* + (z_k - x^*)) \\ &= f(x^*) + (z_k - x^*)^T \nabla f(x^*) + \frac{1}{2}(z_k - x^*)^T \nabla^2 f(\hat{z}_k)(z_k - x^*) \end{aligned}$$

where \hat{z}_k is a point on the line segment joining x^* and z_k . This implies that

$$0 \leq \left(\frac{z_k - x^*}{t_k} \right)^T \nabla f(x^*) + \frac{1}{2} (z_k - x^*)^T \nabla^2 f(\hat{z}_k) \left(\frac{z_k - x^*}{t_k} \right)$$

Letting $k \rightarrow \infty$ gives $d^T \nabla f(x^*) \geq 0$ □

A.2 Solution of Unconstrained and Constrained Optimization Problems

A.2.1 Proof of Equation 2.3

Proof. Firstly, for any optimal y , according to the first-order optimality condition, we have

$$\frac{df(x, y)}{dy} = \mathbf{0} \in \mathbb{R}^{1 \times m}$$

Then from the implicit function theorem, rearranging and differentiating both sides we have

$$\begin{aligned} D\left(\frac{df(x, y)}{dy}\right)^T &= \mathbf{0} \in \mathbb{R}^{m \times n} \\ &= \frac{\partial^2}{\partial x \partial y} f(x, y) + \frac{\partial^2}{\partial y^2} f(x, y) \frac{dy(x)}{dx} \\ \frac{dy(x)}{dx} &= -\left[\left(\frac{\partial^2}{\partial y^2}\right) f(x, y)\right]^{-1} \left(\frac{\partial^2}{\partial x \partial y}\right) f(x, y) \end{aligned}$$

□

A.2.2 Proof of Equation 2.4 [Gould et al., 2019]

Proof. According to the definition of Lagrange multipliers, we can define the Lagrangian:

$$\mathcal{L}(x, y, \lambda) = f(x, y) - \sum_{i=1}^p \lambda_i (A_i y_i - b_i)$$

We are going to find the stationary point (y, λ) for this lagrangian. Therefore, we calculate the derivative of \mathcal{L} with respect to y and λ separately:

$$\frac{\partial}{\partial y} f(x, y) - \sum_{i=1}^p \lambda_i \frac{\partial}{\partial y} (A_i y_i - b_i) = 0 \quad (1)$$

$$A y - b = 0 \quad (2)$$

Since y is the optimal point, we have $\frac{\partial}{\partial y} f(x, y) = 0$, which can be an unconstrained problem or it is orthogonal to the constraint surface. For unconstrained cases, we can

set $\lambda = 0$ directly. For the orthogonal case, from Equation 1, we have

$$\frac{\partial}{\partial y} f(x, y) = \sum_{i=1}^p \lambda_i \frac{\partial}{\partial y} (A_i y_i - b_i) = \lambda^T A$$

Now we are going to calculate the derivative of the Lagrangian with respect to x for both 1 and 2:

$$\frac{\partial^2}{\partial x \partial y} f(x, y) + \frac{\partial^2}{\partial y^2} f(x, y) Dy - \frac{\partial}{\partial y} (A y - b)^T D \lambda = 0 \quad (3)$$

$$\frac{\partial}{\partial x} (A y - b) + \frac{\partial}{\partial y} (A y - b) Dy = 0 \quad (4)$$

Solving 3 and 4, we get:

$$Dy(x) = \left(H^{-1} A^T \left(A H^{-1} A^T \right)^{-1} A H^{-1} - H^{-1} \right) B$$

where

$$H = \frac{\partial^2}{\partial y^2} f(x, y), \quad B = \frac{\partial^2}{\partial x \partial y} f(x, y)$$

□