

Assignment 1

Sujal - AI20BTECH11020

Download all python codes from

https://github.com/sujal100/Probability_and_Random_variable/tree/main/exercise_1/codes

and latex codes from

https://github.com/sujal100/Probability_and_Random_variable/blob/main/exercise_1/exercise_1_main_tex.tex

$A \cap B$ = the number appearing is even multiple of 3 = (3)

So,

$$\Pr(A \cap B) = \frac{1}{6} \quad (2.0.8)$$

Now,

$$\Pr(A) \cdot \Pr(B) = \frac{1}{3} \cdot \frac{1}{2} = \frac{1}{6} \quad (2.0.9)$$

Since,

$$\Pr(A \cap B) = \Pr(A) \cdot \Pr(B) \quad (2.0.10)$$

Therefore A and B are independent events.

1 PROBLEM

A die is thrown. If E is the event "the number appearing is a multiple of 3" and F be the event "the number appearing is even" then find whether E and F are independent ?

2 SOLUTION

Two event A and B are independent if

$$P(A \cap B) = P(A) \cdot P(B) \quad (2.0.1)$$

A die is thrown.

We know that the sample space is

$$S = (1, 2, 3, 4, 5, 6) \quad (2.0.2)$$

Let two events be

A : the number appear is a multiple of 3 .

B : the number appearing is even.

$$A : (3, 6) \quad (2.0.3)$$

$$\Pr(A) = \frac{2}{6} = \frac{1}{3} \quad (2.0.4)$$

$$B : (2, 4, 6) \quad (2.0.5)$$

$$\Pr(B) = \frac{3}{6} = \frac{1}{2} \quad (2.0.6)$$

$$(2.0.7)$$