Problem 12.217

Sujal Rajani

October 12, 2025

Question : Given the surface

$$x^2 + \frac{y^2}{4} + \frac{z^2}{9} = 3,$$

find the directional derivative at $P(1, 2, 3)$ in the direction of the vector **P**, where O is the origin.

Solution

SOLUTION

we represent the partial derivative as :

$$\frac{\partial f}{\partial x} = f_x(x, y)$$

the directional derivative of $f(x, y)$ in the direction of the unit vector $\mathbf{u} = \begin{pmatrix} a \\ b \end{pmatrix}$ is

$$D_{\mathbf{u}} f(x, y) = f_x(x, y)a + f_y(x, y)b, \tag{1.1}$$

and in three dimensions,

$$D_{\mathbf{u}} F(x, y, z) = \nabla \mathbf{F}(\mathbf{x}, \mathbf{y}, \mathbf{z})^{\top} \mathbf{u}, \tag{1.2}$$

where $\nabla \mathbf{F}$ is the gradient of F.

Step-by-Step Calculation

Let

$$F(x, y, z) = x^2 + \frac{y^2}{4} + \frac{z^2}{9}. \tag{1.3}$$

Compute the gradient:

$$\nabla \mathbf{F} = \left(\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y}, \frac{\partial F}{\partial z} \right)^\top = \begin{pmatrix} 2x & \frac{y}{2} & \frac{2z}{9} \end{pmatrix}^\top \tag{1.4}$$

At $P(1, 2, 3)$:

$$\nabla \mathbf{F}(\mathbf{1}, \mathbf{2}, \mathbf{3}) = \begin{pmatrix} 2 & 1 & \frac{2}{3} \end{pmatrix}^\top \tag{1.5}$$

Step-by-Step Calculation

Step 3: Unit Direction Vector

The position vector from the origin to the point $P(1, 2, 3)$ is

$$\mathbf{P} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \tag{1.6}$$

To get the unit vector,

$$||\mathbf{P}||^2 = \mathbf{P}^\top \mathbf{P} = 1^2 + 2^2 + 3^2 = 14 \tag{1.7}$$

$$\hat{\mathbf{P}} = \frac{\mathbf{P}}{||\mathbf{P}||} \tag{1.8}$$

So,

$$\hat{\mathbf{P}} = \left(\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}} \right)^\top$$

Step-by-Step Calculation

By the theorem :

$$D_{\mathbf{u}}F = \nabla \mathbf{F}^{\top}\hat{\mathbf{P}} \tag{1.9}$$

$$= \frac{2}{\sqrt{14}} + \frac{2}{\sqrt{14}} + \frac{2}{\sqrt{14}} \tag{1.10}$$

$$= \frac{6}{\sqrt{14}} \tag{1.11}$$

$$\boxed{\frac{6}{\sqrt{14}}}$$

3D Plot of **P** and ∇F(1, 2, 3)