

Diabetes Hospital Readmission Analysis

Key Patient, Treatment and Encounter Characteristics Associated with Readmission Risk

Total Encounters	Unique Patients	30 Day Readmission Rate	Average Hospital Stay
102K	72K	0.11	4.40

Average Age by Readmission Status






Observation

Older patients show a slightly higher average age among those who were readmitted. However, the difference is minimal, suggesting that age alone is not a strong indicator of 30-day readmission risk.


Average Age by Readmission Status



Observation

Patients readmitted within 30 days had the longest hospital stay (4.77 days), compared with 4.50 days for patients readmitted after 30 days and 4.25 days for patients not readmitted.

Readmission Rate by Admission Type




Observation

Emergency admissions have the highest 30-day readmission rate at 11.52%, followed by urgent admissions at 11.18%. Elective admissions have a lower rate of 10.39%. This suggests that patients admitted through emergency or urgent care may have a slightly higher risk of being readmitted within 30 days, although the differences are relatively small.

Average Number of Medications by Readmission Status







Observation

Patients readmitted within 30 days had the highest average number of medications (16.90), compared with 16.28 for patients readmitted after 30 days and 15.67 for patients who were not readmitted. This suggests that patients taking more medications may have a slightly higher likelihood of readmission, although the differences are relatively small.

Insulin Use and Readmission Rate






Observation

Patients who took insulin (Up or Steady) had a higher readmission rate, with 43.17% for patients readmitted within 30 days, 41.24% for patients readmitted after 30 days and 36.45% for those not readmitted. This suggests that insulin use may be associated with a higher risk of readmission.

Medication Change and Prior Use (for <30 Day Readmission)



Observation

Patients readmitted within 30 days had the highest average number of medications (16.90), compared with 16.28 for patients readmitted after 30 days and 15.67 for patients who were not readmitted. This suggests that patients taking more medications may have a slightly higher likelihood of readmission, although the differences are relatively small.