

9:41

CropGuard

AI DISEASE ANALYST

Welcome, Farmer

Sign in to instantly diagnose crop health and access localized prevention guides.

Full Name

e.g. John Doe

Phone Number

e.g. +1 555 123 4567

Login to CropGuard

Need help joining? [Contact Support](#)

9:41

Good morning, Farmer John

Region North

Diagnose Crop

Upload a clear close-up photo of the affected plant leaf or stem to detect diseases in seconds.

Take or Upload Photo

Tap to open camera or drag & drop leaf picture here

Make sure lighting is bright

Analyze Crop Health

9:41

Diagnosis Report

HIGH SEVERITY

DETECTED DISEASE

Early Blight

Target Crop: Tomato

Confidence Level: 94.2% Accuracy

Recommended Precautions

Early Blight is caused by the fungus Alternaria solani. It initiates brown spots on older leaves. Follow these immediate biological and chemical treatment steps to protect your total harvest yield:

Prune Infected Foliage

Cut away lower affected leaves carefully. Clean shears with alcohol between cuts to prevent spreading spores.

Apply Organic Fungicide

Spray copper-based organic fungicides or Bacillus subtilis thoroughly on both sides of remaining healthy leaves.

Improve Soil & Watering

Switch to drip irrigation. Avoid overhead watering completely to keep foliage dry. Mulch around stem base.

Diagnose Another Crop

AGRICULTURAL DISEASE SURVEILLANCE SYSTEM

Official Monitoring & Response Dashboard • Ministry of Agriculture

2026-03-31 14:45:22 UTC

SYSTEM ONLINE

Officer J. Kiptoo
National Surveillance Unit

Dashboard

Disease Reports

Active Alerts

GIS Map View

Agri-Weather Data

Farmer Database

Threshold Settings

DISEASE RESPONSE PIPELINE TRACKING

STAGE 01
Farmer Intake

STAGE 02
AI Diagnosis

STAGE 03
Observation DB

STAGE 04
Priority Filter

STAGE 05
Threshold Det.

STAGE 06
GIS Analysis

STAGE 07
Weather Corr.

STAGE 08
Risk Scoring

STAGE 09
Human Verification

STAGE 10
Alert Decision

CASE ID: #WSR-2026-0982

PRIORITY: HIGH RISK

Intake: 2 hours ago (2026-03-31 12:30)

Submitted Image: Nakuru Field B3

Wheat Stem Rust (Puccinia graminis)

Highly infectious fungal pathogen capable of causing complete crop failure.

AI Confidence

87 %

Detection Density

14 farms

+3 in last 24h

Affected Location

Nakuru County

AI AUTOMATED ANALYSIS SUMMARY

Multiple alerts matched within a 15km radius. Spore propagation probability is elevated due to consistent moderate winds blowing South-East towards the high-yield grain belt. Pathogen markers correlate with the Ug99 lineage. Action is recommended within 24-48 hours.

AGRI-WEATHER CORRELATION

Temperature

24 °C

Relative Humidity

78%

Rainfall (48h)

12.4mm

FAVORABLE FOR FUNGAL SPREAD

30-DAY OUTBREAK TREND

T-30 Days

Peak Case Velocity

Active

OFFICER DECISION REQUIRED

Review the surveillance synthesis above before executing response actions.

Target Alert Severity Level

Level 3 - County Wide Broadcast & Quarantines Recommended

Officer Triage Notes & Strategic Directives

AI diagnosis confirmed via image evaluation of rust pustule density. High probability Ug99. Regional Extension Officers must deploy immediate copper-based fungicide advisories to Ward 3 and neighboring buffer Ward 4.

Confirm visual verification and authorize system state broadcast under Ministerial Act 14.

EXECUTE & ISSUE SYSTEM ALERT

DISMISS / PASSIVE MONITOR

Collapse Navigation

ADSS V4.2.1-PROD

OPERATIONAL LOGS:

[14:41] GIS layer updated for Nakuru County B3 • [14:38] Agri-weather pipeline retrieved current wind matrices • [14:15] Outbreak pattern detection executed successfully

3 PENDING VERIFICATIONS