

PATTERN RECOGNITION AND MACHINE LEARNING

CHAPTER 8: GRAPHICAL MODELS

Bayesian Networks

Directed Acyclic Graph (DAG)




$$p(a, b, c) = p(c|a, b)p(a, b) = p(c|a, b)p(b|a)p(a)$$

$$p(x_1, \dots, x_K) = p(x_K|x_1, \dots, x_{K-1}) \dots p(x_2|x_1)p(x_1)$$

Bayesian Networks


$$p(x_1, \dots, x_7) = p(x_1)p(x_2)p(x_3)p(x_4|x_1, x_2, x_3) \\ p(x_5|x_1, x_3)p(x_6|x_4)p(x_7|x_4, x_5)$$



General Factorization

$$p(\mathbf{x}) = \prod_{k=1}^K p(x_k | \text{pa}_k)$$

Bayesian Curve Fitting (1)




Polynomial

$$y(x, \mathbf{w}) = \sum_{j=0}^M w_j x^j$$

$$p(\mathbf{t}, \mathbf{w}) = p(\mathbf{w}) \prod_{n=1}^N p(t_n | y(\mathbf{w}, x_n))$$

Bayesian Curve Fitting (2)


$$p(\mathbf{t}, \mathbf{w}) = p(\mathbf{w}) \prod_{n=1}^N p(t_n | y(\mathbf{w}, x_n))$$



Bayesian Curve Fitting (3)

Input variables and explicit hyperparameters


$$p(\mathbf{t}, \mathbf{w} | \mathbf{x}, \alpha, \sigma^2) = p(\mathbf{w} | \alpha) \prod_{n=1}^N p(t_n | \mathbf{w}, x_n, \sigma^2).$$



Bayesian Curve Fitting—Learning

Condition on data

$$p(\mathbf{w}|\mathbf{t}) \propto p(\mathbf{w}) \prod_{n=1}^N p(t_n|\mathbf{w})$$




Bayesian Curve Fitting—Prediction

Predictive distribution: $p(\hat{t}|\hat{x}, \mathbf{x}, \mathbf{t}, \alpha, \sigma^2) \propto \int p(\hat{t}, \mathbf{t}, \mathbf{w}|\hat{x}, \mathbf{x}, \alpha, \sigma^2) d\mathbf{w}$


where

$$p(\hat{t}, \mathbf{t}, \mathbf{w}|\hat{x}, \mathbf{x}, \alpha, \sigma^2) = \left[\prod_{n=1}^N p(t_n|x_n, \mathbf{w}, \sigma^2) \right] p(\mathbf{w}|\alpha) p(\hat{t}|\hat{x}, \mathbf{w}, \sigma^2)$$



Generative Models

Causal process for generating images



Conditional Independence

a is independent of b given c

$$p(a|b, c) = p(a|c)$$


Equivalently

$$\begin{aligned} p(a, b|c) &= p(a|b, c)p(b|c) \\ &= p(a|c)p(b|c) \end{aligned}$$

Notation

$$a \perp\!\!\!\perp b \mid c$$

Conditional Independence: Example 1




$$p(a, b, c) = p(a|c)p(b|c)p(c)$$

$$p(a, b) = \sum_c p(a|c)p(b|c)p(c)$$

$$a \not\perp b \mid \emptyset$$


Conditional Independence: Example 1



$$\begin{aligned} p(a, b|c) &= \frac{p(a, b, c)}{p(c)} \\ &= p(a|c)p(b|c) \end{aligned}$$

$$a \perp\!\!\!\perp b \mid c$$

Conditional Independence: Example 2




$$p(a, b, c) = p(a)p(c|a)p(b|c)$$

$$p(a, b) = p(a) \sum_c p(c|a)p(b|c) = p(a)p(b|a)$$

$$a \not\perp b \mid \emptyset$$


Conditional Independence: Example 2



$$\begin{aligned} p(a, b|c) &= \frac{p(a, b, c)}{p(c)} \\ &= \frac{p(a)p(c|a)p(b|c)}{p(c)} \\ &= p(a|c)p(b|c) \end{aligned}$$

$$a \perp\!\!\!\perp b \mid c$$

Conditional Independence: Example 3




$$p(a, b, c) = p(a)p(b)p(c|a, b)$$

$$p(a, b) = p(a)p(b)$$

$$a \perp\!\!\!\perp b \mid \emptyset$$

Note: this is the opposite of Example 1, with c unobserved.

Conditional Independence: Example 3



$$\begin{aligned} p(a, b|c) &= \frac{p(a, b, c)}{p(c)} \\ &= \frac{p(a)p(b)p(c|a, b)}{p(c)} \end{aligned}$$

$$a \not\perp b \mid c$$

Note: this is the opposite of Example 1, with c observed.


“Am I out of fuel?”

$$p(G = 1|B = 1, F = 1) = 0.8$$

$$p(G = 1|B = 1, F = 0) = 0.2$$

$$p(G = 1|B = 0, F = 1) = 0.2$$

$$p(G = 1|B = 0, F = 0) = 0.1$$



$$p(B = 1) = 0.9$$

$$p(F = 1) = 0.9$$

and hence


$$p(F = 0) = 0.1$$

B = Battery (0=flat, 1=fully charged)

F = Fuel Tank (0=empty, 1=full)

G = Fuel Gauge Reading
(0=empty, 1=full)


“Am I out of fuel?”



$$\begin{aligned} p(F = 0|G = 0) &= \frac{p(G = 0|F = 0)p(F = 0)}{p(G = 0)} \\ &\simeq 0.257 \end{aligned}$$

Probability of an empty tank increased by observing $G = 0$.

“Am I out of fuel?”




$$\begin{aligned} p(F = 0 | G = 0, B = 0) &= \frac{p(G = 0 | B = 0, F = 0)p(F = 0)}{\sum_{F \in \{0,1\}} p(G = 0 | B = 0, F)p(F)} \\ &\simeq 0.111 \end{aligned}$$

Probability of an empty tank reduced by observing $B = 0$.
This referred to as “explaining away”.


D-separation

- A , B , and C are non-intersecting subsets of nodes in a directed graph.
 - A path from A to B is blocked if it contains a node such that either
 - a) the arrows on the path meet either head-to-tail or tail-to-tail at the node, and the node is in the set C , or
 - b) the arrows meet head-to-head at the node, and neither the node, nor any of its descendants, are in the set C .
 - If all paths from A to B are blocked, A is said to be d-separated from B by C .
 - If A is d-separated from B by C , the joint distribution over all variables in the graph satisfies $A \perp\!\!\!\perp B \mid C$.
-

D-separation: Example




$$a \not\perp b \mid c$$



$$a \perp b \mid f$$

D-separation: I.I.D. Data



$$p(\mathcal{D}|\mu) = \prod_{n=1}^N p(x_n|\mu)$$

$$p(\mathcal{D}) = \int_{-\infty}^{\infty} p(\mathcal{D}|\mu)p(\mu) \, d\mu \neq \prod_{n=1}^N p(x_n)$$
