

ADR Identification Basics

Assignment Report – Task 1

Task ID	task1
Subject	Adverse Drug Reaction Identification
Topic	Identification of a suspected drug-related reaction
Prepared by	Student

This report explains the basic process used to identify a possible Adverse Drug Reaction (ADR) by reviewing symptoms, medication history and the time relationship between medicine use and the reaction.

1. Introduction

An Adverse Drug Reaction (ADR) is a harmful and unintended response to a medicine when it is used at normal doses. ADR identification is an important part of clinical pharmacy and pharmacovigilance. A suspected ADR can be identified by checking the patient's symptoms, medicines, time of onset and response after stopping the suspected medicine.

2. Aim

To understand how a possible drug-related adverse reaction is identified from a patient case and how the suspected medicine is documented.

3. Objectives

- To recognize the signs and symptoms of a possible ADR.
- To review the patient's medication history.
- To compare the time of medicine administration with the time of symptom onset.
- To identify the suspected drug responsible for the reaction.
- To understand the importance of reporting and documenting ADRs.

4. Basic Steps for ADR Identification

Step	What is checked?
1	Patient details: age, sex, diagnosis and relevant medical history.
2	Medicine history: name of drug, dose, route, frequency and starting date.
3	Symptoms: type, severity and date/time of onset.
4	Time relationship: whether symptoms appeared after the medicine was started.
5	Dechallenge: whether symptoms improve after withdrawal of the suspected drug.
6	Other causes: disease, food allergy, infection or another medicine.
7	Final assessment: identify the suspected drug and document the reaction.

5. Illustrative Patient Case

A 25-year-old male patient was prescribed Amoxicillin 500 mg orally for a bacterial throat infection. The patient took the medicine as directed. After about two days, he developed red skin rashes and itching over the arms and chest. He had not experienced these symptoms before starting the medicine. After medical advice, the suspected medicine was stopped and the symptoms gradually improved.

6. Case Analysis

Parameter	Finding
Patient	25-year-old male
Medicine	Amoxicillin 500 mg
Indication	Bacterial throat infection
Observed symptoms	Red skin rash and itching
Time of onset	Approximately 2 days after starting the medicine
Dechallenge	Symptoms gradually improved after withdrawal
Suspected reaction	Possible allergic skin reaction
Suspected drug	Amoxicillin

7. Timeline

- Day 0: Amoxicillin was started.
- Day 1: Patient continued the prescribed medicine.
- Day 2: Red rash and itching appeared.
- After withdrawal: Symptoms gradually reduced.

8. Assessment and Interpretation

The symptoms appeared after the medicine was started, and they improved after withdrawal. This time relationship supports Amoxicillin as the suspected drug. The findings are consistent with a possible allergic skin reaction. However, a definite diagnosis requires clinical evaluation and exclusion of other possible causes.

9. Conclusion

ADR identification is performed by reviewing symptoms, medication history, time of onset, possible alternative causes and improvement after withdrawal. In the illustrative case, Amoxicillin was identified as the suspected drug associated with a possible allergic skin reaction. Proper documentation and reporting of suspected ADRs help improve patient safety.

10. Note

This is a student-prepared educational report based on the task description. The patient case used in this report is illustrative and is not a real patient record. It should not be used for diagnosis or treatment decisions.

11. References

- World Health Organization (WHO): Pharmacovigilance and adverse drug reaction monitoring.
- Standard textbooks of Pharmacology and Clinical Pharmacy.
- General principles of adverse drug reaction reporting and patient safety.