Statistics for the severity of head injuries in children

To support our project with numbers, we provided a questionnaire to the community with a set of questions. The questionnaire aimed to gather data on the frequency of falls and head injuries among children in the community. By collecting this information, we can better understand the potential risks and benefits of helmet usage Furthermore, by involving the community in this research, we can foster a senso of ownership and collaboration, ultimately leading to a safer environment for children to thrive in.



Figure 2: Age of children had fall in their head.



Figure 1: Symptoms has occurred to children due to fall.

As you see, 32% of the children who fell needed to visit the hospital, and most of them are 1-3 years old, which is a huge number of young children at risk for serious injuries. These statistics highlight the risk of fall-related injuries among children. Additionally, according to the statistics, children who had fallen in accidents most frequently experienced dizziness and a need for sleep.

The statistics were collected by Faisal Saud Al Harbi, Sultanah Al Mutairi, Wajd Suliman Al Mehaileb

Together they account for 29% of the symptoms. Many participants selected the option Other, and they listed the following symptoms: swelling on the side of the head like water; hallucinations; momentary amnesia; swallowing the tongue; vomiting.



Figure 3: Number of children admitted into the hospital due to fall.

Furthermore, we discovered that 8% of children had strabismus and 45% of children had contusions as a result of a fall accident, while others chose the option" Other" which includes symptoms such as difficulty concentrating, nose fractures, and blood mingling with water.



Figure 4: Statistics of negative results that occurred to children after fall on the head .

The statistics were collected by Faisal Saud Al Harbi, Sultanah Al Mutairi, Wajd Suliman Al Mehaileb



Figure 5: The percentage of the desire of the parents of children for a helmet that combines protection and therapeutic.

As a distinctive addition to our design, we decided to make it not only protective but also therapeutic, and to make sure that our project is effective and desired by the community, we asked in the questionnaire if they prefer to use a helmet that protects the child's head and at the same time treats it or not. The results of the questionnaire revealed that a majority of respondents expressed a strong preference for a helmet that offers both protection and therapeutic benefits for their child's head.

In conclusion, the results of the questionnaire made a big difference in our progress on our project. The positive response from the majority of respondents indicates that there is a clear market need for a helmet that combines protection and therapeutic benefits.

The statistics were collected by Faisal Saud Al Harbi, Sultanah Al Mutairi, Wajd Suliman Al Mehaileb