

UNIVERSITÉ LIBRE DE BRUXELLES

INFO-F-409 - LEARNING DYNAMICS

Assignment One

Raymond Lochner - 000443637

raymond.lochner@ulb.ac.be

Contents

1	The Hawk-Dove game	1
2	Which social dilemma?	1
3	Sequential truel	2
	References	3

October 16, 2017

1 The Hawk-Dove game

The Hawk-Dove game was first formulated by John Maynard Smith and Georg Prince in 1973 [1]. The aim of the game is to gain a better understanding of conflicts in the animal kingdom. It consists of two players {Player One, Player Two} who have each two actions {Hawk, Dove}. The resulting payoff matrix can be seen in Table 1 where:

- V = fitness value of winning resources in fight
- D = fitness costs of injury
- T = fitness costs of wasting time

and we assume that $V, D, T \geq 0$.

Table 1: Hawk-Dove Payoff Matrix

		Player Two	
		Hawk	Dove
Player One	Hawk	$(V-D)/2$ / $(V-D)/2$	0 / V
	Dove	0 / V	$V/2-T$ / $V/2-T$

In a mixed strategy game, we consider each player performing his action with a certain probability p , which results in the following payoff matrix displayed in Table 2.

Table 2: Hawk-Dove Probability Payoff Matrix

		Player Two	
		$P(\text{Hawk}) = q$	$P(\text{Dove}) = 1-q$
Player One	$P(\text{Hawk}) = p$	$(V-D)/2$ / $(V-D)/2$	0 / V
	$P(\text{Dove}) = 1-p$	0 / V	$V/2-T$ / $V/2-T$

2 Which social dilemma?

Player A is confronted with one of three social dilemma's - the corresponding payoff matrix is shown in tables 3, 4 and 5. The player has to decide whether to cooperate (C) or to defect (D) without knowing which game he is actually facing. Each dilemma has the same probability $1/3$ of being played. Opponent B knows the game.

Table 3: Prisoners dilemma

	C	D
C	2 / 2	5 / 0
D	0 / 5	1 / 1

Table 4: Stag-Hunt game

	C	D
C	5 / 5	2 / 0
D	0 / 2	1 / 1

Table 5: Snowdrift game

	C	D
C	2 / 2	5 / 1
D	1 / 5	0 / 0

with this information we can calculate the expected payoff for player A for every possible strategy of Player B - Table 6.

From this table we can select the best response for Player A for each strategy of Player B - cells marked red in Table 7.

Table 6: Expected payoff for Player A

		Player B							
		(C,C,C)	(C,C,D)	(C,D,C)	(C,D,D)	(D,C,C)	(D,C,D)	(D,D,C)	(D,D,D)
Player A	C	9/3	8/3	4/3	3/3	7/3	6/3	2/3	1/3
	D	12/3	7/3	11/3	6/3	8/3	3/3	7/3	2/3

Table 7: Best responses for Player A

		Player B							
		(C,C,C)	(C,C,D)	(C,D,C)	(C,D,D)	(D,C,C)	(D,C,D)	(D,D,C)	(D,D,D)
Player A	C	9/3	8/3	4/3	3/3	7/3	6/3	2/3	1/3
	D	12/3	7/3	11/3	6/3	8/3	3/3	7/3	2/3

Now we have to determine the best responses of Player B against Player A of the three different strategies - marked by green cells in Tables 8, 9 and 10.

Table 8: Prisoners dilemma

	C	D
C	2	5
D	0	1

Table 9: Stag-Hunt game

	C	D
C	5	2
D	0	1

Table 10: Snowdrift game

	C	D
C	2	5
D	1	0

The pure strategy Nash Equilibria can now be determined by matching these results. We find two Nash Equilibria at $\{C,(D,C,D)\}$ and $\{D,(D,D,C)\}$.

3 Sequential truel

References

- [1] J. Maynard Smith and G. R. Price. The logic of animal conflict. *Nature*, 246(5427):15–18, 1973.