

MASARYKOVA UNIVERZITA
PŘÍRODOVĚDECKÁ FAKULTA
ÚSTAV TEORETICKÉ FYZIKY A ASTROFYZIKY

Bakalářská práce

BRNO 2018

MILAN SUK

Aspekty přechodu od kvantové mechaniky ke kvantové teorii pole

Bakalářská práce

Milan Suk

Bibliografický záznam

Autor:	Milan Suk Přírodovědecká fakulta, Masarykova univerzita Ústav teoretické fyziky a astrofyziky
Název práce:	Aspekty přechodu od kvantové mechaniky ke kvantové teorii pole
Studijní program:	Fyzika
Studijní obor:	Fyzika
Vedoucí práce:	doc. Klaus Bering Larsen, Ph.D
Akademický rok:	2017/2018
Počet stran:	?? + ??
Klíčová slova:	kvantová mechanika, kvantová teorie pole, relativistická kvantová mechanika

Bibliographic Entry

Author:	Milan Suk Faculty of Science, Masaryk University Department of physics and astrophysics
Title of Thesis:	Aspects of the transition from quantum mechanics to quantum field theory
Degree Programme:	Physics
Field of Study:	Physics
Supervisor:	doc. Klaus Bering Larsen, Ph.D
Academic Year:	2017/2018
Number of Pages:	?? + ??
Keywords:	quantum mechanics; quantum field theory, rela- tivistic quantum mechanics

Abstrakt

V této bakalářské/diplomové/rigorózní práci se věnujeme ...

Abstract

In this thesis we study ...

Místo tohoto listu vložte kopii oficiálního (podepsaného) zadání práce.

Poděkování

Na tomto místě bych chtěl(-a) poděkovat ...

Prohlášení

Prohlašuji, že jsem svoji bakalářskou/diplomovou/rigorózní práci vypracoval(-
a) samostatně s využitím informačních zdrojů, které jsou v práci citovány.

Brno 1. května 2018

.....
Milan Suk

Contents

Nomenclature	xv
Introduction	1
Kapitola 1 Relativistic quantum mechanics	3
1.1 Canonical quantization	3
1.2 Klein-Gordon equation	4
1.3 Dirac equation	5
Kapitola 2 Failures of the Relativistic Quantum Mechanics	7
2.1 Probability density	7
2.2 Causality	8
Kapitola 3 Quantum field theory	9
3.1 Scalar field	9
3.2 Second quantization	9
3.3 Causality problem	9
Conclusion	11
Bibliography	13

Nomenclature

- \mathbb{C} množina všech komplexních čísel
- \mathbb{R} množina všech reálných čísel
- \mathbb{Z} množina všech celých čísel

Introduction

The Quantum Mechanics (QM) is a fundamental theory in physics which aims to describe the world at the lowest scales of measured units. It had a great success of explaining a lot of new observations in the very beginning of 20th century, e.g. spectra of atoms, the black-body radiation or photoelectric effect. In 1905, Albert Einstein published his famous article *On the Electrodynamics of Moving Bodies* and proposed the Special theory of relativity (STR). STR corrects the classical mechanics in a way that it can predict a state of a physical system having *relativistic velocities* and setups a new restrictions for mathematical formulations of physical theories.

A logical step is to develop a physical theory which preserves STR and is capable of describing particles in quantum mechanics. This theory is known as Relativistic Quantum Mechanics (RQM) and it has been successful in prediction of interesting phenomena in physics like antimatter or spin. Nevertheless, RQM can't be introduced without formal inconsistencies and it cant deal with varying number of particles. The modern framework which consistently combines QM and STR is the Quantum Field Theory (QFT).

The thesis consists of three main chapters. In the first one we will discuss the formulation of RQM, i.e. we will construct two relativistic quantum mechanical eqautions - the Klein-Gordon equation and the Dirac equation. In the second chapter we will focus on the failures of these formulations and we show the need for a new theory. In the last chapter we will briefly introduce the new framework of QFT. In the terms of Quantum scalar fields we will try to check whether the new formulation solves the problems described in the second chapter.

Chapter 1

Relativistic quantum mechanics

In this chapter (TODO)

1.1 Canonical quantization

Canonical quantization is a process of transition from a classical theory to the quantum one, i.e. procedure for quantizing the theory. The term *canonical* comes from (TODO).

Definition 1. Given two functions $f(\vec{q}, \vec{p}, t)$ and $g(\vec{q}, \vec{p}, t)$, the binary operation $\{f, g\}$ is called the **Poisson bracket** of these functions and takes the form

$$\{f, g\} = \sum_{i=1}^N \left(\frac{\partial f}{\partial q_i} \frac{\partial g}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q_i} \right) \quad (1.1)$$

Theorem 1. The Poisson brackets of canonical coordinates q_i and p_i are

$$\begin{aligned} \{q_i, q_j\} &= 0 \\ \{p_i, p_j\} &= 0 \\ \{q_i, p_j\} &= \delta_{ij} \end{aligned} \quad (1.2)$$

Proof. The proof of 1 can be easily done from the definition 1. The first two equations are manifestly zero because $\frac{\partial q_i}{\partial p_j} = \frac{\partial p_i}{\partial q_j} = 0$ and the last can one be proved as follows.

$$\begin{aligned} \{q_i, p_j\} &= \sum_{k=1}^N \left(\frac{\partial q_i}{\partial q_k} \frac{\partial p_j}{\partial p_k} - \frac{\partial q_i}{\partial p_k} \frac{\partial p_j}{\partial q_k} \right) = \sum_{k=1}^N \frac{\partial q_i}{\partial q_k} \frac{\partial p_j}{\partial p_k} = \begin{cases} 1 & \text{if } i = j = k \\ 0 & \text{otherwise} \end{cases} \\ &\implies \{q_i, p_j\} = \delta_{ij} \end{aligned} \quad (1.3)$$

□

Dirac has formulated a technique for generating quantum operators from classical functions. The famous statement goes as follows.

$$\{A, B\} \rightarrow \frac{1}{i\hbar}[\hat{A}, \hat{B}] \quad (1.4)$$

For the next discussion let's define \mathcal{O} map which takes a function from the configuration space and maps it to the Hilbert space.

Definition 2. Let \mathcal{Q} be a configuration space and H a Hilbert space. $End(V)$ is an endomorphism of space V , i.e. a set of linear operators $\{L : V \rightarrow V \mid L \text{ linear}\}$.

$$\begin{aligned} \mathcal{O} : \mathcal{Q} &\rightarrow End(H) \\ \mathcal{O}(\{f, g\}) &= \frac{1}{i\hbar}[\hat{A}, \hat{B}] \end{aligned} \quad (1.5)$$

Where \hat{A} is the operator corresponding the the classical value f and B corresponds to g .

Now we will use the fact that $[\hat{q}_i, \hat{p}_j] = i\hbar\delta_{ij}$ to show that $\mathcal{O}(q_i) = \hat{q}_i$. Firstly, we will rewrite q using the Poisson bracket. The obvious choice is $q_i = \{\frac{q_i^2}{2}, p_i\}$. And from the definition 2 we see that $\mathcal{O}(\{\frac{q_i^2}{2}, p_i\}) = \frac{1}{i\hbar}[\frac{\hat{q}_i^2}{2}, \hat{p}_i]$. Using the identity for commutation relation $[AB, C] = [A, B]C + B[A, C]$ ¹ the commutator on the right-hand side can be written as follows.

$$\mathcal{O}(x_i) = \frac{1}{i\hbar} \left[\frac{\hat{x}_i^2}{2}, \hat{p}_i \right] = \frac{1}{2i\hbar} (\hat{x}_i[\hat{x}_i, \hat{p}_i] + [\hat{x}_i, \hat{p}_i]\hat{x}_i) = \frac{1}{2i\hbar} (\hat{x}_i i\hbar + i\hbar \hat{x}_i) = \hat{x}_i$$

The proof of $\mathcal{O}(p) = \hat{p}$ is very similar.

$$\mathcal{O}(p_i) = \frac{1}{i\hbar} \left[\hat{x}_i, \frac{\hat{p}_i^2}{2} \right] = \frac{1}{2i\hbar} ([\hat{x}_i, \hat{p}_i]\hat{p}_i + \hat{p}_i[\hat{x}_i, \hat{p}_i]) = \frac{1}{2i\hbar} (i\hbar\hat{p}_i + \hat{p}_i i\hbar) = \hat{p}_i$$

In the x -representation the explicit form of \hat{x} and \hat{p} is $\hat{x} = x$ and $\hat{p}_x = -i\hbar\frac{\partial}{\partial x}$.
TODO: study $\mathcal{O}(\{f(\vec{q}, \vec{p})\})$ and $\mathcal{O}(\{f + g, h\})$. How to get $\mathcal{O}(E)$?

1.2 Klein-Gordon equation

Klein-Gordon equation is relativistic quantum mechanical equation named after Oskar Klein and Walter Gordon and it describes relativistic spinless particles. It can be found using quantization of the relativistic energy equation. We need to apply the \mathcal{O} map on each side while taking into account that $\mathcal{O}(E^2) = -\hbar^2 \frac{\partial^2}{\partial t^2}$ and $\mathcal{O}(\vec{p}^2) = -\hbar^2 \Delta$.

¹Proof by directly using the definition of the commutator: $[AB, C] = ABC - CAB = ABC - ACB + ACB - CAB = A[B, C] + [A, C]B$

$$E^2 = \vec{p}^2 c^2 + m^2 c^4$$

$$-\hbar^2 \frac{\partial^2}{\partial t^2} \psi = -\hbar^2 c^2 \Delta \psi + m^2 c^4 \psi$$

After a little rearranging and setting $\frac{m^2 c^2}{\hbar^2} = \mu^2$, we will finally get the famous form 1.6.

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \Delta + \mu^2 \right) \psi = 0 \quad (1.6)$$

Let's rewrite the equation in a form so we can see it is *lorentz invariant*. We use the fact that *4-gradient* transforms the same way as any other 4-vector and then express the Klein-Gordon equation using 4-gradient dot products. Using the metric $\eta^{\mu\nu} = \text{diag}(1, -1, -1, -1)$ and the *4-gradient* $\partial_\mu = (\frac{\partial}{\partial t}, \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z})$ it is possible to write the term $\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \Delta$ as follows.

$$\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} = \eta^{\mu\nu} \partial_\mu \partial_\nu = \partial_\mu \partial^\mu$$

$$\left(\partial_\mu \partial^\mu + \mu^2 \right) \psi = 0 \quad (1.7)$$

1.3 Dirac equation

One important fact about the Klein-Gordon equation is that it is second-order in time. Thus we need one more initial condition compared to the Schrodinger Equation. Dirac tried to find an equation that is first order in both space and time by finding the *square root* of the Klein-Gordon equation. Let's start with the equation 1.6.

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \Delta + \mu^2 \right) \psi = 0$$

Chapter 2

Failures of the Relativistic Quantum Mechanics

In this chapter we will discuss the problems of relativistic quantum mechanics. We will start with the problem of probability density within the Klein-Gordon equation.

2.1 Probability density

In the non-relativistic case we can find the continuity equation by taking the complex conjugate of the Schrodinger's equation.

$$\begin{aligned}i\hbar \frac{\partial}{\partial t} \psi &= \hat{H} \psi \\ -i\hbar \frac{\partial}{\partial t} \psi^* &= (\hat{H} \psi)^*\end{aligned}$$

Then we multiply the first equation by ψ^* and the second one by ψ .

$$\begin{aligned}i\hbar \psi^* \frac{\partial}{\partial t} \psi &= \psi^* \hat{H} \psi \\ i\hbar \psi \frac{\partial}{\partial t} \psi^* &= \psi (\hat{H} \psi)^*\end{aligned}$$

If we subtract these two equations from each other and use the formula for derivative $(a \cdot b)' = a' \cdot b + a \cdot b'$ we will get a form from which it will be easy to identify the continuity equation $\frac{\partial \rho}{\partial t} + \text{div} \vec{j} = 0$.

$$i\hbar \frac{\partial |\psi|^2}{\partial t} = \psi^* \hat{H} \psi - \psi (\hat{H} \psi)^*$$

We assume the potential V in $\hat{H} = \frac{\hat{p}^2}{2m} + V$ to be real-valued thus the right-hand side will simplify to

$$\psi^* \frac{\hbar^2}{2m} \nabla^2 \psi - \psi \frac{\hbar^2}{2m} \nabla^2 \psi^* = \frac{\hbar^2}{2m} \left(\psi^* \nabla^2 \psi - \psi \nabla^2 \psi^* \right)$$

and the desired form is

$$\frac{\partial |\psi|^2}{\partial t} + \text{div} \left[\frac{\hbar}{2mi} \left(\psi \nabla \psi^* - \psi^* \nabla \psi \right) \right] = 0$$

Now we will do the same for the Klein-Gordon equation 1.6. Let's do the same procedure here and find the complex conjugate, multiply both equations by ψ and ψ^* and then subtract them. After that let's focus on the term which contains the time derivative and thus will represent the *probability density*.

$$\frac{\partial \rho}{\partial t} \sim \frac{1}{2i} \left[\psi^* \frac{\partial^2}{\partial t^2} \psi - \psi \frac{\partial^2}{\partial t^2} \psi^* \right]$$

Now we can find what ρ is proportional to.

$$\rho \sim \frac{1}{2i} \left[\psi^* \frac{\partial}{\partial t} \psi - \psi \frac{\partial}{\partial t} \psi^* \right] \quad (2.1)$$

Let's divide the wave function ψ into the real and imaginary part.

$$\begin{aligned} \psi &= i\psi_1 + \psi_2 \\ \rho &\sim \left[\psi_2 \frac{\partial}{\partial t} \psi_1 - \psi_1 \frac{\partial}{\partial t} \psi_2 \right] \end{aligned}$$

Now it is easy to see the problem. In the non-relativistic case the probability density was simply $|\psi|^2$ which is non-negative for any $\psi : \mathbb{R} \rightarrow \mathbb{C}$. In the case of Klein-Gordon equation, we can see the probability density is zero whenever the wave function is either exclusively real or imaginary. Moreover, since both $\frac{\partial \psi_1}{\partial t}$ and $\frac{\partial \psi_2}{\partial t}$ are arbitrary, the value can be negative. This is in contradiction with the interpretation of ρ as the *probability density of measuring the system within a given state ψ* .

2.2 Causality

Chapter 3

Quantum field theory

3.1 Scalar field

In non-relativistic case we can find the continuity equation by

3.2 Second quantization

3.3 Causality problem

Conclusion

Bibliography

- [1] S. J. Monaquel a K. M. Schmidt, *On M -functions and operator theory for non-self-adjoint discrete Hamiltonian systems*
- [2] Shewell, J. R. (1959). *On the Formation of Quantum-Mechanical Operators*. *American Journal of Physics*

