

# UV Spectral Characterization of Low-Mass Stars With AstroSat UVIT for Exoplanet Applications: HIP 23309 Case Study

SUKRIT RANJAN,<sup>1</sup> PRASANTA K. NAYAK,<sup>2</sup> J. SEBASTIAN PINEDA,<sup>3</sup> AND MAYANK NARANG<sup>4,2</sup>

<sup>1</sup>*University of Arizona, Lunar and Planetary Laboratory/Department of Planetary Sciences, Tucson, 85721, USA\**

<sup>2</sup>*Department of Astronomy and Astrophysics, Tata Institute for Fundamental Research, Mumbai, 400005, India*

<sup>3</sup>*University of Colorado-Boulder, Boulder, 80309, USA*

<sup>4</sup>*Academia Sinica Institute of Astronomy & Astrophysics, 11F of Astro-Math Bldg., No.1, Sec. 4, Roosevelt Rd., Taipei 10617, Taiwan, R.O.C.*

(Received January 14, 2023)

Submitted to AJ

## ABSTRACT

Characterizing rocky exoplanet atmospheres is a key goal of exoplanet science, but interpreting such observations will require understanding the stellar UV irradiation incident on the planet from its host star. Stellar UV mediates atmospheric escape, photochemistry, and planetary habitability, and observations of rocky exoplanets can only be understood in the context of the UV SED of their host stars. Particularly important are SEDs from observationally favorable but poorly understood low-mass M-dwarf stars, which are the only plausible targets for rocky planet atmospheric characterization for the next 1-2 decades. In this work, we explore the utility of ASTROSAT UVIT for the characterization of the UV SEDs of low-mass stars. We present observations of the nearby M0 star HIP 23309 in the FUV and NUV grisms of UVIT. Our FUV spectra are consistent with contemporaneous HST data and our NUV spectra are stable between orbits, suggesting UVIT is a viable tool for the characterization of the SEDs of low-mass stars. We apply our measured spectra to simulations of photochemistry and habitability for a hypothetical rocky planet orbiting HIP 23309 and elucidate the utility and limitations of UVIT in deriving UV SEDs of M-dwarf exoplanet hosts. Our work validates UVIT as a tool to complement HST in the characterization of exoplanet host stars and carries implications for its successor missions like INSIST.

*Keywords:* M dwarf stars (982), Exoplanet atmospheric composition (2021), Ultraviolet telescopes (1743), Astrobiology (74), Habitable planets (695), Spectral energy distribution (2129)

## 1. INTRODUCTION

Temperate, rocky planets orbiting other stars are abundant, and upcoming facilities will have the ability to characterize their atmospheres (Dressing & Charbonneau 2015; Bryson et al. 2021; Rodler & López-Morales 2014; Morley et al. 2017; Lustig-Yaeger et al. 2019). Characterization of rocky exoplanet atmospheres will probe their surface processes including potentially constraining the presence or absence of life (Schwieterman et al. 2018), and such constraints are high priorities for the astronomical community (National Academies of Sciences Engineering and Medicine 2018; of Sciences Engineering & Medicine 2021). However, interpreting measurements of rocky exo-atmospheres requires understanding the UV spectral energy distribution (SED) of their host stars. Stellar UV is a first-order control on the surface-atmosphere system of rocky planets, and mediates atmospheric escape, photochemistry, and habitability, meaning that observations of rocky exo-atmospheres can only be understood in the context of the SED of their host

Corresponding author: Sukrit Ranjan  
sukrit@arizona.edu

\* Blue Marble Space Institute of Science, Seattle, 98154, USA

**Table 1.** HIP 23309 Stellar Parameters

Parameter	Value	Reference
Spectral Type	M0	Torres et al. (2006); Pineda et al. (2021)
Effective Temperature	$3886^{+28}_{-27}$ K	Pineda et al. (2021)
Mass	$0.785^{+0.009}_{-0.010} M_{\odot}$	Pineda et al. (2021)
Radius	$0.932 \pm 0.014 R_{\odot}$	Pineda et al. (2021)
Luminosity	$6.82^{+0.12}_{-1.1} \times 10^{32}$ erg s <sup>-1</sup>	Pineda et al. (2021)

star (Teal et al. 2022). This observation has motivated observational and theoretical work aimed at constraining the high-energy emission of exoplanet host stars (e.g., France et al. 2013; Viswanath et al. 2020).


Characterization of the UV emission of low-mass ( $0.08 - 0.6 M_{\odot}$ ) M-dwarf stars is of particular importance to the quest to characterize rocky exo-atmospheres. Most stars are M-dwarfs, and planets orbiting M-dwarfs are uniquely accessible to characterization (Scalo et al. 2007; Shields et al. 2016). Consequently, M-dwarf planets are the *only* plausible targets for in-depth atmospheric characterization and biosignature search with near-term facilities such as JWST and the ELTs (Cowan et al. 2015; Morley et al. 2017; Batalha et al. 2018; Tremblay et al. 2020). However, M-dwarf UV emission is very different from the UV emission of higher-mass stars. These differences are predicted to impact atmospheric escape, atmospheric photochemistry, and habitability relative to Sun-like stars (Segura et al. 2005; Ramirez & Kaltenegger 2014; Luger & Barnes 2015; Rugheimer et al. 2015; Dong et al. 2017; Rimmer et al. 2018; Wordsworth et al. 2018; Harman et al. 2018; Tilley et al. 2019; Schwieterman et al. 2022)), motivating observational efforts to characterize the SEDs of M-dwarf stars (e.g., France et al. 2016; Peacock et al. 2019a; Pineda et al. 2021). Such observations must be done from space due to telluric absorption, and to-date are primarily executed on the Hubble Space Telescope (HST) (e.g., France et al. 2016). However, the energetic emission of M-dwarfs remains underconstrained, in part because of the need to use extremely scarce HST time, and in part, because the considerable diversity of M-dwarfs necessitates a large number of observations to constrain their emission as a function of parameters such as mass and age (Lloyd et al. 2021; Pineda et al. 2021). Efforts have been made to predict M-dwarf UV SEDs from more tractably-measured emission lines (Youngblood et al. 2017; Melbourne et al. 2020), but the use of such approximations may lead to significant inaccuracies in models of rocky exoplanet atmospheres (Teal et al. 2022). It remains preferred to characterize the UV SEDs of planet-hosting M-dwarf stars to accurately model the atmospheres of their orbiting planets (Teal et al. 2022).

The UltraViolet Imaging Telescope (UVIT) instrument on the ISRO Astronomy Satellite (AstroSat) mission is in principle well-suited to characterization of the UV SEDs of M-dwarf stars. UVIT is composed of two 38-cm telescopes with a field of view of  $28'$  in diameter. One telescope is dedicated for observations in the FUV region (130-180 nm), while the other one is for NUV (200-300 nm) and visible regions (320-550 nm). The UVIT operates in photon counting mode and observes simultaneously with both the telescopes with a spatial resolution (FWHM)  $< 1.''5$  (Tandon et al. 2020), better than the resolution of GALEX ( $\sim 5''$ ). Both FUV and NUV channels incorporate multiple narrow to broad-band filters. UVIT also has a facility for low-resolution slit-less spectroscopy in both NUV and FUV bands: one grating is mounted in the NUV wheel (now inoperative) and two gratings (with orthogonal dispersions axis) in the FUV wheel. The maximum efficiency is achieved in the  $-2$  order of the FUV gratings with a spectral resolution of  $15\text{\AA}$  and  $-1$  order of the NUV grating with a spectral resolution of  $33\text{\AA}$  (Tandon et al. 2020; Dewangan 2021).

In this paper, we explore the utility of UVIT to constrain the SEDs of M-dwarf stars using HIP 23309 as a case-study. HIP 23309 is a young, rapidly rotating, early-type M-dwarf star (Pineda et al. 2021; Table 1). We secure spectrally resolved observations of HIP 23309 in the UVIT FUV (130 – 180 nm) and NUV (200 – 300 nm) grisms. Standard UVIT calibration pipelines are able to extract resolved spectra of HIP 23309 with no modification (Dewangan 2021). We compare our FUV data to contemporaneous HST measurements, and show excellent agreement. We perform atmospheric and surficial photochemical modelling to elucidate the utilities and limitations of such measurements for studying the atmospheres of rocky planets and to assessing their habitability. We close by commenting on the potential of UVIT for ongoing studies of M-dwarfs and their exoplanets, and recommendation for future facilities like INSIST and UV-SCOPE.

## 2. OBSERVATIONS AND DATA REDUCTION

We observed HIP 23309 with the FUV-Grating1 (FUV-G1) and NUV-Grating of UVIT as a part of AO-1 proposal (PI: Lalitha Sairam) on 24-11-2017 with an exposure time of  $\sim 12$  ksec in each channel. The observations were completed in multiple orbits. We applied corrections for spacecraft drift, flat-field and distortion, using the software CCDLAB (Postma & Leahy 2017) and created images for each orbit. Then, the orbit-wise images were co-aligned and combined to generate final science ready images in each grating. Astrometry on the images were also performed using the CCDLAB by comparing the Gaia-DR2 source catalogue. The zoomed-in FUV and NUV grating images are shown in the Figure 1, with boxes indicating the different spectral orders. We extracted the source spectrum of  $-2$  order from FUV-G1 image and  $-1$  order from the NUV image with a cross-dispersion width of 50 pixels (1 pixel  $\sim 0.''41$ ). Background regions are selected exactly in the same pixel range above the source spectrum along the dispersion axis for both the gratings. We used the UVITTools package written in the Julia language by Dewangan 2021 to extract spectral files for a given grating order and cross-dispersion width. Wavelength and flux calibration to the spectra are done using standard UVIT source calibrators: HZ4 and NGC40. We show the NUV and FUV spectra of HIP 23309 in Figure 1.



**Figure 1.** Images of HIP 23309 in FUV-Grating1 (left) and NUV-Grating (right) are shown here. Different grating order are marked in green.

## 3. MEASURED SPECTRA OF HIP 23309

### 3.1. UVIT Measurements of HIP23309

In the NUV spectra (Fig.2), we can clearly identify three peaks: one at  $2800 \text{ \AA}$  due to strong Mg II *h* and *k* lines, and the other two peaks at  $2400 \text{ \AA}$  and  $2600 \text{ \AA}$  due to strong Fe II lines. The strong Mg II *h* and *k* lines emissions indicate that the star is chromospherically active (Blanco et al. 1979; Doherty 1985). In the FUV spectra (Fig.2), comparative poor signal-to-noise (SNR) makes it difficult to identify emission lines. We can clearly identify the strong C IV line at  $1560 \text{ \AA}$  despite low SNR, however, the identification of He II and O III are less reliable for further analysis.


### 3.2. Robustness of UVIT M-dwarf Spectra

We next assess the accuracy of our spectra of HIP 23309 from UVIT. For the far-UV wavelength regime, we are able to directly compare our UVIT observations to contemporaneous FUV measurements collected by HST within 1 day of this work. (Table 2). For the near-UV data, we evaluate the orbit-by-orbit stability of our measured spectrum.

#### 3.2.1. FUV Spectrum: Comparison to HST Data

We coordinated our UVIT observations of HIP 23309 with HST STIS observations of HIP 23309 collected as part of the FUMES survey (PI: J. S. Pineda; Pineda et al. 2021; Youngblood et al. 2021). The spectral coverage of the UVIT FUV grism and the HST STIS overlap from 130-170 nm, enabling us to compare spectra derived from the two facilities.

The FUV UVIT and STIS measurements of HIP 23309 broadly agree to within a factor of  $\sim 2$ , but the UVIT measurements systematically exceed the STIS measurements (Figure 3). The 130-170 nm flux measured by UVIT is  $(5.62 \pm 0.34) \times 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$ , compared to  $(4.02 \pm 0.04) \times 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$ . This corresponds to a 40%



**Figure 2.** UVIT FUV (left) and NUV (right) spectra of HIP23309. Multiple NUV lines are detected; detections of FUV lines are much weaker, with only C IV being clearly detected.

**Table 2.** Summary of HIP 23309 Observations Discussed in this Work

Facility	Instrument	Grism	Orbits	Observation Start Time (UT)	Total Integration Time	Reference
HST	STIS	G140L (115-173 nm)	1	2017-11-24 01:15:22	2350 s	Pineda et al. (2021)
AstroSat	UVIT	FUV (130-180 nm) <sup>a</sup>	7	2017-11-24 10:16:44	12027 s	This Work
AstroSat	UVIT	NUV (200-300 nm) <sup>a</sup>	7	2017-11-24 10:16:44	12027 s	This Work

<sup>a</sup>Tandon et al. (2017)


overestimate by UVIT and  $4.63\sigma$  tension relative to STIS. To compare the STIS data to the UVIT data `spectrally`, we (1) convolve the STIS data against a Gaussian with full-width at half-max of 14.63, corresponding to the resolution of the UVIT FUV grating (Dewangan 2021), and (2) rebin the smoothed spectrum to the same wavelength scale as the UVIT data using the `coronagraph` package and conserving total SNR in each bin (Robinson et al. 2016; Lustig-Yaeger et al. 2019). The synthetic spectrum derived from the HST data is in tension with the UVIT spectrum by an average of  $1.46\sigma$  (Figure 3). The disagreement is strongest in the continuum region between lines, where HIP 23309 emits very little flux ( $3.10 \pm 0.09 \times 10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$  in FUV; Pineda et al. 2021). For example, integrated from 1410-1510 Å, the flux measured by UVIT is  $5.5 \pm 1 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$ , whereas the flux measured by HST is  $3.39 \pm 0.1 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$ , a tension of  $1.9\sigma$ . The agreement between HST and UVIT is better for major lines, i.e. CIV. The CIV line emission from UVIT is  $9.8 \pm 3.5 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ , compared to  $6.9 \pm 0.2 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$  (Pineda et al. 2021). We conclude that the FUV UVIT data broadly reproduce the HST data to within a factor of 2, but that there is the potential for uncorrected systematic noise in the UVIT data, including potentially a noise floor of  $\sim 2 \times 10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$ .

### 3.2.2. NUV Spectrum: Stability of Measurement


The SNR of the NUV data is high enough to support orbit-by-orbit decomposition (Figure 4). The band-integrated (220-290 nm) orbit-by-orbit NUV data are consistent with the mean to within 10%, but vary from it by as much as  $8.4\sigma$ . This suggests that the measurement errors are underestimated, e.g. due to an uncharacterized source of red noise. Indeed, the orbit-by-orbit band-integrated fluxes display a weak increase with orbit number. Comparing the orbit-by-orbit data to the mean on a spectral basis, we find that the average deviation from the mean  $A$ :

$$A = \frac{1}{N} \sum_{i=0}^N \frac{F_i - \bar{F}_i}{\sqrt{\sigma_i^2 + \bar{\sigma}_i^2}} \quad (1)$$

has values ranging from 1.83 to 3.74. This also indicates that the errors are underestimated by a factor of a few. Alternately, our measurements may reflect variations in stellar flux due to activity. We do not detect evidence of large flares (variation in total flux of  $\geq 2\times$ ).



**Figure 3.** Comparison between HST and AstroSat-measured SEDs of HIP 23309 in the FUV (1300-1700 Å). The cyan and black data correspond to the unbinned HST and AstroSat data, respectively. The blue points correspond to HST data degraded to the approximate quality of the AstroSat data, to enable a closer comparison. The green points correspond to the AstroSat data binned down by a factor of 5, to improve SNR and enable better visual comparison. We assumed Gaussian uncorrelated noise when rebinning the AstroSat data.




**Figure 4.** Orbit-by-orbit NUV spectral (left) and band-integrated (right) flux of HIP23309 as measured by UVIT. The band-integrated orbit-by-orbit fluxes agree with the mean flux to within 10%, but are in statistically significant tension with the mean. This indicates that either our errors are underestimated, or that there is stellar variability at the  $\sim 10\%$  level.

### 3.3. HIP23309 SED in Context



We compare our FUV and NUV spectra of HIP23309 to spectra of other low-mass stars to assess whether our measurements are sensible in context (Figure 5). We specifically compare the spectral flux at the stellar surface of HIP23309 to MUSCLES M-dwarf and K-dwarf spectra (v2.2, France et al. 2016; Youngblood et al. 2016; Loyd et al. 2016), and to StarCAT measurements of AU Mic in the FUV (Ayres 2010). The MUSCLES stars are mature, whereas AU Mic is a young pre-main sequence (22 Myr, Mamajek & Bell 2014; Plavchan et al. 2020) early M-dwarf star similar to HIP 23309. HIP23309’s FUV emission is similar to AU Mic’s, which is sensible in that AU Mic is of similar spectral type and age to HIP 23309. HIP23309 is brighter in the FUV and NUV compared to main-sequence M-dwarfs of similar spectral type, consistent with the finding that pre-main sequence M-dwarfs display elevated UV emission compared to main-sequence M-dwarfs due to enhanced stellar activity and larger radii (Shkolnik & Barman 2014; Luger & Barnes 2015). At longer wavelengths in the NUV, the HIP 23309 stellar flux begins to converge with main-sequence stars, which is consistent with the increasing importance of photospheric emission at longer wavelengths. We conclude that to zeroth order, our measured NUV and FUV spectra of HIP 23309 are sensible in the context of other measured spectra of low-mass stars.



**Figure 5.** Stellar flux of HIP23309 at the stellar surface ( $F_{\star} \times \frac{d_{\star}^2}{r^2}$ ) in the FUV (left) and NUV (right), compared to other low-mass stars. HIP23309’s stellar flux is much higher than mature stars of similar spectral type at short wavelengths due to enhanced activity because of its young age, but converges with them at longer wavelengths where photospheric emission dominates. HIP 23309’s flux in the FUV is similar to AU Mic’s, another extremely young M-dwarf of similar spectral class. Overall, our spectra of HIP23309 appears sensible in the context of other measured spectra of low-mass stars.

#### 4. APPLICATIONS OF UV SEDS TO ROCKY EXOPLANET MODELING

In this section, we explore the implications of our measured SED of HIP 23309 for a hypothetical temperate terrestrial planet orbiting HIP 23309. Our intent is to illustrate the utility of M-dwarf SEDs for studying rocky exoplanets, and particularly for questions of astrobiological interest.

##### 4.1. Implications of HIP23309 SED for UV-Dependent Prebiotic Chemistry

UV light has been proposed necessary to the origin of life, but the surfaces of habitable M-dwarf planets are low-UV environments (Ranjan et al. 2017). Rimmer et al. (2018) showed that HCN photohomologation chemistry used in proposed prebiotic pathways for ribonucleotide synthesis (Xu et al. 2018; Xu et al. 2020; Rimmer et al. 2021b) require  $F_{200-280nm} \geq (6.8 \pm 3.6) \times 10^{10} \text{ cm}^{-2} \text{ s}^{-1}, \text{ nm}^{-1}$  to function, where  $F_{200-280nm}$  is the surface flux integrated from 200-280 nm. This means that if HCN photohomologation is required for the emergence of life, then life cannot emerge on planets which have access to less than this critical flux of UV. Rimmer et al. (2018) find that the quiescent emission from known M-dwarfs is insufficient to meet this criterion, and that HCN photohomologation chemistry can only proceed on planets orbiting M-dwarfs if enabled by transiently enhanced UV from flares. Efforts to experimentally

determine if flare UV can substitute for quiescent UV are now under way<sup>1</sup>; if they cannot, then life search on exoplanets may constrain the pathways towards origin of life (Ranjan et al. 2017; Rimmer et al. 2018).

We consider whether the UV emission from HIP 23309 meets the Rimmer et al. (2018) criterion. We assembled an FUV-NIR SED of HIP 23309 by concatenating our UVIT-measured FUV and NUV fluxes to the low-resolution GAIA spectrum of HIP 23309 (Gaia Collaboration et al. 2022). We scale these observed fluxes to the flux that would be experienced by a hypothetical Earth-like planet orbiting HIP 23309 by multiplying them by  $(\frac{d_*}{a})^2$ , where  $d_* = 26.8$  pc is the distance to HIP 23309 (Malo et al. 2014),  $a = 1\text{AU} \times (\frac{L_*}{L_\odot * 0.9})^{0.5}$  is the distance such that the bolometric flux received at distance  $a$  is equivalent to the bolometric flux received by Earth, corrected by  $0.9\times$  to account for the redder emission from M-dwarf stars (Segura et al. 2005), and  $L_* = 6.82 \times 10^{32} \text{ erg s}^{-1}$  (Pineda et al. 2021). We then calculated the hemispherically-integrated surface actinic flux ("surface radiance") for an early Earth-like planet with a 0.9 bar N<sub>2</sub>, 0.1 bar CO<sub>2</sub> atmosphere as in Ranjan et al. (2017). We find that HIP 23309 comfortably meets the Rimmer et al. (2018) criterion, with  $F_{200-280\text{nm}} = 3 \times 10^{11} \text{ cm}^{-2} \text{ s}^{-1} \text{ nm}^{-1}$ . Interestingly, we also find that the quiescent flux from the extremely active M-dwarf AD Leo also meets the Rimmer et al. (2018) criterion ( $F_{200-280\text{nm}} = 2 \times 10^{11} \text{ cm}^{-2} \text{ s}^{-1} \text{ nm}^{-1}$ ), and that quiescent flux from early (high-mass) M-dwarfs like GJ 832 also marginally meet this criterion. This apparent contradiction with Rimmer et al. (2018) is due to their assignment of single-scattering albedo  $\omega_0 = 0.9 < 1$  when calculating Rayleigh scattering, which underestimates surface UV in the unique radiative transfer regime unveiled on anoxic early Earth-like planets (Ranjan & Sasselo 2017; Rimmer et al. 2021b; P. Rimmer, personal communication, 12/9/2020). We instead find that planets orbiting young or active M-dwarf stars, as well as some quiescent early M-dwarf stars, can meet the Rimmer et al. (2018) criterion, illustrating the necessity of measuring and constraining the evolution of NUV SEDs of individual planet-hosting M-dwarfs to understand the habitability of their worlds (Rimmer et al. 2021a).


#### 4.2. Implications of HIP23309 SED for Atmospheric Photochemistry

In this section, we explore the utility and limitations of ASTROSAT UVIT for constraining the UV SEDs of exoplanet host stars for atmospheric photochemistry applications. UV photochemistry controls the trace gas composition of temperate terrestrial planet atmospheres, and accurate UV SEDs are necessary for accurate predictions of atmospheric composition and inferences derived thereof (Rugheimer et al. 2015; Teal et al. 2022). We assess the impact of UV SEDs on predicted atmospheric trace gas concentrations using a procedure established first by Kasting et al. (1997) and subsequently extensively deployed (e.g., Segura et al. 2005; Rauer et al. 2011; Rugheimer et al. 2015; Teal et al. 2022): We run a photochemical model with a fixed planetary scenario, varying only the UV irradiation between model runs, and examine the implications for predicted atmospheric composition. Our calculations are solely sensitivity tests aimed at constraining the utility and limitations of using ASTROSAT UVIT to constrain M-dwarf SEDs for photochemical modeling applications, as opposed to self-consistent predictions of the atmospheric composition of temperate terrestrial planets orbiting young M-dwarf stars.

To conduct this sensitivity test, we use the MIT Exoplanet Atmospheric Chemistry model (MEAC, Hu et al. 2012), as updated by Ranjan et al. (2022). We apply MEAC to the N<sub>2</sub>-dominated abiotic temperate terrestrial planet benchmark scenario from Hu et al. (2012), which simulates a planet geologically similar to Earth in the absence of life. We explore the effects of different estimates of the HIP23309 SED on the abundances of key HCO species which dominate the atmosphere and whose remote detection could constrain surfaces processes like the carbon cycle and the presence or absence of life (Bean et al. 2017; Schwieterman et al. 2019; Wogan & Catling 2020; Lehmer et al. 2020; Thompson et al. 2022).

We begin by showing that the UVIT-measured SED alone can achieve order-of-magnitude accuracy at best in photochemical modeling of rocky exoplanet atmospheres. We formulate a UVIT-only input SED for HIP 23309 as in Section 4.1, except that we do not include the luminosity correction factor of 0.9 to maintain consistency with similar past sensitivity tests conducted with MEAC (e.g., Seager et al. 2013). We formulate a second SED as the UVIT SED, except that we replace the FUV UVIT measurements with the FUMES HST measurements (Figure 7). Calculating the trace gas composition under irradiation by each of these SEDs (Figure 8), we find that the use of the UVIT-only SED leads to overpredictions of CH<sub>4</sub> by  $7\times$  and underprediction of CO by  $6\times$  relative to the UVIT+HST SED (Table 3). Two factors drive model inaccuracy when using the UVIT-only SED. First, the UVIT SED systematically overestimates the continuum flux due to its lower sensitivity. This leads to overestimates of H<sub>2</sub>O photolysis, and

<sup>1</sup> <https://www.mrao.cam.ac.uk/~pbr27/resources.html>, accessed 12/7/2022




**Figure 6.** Surface radiances on early Earth-like planets orbiting different M-dwarfs including HIP23309, computed as in [Ranjan et al. \(2017\)](#). The [Rimmer et al. \(2018\)](#) minimum flux criterion for HCN homologation towards ribonucleotide synthesis is shown in pink. Extremely active stars like HIP23309 meet the [Rimmer et al. \(2018\)](#) criterion, illustrating the need to constrain UV emission of M-dwarfs over the course of their lifetimes to understand the potential of life search on M-dwarf planets to test theories of abiogenesis ([Rimmer et al. 2021a](#)).

accounts for the variation in  $p\text{CO}$  and some of the variation in  $p\text{CH}_4$ . The effects of this overestimate are amplified by the large wavelength gap between reliable measurements in the FUV and NUV grism, which is interpolated in the model. The importance of the UV continuum has been affirmed in other studies ([Teal et al. 2022](#); [Peacock et al. 2022](#)). Second, the UVIT SED does not reliably cover the Lyman-alpha line, where M-dwarfs emit a substantial fraction of their flux and which is particularly relevant to  $\text{CH}_4$  photolysis in anoxic atmospheres ([France et al. 2013](#); [Ranjan et al. 2022](#)). [Peacock et al. \(2022\)](#) reported predicted atmospheric composition of M-dwarf planets to be insensitive to the degree of self-reversal in the stellar  $\text{Ly}\alpha$  line, and we reproduced this finding with our model. However, omitting  $\text{Ly}\alpha$  entirely results in a much bigger change in irradiating flux than variations in the degree of self-reversal in the  $\text{Ly}\alpha$  line, and we find this omission to result in significant variations in predicted  $p\text{CH}_4$ .

We next consider how the UVIT-only SED performs in comparison to proxy SEDs from stars of similar spectral type, which are sometimes employed when accurate SEDs of planet-hosting stars are not available (e.g., [Hu et al. 2020, 2021](#)). We consider SEDs from GJ832 (M1.5) and HD 85512 (K6), which are well-characterized by the MUSCLES survey (v2.2, [France et al. 2016](#); [Lloyd et al. 2016](#); [Youngblood et al. 2016](#)), as proxies for HIP23309 (M0). We normalize their SEDs to the solar constant as for HIP23309, and apply these SEDs to our planetary scenario (Figure 7). We find that use of proxy SEDs matches the HST+UVIT SED calculation better than the UVIT-only SED (Figure 8, Table 3).






**Figure 7.** Input UV SEDs used for modeling the atmospheric composition (Figure 8) of a hypothetical abiotic Earth-like planet orbiting HIP23309.

**Table 3.** Surface Mixing Ratios Predicted in MEAC  
Abiotic N<sub>2</sub>-dominated Benchmark Scenario

SED	$r_{CH_4}(z=0)$	$r_{CO}(z=0)$
HIP 23309 HST FUV	$1 \times 10^{-6}$	$6 \times 10^{-8}$
HIP 23309 UVIT FUV	$7 \times 10^{-6}$	$1 \times 10^{-8}$
GJ 832 (MUSCLES)	$4 \times 10^{-7}$	$3 \times 10^{-7}$
HD85512 (MUSCLES)	$4 \times 10^{-7}$	$3 \times 10^{-7}$
GJ 832 (Scaled)	$6 \times 10^{-7}$	$1 \times 10^{-7}$
HD85512 (Scaled)	$4 \times 10^{-7}$	$1 \times 10^{-7}$

We finally consider whether UVIT data can be used to calibrate other techniques for estimating M-dwarf UV SEDs. For example, partial SED measurements have been used to calibrate full-SED estimates derived from stellar models (Peacock et al. 2019a,b, 2020). Such sophisticated estimates are beyond the scope of this work; as proof-of-concept, we instead utilize our UVIT measurements to crudely calibrate our use of similar stellar spectra as proxies. Specifically, we scale our GJ832 and HD85512 SEDs by a constant factor, such that they have the same flux integrated from 130-170 nm as our UVIT-only HIP23309 SED (Figure 7). We do not utilize the NUV data because the UVIT NUV grism is no longer operative, and so it is useless to determine its utility at calibrating other estimates. We find that use of scaled proxy SEDs to marginally improve agreement with the HIP23309 UVIT+HST SED compared to unscaled proxy SEDs, in the case of GJ832 reproducing the predicted surface concentrations of CO and CH<sub>4</sub> to within a factor



**Figure 8.** Predicted atmospheric composition of a hypothetical abiotic Earth-like planet orbiting HIP23309, subject to varying estimates of its UV irradiation (Figure 7). The different colors correspond to different chemical species. The left column corresponds to chemical species emitted by the surface, while the right column corresponds to photochemically generated species. In each row, the different linestyles correspond to different SEDs. The UVIT-only SED alone is insufficient to achieve highly precise photochemical modeling (top row), but can be used to calibrate other estimates of the stellar SED (middle row) to achieve higher precision, as done crudely here (bottom row) and with vastly more sophistication by, e.g., (Peacock et al. 2019a,b))

of 2 (Figure 8, Table 3). That even an extremely crude calibration with UVIT can marginally improve agreement suggests that it is possible that more sophisticated calibrations with UVIT can significantly improve agreement; we advocate such work.

## 5. DISCUSSION & CONCLUSIONS

Constraining the UV SEDs of low-mass stars is crucial to interpreting atmospheric measurements of the exoplanets they host, and to understanding the habitability and **urability** of those planets. The primary instrument used for such studies, HST, is oversubscribed and well past its designed lifetime. Here, we have examined the potential of ASTROSAT UVIT to carry out such studies through spectrally resolved NUV and FUV grism observations of the young M0 star HIP 23309. We robustly detect stellar lines of C-IV in the FUV and Fe-II and Mg-II in the NUV. We find UVIT can reproduce coterminous HST measurements in the FUV to within a factor of 2 and that its orbit-by-orbit NUV measurements are stable to 10%, but that its errors may be underestimated by a factor of a few. The

UVIT-measured HIP23309 SED displays trends in context of other measured M-dwarf SEDs that are consistent with first-order expectations. Taken together, the example of HIP 23309 suggests ASTROSAT UVIT is a viable tool for characterizing the UV SEDs of bright, nearby low-mass stars to within a factor of a few.

We have demonstrated the utility of such measurements for planetary applications using HIP23309 as a case study. Measurements from the UVIT NUV grism can inform studies of planetary habitability, because it constrains emission at the  $\geq 190$  nm wavelength range relevant to habitable planet surfaces (Ranjan & Sassellov 2017). For example, HIP23309 meets the Rimmer et al. (2018) criterion for sufficient UV output to power HCN photohomologation in its habitable zone, demonstrating the necessity of characterizing the detailed evolution of M-dwarf SEDs to understand the potential of biosignature search on M-dwarf planets to test theories of the origin of life. Our FUV and NUV spectra of HIP23309 on their own permit limited accurate photochemical modeling of planetary atmospheres, because the UVIT FUV grism does not reliably cover the crucial Lyman alpha line and because the limited sensitivity and wavelength coverage of the FUV grism lead to overestimates of the continuum flux. However, our sensitivity tests suggest that UVIT FUV observations can be used to calibrate estimates of M-dwarf SEDs, thereby supporting photochemical modeling. Our work suggests a role for archival NUV observations and archival and future FUV observations in supporting studies of exoplanet habitability and photochemistry.

Our study bears implications for the proposed Indian Spectroscopic and Imaging Space Telescope (INSIST; Subramaniam 2022). The INSIST mission design features a 1m mirror ( $3\times$  the size of UVIT), suggesting a factor of 3 improvement in signal-to-noise relative to UVIT, and continuous wavelength coverage from the FUV to the NUV (150-300 nm). This means that INSIST would likely outperform UVIT at characterizing the important UV continuum. However, termination of its wavelength coverage at 150 nm means that INSIST would not be able to measure the 121.6 nm Lyman alpha line that can be important to temperate exoplanet photochemistry. We advocate for the extension of INSIST's spectral coverage to 120 nm or shorter, to enable direct characterization of the full wavelength range relevant to exoplanet photochemistry. However, even if INSIST is unable to directly measure full exoplanet host star UV SEDs, measurements from INSIST will still be able to calibrate other estimates of M-dwarf SEDs like stellar models (Peacock et al. 2019a,b), and its coverage of NUV wavelengths will enable planetary habitability studies (e.g., Rugheimer et al. 2015; O'Malley-James & Kaltenegger 2019; Rimmer et al. 2018). The UV-SCOPE mission concept (Ardila et al. 2022) covers 100-400 nm and is ideally suited to studies of the type discussed here.

## ACKNOWLEDGMENTS

We thank Sarah Peacock and Edward Schwieterman for sharing TRAPPIST-1 spectra from Peacock et al. (2022) as an additional validation test. SR thanks Sujan Sengupta for mentorship in developing the project idea, and the Indian Institute of Astrophysics, Northwestern University, and University of Arizona for support while carrying out this project.

*Author contribution statement:* SR conceived the project, served as co-I on the UVIT proposal, assessed the reduced data, performed the modeling, and wrote the paper. PN led reduction of the UVIT data. JSP served as co-I on the UVIT proposal, advised on data reduction, and supplied pointers to other M-dwarf spectra utilized in this paper. MN assisted PN in reducing the UVIT data. All authors reviewed and edited the paper.

*Facilities:* HST(STIS), AstroSat (UVIT)

*Software:* MEAC (Hu et al. 2012; Ranjan et al. 2022)

## REFERENCES

- |  |  |
|--|--|
| <p>Ardila, D. R., Shkolnik, E., Ziemer, J., et al. 2022, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 12181, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, ed. J.-W. A. den Herder, S. Nikzad, &amp; K. Nakazawa, 1218104, doi: <a href="https://doi.org/10.1117/12.2629000">10.1117/12.2629000</a></p> | <p>Ayres, T. R. 2010, ApJS, 187, 149, doi: <a href="https://doi.org/10.1088/0067-0049/187/1/149">10.1088/0067-0049/187/1/149</a></p> <p>Batalha, N. E., Lewis, N. K., Line, M. R., Valenti, J., &amp; Stevenson, K. 2018, ApJL, 856, L34, doi: <a href="https://doi.org/10.3847/2041-8213/aab896">10.3847/2041-8213/aab896</a></p> |
|--|--|

- Bean, J. L., Abbot, D. S., & Kempton, E. M. R. 2017, *ApJL*, 841, L24, doi: [10.3847/2041-8213/aa738a](https://doi.org/10.3847/2041-8213/aa738a)
- Blanco, C., Catalano, S., & Marilli, E. 1979, *Nature*, 280, 661, doi: [10.1038/280661a0](https://doi.org/10.1038/280661a0)
- Bryson, S., Kunimoto, M., Kopparapu, R. K., et al. 2021, *AJ*, 161, 36, doi: [10.3847/1538-3881/abc418](https://doi.org/10.3847/1538-3881/abc418)
- Cowan, N. B., Greene, T., Angerhausen, D., et al. 2015, *PASP*, 127, 311, doi: [10.1086/680855](https://doi.org/10.1086/680855)
- Dewangan, G. C. 2021, *Journal of Astrophysics and Astronomy*, 42, 49, doi: [10.1007/s12036-021-09691-w](https://doi.org/10.1007/s12036-021-09691-w)
- Doherty, L. R. 1985, *MNRAS*, 217, 41, doi: [10.1093/mnras/217.1.41](https://doi.org/10.1093/mnras/217.1.41)
- Dong, C., Lingam, M., Ma, Y., & Cohen, O. 2017, *ApJL*, 837, L26, doi: [10.3847/2041-8213/aa6438](https://doi.org/10.3847/2041-8213/aa6438)
- Dressing, C. D., & Charbonneau, D. 2015, *Astrophysical Journal*, 807, 45, doi: [10.1088/0004-637X/807/1/45](https://doi.org/10.1088/0004-637X/807/1/45)
- France, K., Froning, C. S., Linsky, J. L., et al. 2013, *The Astrophysical Journal*, 763, 149
- France, K., Parke Loyd, R. O., Youngblood, A., et al. 2016, *Astrophysical Journal*, 820, 89, doi: [10.3847/0004-637X/820/2/89](https://doi.org/10.3847/0004-637X/820/2/89)
- Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2022, arXiv e-prints, arXiv:2208.00211. <https://arxiv.org/abs/2208.00211>
- Harman, C. E., Felton, R., Hu, R., et al. 2018, *ApJ*, 866, 56, doi: [10.3847/1538-4357/aadd9b](https://doi.org/10.3847/1538-4357/aadd9b)
- Hu, R., Damiano, M., Scheucher, M., et al. 2021, *ApJL*, 921, L8, doi: [10.3847/2041-8213/ac1f92](https://doi.org/10.3847/2041-8213/ac1f92)
- Hu, R., Peterson, L., & Wolf, E. T. 2020, *ApJ*, 888, 122, doi: [10.3847/1538-4357/ab5f07](https://doi.org/10.3847/1538-4357/ab5f07)
- Hu, R., Seager, S., & Bains, W. 2012, *Astrophysical Journal*, 761, 166, doi: [10.1088/0004-637X/761/2/166](https://doi.org/10.1088/0004-637X/761/2/166)
- Kasting, J. F., Whittet, D. C. B., & Sheldon, W. R. 1997, *Origins of Life and Evolution of the Biosphere*, 27, 413, doi: [10.1023/A:1006596806012](https://doi.org/10.1023/A:1006596806012)
- Lehmer, O. R., Catling, D. C., & Krissansen-Totton, J. 2020, *Nature Communications*, 11, 6153, doi: [10.1038/s41467-020-19896-2](https://doi.org/10.1038/s41467-020-19896-2)
- Loyd, R. O. P., France, K., Youngblood, A., et al. 2016, *Astrophysical Journal*, 824, 102, doi: [10.3847/0004-637X/824/2/102](https://doi.org/10.3847/0004-637X/824/2/102)
- Loyd, R. O. P., Shkolnik, E. L., Schneider, A. C., et al. 2021, *ApJ*, 907, 91, doi: [10.3847/1538-4357/abd0f0](https://doi.org/10.3847/1538-4357/abd0f0)
- Luger, R., & Barnes, R. 2015, *Astrobiology*, 15, 119, doi: [10.1089/ast.2014.1231](https://doi.org/10.1089/ast.2014.1231)
- Lustig-Yaeger, J., Meadows, V. S., & Lincowski, A. P. 2019, *AJ*, 158, 27, doi: [10.3847/1538-3881/ab21e0](https://doi.org/10.3847/1538-3881/ab21e0)
- Lustig-Yaeger, J., Robinson, T. D., & Arney, G. 2019, *Journal of Open Source Software*, 4, 1387, doi: [10.21105/joss.01387](https://doi.org/10.21105/joss.01387)
- Malo, L., Doyon, R., Feiden, G. A., et al. 2014, *ApJ*, 792, 37, doi: [10.1088/0004-637X/792/1/37](https://doi.org/10.1088/0004-637X/792/1/37)
- Mamajek, E. E., & Bell, C. P. M. 2014, *MNRAS*, 445, 2169, doi: [10.1093/mnras/stu1894](https://doi.org/10.1093/mnras/stu1894)
- Melbourne, K., Youngblood, A., France, K., et al. 2020, *AJ*, 160, 269, doi: [10.3847/1538-3881/abbf5c](https://doi.org/10.3847/1538-3881/abbf5c)
- Morley, C. V., Kreidberg, L., Rustamkulov, Z., Robinson, T., & Fortney, J. J. 2017, *ApJ*, 850, 121, doi: [10.3847/1538-4357/aa927b](https://doi.org/10.3847/1538-4357/aa927b)
- National Academies of Sciences Engineering and Medicine. 2018, *Exoplanet Science Strategy* (Washington, DC: The National Academies Press), doi: [10.17226/25187](https://doi.org/10.17226/25187)
- of Sciences Engineering, N. A., & Medicine. 2021, *Pathways to Discovery in Astronomy and Astrophysics for the 2020s* (Washington, DC: The National Academies Press), doi: [10.17226/26141](https://doi.org/10.17226/26141)
- O'Malley-James, J. T., & Kaltenegger, L. 2019, *MNRAS*, 485, 5598, doi: [10.1093/mnras/stz724](https://doi.org/10.1093/mnras/stz724)
- Peacock, S., Barman, T., Shkolnik, E. L., Hauschildt, P. H., & Baron, E. 2019a, *ApJ*, 871, 235, doi: [10.3847/1538-4357/aaf891](https://doi.org/10.3847/1538-4357/aaf891)
- Peacock, S., Barman, T., Shkolnik, E. L., et al. 2019b, *ApJ*, 886, 77, doi: [10.3847/1538-4357/ab4f6f](https://doi.org/10.3847/1538-4357/ab4f6f)
- . 2020, *ApJ*, 895, 5, doi: [10.3847/1538-4357/ab893a](https://doi.org/10.3847/1538-4357/ab893a)
- Peacock, S., Barman, T. S., Schneider, A. C., et al. 2022, *ApJ*, 933, 235, doi: [10.3847/1538-4357/ac77f2](https://doi.org/10.3847/1538-4357/ac77f2)
- Pineda, J. S., Youngblood, A., & France, K. 2021, *ApJ*, 911, 111, doi: [10.3847/1538-4357/abe8d7](https://doi.org/10.3847/1538-4357/abe8d7)
- Plavchan, P., Barclay, T., Gagné, J., et al. 2020, *Nature*, 582, 497, doi: [10.1038/s41586-020-2400-z](https://doi.org/10.1038/s41586-020-2400-z)
- Postma, J. E., & Leahy, D. 2017, *PASP*, 129, 115002, doi: [10.1088/1538-3873/aa8800](https://doi.org/10.1088/1538-3873/aa8800)
- Ramirez, R. M., & Kaltenegger, L. 2014, *Astrophysical Journal*, 797, L25, doi: [10.1088/2041-8205/797/2/L25](https://doi.org/10.1088/2041-8205/797/2/L25)
- Ranjan, S., & Sasselov, D. D. 2017, *Astrobiology*, 17, 169, doi: [10.1089/ast.2016.1519](https://doi.org/10.1089/ast.2016.1519)
- Ranjan, S., Seager, S., Zhan, Z., et al. 2022, *ApJ*, 930, 131, doi: [10.3847/1538-4357/ac5749](https://doi.org/10.3847/1538-4357/ac5749)
- Ranjan, S., Wordsworth, R., & Sasselov, D. D. 2017, *The Astrophysical Journal*, 843, 110. <http://stacks.iop.org/0004-637X/843/i=2/a=110>
- Rauer, H., Gebauer, S., Paris, P. V., et al. 2011, *A&A*, 529, A8, doi: [10.1051/0004-6361/201014368](https://doi.org/10.1051/0004-6361/201014368)
- Rimmer, P. B., Ranjan, S., & Rugheimer, S. 2021a, *Elements*, 17, 265. <https://arxiv.org/abs/2108.08388>
- Rimmer, P. B., Thompson, S. J., Xu, J., et al. 2021b, *Astrobiology*, 21, 1099, doi: [10.1089/ast.2020.2335](https://doi.org/10.1089/ast.2020.2335)
- Rimmer, P. B., Xu, J., Thompson, S. J., et al. 2018, *Science Advances*, 4, eaar3302, doi: [10.1126/sciadv.aar3302](https://doi.org/10.1126/sciadv.aar3302)

- Robinson, T. D., Stapelfeldt, K. R., & Marley, M. S. 2016, *PASP*, 128, 025003, doi: [10.1088/1538-3873/128/960/025003](https://doi.org/10.1088/1538-3873/128/960/025003)
- Rodler, F., & López-Morales, M. 2014, *Astrophysical Journal*, 781, 54, doi: [10.1088/0004-637X/781/1/54](https://doi.org/10.1088/0004-637X/781/1/54)
- Rugheimer, S., Kaltenegger, L., Segura, A., Linsky, J., & Mohanty, S. 2015, *The Astrophysical Journal*, 809, 57
- Rugheimer, S., Segura, A., Kaltenegger, L., & Sasselov, D. 2015, *Astrophysical Journal*, 806, 137, doi: [10.1088/0004-637X/806/1/137](https://doi.org/10.1088/0004-637X/806/1/137)
- Scalo, J., Kaltenegger, L., Segura, A. G., et al. 2007, *Astrobiology*, 7, 85, doi: [10.1089/ast.2006.0125](https://doi.org/10.1089/ast.2006.0125)
- Schwieterman, E. W., Reinhard, C. T., Olson, S. L., et al. 2019, *ApJ*, 874, 9, doi: [10.3847/1538-4357/ab05e1](https://doi.org/10.3847/1538-4357/ab05e1)
- Schwieterman, E. W., Kiang, N. Y., Parenteau, M. N., et al. 2018, *Astrobiology*, 18, 663, doi: [10.1089/ast.2017.1729](https://doi.org/10.1089/ast.2017.1729)
- Schwieterman, E. W., Olson, S. L., Pidhorodetska, D., et al. 2022, *ApJ*, 937, 109, doi: [10.3847/1538-4357/ac8cfb](https://doi.org/10.3847/1538-4357/ac8cfb)
- Seager, S., Bains, W., & Hu, R. 2013, *ApJ*, 777, 95, doi: [10.1088/0004-637X/777/2/95](https://doi.org/10.1088/0004-637X/777/2/95)
- Segura, A., Kasting, J. F., Meadows, V., et al. 2005, *Astrobiology*, 5, 706, doi: [10.1089/ast.2005.5.706](https://doi.org/10.1089/ast.2005.5.706)
- Shields, A. L., Ballard, S., & Johnson, J. A. 2016, *ArXiv e-prints*. <https://arxiv.org/abs/1610.05765>
- Shkolnik, E. L., & Barman, T. S. 2014, *AJ*, 148, 64, doi: [10.1088/0004-6256/148/4/64](https://doi.org/10.1088/0004-6256/148/4/64)
- Subramaniam, A. 2022, *Journal of Astrophysics and Astronomy*, 43, 80, doi: [10.1007/s12036-022-09870-3](https://doi.org/10.1007/s12036-022-09870-3)
- Tandon, S. N., Subramaniam, A., Girish, V., et al. 2017, *AJ*, 154, 128, doi: [10.3847/1538-3881/aa8451](https://doi.org/10.3847/1538-3881/aa8451)
- Tandon, S. N., Postma, J., Joseph, P., et al. 2020, *AJ*, 159, 158, doi: [10.3847/1538-3881/ab72a3](https://doi.org/10.3847/1538-3881/ab72a3)
- Teal, D. J., Kempton, E. M. R., Bastelberger, S., Youngblood, A., & Arney, G. 2022, *arXiv e-prints*, arXiv:2201.08805. <https://arxiv.org/abs/2201.08805>
- Thompson, M. A., Krissansen-Totton, J., Wogan, N., Telus, M., & Fortney, J. J. 2022, *Proceedings of the National Academy of Science*, 119, e2117933119, doi: [10.1073/pnas.2117933119](https://doi.org/10.1073/pnas.2117933119)
- Tilley, M. A., Segura, A., Meadows, V., Hawley, S., & Davenport, J. 2019, *Astrobiology*, 19, 64, doi: [10.1089/ast.2017.1794](https://doi.org/10.1089/ast.2017.1794)
- Torres, C. A. O., Quast, G. R., da Silva, L., et al. 2006, *A&A*, 460, 695, doi: [10.1051/0004-6361:20065602](https://doi.org/10.1051/0004-6361:20065602)
- Tremblay, L., Line, M. R., Stevenson, K., et al. 2020, *AJ*, 159, 117, doi: [10.3847/1538-3881/ab64dd](https://doi.org/10.3847/1538-3881/ab64dd)
- Viswanath, G., Narang, M., Manoj, P., Mathew, B., & Kartha, S. S. 2020, *AJ*, 159, 194, doi: [10.3847/1538-3881/ab7d3b](https://doi.org/10.3847/1538-3881/ab7d3b)
- Wogan, N. F., & Catling, D. C. 2020, *ApJ*, 892, 127, doi: [10.3847/1538-4357/ab7b81](https://doi.org/10.3847/1538-4357/ab7b81)
- Wordsworth, R. D., Schaefer, L. K., & Fischer, R. A. 2018, *AJ*, 155, 195, doi: [10.3847/1538-3881/aab608](https://doi.org/10.3847/1538-3881/aab608)
- Xu, J., Ritson, D. J., Ranjan, S., et al. 2018, *Chemical Communications*
- Xu, J., Chmela, V., Green, N. J., et al. 2020, *Nature*, 582, 60, doi: [10.1038/s41586-020-2330-9](https://doi.org/10.1038/s41586-020-2330-9)
- Youngblood, A., Pineda, J. S., & France, K. 2021, *ApJ*, 911, 112, doi: [10.3847/1538-4357/abe8d8](https://doi.org/10.3847/1538-4357/abe8d8)
- Youngblood, A., France, K., Loyd, R. O. P., et al. 2016, *ApJ*, 824, 101, doi: [10.3847/0004-637X/824/2/101](https://doi.org/10.3847/0004-637X/824/2/101)
- . 2017, *ApJ*, 843, 31, doi: [10.3847/1538-4357/aa76dd](https://doi.org/10.3847/1538-4357/aa76dd)