**IEE(Main) 2023** Department of Higher Education Ministry of Education Government of India Session 1

QuestionID Correct Option(s)/

3666941171 | 3666943514

3666941172 | 3666943518

3666941173 3666943521

3666941174 3666943526

3666941175 | 3666943528

3666941176 3666943532

3666941177 | 3666943538

3666941178 3666943540

3666941179 3666943543

3666941180 3666943547

3666941181 3666943553

3666941182 3666943557

3666941183 3666943559

3666941184 3666943564

3666941185 3666943567

3666941186 3666943573

3666941187 3666943577

3666941188 3666943581

3666941189 3666943586

3666941190 3666943588

3666941191 27

3666941192 | 52

3666941193 | 100

3666941194 10

3666941195 | 45

3666941196 18

3666941197 5

3666941198 | 17

3666941199 2

3666941200 4

3666941201 | 3666943602

3666941202 3666943608

3666941203 | 3666943611

3666941204 | 3666943616

3666941205 3666943620

3666941206 3666943623

3666941207 | 3666943628

3666941208 | 3666943630

3666941209 | 3666943634

3666941210 3666943639

3666941211 3666943643

3666941212 | 3666943648

3666941213 | 3666943651

3666941214 3666943655

3666941215 3666943659

3666941216 3666943663

3666941217 3666943668

3666941218 3666943671

3666941219 3666943674

3666941220 | 3666943677

3666941221 12

3666941222 6

3666941223 | 360

3666941224 41500

3666941225 9079

3666941226 10

3666941227 | 60

3666941228 2

3666941229 4

3666941230 42

3666941231 3666943692

3666941232 3666943695

3666941233 3666943700

3666941234 3666943706

3666941235 3666943709

3666941236 3666943711

3666941237 3666943717

3666941238 | 3666943719

3666941239 | 3666943725

3666941240 | 3666943729

3666941241 3666943731

3666941242 | 3666943736

3666941243 3666943741

3666941244 3666943744

3666941245 3666943750

3666941246 3666943753

3666941247 3666943758

3666941248 3666943762

3666941249 | 3666943763

3666941250 3666943769

3666941251 2039

3666941252 25

3666941253 43

3666941254 | 1080

3666941255 | 495

3666941256 216

3666941257 120

3666941258 | 600

3666941259 9

3666941260 2

This site is designed, developed and hosted by CNR Division, NIC and the contents are provided by NTA. For any

Choose file No file chosen

In case the candidates want to submit documents in support of challenge of answer key, they should upload the PDF file.

**Submit Your Claims** 

Answers

Change Password Register Query Logout ♣ Home

**Candidate Details** 

Application Number:

Candidate's Name:

**Claimed Answer Key List** 

**B** TECH - Physics Section A

B TECH - Physics Section A

**B** TECH - Physics Section **B** 

■ TECH - Physics Section ■

**B** TECH - Physics Section **B** 

**B** TECH - Physics Section **B** 

**B** TECH - Physics Section **B** 

BTECH - Chemistry Section A

B TECH - Chemistry Section A Objective

BTECH - Chemistry Section A Objective

B TECH - Chemistry Section A Objective

BTECH - Chemistry Section A Objective

B TECH - Chemistry Section A Objective

BTECH - Chemistry Section A Objective

BTECH - Chemistry Section A Objective

B TECH - Chemistry Section A Objective

B TECH - Chemistry Section A Objective

B TECH - Chemistry Section B Numerical

Numerical

Numerical

Numerical

Numerical

Numerical

Numerical

Objective

bjective

Objective

Objective

Objective

Objective

Objective

Objective

Objective

Numerical

Claimed Answer Key ListUpload Document:

further information, please contact to NTA. (Ver 1.0.43.2.0)

Claimed Answer Key ListUpload Document

**B** TECH - Chemistry Section B

BTECH - Chemistry Section B

B TECH - Chemistry Section B

BTECH - Chemistry Section B

BTECH - Chemistry Section B

BTECH - Chemistry Section B

**BTECH** - Mathematics

**B TECH - Mathematics** 

**BTECH** - Mathematics

■ TECH - Mathematics

**B** TECH - Mathematics

■ TECH - Mathematics

**B** TECH - Mathematics

**B** TECH - Mathematics

B TECH - Mathematics

**B TECH - Mathematics** 

**B TECH - Mathematics** 

Section A

Section B

Disclaimer:

Question

Objective

Numer cal

Numerical

Numerical

Numerical

Numerical

Numerical

Numerical

Numerical

Numerical

Numerical

Objective

Type

Father's Name:

Paper

**Challenges regarding Answer Key** 

Roll Number:

Date of Birth:

Mother's Name:

3666943511

3666943515

3666943519

3666943523

3666943527

3666943531

3666943535

3666943539

3666943543

3666943547

3666943551

3666943555

3666943559

3666943563

3666943567

3666943571

3666943575

3666943579

3666943583

3666943587

3666943601

3666943605

3666943609

3666943613

3666943617

3666943621

3666943625

3666943629

3666943633

3666943637

366 69 43 641

3666943645

3666943649

3666943653

3666943657

3666943661

3666943665

3666943669

3666943673

3666943677

3666943691

3666943695

3666943699

3666943703

3666943707

3666943711

3666943715

3666943719

3666943723

3666943727

3666943731

3666943735

3666943739

3666943743

3666943747

3666943751

3666943755

3666943759

3666943763

3666943767

Option(s) ID for Challenge

3666943512

3666943516

3666943520

3666943524

3666943528

3666943532

3666943536

3666943540

3666943544

3666943548

3666943552

3666943556

3666943560

3666943564

3666943568

3666943572

3666943576

3666943580

3666943584

3666943588

3666943602

3666943606

3666943610

3666943614

3666943618

3666943622

3666943626

3666943630

3666943634

3666943638

3666943642

3666943646

3666943650

3666943654

3666943658

3666943662

3666943666

3666943670

3666943674

3666943678

3666943692

3666943696

3666943700

3666943704

3666943708

3666943712

3666943716

3666943720

3666943724

3666943728

3666943732

3666943736

3666943740

3666943744

3666943748

3666943752

3666943756

3666943760

3666943764

3666943768

Name:

13-05-2006

3666943513

3666943517

3666943521

3666943525

3666943529

3666943533

3666943537

3666943541

3666943545

3666943549

3666943553

3666943557

3666943561

3666943565

3666943569

3666943573

3666943577

3666943581

3666943585

3666943589

राष्ट्रीय परीक्षा एजेंसी **National Testing Agency** 

**Application No:** 

3666943514

3666943518

3666943522

3666943526

3666943530

3666943534

3666943538

3666943542

3666943546

3666943550

3666943554

3666943558

3666943562

3666943566

3666943570

3666943574

3666943578

3666943582

3666943586

3666943590

3666943604

3666943608

3666943612

3666943616

3666943620

3666943624

3666943628

3666943632

3666943636

3666943640

3666943644

3666943648

3666943652

3666943656

3666943660

3666943664

3666943668

3666943672

3666943676

3666943680

3666943694

3666943698

3666943702

3666943706

3666943710

3666943714

3666943718

3666943722

3666943726

3666943730

3666943734

3666943738

3666943742

3666943746

3666943750

3666943754

3666943758

3666943762

3666943766

3666943770

3666943693

3666943697

3666943701

3666943705

3666943709

3666943713

3666943717

3666943721

3666943725

3666943729

3666943733

3666943737

3666943741

3666943745

3666943749

3666943753

3666943757

3666943761

3666943765

3666943769

None of These

3666943603

3666943607

3666943611

3666943615

3666943619

3666943623

3666943627

3666943631

3666943635

3666943639

3666943643

3666943647

3666943651

3666943655

3666943659

3666943663

3666943667

3666943671

3666943675

3666943679

None of These

## **JEE 2023 Session-1 24th Jan to 1st Feb 2023**

Application No	
Candidate Name	
Roll No	
Test Date	25/01/2023
Test Time	9:00 AM - 12:00 PM
Subject	В ТЕСН

Section: Physics Section A

- Q.1 The root mean square velocity of molecules of gas is Options
  - 1. Inversely proportional to square root of temperature



- 2. Proportional to square root of temperature  $(\sqrt{T})$
- 3. Proportional to square of temperature  $(T^2)$
- 4. Proportional to temperature (T)

Question Type: MCQ

Question ID: 3666941178 Option 1 ID: 3666943542 Option 2 ID: 3666943540

Option 3 ID: 3666943541 Option 4 ID: 3666943539

Status: Answered

Chosen Option: 2

## Q.2 Match List I with List II

List I	List II
A. Surface tension	I. kg m <sup>-1</sup> s <sup>-1</sup>
B. Pressure	II. kg ms <sup>-1</sup>
C. Viscosity	III. kg m <sup>-1</sup> s <sup>-2</sup>
D. Impulse	IV. kg s <sup>-2</sup>

Choose the correct answer from the options given below:

Options 1. A-IV , B-III , C-II , D-I

2. A-II, B-I, C-III, D-IV

3. A-III, B-IV, C-I, D-II

4. A-IV, B-III, C-I, D-II

Question Type : MCQ

Question ID: 3666941171 Option 1 ID: 3666943511

Option 2 ID: 3666943513 Option 3 ID: 3666943512 Option 4 ID: 3666943514

Status: Answered

Q.3 Match List I with List II

List I (Current configuration	))		List II (Magnitude of Magnetic Field at point O)
$A. \longrightarrow (?)$	) I.		$B_0 = \frac{\mu_0 I}{4\pi r} [\pi + 2]$
B. I 0 1	11	I.	$B_0 = \frac{\mu_0}{4} \; \frac{I}{r}$
$\mathbf{c}. \qquad \qquad \begin{matrix} \downarrow I \\ \downarrow 0 \\ r \end{matrix}$		П.	$B_0 = \frac{\mu_0 I}{2\pi r} \big[\pi - 1\big]$
D. I r	Г	V.	$B_0 = \frac{\mu_0 I}{4\pi r} \left[\pi + 1\right]$

Choose the correct answer from the options given below:

Options 1. A-I, B-III, C-IV, D-II

- 2. A-III, B-IV, C-I, D-II
- 3. A-III, B-I, C-IV, D-II
- 4. A-II, B-I, C-IV, D-III

Question Type : MCQ

Question ID: 3666941182 Option 1 ID: 3666943556 Option 2 ID: 3666943555 Option 3 ID: 3666943557

Option 4 ID: 3666943558 Status : **Answered** 

Chosen Option: 3

Assume that the earth is a solid sphere of uniform density and a tunnel is dug along its diameter throughout the earth. It is found that when a particle is released in this tunnel, it executes a simple harmonic motion. The mass of the particle is 100 g. The time period of the motion of the particle will be (approximately)

(Take  $g=10 \text{ m s}^{-2}$ , radius of earth = 6400 km)

Options 1. 24 hours

- 2. 1 hour 40 minutes
- 3. 12 hours
- 4. 1 hour 24 minutes

Question Type: MCQ

Question ID: 3666941179

Option 1 ID: 3666943546

Option 2 ID: 3666943544

Option 3 ID: 3666943545

Option 4 ID: 3666943543

Status: Answered Chosen Option: 4

A message signal of frequency 5 kHz is used to modulate a carrier signal of frequency 2 MHz. The bandwidth for amplitude modulation is:

Options 1. 10 kHz

- 2. 20 kHz
- 3. 5 kHz
- 4. 2.5 kHz

Question Type: MCQ

Question ID: 3666941190 Option 1 ID: 3666943588 Option 2 ID: 3666943590 Option 3 ID: 3666943587 Option 4 ID: 3666943589 Status: Answered

Chosen Option: 2

Given below are two statements: one is labelled as Assertion A and the other is labelled as Reason R

Assertion A: Photodiodes are used in forward bias usually for measuring the light intensity.

Reason R: For a p-n junction diode, at applied voltage V the current in the forward bias is more than the current in the reverse bias for  $|V_z| \ge \pm V \ge |V_0|$  where  $V_0$  is the threshold voltage and Vz is the breakdown voltage.

In the light of the above statements, choose the correct answer from the options given below

Options 1. Both A and R are true and R is correct explanation A

- 2. A is false but R is true
- 3. A is true but R is false
- 4. Both A and R are true but R is NOT the correct explanation A

Question Type : MCQ

Question ID: 3666941189 Option 1 ID: 3666943583 Option 2 ID: 3666943586 Option 3 ID: 3666943585 Option 4 ID: 3666943584 Status: Answered

Chosen Option: 4

A Carnot engine with efficiency 50% takes heat from a source at 600 K. In order to increase the efficiency to 70%, keeping the temperature of sink same, the new temperature of the source will be:

Options 1. 1000 K

- 2. 360 K
- 3. 300 K
- 4.900 K

Question Type: MCQ

Question ID: 3666941177 Option 1 ID: 3666943538 Option 2 ID: 3666943536 Option 3 ID: 3666943535 Option 4 ID: 3666943537

Status: Answered

In an LC oscillator, if values of inductance and capacitance become twice and eight times, respectively, then the resonant frequency of oscillator becomes x times its initial resonant frequency  $\omega_0$ . The value of x is:

Options 1. 1/16

- 2.16
- 3. 4
- 4. 1/4

Question Type: MCQ

Question ID: 3666941184

Option 1 ID: 3666943565 Option 2 ID: 3666943566

Option 3 ID: 3666943563 Option 4 ID: 3666943564

> Not Attempted and Status: Marked For Review

Chosen Option: --

A car is moving with a constant speed of 20 m/s in a circular horizontal track of radius 40 m. A bob is suspended from the roof of the car by a massless string. The angle made by the string with the vertical will be : (Take  $g = 10 \text{ m/s}^2$ )

Options

- 3

Question Type: MCQ

Question ID: 3666941173 Option 1 ID: 3666943520

Option 2 ID: 3666943521 Option 3 ID: 3666943519

Option 4 ID: 3666943522 Status: Answered

Chosen Option: 1

Q.10 The ratio of the density of oxygen nucleus  $\binom{16}{8}0$  and helium nucleus  $\binom{4}{2}\text{He}$  is

Options 1. 1:1

- 2. 8:1
- 3. 2:1
- 4. 4:1

Question Type: MCQ

Question ID: 3666941186 Option 1 ID: 3666943573


Option 2 ID: 3666943571 Option 3 ID: 3666943572

Option 4 ID: 3666943574 Status: Answered

- Q.11 A uniform metallic wire carries a current 2 A, when 3.4 V battery is connected across it . The mass of uniform metallic wire is  $8.92 \times 10^{-3}$ kg, density is  $8.92 \times 10^3$  kg/m³ and resistivity is  $1.7 \times 10^{-8}$   $\Omega$  -m . The length of wire is :
- Options 1. l = 6.8 m
  - 2. l = 5 m
  - 3. l = 100 m
  - 4. l = 10 m

- Question Type: MCQ
  - Question ID: 3666941181
  - Option 1 ID: 3666943554
  - Option 2 ID: 3666943552
  - Option 3 ID: 3666943551
  - Option 4 ID: 3666943553
    - Status: Answered
- Chosen Option: 4
- Q.12 A solenoid of 1200 turns is wound uniformly in a single layer on a glass tube 2 m long and 0.2 m in diameter. The magnetic intensity at the center of the solenoid when a current of 2 A flows through it is:
- Options 1. 1.2×10<sup>3</sup> A m<sup>-1</sup>
  - <sup>2.</sup> 1 A m<sup>-1</sup>
  - $3.2.4 \times 10^3 \text{A m}^{-1}$
  - 4. 2.4×10<sup>-3</sup> A m<sup>-1</sup>

- Question Type: MCQ
- Question ID: 3666941183
- Option 1 ID: 3666943559
- Option 2 ID: 3666943561
- Option 3 ID: 3666943560
- Option 4 ID: 3666943562
- Status: Answered Chosen Option: 1
- Q.13 An object of mass 8 kg is hanging from one end of a uniform rod CD of mass 2 kg and length 1 m pivoted at its end C on a vertical wall as shown in figure. It is supported by a cable AB such that the system is in equilibrium. The tension in the cable is : (Take  $g = 10 \text{ m/s}^2$ )



- Options 1. 30 N
  - 2. 240 N
  - 3. 300 N
  - 4. 90 N

Question Type: MCQ

Question ID: 3666941174

Option 1 ID: 3666943523

Option 2 ID: 3666943525

Option 3 ID: 3666943526

Option 4 ID: 3666943524

Status: Not Answered

Q.14 A parallel plate capacitor has plate area 40 cm<sup>2</sup> and plates separation 2 mm. The space between the plates is filled with a dielectric medium of a thickness 1 mm and dielectric constant 5. The capacitance of the system is:

Options

$$\frac{10}{3} \epsilon_0 F$$

- 2.  $10\,\epsilon_0\,F$
- 3. 24 ε<sub>0</sub> F
- $4.\frac{3}{10}\varepsilon_0$  F

Question Type : MCQ

Question ID: 3666941180 Option 1 ID: 3666943547 Option 2 ID: 3666943549 Option 3 ID: 3666943550 Option 4 ID: 3666943548

Status: Answered

Chosen Option: 1

Q.15 Electron beam used in an electron microscope, when accelerated by a voltage of 20 kV, has a de-Broglie wavelength of  $\lambda_0$ . If the voltage is increased to 40 kV, then the de-Broglie wavelength associated with the electron beam would be:

Options

- 3. <mark>9 λ</mark>ο
- 4. 3 h

Question Type : MCQ

Question ID: 3666941188 Option 1 ID: 3666943581 Option 2 ID: 3666943582 Option 3 ID: 3666943580 Option 4 ID: 3666943579 Status: Answered

Chosen Option: 1

T is the time period of simple pendulum on the earth's surface. Its time period becomes x T when taken to a height R (equal to earth's radius) above the earth's surface. Then, the value of x will be:

**Options** 

Question Type: MCQ

Question ID: 3666941175 Option 1 ID: 3666943529 Option 2 ID: 3666943528 Option 3 ID: 3666943530 Option 4 ID: 3666943527

Status: Answered

Q.17 In Young's double slits experiment, the position of 5th bright fringe from the central maximum is 5 cm. The distance between slits and screen is 1 m and wavelength of used monochromatic light is 600 nm. The separation between the slits is:

Options 1. 48 µm

- 2. 36 µm
- 3. 60 µm
- 4. 12 µm

Question Type: MCQ

Question ID: 3666941187

Option 1 ID: 3666943576

Option 2 ID: 3666943575 Option 3 ID: 3666943577

Option 4 ID: 3666943578

Status: Answered

Chosen Option: 1

Q.18 A car travels a distance of 'x' with speed  $v_1$  and then same distance 'x' with speed  $v_2$  in the same direction. The average speed of the car is:

Options

$$1. \frac{v_1 v_2}{2(v_1 + v_2)}$$

2. 
$$\frac{v_1 + v_2}{2}$$

$$3. \frac{2 v_1 v_2}{v_1 + v_2}$$

4. 
$$\frac{2x}{v_1 + v_2}$$

Question Type: MCQ

Question ID: 3666941172

Option 1 ID: 3666943517

Option 2 ID: 3666943516

Option 3 ID: 3666943518 Option 4 ID: 3666943515

Status: Answered

Chosen Option: 3

Q.19 A bowl filled with very hot soup cools from 98°C to 86°C in 2 minutes when the room temperature is 22°C. How long it will take to cool from 75°C to 69°C?

Options 1. 1 minute

- 2. 0.5 minute
- 3. 2 minutes
- 4. 1.4 minutes

Question Type: MCQ

Question ID: 3666941176

Option 1 ID: 3666943533

Option 2 ID: 3666943534

Option 3 ID: 3666943531

Option 4 ID: 3666943532

Status: Answered

- Q.20 An electromagnetic wave is transporting energy in the negative z direction. At a certain point and certain time the direction of electric field of the wave is along positive y direction. What will be the direction of the magnetic field of the wave at that point and instant?
- Options 1. Positive direction of x
  - 2. Negative direction of y
  - 3. Positive direction of z
  - 4. Negative direction of x


Question Type : MCQ

Question ID: 3666941185 Option 1 ID: 3666943567 Option 2 ID: 3666943570 Option 3 ID: 3666943569 Option 4 ID: 3666943568 Status: Answered

Chosen Option: 4

Section: Physics Section B

Q.21 A uniform electric field of 10 N/C is created between two parallel charged plates (as shown in figure). An electron enters the field symmetrically between the plates with a kinetic energy 0.5 eV. The length of each plate is 10 cm. The angle ( $\theta$ ) of deviation of the path of electron as it comes out of the field is degree).



Given --Answer:

> Question Type: SA Question ID: 3666941195 Status: Not Answered

A ray of light is incident from air on a glass plate having thickness  $\sqrt{3}$  cm and Q.22 refractive index  $\sqrt{2}$ . The angle of incidence of a ray is equal to the critical angle for glass-air interface. The lateral displacement of the ray when it passes through the plate is  $\times 10^{-2}$  cm. (given sin 15° = 0.26)

Given --Answer:

> Question Type: SA Question ID: 3666941192 Status : Not Answered

Q.23 An LCR series circuit of capacitance 62.5 nF and resistance of 50  $\Omega$ , is connected to an A.C. source of frequency 2.0 kHz. For maximum value of amplitude of current in circuit, the value of inductance is mH.

(Take  $\pi^2 = 10$ )

Given --Answer:

> Question Type: SA Question ID: 3666941193 Status: Not Answered

Q.24 As shown in the figure, in an experiment to determine Young's modulus of a wire, the extension-load curve is plotted. The curve is a straight line passing through the origin and makes an angle of  $45^{\circ}$  with the load axis. The length of wire is 62.8 cm and its diameter is 4 mm. The Young's modulus is found to be  $x \times 10^4$  Nm<sup>-2</sup>.

The value of x is \_\_\_\_\_\_\_ But the value of x is \_\_\_\_\_\_\_ A50

Given --Answer :

Question Type : SA

Question ID : 3666941197

Status : Not Answered

Q.25 The wavelength of the radiation emitted is  $\lambda_0$  when an electron jumps from the second excited state to the first excited state of hydrogen atom. If the electron jumps from the third excited state to the second orbit of the hydrogen atom, the wavelength of the radiation emitted will be  $\frac{20}{x}\lambda_0$ . The value of x is \_\_\_\_\_.


Given --Answer :

Question Type : SA

Question ID : 3666941191

Status : Not Answered

Q.26 An object of mass 'm' initially at rest on a smooth horizontal plane starts moving under the action of force F = 2N. In the process of its linear motion, the angle  $\theta$  (as shown in figure) between the direction of force and horizontal varies as  $\theta = kx$ , where k is a constant and x is the distance covered by the object from its initial position. The expression of kinetic energy of the object will be  $E = \frac{n}{k} \sin \theta$ , The value of n is \_\_\_\_\_\_.



Smooth horizontal surface

Given --Answer :

Question Type : SA

Question ID : 3666941199


Status : Not Answered

Q.27 If  $\vec{P} = 3\hat{i} + \sqrt{3} \hat{j} + 2\hat{k}$  and  $\vec{Q} = 4\hat{i} + \sqrt{3} \hat{j} + 2.5\hat{k}$  then, The unit vector in the direction of  $\vec{P} \times \vec{Q}$  is  $\frac{1}{x} (\sqrt{3}i + \hat{j} - 2\sqrt{3}\hat{k})$ . The value of x is

Given --Answer :

Question Type : SA
Question ID : 3666941200
Status : Not Answered

Q.28	In the given circuit,	the equivalent resistan	ice between the ten	ninal A and B is
	0			



Given --Answer:

> Question Type : SA Question ID: 3666941194 Status: Not Answered

Q.29 The distance between two consecutive points with phase difference of 60° in a wave of frequency 500 Hz is 6.0 m. The velocity with which wave is traveling is

Given --Answer:

> Question Type : SA Question ID: 3666941196 Status: Not Answered

I<sub>CM</sub> is the moment of inertia of a circular disc about an axis (CM) passing through Q.30 its center and perpendicular to the plane of disc. IAB is it's moment of inertia about an axis AB perpendicular to plane and parallel to axis CM at a distance  $\frac{2}{3}$ R from center. Where R is the radius of the disc. The ratio of  $I_{\mbox{\footnotesize{AB}}}$  and  $I_{\mbox{\footnotesize{CM}}}$  is x : 9. The value of x is \_\_\_\_\_



Given --Answer:

> Question Type: SA Question ID: 3666941198

Status: Not Answered

Section: Chemistry Section A

Q.31 The correct order in aqueous medium of basic strength in case of methyl substituted amines

Options 1.  $Me_3N > Me_2NH > MeNH_2 > NH_3$ 

- 2. Me<sub>2</sub>NH > Me<sub>3</sub>N > MeNH<sub>2</sub> > NH<sub>3</sub>
- 3. NH<sub>3</sub>> Me<sub>3</sub>N> MeNH<sub>2</sub>> Me<sub>2</sub>NH
- 4.  $Me_2NH > MeNH_2 > Me_3N > NH_3$

Question Type : MCQ

Question ID: 3666941217 Option 1 ID: 3666943666 Option 2 ID: 3666943665 Option 3 ID: 3666943667 Option 4 ID: 3666943668

Status: Answered Chosen Option: 4

Q.32 Inert gases have positive electron gain enthalpy. Its correct order is

Options 1. He  $\leq Xe \leq Kr \leq Ne$ 

- 2. He < Ne < Kr < Xe
- 3. He < Kr < Xe < Ne
- 4. Xe < Kr < Ne < He


Question Type: MCQ

Question ID: 3666941204 Option 1 ID: 3666943616 Option 2 ID: 3666943613 Option 3 ID: 3666943615 Option 4 ID: 3666943614

Status: Answered Chosen Option: 4

Q.33 In the cumene to phenol preparation in presence of air, the intermediate is

Options



Question Type: MCQ

Question ID: 3666941214 Option 1 ID: 3666943655 Option 2 ID: 3666943654 Option 3 ID: 3666943656 Option 4 ID: 3666943653 Status : **Answered** 

Chosen Option: 2

Q.34 Which of the following statements is incorrect for antibiotics?

Options 1.

An antibiotic is a synthetic substance produced as a structural analogue of naturally occurring

- 2. An antibiotic should promote the growth or survival of microorganisms.
- 3. An antibiotic should be effective in low concentrations.
- 4. An antibiotic must be a product of metabolism.


Question Type: MCQ


Option 1 ID: 3666943672 Option 2 ID: 3666943671 Option 3 ID: 3666943670


Question ID: 3666941218


Option 4 ID: 3666943669 Status : Answered

# Q.35 The variation of the rate of an enzyme catalyzed reaction with substrate concentration is correctly represented by graph









Options 1. (a)

- 2. (d)
- 3. (c)
- 4. (b)

Question Type : MCQ

Question ID: 3666941203

Option 1 ID: 3666943609

Option 2 ID: 3666943612 Option 3 ID: 3666943611

Option 4 ID: 3666943610

Status: Answered

Chosen Option: 2

### Q.36 Match List I with List II

LIST I Elements		c	LIST II  Colour imparted to the flame		
A.	K	I.	Brick Red		
B.	Ca	II.	Violet		
C.	Sr	III.	Apple Green		
D.	Ba	IV.	Crimson Red		

Choose the correct answer from the options given below:

Options 1. A-II, B-IV, C-I, D-III

- 2. A-II, B-I, C-IV, D-III
- 3. A-IV, B-III, C-II, D-I
- 4. A-II, B-I, C-III, D-IV

Question Type : MCQ

Question ID: 3666941209

Option 1 ID: 3666943636

Option 2 ID: 3666943634

Option 3 ID: 3666943635

Option 4 ID: 3666943633

Status: Answered

The correct sequence of reagents for the preparation of Q and R is :

(i) CrO<sub>2</sub>Cl<sub>2</sub>, H<sub>3</sub>O<sup>+</sup>; (ii) Cr<sub>2</sub>O<sub>3</sub>, 770 K, 20 atm; Options

- (iii) NaOH; (iv) H3O+
- 2. (i) KMnO<sub>4</sub>, OH<sup>-</sup>; (ii) Mo<sub>2</sub>O<sub>3</sub>, Δ; (iii) NaOH; (iv) H<sub>3</sub>O<sup>+</sup>
- (i)  $Cr_2O_3$ , 770 K, 20 atm; (ii)  $CrO_2Cl_2$ ,  $H_3O^+$ ;
  - (iii) NaOH; (iv) H3O+
- 4. (i) Mo<sub>2</sub>O<sub>3</sub>, Δ; (ii) CrO<sub>2</sub>Cl<sub>2</sub>, H<sub>3</sub>O<sup>+</sup>; (iii) NaOH; (iv) H<sub>3</sub>O<sup>+</sup>

Question Type : MCQ

Question ID: 3666941212

Option 1 ID: 3666943646

Option 2 ID: 3666943645

Option 3 ID: 3666943648

Option 4 ID: 3666943647

Status: Answered

Chosen Option: 2

- Q.38 Compound A reacts with NH<sub>4</sub>Cl and forms a compound B. Compound B reacts with H2O and excess of CO2 to form compound C which on passing through or reaction with saturated NaCl solution forms sodium hydrogen carbonate. Compound A, B and C, are respectively.
- Options 1.  $Ca(OH)_2$ ,  $NH_4^{\oplus}$ ,  $(NH_4)_2CO_3$ 
  - 2. Ca(OH)<sub>2</sub>, NH<sub>3</sub>, NH<sub>4</sub>HCO<sub>3</sub>
  - 3. CaCl2, NH3, NH4HCO3
  - 4. CaCl<sub>2.</sub> NH<sub>4</sub><sup>⊕</sup>, (NH<sub>4</sub>)<sub>2</sub>CO<sub>3</sub>

Question Type: MCQ

Question ID: 3666941207

Option 1 ID: 3666943626

Option 2 ID: 3666943628

Option 3 ID: 3666943625

Option 4 ID: 3666943627

Status: Answered

Q.39 The radius of the  $2^{nd}$  orbit of  $Li^{2+}$  is x. The expected radius of the  $3^{rd}$  orbit of  $Be^{3+}$  is

Question Type : MCQ

Question ID: 3666941202 Option 1 ID: 3666943606 Option 2 ID: 3666943608 Option 3 ID: 3666943607 Option 4 ID: 3666943605

Status: Answered Chosen Option: 3

Q.40 The compound which will have the lowest rate towards nucleophilic aromatic substitution on treatment with OH is

Options

NO2

Question Type : MCQ

Question ID: 3666941213 Option 1 ID: 3666943649 Option 2 ID: 3666943652 Option 3 ID: 3666943651 Option 4 ID: 3666943650 Status: Answered

#### Q.41 Identify the product formed (A and E)

Options

2. 
$$A = \bigcup_{NO_2}^{Me} Br$$
 ,  $E = \bigcup_{OH}^{COOH} Br$ 

$$A = Br$$
 $Br$ 
 $Br$ 
 $Br$ 
 $Br$ 

$$A = \begin{cases} Me \\ Br \\ NO_2 \end{cases}$$

$$E = \begin{cases} Me \\ Br \\ Br \\ Br \end{cases}$$

Question Type: MCQ

Question ID: 3666941216 Option 1 ID: 3666943663 Option 2 ID: 3666943664 Option 3 ID: 3666943661 Option 4 ID: 3666943662 Status: Answered

Chosen Option : 1

Q.42 Reaction of thionyl chloride with white phosphorus forms a compound [A], which on hydrolysis gives [B], a dibasic acid. [A] and [B] are respectively

Options 1. PCl<sub>3</sub> and H<sub>3</sub>PO<sub>3</sub>

2. P4O6 and H3PO3

3. PCl<sub>5</sub> and H<sub>3</sub>PO<sub>4</sub>

4. POCl<sub>3</sub> and H<sub>3</sub>PO<sub>4</sub>

Question Type: MCQ

Question ID: 3666941208 Option 1 ID: 3666943630 Option 2 ID: 3666943629 Option 3 ID: 3666943632

Option 4 ID: 3666943631 Status: Answered

Given below are two statements: one is labelled as Assertion A and the other is labelled as Reason R:

Assertion A: Acetal / Ketal is stable in basic medium.

Reason R: The high leaving tendency of alkoxide ion gives the stability to acetal / ketal in basic medium.

In the light of the above statements, choose the correct answer from the options given below:

Options 1. Both A and R are true but R is NOT the correct explanation of A

- 2. Both A and R are true and R is the correct explanation of A
- 3. A is false but R is true
- 4. A is true but R is false

Question Type: MCQ

Question ID: 3666941215 Option 1 ID: 3666943658 Option 2 ID: 3666943657 Option 3 ID: 3666943660 Option 4 ID: 3666943659

Status: Answered

Chosen Option: 2

### Q.44 '25 volume' hydrogen peroxide means

Options 1. 1 L marketed solution contains 75 g of H<sub>2</sub>O<sub>2</sub>.

- 2. 100 mL marketed solution contains 25 g of H2O2.
- 3. 1 L marketed solution contains 250 g of H<sub>2</sub>O<sub>2</sub>.
- 4. 1 L marketed solution contains 25 g of H<sub>2</sub>O<sub>2</sub>.

Question Type: MCQ

Chosen Option: 2

Question ID: 3666941206 Option 1 ID: 3666943623 Option 2 ID: 3666943622 Option 3 ID: 3666943624 Option 4 ID: 3666943621 Status: Answered

Q.45 Which one of the following reactions does not occur during extraction of copper?

Options 1. CaO + SiO<sub>2</sub> → CaSiO<sub>3</sub>

2. 2 FeS + 3  $O_2 \rightarrow$  2 FeO + 2  $SO_2$ 

3.  $2 \text{ Cu}_2\text{S} + 3 \text{ O}_2 \rightarrow 2 \text{ Cu}_2\text{O} + 2 \text{ SO}_2$ 

4. FeO + SiO<sub>2</sub> → FeSiO<sub>3</sub>

Question Type : MCQ

Question ID: 3666941205 Option 1 ID: 3666943620 Option 2 ID: 3666943618 Option 3 ID: 3666943619

Option 4 ID: 3666943617

Status: Answered

Q.46 A cubic solid is made up of two elements X and Y. Atoms of X are present on every alternate corner and one at the center of cube. Y is at  $\frac{1}{2}$ rd of the total faces. The empirical formula of the compound is

Options 1.  $X_2Y_{1.5}$ 

- 2. XY<sub>2.5</sub>
- 3. X<sub>2.5</sub>Y
- 4. X<sub>1.5</sub>Y<sub>2</sub>

Question Type: MCQ

Question ID: 3666941201 Option 1 ID: 3666943603 Option 2 ID: 3666943601 Option 3 ID: 3666943602

Option 4 ID: 3666943604 Status: Answered

Chosen Option: 4

#### Q.47 Match the List-I with List-II:

List-I	List-II		
Cations	Group reagents		
$A \rightarrow Pb^{2+}, Cu^{2+}$	i) H <sub>2</sub> S gas in presence of dilute HCl		
$B \rightarrow A1^{3+}$ , $Fe^{3+}$	ii) (NH <sub>4</sub> ) <sub>2</sub> CO <sub>3</sub> in presence of NH <sub>4</sub> OH		
$C \rightarrow Co^{2+}, Ni^{2+}$	iii) NH <sub>4</sub> OH in presence of NH <sub>4</sub> Cl		
$D \rightarrow Ba^{2+}, Ca^{2+}$	iv) H <sub>2</sub> S in presence of NH <sub>4</sub> OH		

Correct match is -

Options 1.  $A \rightarrow iv$ ,  $B \rightarrow ii$ ,  $C \rightarrow iii$ ,  $D \rightarrow i$ 

- 2.  $A \rightarrow i$ ,  $B \rightarrow iii$ ,  $C \rightarrow iv$ ,  $D \rightarrow ii$
- 3.  $A \rightarrow iii$ ,  $B \rightarrow i$ ,  $C \rightarrow iv$ ,  $D \rightarrow ii$
- 4.  $A \rightarrow i$ ,  $B \rightarrow iii$ ,  $C \rightarrow ii$ ,  $D \rightarrow iv$

Question Type :  $\boldsymbol{\mathsf{MCQ}}$ 

Question ID: 3666941220 Option 1 ID: 3666943680

Option 2 ID: 3666943677 Option 3 ID: 3666943678

Option 4 ID: 3666943679

Status: Answered

Q.48 Some reactions of NO2 relevant to photochemical smog formation are

$$\begin{array}{c} NO_2 \xrightarrow{\text{sunlight}} X + Y \\ \downarrow A \\ B \end{array}$$

Identify A, B, X and Y

Options 1. 
$$X = [O]$$
,  $Y = NO$ ,  $A = O_2$ ,  $B = O_3$ 

2. 
$$X = \frac{1}{2}O_2$$
,  $Y = NO_2$ ,  $A = O_3$ ,  $B = O_2$ 

3. 
$$X = N_2O$$
,  $Y = [O]$ ,  $A = O_3$ ,  $B = NO$ 

4. 
$$X = NO$$
,  $Y = [O]$ ,  $A = O_2$ ,  $B = N_2O_3$ 

Question Type: MCQ

Question ID: 3666941210


Option 1 ID: 3666943639

Option 2 ID: 3666943640

Option 3 ID: 3666943637

Option 4 ID: 3666943638

Status: Answered



#### Q.50 Match items of Row I with those of Row II.

Row I:

Row II:

- (i) α-D-(-)-Fructofuranose,
- (ii) β-D-(-)-

- Fructofuranose
- (iii) α-D-(-) Glucopyranose,
- (iv) β-D-(-)-Glucopyranose

Correct match is

Options 1.  $A \rightarrow i$ ,  $B \rightarrow ii$ ,  $C \rightarrow iii$ ,  $D \rightarrow iv$ 

- 2.  $A \rightarrow iv$ ,  $B \rightarrow iii$ ,  $C \rightarrow i$ ,  $D \rightarrow ii$
- 3.  $A \rightarrow iii$ ,  $B \rightarrow iv$ ,  $C \rightarrow i$ ,  $D \rightarrow ii$
- 4. A  $\rightarrow$  iii, B  $\rightarrow$  iv, C  $\rightarrow$  ii, D  $\rightarrow$  i

Question Type : MCQ

Question ID: 3666941219 Option 1 ID: 3666943673 Option 2 ID: 3666943676 Option 3 ID: 3666943674 Option 4 ID: 3666943675

Status : Not Answered

Chosen Option: --

Section : Chemistry Section B

Q.51 How many of the following metal ions have similar value of spin only magnetic moment in gaseous state?

(Given: Atomic number : V, 23; Cr, 24; Fe, 26; Ni, 28)

Given 2 Answer:

Question Type : SA

Question ID : **3666941228** Status : **Answered** 

Q.52 An athlete is given 100 g of glucose (C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>) for energy. This is equivalent to 1800kJ of energy. The 50% of this energy gained is utilized by the athlete for sports activities at the event. In order to avoid storage of energy, the weight of extra water he would need to perspire is \_\_\_\_\_\_ g (Nearest integer)

Assume that there is no other way of consuming stored energy.

Given: The enthalpy of evaporation of water is 45 kJ mol<sup>-1</sup>

Molar mass of C, H & O are 12, 1 and 16 g mol<sup>-1</sup>.

Given --Answer :


Question Type : SA

Question ID : **3666941223**Status : **Not Answered** 

Q.53 The density of a monobasic strong acid (Molar mass 24.2 g/mol) is 1 its solution required for the complete neutralization of	.21 kg/L. The volume of
25 mL of 0.24 M NaOH is × 10 <sup>-2</sup> mL (Nearest integer)	
Given	
Answer:	
	Outseties Ture : CA
	Question Type : <b>SA</b> Question ID : <b>3666941221</b>
	Status: Not Answered
Q.54 The total number of lone pairs of electrons on oxygen atoms of	of ozone is
Given 1	
Answer:	
	Question Type : SA
	Question ID : 3666941222
	Status : Answered
0.471	20.000.010.700000.111.000
Q.55 In sulphur estimation, 0.471 g of an organic compound gave 1.44  The percentage of sulphur in the compound is(N	
(Given: Atomic mass Ba: 137 u, S: 32 u, O: 16 u)	
Given	
Answer:	
	Question Type : SA
	Question ID : 3666941230
	Status : Not Answered
Q.56 Consider the cell	
Q.56 Consider the cell	
$Pt(s) \mid H_2(g) (1 \text{ atm}) \mid H^+ (aq, [H^+] = 1) \parallel Fe^{3+} (aq), Fe^{3+} (aq) \mid H^+ (aq, [H^+] = 1) \mid H^+ (a$	2+(ag)   Pt(s)
1 ((3)   11½(g) (1 am)   11 (aq, [11 ] - 1)    10 (aq), 10	(44) (11(3)
Given $E_{Fe}^{3+}/F_{e}^{2+} = 0.771 \text{ V}$ and $E_{H}^{+}/_{\frac{1}{2}}/H_{2}^{2} = 0 \text{ V}$ , T	- 208 K
Given by Fe Fe -0.771 v and b H /1/2 H2 -0 v . 1	- 270 K
If the potential of the cell is 0.712 V, the ratio of conce	ntration of
Fe <sup>2+</sup> to Fe <sup>3+</sup> is (Nearest integer)	
Given	
Answer:	
	Question Type : <b>SA</b> Question ID : <b>3666941226</b>
	Status : Not Answered
Q.57 For the first order reaction $A \rightarrow B$ , the half life is 30 min. The time to	aken for 75% completion
of the reaction is min. (Nearest integer)	
Given: log 2 = 0.3010	
$\log 3 = 0.4771$	
log 5 = 0.6989	
Given Answer :	
	Question Type : SA
	Question ID : 3666941227 Status : Not Answered
	Status : NOT Allswelled

Q.58 The osmotic pressure of solutions of PVC in cyclohexanone at 300 K are plotted on the graph.

The molar mass of PVC is g mol<sup>-1</sup> (Nearest integer)



(Given :  $R = 0.083 L atm K^{-1} mol^{-1}$ )

Given --Answer :

Question Type : SA

Question ID : 3666941224

Status : Not Answered

Q.59 A litre of buffer solution contains 0.1 mole of each of NH<sub>3</sub> and NH<sub>4</sub>Cl. On the addition of

0.02 mole of HCl by dissolving . gaseous HCl, the pH of the solution is found to be  $\times$  10<sup>-3</sup> (Nearest integer)

[Given:  $pK_b(NH_3) = 4.745$ 

log 2 = 0.301 log 3 = 0.477 T = 298 K ]

Given --Answer :

Question Type : SA

Question ID : 3666941225

Status : Not Answered

Q.60 The number of paramagnetic species from the following is \_\_\_\_\_\_.

[Ni(CN)<sub>4</sub>]<sup>2-</sup>, [Ni(CO)<sub>4</sub>], [NiCl<sub>4</sub>]<sup>2-</sup>,

 $[Fe(CN)_6]^{4-}$ ,  $[Cu(NH_3)_4]^{2+}$ 

 $[Fe(CN)_6]^{3-}$  and  $[Fe(H_2O)_6]^{2+}$ 

Given --Answer :

Question Type : SA

Question ID : 3666941229

Status : Not Answered

Section: Mathematics Section A

Q.61 The distance of the point P (4, 6, -2) from the line passing through the point (-3, 2, 3) and parallel to a line with direction ratios 3, 3, -1 is equal to:

Options 1.  $2\sqrt{3}$ 

- 2. 3
- 3.  $\sqrt{14}$
- 4.  $\sqrt{6}$

Question Type : MCQ

Question ID: 3666941245 Option 1 ID: 3666943747

Option 2 ID: 3666943748 Option 3 ID: 3666943750

Option 4 ID: 3666943749 Status: Not Answered

Chosen Option: --

Q.62 The mean and variance of the marks obtained by the students in a test are 10 and 4 respectively. Later, the marks of one of the students is increased from 8 to 12. If the new mean of the marks is 10.2, then their new variance is equal to:

Options 1. 3.96

- 2. 3.92
- 3. 4.04
- 4.4.08

Question Type: MCQ

Question ID: 3666941247 Option 1 ID: 3666943758 Option 2 ID: 3666943755 Option 3 ID: 3666943756

Option 4 ID: 3666943757 Status: Answered

Chosen Option: 3

Q.63 The statement  $(p \land (\sim q)) \Rightarrow (p \Rightarrow (\sim q))$  is

Options 1. a contradiction

- 2. equivalent to  $p \vee q$
- 3. a tautology
- 4. equivalent to  $(-p)\vee(-q)$

Question Type : MCQ

Question ID: 3666941250 Option 1 ID: 3666943770 Option 2 ID: 3666943767 Option 3 ID: 3666943769 Option 4 ID: 3666943768 Status: Answered

Q.64 Let  $S_1$  and  $S_2$  be respectively the sets of all  $a \in \mathbb{R} - \{0\}$  for which the system of linear equations

$$ax + 2ay - 3az = 1$$

$$(2a+1)x + (2a+3)y + (a+1)z = 2$$

$$(3a+5)x+(a+5)y+(a+2)z=3$$

has unique solution and infinitely many solutions. Then

Options 1.  $n(S_1) = 2$  and  $S_2$  is an infinite set

2. 
$$S_1 = \mathbb{R} - \{0\}$$
 and  $S_2 = \Phi$ 

3. 
$$S_1$$
 is an infinite set and  $n(S_2) = 2$ 

4. 
$$S_1 = \Phi$$
 and  $S_2 = \mathbb{R} - \{0\}$ 

Question Type : MCQ

Question ID: 3666941232

Option 1 ID: 3666943697

Option 2 ID : **3666943695** 

Option 3 ID : **3666943698** 

Option 4 ID : 3666943696

Status: Not Answered

Chosen Option : --

Q.65 Let  $x, y, z \ge 1$  and  $A = \begin{bmatrix} 1 & \log_x y & \log_x z \\ \log_y x & 2 & \log_y z \\ \log_z x & \log_z y & 3 \end{bmatrix}$ . Then |adj (adj A<sup>2</sup>)| is equal to

Options 1. 48

2. 24

3. 28

4. 64

Question Type : MCQ

Question ID: 3666941233

Option 1 ID: 3666943701

Option 2 ID : **3666943699** 

Option 3 ID: 3666943700

Option 4 ID: 3666943702

Status: Not Answered

Q.66 Let x = 2 be a local minima of the function  $f(x) = 2x^4 - 18x^2 + 8x + 12$ ,  $x \in (-4, 4)$ . If M is local maximum value of the function f in (-4, 4), then M = (-4, 4).

Options

1. 
$$18\sqrt{6} - \frac{33}{2}$$

2. 
$$12\sqrt{6} - \frac{31}{2}$$

3. 
$$18\sqrt{6} - \frac{31}{2}$$

4. 
$$12\sqrt{6} - \frac{33}{2}$$

Question Type : MCQ

Question ID : 3666941237 Option 1 ID : 3666943715 Option 2 ID : 3666943716

Option 3 ID : **3666943718** Option 4 ID : **3666943717** 

Status : Not Answered

Chosen Option: --

Q.67 Let 
$$f(x) = \int \frac{2x}{(x^2 + 1)(x^2 + 3)} dx$$
. If  $f(3) = \frac{1}{2} (\log_e 5 - \log_e 6)$ , then  $f(4)$  is equal to

Options 1.  $\log_e 17 - \log_e 18$ 

$$2.\frac{1}{2}(\log_e 19 - \log_e 17)$$

3. 
$$\frac{1}{2} (\log_e 17 - \log_e 19)$$

Question Type : MCQ

Question ID: 3666941239 Option 1 ID: 3666943726 Option 2 ID: 3666943724 Option 3 ID: 3666943725

Option 4 ID: **3666943723** Status: **Not Answered** 

Chosen Option : --

Q.68 The vector  $\vec{a} = -\hat{i} + 2\hat{j} + \hat{k}$  is rotated through a right angle, passing through the y-axis in its way and the resulting vector is  $\vec{b}$ . Then the projection of  $3\vec{a} + \sqrt{2}\vec{b}$  on  $\vec{c} = 5\hat{i} + 4\hat{j} + 3\hat{k}$  is:

Options 1.  $2\sqrt{3}$ 

2. 3√2

3 1

4.  $\sqrt{6}$ 

Question Type: MCQ

Question ID: 3666941246 Option 1 ID: 3666943754 Option 2 ID: 3666943753 Option 3 ID: 3666943751

Option 4 ID : **3666943752** Status : **Not Answered** 

```
Q.69 The points of intersection of the line ax + by = 0, (a \ne b) and the circle x^2 + y^2 - 2x = 0 are A (\alpha, 0) and B (1, \beta). The image of the circle with AB as a diameter in the line x + y + 2 = 0 is:
```

Options 1. 
$$x^2 + y^2 + 3x + 5y + 8 = 0$$

2. 
$$x^2 + y^2 - 5x - 5y + 12 = 0$$

3. 
$$x^2 + y^2 + 3x + 3y + 4 = 0$$

4. 
$$x^2 + y^2 + 5x + 5y + 12 = 0$$

Question Type : MCQ

Question ID: 3666941244 Option 1 ID: 3666943745 Option 2 ID: 3666943743 Option 3 ID: 3666943746 Option 4 ID: 3666943744

Status: Not Answered

Chosen Option : --

Q.70

Let  $f:(0,1) \to \mathbb{R}$  be a function defined by  $f(x) = \frac{1}{1 - e^{-x}}$ , and

g(x) = (f(-x) - f(x)). Consider two statements

(I) g is an increasing function in (0, 1)

(II) g is one-one in (0, 1)

Then,

Options 1. Only (II) is true

- 2. Both (I) and (II) are true
- 3. Only (I) is true
- 4. Neither (I) nor (II) is true

Question Type :  $\boldsymbol{\mathsf{MCQ}}$ 

Option 1 ID: 3666943740
Option 2 ID: 3666943741
Option 3 ID: 3666943739
Option 4 ID: 3666943742
Status: Not Answered

Question ID: 3666941243

Chosen Option : --

Q.71 Let 
$$y(x) = (1+x)(1+x^2)(1+x^4)(1+x^8)(1+x^{16})$$
. Then  $y' - y''$  at  $x = -1$  is equal to:

Options 1. 976

2. 464

3.496

4. 944

Question Type :  $\ensuremath{\mathsf{MCQ}}$ 

Question ID: 3666941238 Option 1 ID: 3666943721 Option 2 ID: 3666943720 Option 3 ID: 3666943719

Option 4 ID : 3666943722 Status : Not Answered

Options

Question Type: MCQ

Question ID: 3666941248 Option 1 ID: 3666943760 Option 2 ID: 3666943759

Option 3 ID: 3666943761 Option 4 ID: 3666943762

Status: Not Answered

Chosen Option: --

Q.73 The distance of the point  $(6, -2\sqrt{2})$  from the common tangent y = mx + c, m > 0, of the curves  $x = 2y^2$  and  $x = 1 + y^2$  is:

Options 1.

3. 5√3

4. 5

Question Type: MCQ

Question ID: 3666941242

Option 1 ID: 3666943737 Option 2 ID: 3666943735

Option 3 ID: 3666943738

Option 4 ID: 3666943736

Status: Not Answered

Q.74 Consider the lines  $L_1$  and  $L_2$  given by

$$L_1: \frac{x-1}{2} = \frac{y-3}{1} = \frac{z-2}{2}$$

$$L_2: \frac{x-2}{1} = \frac{y-2}{2} = \frac{z-3}{3}$$
.

A line  $L_3$  having direction ratios 1, -1, -2, intersects  $L_1$  and  $L_2$  at the points P and Q respectively. Then the length of line segment PQ is

Options 1.  $2\sqrt{6}$ 

- 2. 4
- 3.  $3\sqrt{2}$
- 4.  $4\sqrt{3}$

Question Type: MCQ

Question ID: 3666941236

Option 1 ID: 3666943711

Option 2 ID: 3666943714

Option 3 ID: 3666943713

Option 4 ID: 3666943712

Status: Not Answered

Chosen Option: --

The value of  $\lim_{n\to\infty} \frac{1+2-3+4+5-6+\dots+(3n-2)+(3n-1)-3n}{\sqrt{2n^4+4n+3}-\sqrt{n^4+5n+4}}$  is: Q.75

Options 1.  $\frac{\sqrt{2}+1}{2}$ 

- 2. 3  $(\sqrt{2}+1)$
- $3. \frac{3}{2\sqrt{2}}$
- 4.  $\frac{3}{2}(\sqrt{2}+1)$

Question Type: MCQ

Question ID: 3666941235

Option 1 ID: 3666943707

Option 2 ID: 3666943710

Option 3 ID: 3666943708

Option 4 ID: 3666943709 Status: Not Answered

Q.76 Let y = y(x) be the solution curve of the differential equation

$$\frac{dy}{dx} = \frac{y}{x} \left( 1 + xy^2 \left( 1 + \log_e x \right) \right), x > 0, \quad y(1) = 3 \quad \text{Then} \quad \frac{y^2(x)}{9} \quad \text{is equal to} :$$

Ontions

$$\int_{1}^{3} \frac{x^2}{7 - 3x^3(2 + \log_e x^2)}$$

$$2. \frac{x^2}{5 - 2x^3(2 + \log_e x^3)}$$

$$3. \frac{x^2}{3x^3(1 + \log_e x^2) - 2}$$

$$4. \frac{x^2}{2x^3(2 + \log_e x^3) - 3}$$

Question Type : MCQ

Question ID : 3666941241

Option 1 ID: 3666943732

Option 2 ID: 3666943731

Option 3 ID: 3666943733

Option 4 ID: 3666943734

Status: Not Answered

Chosen Option: --

Q.77 If  $a_r$  is the coefficient of  $x^{10-r}$  in the Binomial expansion of  $(1+x)^{10}$ , then

$$\sum_{r=1}^{10} r^3 \left( \frac{a_r}{a_{r-1}} \right)^2$$
 is equal to

Options 1. 1210

2. 5445

3. 3025

4. 4895

Question Type : MCQ

Question ID: 3666941234

Option 1 ID: 3666943706

Option 2 ID: 3666943703

Option 3 ID: 3666943705

Option 4 ID: 3666943704

Status: Not Answered

Q.78 Let 
$$z_1 = 2 + 3i$$
 and  $z_2 = 3 + 4i$ . The set

$$S = \left\{ z \in \mathbb{C} : \left| z - z_1 \right|^2 - \left| z - z_2 \right|^2 = \left| z_1 - z_2 \right|^2 \right\} \text{ represents a}$$

Options 1.

straight line with the sum of its intercepts on the coordinate axes equals 14

straight line with the sum of its intercepts on the coordinate axes equals -18

- 3. hyperbola with eccentricity 2
- 4. hyperbola with the length of the transverse axis 7

Question Type: MCQ

Question ID: 3666941231 Option 1 ID: 3666943692 Option 2 ID: 3666943694 Option 3 ID: 3666943691 Option 4 ID: 3666943693 Status: Answered

Chosen Option: 2

Q.79

Let  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  be three non zero vectors such that  $\vec{b} \cdot \vec{c} = 0$  and  $\vec{a} \times (\vec{b} \times \vec{c}) = \frac{\vec{b} - \vec{c}}{2}$ .

If  $\vec{d}$  be a vector such that  $\vec{b} \cdot \vec{d} = \vec{a} \cdot \vec{b}$ , then  $(\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d})$  is equal to

Options

Question Type : MCQ

Question ID: 3666941249 Option 1 ID: 3666943766 Option 2 ID: 3666943763 Option 3 ID: 3666943765

Option 4 ID: 3666943764

Status: Not Answered

The minimum value of the function  $f(x) = \int_{0}^{2} e^{|x-t|} dt$  is:

Options 1. 2e-1

- 2. 2
- 3. 2(e-1)
- 4. e(e-1)

Question Type : MCQ

Question ID: 3666941240 Option 1 ID: 3666943727 Option 2 ID: 3666943730 Option 3 ID: 3666943729

Option 4 ID : 3666943728
Status : Not Answered

Chosen Option : --

Section: Mathematics Section B

Q.81 Let x and y be distinct integers where  $1 \le x \le 25$  and  $1 \le y \le 25$ . Then, the number of ways of choosing x and y, such that x + y is divisible by 5, is \_\_\_\_\_.

Given --Answer :

Question Type : SA

Question ID : **3666941257**Status : **Not Answered** 

Q.82 Let  $S = \left\{ \alpha : \log_2(9^{2\alpha - 4} + 13) - \log_2\left(\frac{5}{2} \cdot 3^{2\alpha - 4} + 1\right) = 2 \right\}$ . Then the maximum

value of  $\beta$  for which the equation  $x^2 - 2\left(\sum_{\alpha \in S} \alpha\right)^2 x + \sum_{\alpha \in S} (\alpha + 1)^2 \beta = 0$  has real

roots, is \_\_\_\_\_

Given --Answer :

Question Type : SA

Question ID : 3666941252 Status : Not Answered

Q.83 It the area enclosed by the parabolas  $P_1: 2y = 5x^2$  and  $P_2: x^2 - y + 6 = 0$  is equal to the area enclosed by  $P_1$  and  $y = \alpha x$ ,  $\alpha > 0$ , then  $\alpha^3$  is equal to \_\_\_\_\_.

Given --Answer :

Question Type : SA

Question ID : **3666941258** Status : **Not Answered** 

Q.84	Let the equation of the plane passing through the line
	x-2y-z-5=0=x+y+3z-5 and parallel to the line
	x + y + 2z - 7 = 0 = 2x + 3y + z - 2 be $ax + by + cz = 65$ . Then the distance of the
	point (a, b, c) from the plane $2x + 2y - z + 16 = 0$ is

Given --Answer :

Question Type : **SA**Question ID : **3666941259**Status : **Not Answered** 

The constant term in the expansion of  $\left(2x + \frac{1}{x^7} + 3x^2\right)^5$  is \_\_\_\_\_.

Given --Answer :

Question Type : SA

Question ID : 3666941254

Status : Not Answered

Q.86 The vertices of a hyperbola H are  $(\pm 6, 0)$  and its eccentricity is  $\frac{\sqrt{5}}{2}$ . Let N be the normal to H at a point in the first quadrant and parallel to the line  $\sqrt{2}x + y = 2\sqrt{2}$ .

If d is the length of the line segment of N between H and the y-axis then  $d^2$  is equal to

Given --Answer :

Question Type : SA

Question ID : 3666941256

Status : Not Answered

Q.87 If the sum of all the solutions of  $\tan^{-1}\left(\frac{2x}{1-x^2}\right) + \cot^{-1}\left(\frac{1-x^2}{2x}\right) = \frac{\pi}{3}$ , -1 < x < 1,  $x \ne 0$ , is  $\alpha - \frac{4}{\sqrt{3}}$ , then  $\alpha$  is equal to \_\_\_\_\_.

Given --Answer :

Question Type : **SA**Question ID : **3666941260**Status : **Not Answered** 

Q.88 Let  $S = \{1, 2, 3, 5, 7, 10, 11\}$ . The number of non-empty subsets of S that have the sum of all elements a multiple of 3, is

Given --Answer :

Question Type : SA

Question ID : 3666941253

Status : Not Answered

Q.89 Let  $A_1, A_2, A_3$  be the three A.P. with the same common difference d and having their first terms as A, A+1, A+2, respectively. Let a, b, c be the 7<sup>th</sup>, 9<sup>th</sup>, 17<sup>th</sup> terms of  $A_1, A_2, A_3$ , respectively such that  $\begin{vmatrix} a & 7 & 1 \\ 2b & 17 & 1 \\ c & 17 & 1 \end{vmatrix} + 70 = 0.$ 

If a = 29, then the sum of first 20 terms of an AP whose first term is c-a-b and common difference is  $\frac{d}{12}$ , is equal to\_\_\_\_\_.

Given --Answer :

Question Type : **SA**Question ID : **3666941255** 

Status: Not Answered

Q.90 For some a, b,  $c \in \mathbb{N}$ , let f(x) = ax - 3 and  $g(x) = x^b + c$ ,  $x \in \mathbb{R}$ . If  $(f \circ g)^{-1}(x) = \left(\frac{x-7}{2}\right)^{1/3}$ , then  $(f \circ g)(ac) + (g \circ f)(b)$  is equal to \_\_\_\_\_\_

Given --Answer :

Question Type :  $\ensuremath{\mathbf{SA}}$ 

Question ID : **3666941251**Status : **Not Answered**