

Methylome Plasticity as a Biomarker of Treatment Response in Small Cell Lung Cancer

Danielle B. Sacdalan^{1,3*}; **Sami Ul Haq**^{1,2*}; Zhan, LJ¹; Li, JJ^{1,4}; Philip, V¹; Bratman, SV^{1,4}; Liu, G^{1,4}; Lok, BH^{1,3,4}

1. Princess Margaret Cancer Centre; 2. Schulich School of Medicine and Dentistry, Western University; 3. Institute of Medical Science, Temerty Faculty of Medicine, University of Toronto; 4. Department of Medical Biophysics, Temerty of Faculty of Medicine, University of Toronto; *Contributed Equally to the Work

INTRODUCTION

Small cell lung cancer (SCLC) is an aggressive disease with poor treatment outcomes, in part due to epigenetic mechanisms driving tumour growth and resistance.




Figure 1. Methylome-defined prognostic clusters identified using cfMeDIP-seq. Median OS (mOS) low-risk cluster = 20mo high-risk cluster = 11mo

HYPOTHESIS

We hypothesize **changes in the cell free methylome** of SCLC **influence disease response** to therapy and **drive treatment resistance** via dysregulation of cancer-associated pathways.

METHODS




Figure 2. cfMeDIP-seq on paired pre-treatment and relapse SCLC liquid biopsies (N=37).




Figure 3. Change in methylation across pairs ranked. Lowest quartile of change = stable group. Clinical outcomes compared between groups.




Figure 4. Overall workflow for DMR analysis and GSEA approach to pathway identification.

RESULTS

A.

Age at Diagnosis	Median (range)	67 (41-88)
Sex	Male	23 (62%)
Smoking status	Ever smoker	34/37 (92%)
Pack Years	Median (range)	40 (0-100)
VALSG Stage	Extensive stage	26 (70%)
1st-line chemo only	Proportion of ES patients given EP only	20/26 (77%)
Vital Status*	Dead	35/37 (95%)

Figure 5. Patient demographics (A) and distribution of methylome change groups (B). Broken line represents cut-off for lowest quartile (i.e., stable methylome group)




Figure 6. Methylome plasticity with shorter time to relapse from start of 1L treatment. (Med. Diff: 66.5 days, 95% CI 6.0-146.0; $p = 0.041$) and are more likely to be platinum resistant (χ^2 test $p = 0.044$). As for best response to treatment, progressive disease only in plastic group ($n=5$), complete response only in stable group ($n=3$).






Figure 7. Top hallmark pathways between pre-treatment and relapse SCLC accounting for the relationship between promoter methylation and gene expression. Myogenesis and TGF- β signalling pass the FDR cut-off of <0.25 .

RESULTS

A.



B.




Figure 8. Methylome plasticity associated with worse PFS (A) and OS (B). Additional cox regression analysis. PFS: adjusted hazard ratio [aHR] = 2.7, $p = 0.015$. OS: aHR = 2.4, $p = 0.039$.

DISCUSSION




Figure 9. Tumors with plastic methylomes demonstrate shorter periods of disease control with first-line treatment. Changes in methylation in during therapy are an efficient mechanism of response to treatment.

FUTURE DIRECTIONS

Confirmation by knockdown and over-expression of key pathway genes for myogenesis and TGF- β pathways are underway to validate and results of DMR and GSEA analysis are underway.