

# Notes on Finite Differences

Sultan Aitzhan

February 19, 2020

## Contents

<b>1 Time stepping and finite differences: the whole line</b>	<b>1</b>
<b>2 Time stepping and finite differences: the half line</b>	<b>10</b>

## 1 Time stepping and finite differences: the whole line

Recall the equation we obtained for the surface elevation on the whole line:

$$\eta_{tt} = \eta_{xx} + \mu^2 \left( \frac{1}{3} \eta_{xxxx} + \eta_{xt} \int_{-\infty}^x \eta_t \, dx' + \eta_x^2 + \eta_t^2 + \eta \eta_{xx} + \partial_x^2 \left( \int_{-\infty}^x \eta_t \, dx' \right)^2 \right). \quad (1)$$

To do time stepping, introduce

$$u = \eta_t. \quad (2)$$

Also, note that

$$\partial_x^2 \left( \int_{-\infty}^x \eta_t \, dx' \right)^2 = 2(\eta_t^2 + \eta_{tx} \int_{-\infty}^x \eta_t \, dx')$$

Then, combining (2) and (1), we obtain a two-dimensional system:

$$\partial_t \begin{bmatrix} u \\ \eta \end{bmatrix} = \begin{bmatrix} \mu^2 \left( 2u_x \int_{-\infty}^x u \, dx' + 2u^2 \right) + \eta_{xx} + \mu^2 \left( \frac{1}{3} \eta_{xxxx} + \eta_x^2 + \eta \eta_{xx} \right) \\ u \end{bmatrix}. \quad (3)$$

Now, consider (1) on a finite interval  $[a, b]$ , and let partition the interval into  $n + 1$  points  $\{x_k\}_{k=0}^n$ , with  $x_0 = a$  and  $x_n = b$ . This means that the integral terms becomes

$$\int_{-\infty}^x \eta_t \, dx' = \left\{ \int_{-\infty}^a + \int_a^x \right\} \eta_t \, dx' \approx \int_a^x \eta_t \, dx',$$

while assuming that

$$\int_{-\infty}^a \eta_t \, dx'$$

is small enough. Now, we employ forward Euler time stepping. First, observe that

$$\begin{aligned} u_t(x_k, t_j) &= \frac{u(x_k, t_{j+1}) - u(x_k, t_j)}{\Delta t} = f_1(\eta, u, t) & \implies u(x_k, t_{j+1}) &= u(x_k, t_j) + \Delta t f_1(\eta, u, t) \\ \eta_t(x_k, t_j) &= \frac{\eta(x_k, t_{j+1}) - \eta(x_k, t_j)}{\Delta t} = f_2(\eta, u, t) & \implies \eta(x_k, t_{j+1}) &= \eta(x_k, t_j) + \Delta t f_2(\eta, u, t), \end{aligned}$$

where

$$\begin{aligned} f_1(\eta, u, t) &= \eta_{xx} + \mu^2 \left( 2u_x \int_{-\infty}^x u \, dx' + 2u^2 + \frac{1}{3} \eta_{xxxx} + \eta_x^2 + \eta \eta_{xx} \right) \\ f_2(\eta, u, t) &= u(x_k, t_j). \end{aligned}$$

Observe that the highest order derivative is 4, so we need to use five-point stencils. In particular, we use five point midpoint stencil for  $x_2, \dots, x_{n-2}$ , and five point, one-sided stencils at  $x_0, x_1, x_{n-1}, x_n$ . First, consider the system for  $x_2, \dots, x_{n-2}$  :

$$\begin{aligned} f_1(\eta(x_k, t_j), u(x_k, t_j), t) &= \eta(x_k, t_j)_{xx} + \frac{\mu^2}{3} \eta(x_k, t_j)_{xxxx} \\ &\quad + \mu^2 \left( 2u(x_k, t_j)_x \int_{x_0}^{x_k} u \, dx' + 2u(x_k, t_j)^2 + \eta(x_k, t_j)_x^2 + \eta(x_k, t_j) \eta(x_k, t_j)_{xx} \right), \end{aligned}$$

where we have separated the linear and nonlinear terms. Let  $\Delta x = x_k - x_{k-1}$  and recall the finite difference formulas at  $x$  :

$$\begin{aligned} f'(x) &= \frac{f(x - 2\Delta x) - 8f(x - \Delta x) + 8f(x + \Delta x) - f(x + 2\Delta x)}{12\Delta x} + \mathcal{O}((\Delta x)^4) \\ f''(x) &= \frac{-f(x - 2\Delta x) + 16f(x - \Delta x) - 30f(x) + 16f(x + \Delta x) - f(x + 2\Delta x)}{12(\Delta x)^2} + \mathcal{O}((\Delta x)^2) \\ f'''(x) &= \frac{f(x - 2\Delta x) - 4f(x - \Delta x) + 6f(x) - 4f(x + \Delta x) + f(x + 2\Delta x)}{(\Delta x)^4} + \mathcal{O}((\Delta x)^2) \end{aligned}$$

so that

$$\begin{aligned}(\eta_k)_x &= \frac{\eta_{k-2} - 8\eta_{k-1} + 8\eta_{k+1} - \eta_{k+2}}{12\Delta x} + \mathcal{O}((\Delta x)^4) \\(u_k)_x &= \frac{u_{k-2} - 8u_{k-1} + 8u_{k+1} - u_{k+2}}{12\Delta x} + \mathcal{O}((\Delta x)^4) \\(\eta_k)_{xx} &= \frac{-\eta_{k-2} + 16\eta_{k-1} - 30\eta_k + 16\eta_{k+1} - \eta_{k+2}}{12(\Delta x)^2} + \mathcal{O}((\Delta x)^2) \\(\eta_k)_{xxxx} &= \frac{\eta_{k-2} - 4\eta_{k-1} + 6\eta_k - 4\eta_{k+1} + \eta_{k+2}}{(\Delta x)^4} + \mathcal{O}((\Delta x)^2),\end{aligned}$$

where it is assumed  $t = t_j$ . Also, by trapezoidal rule,

$$\int_{x_0}^{x_k} u \, dx' = \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} u \, dx = \frac{\Delta x}{2} \sum_{i=0}^{n-1} u(x_i) + u(x_{i+1}).$$

At  $x = x_0$ , we have

$$\begin{aligned}f'(x) &= \frac{-25f(x) + 48f(x + \Delta x) - 36f(x + 2\Delta x) + 16f(x + 3\Delta x) - 3f(x + 4\Delta x)}{12\Delta x} + \mathcal{O}((\Delta x)^4) \\f''(x) &= \frac{35f(x) - 104f(x + \Delta x) + 114f(x + 2\Delta x) - 56f(x + 3\Delta x) + 11f(x + 4\Delta x)}{12(\Delta x)^2} + \mathcal{O}((\Delta x)^2) \\f'''(x) &= \frac{f(x) - 4f(x + \Delta x) + 6f(x + 2\Delta x) - 4f(x + 3\Delta x) + f(x + 4\Delta x)}{(\Delta x)^4} + \mathcal{O}((\Delta x)^2),\end{aligned}$$

so that

$$\begin{aligned}(\eta_0)_x &= \frac{-25\eta_0 + 48\eta_1 - 36\eta_2 + 16\eta_3 - 3\eta_4}{12\Delta x} \\(u_0)_x &= \frac{-25u_0 + 48u_1 - 36u_2 + 16u_3 - 3u_4}{12\Delta x} \\(\eta_0)_{xx} &= \frac{35\eta_0 - 104\eta_1 + 114\eta_2 - 56\eta_3 + 11\eta_4}{12\Delta x^2} \\(\eta_0)_{xxxx} &= \frac{\eta_0 - 4\eta_1 + 6\eta_2 - 4\eta_3 + \eta_4}{\Delta x^4}\end{aligned}$$

At  $x = x_1$ , we have

$$f'(x) = \frac{-3f(x - \Delta x) - 10f(x) + 18f(x + \Delta x) - 6f(x + 2\Delta x) + f(x + 3\Delta x)}{12\Delta x} + \mathcal{O}((\Delta x)^4)$$

$$f''(x) = \frac{11f(x - \Delta x) - 20f(x) + 6f(x + \Delta x) + 4f(x + 2\Delta x) - f(x + 3\Delta x)}{12(\Delta x)^2} + \mathcal{O}((\Delta x)^2)$$

$$f'''(x) = \frac{f(x - \Delta x) - 4f(x) + 6f(x + \Delta x) - 4f(x + 2\Delta x) + f(x + 3\Delta x)}{(\Delta x)^4} + \mathcal{O}((\Delta x)^2),$$

so that

$$(\eta_1)_x = \frac{-3\eta_0 - 10\eta_1 + 18\eta_2 - 6\eta_3 + \eta_4}{12\Delta x}$$

$$(u_1)_x = \frac{-3u_0 - 10u_1 + 18u_2 - 6u_3 + u_4}{12\Delta x}$$

$$(\eta_1)_{xx} = \frac{11\eta_0 - 20\eta_1 + 6\eta_2 + 4\eta_3 - \eta_4}{12\Delta x^2}$$

$$(\eta_1)_{xxxx} = \frac{\eta_0 - 4\eta_1 + 6\eta_2 - 4\eta_3 + \eta_4}{\Delta x^4}$$

At  $x = x_{n-1}$ , we have

$$f'(x) = \frac{-f(x - 3\Delta x) + 6f(x - 2\Delta x) - 18f(x - \Delta x) + 10f(x) + 3f(x + \Delta x)}{12\Delta x} + \mathcal{O}((\Delta x)^4)$$

$$f''(x) = \frac{-f(x - 3\Delta x) + 4f(x - 2\Delta x) + 6f(x - \Delta x) - 20f(x) + 11f(x + \Delta x)}{12(\Delta x)^2} + \mathcal{O}((\Delta x)^2)$$

$$f'''(x) = \frac{f(x - 3\Delta x) - 4f(x - 2\Delta x) + 6f(x - \Delta x) - 4f(x) + f(x + \Delta x)}{(\Delta x)^4} + \mathcal{O}((\Delta x)^2),$$

so that

$$(\eta_{n-1})_x = \frac{-\eta_{n-4} + 6\eta_{n-3} - 18\eta_{n-2} + 10\eta_{n-1} + 3\eta_n}{12\Delta x}$$

$$(u_{n-1})_x = \frac{-u_{n-4} + 6u_{n-3} - 18u_{n-2} + 10u_{n-1} + 3u_n}{12\Delta x}$$

$$(\eta_{n-1})_{xx} = \frac{-\eta_{n-4} + 4\eta_{n-3} + 6\eta_{n-2} - 20\eta_{n-1} + 11\eta_n}{12\Delta x^2}$$

$$(\eta_{n-1})_{xxxx} = \frac{\eta_{n-4} - 4\eta_{n-3} + 6\eta_{n-2} - 4\eta_{n-1} + \eta_n}{\Delta x^4}$$

At  $x = x_n$ , we have

$$f'(x) = \frac{f(x - 4\Delta x) - 4f(x - 3\Delta x) + 6f(x - 2\Delta x) - 4f(x - \Delta x) + f(x)}{12\Delta x} + \mathcal{O}((\Delta x)^4)$$

$$f''(x) = \frac{11f(x-4\Delta x) - 56f(x-3\Delta x) + 114f(x-2\Delta x) - 104f(x-\Delta x) + 35f(x)}{12(\Delta x)^2} + \mathcal{O}((\Delta x)^2)$$

$$f'''(x) = \frac{f(x-4\Delta x) - 4f(x-3\Delta x) + 6f(x-2\Delta x) - 4f(x-\Delta x) + f(x)}{(\Delta x)^4} + \mathcal{O}((\Delta x)^2),$$

so that

$$(\eta_n)_x = \frac{3\eta_{n-4} - 16\eta_{n-3} + 36\eta_{n-2} - 48\eta_{n-1} + 25\eta_n}{12\Delta x}$$

$$(u_n)_x = \frac{3u_{n-4} - 16u_{n-3} + 36u_{n-2} - 48u_{n-1} + 25u_n}{12\Delta x}$$

$$(\eta_n)_{xx} = \frac{11\eta_{n-4} - 56\eta_{n-3} + 114\eta_{n-2} - 104\eta_{n-1} + 35\eta_n}{12\Delta x^2}$$

$$(\eta_n)_{xxx} = \frac{\eta_{n-4} - 4\eta_{n-3} + 6\eta_{n-2} - 4\eta_{n-1} + \eta_n}{\Delta x^4}.$$

All in all, we obtain:

$$f_1(\eta(x_0, t_j), u(x_0, t_j), t) = \eta(x_0, t_j)_{xx} + \frac{\mu^2}{3}\eta(x_0, t_j)_{xxx} + \mu^2 (2u(x_0, t_j)^2 + \eta(x_0, t_j)_x^2 + \eta(x_0, t_j)\eta(x_0, t_j)_{xx}),$$

$$f_1(\eta(x_1, t_j), u(x_1, t_j), t) = \eta(x_1, t_j)_{xx} + \frac{\mu^2}{3}\eta(x_1, t_j)_{xxx}$$

$$+ \mu^2 \left( 2u(x_1, t_j)_x \int_{x_0}^{x_1} u \, dx' + 2u(x_1, t_j)^2 + \eta(x_1, t_j)_x^2 + \eta(x_1, t_j)\eta(x_1, t_j)_{xx} \right),$$

$$f_1(\eta(x_2, t_j), u(x_2, t_j), t) = \eta(x_2, t_j)_{xx} + \frac{\mu^2}{3}\eta(x_2, t_j)_{xxx}$$

$$+ \mu^2 \left( 2u(x_2, t_j)_x \int_{x_0}^{x_2} u \, dx' + 2u(x_2, t_j)^2 + \eta(x_2, t_j)_x^2 + \eta(x_2, t_j)\eta(x_2, t_j)_{xx} \right),$$

$$\dots$$

$$f_1(\eta(x_k, t_j), u(x_k, t_j), t) = \eta(x_k, t_j)_{xx} + \frac{\mu^2}{3}\eta(x_k, t_j)_{xxx}$$

$$+ \mu^2 \left( 2u(x_k, t_j)_x \int_{x_0}^{x_k} u \, dx' + 2u(x_k, t_j)^2 + \eta(x_k, t_j)_x^2 + \eta(x_k, t_j)\eta(x_k, t_j)_{xx} \right),$$

$$\dots$$

$$f_1(\eta(x_{n-1}, t_j), u(x_{n-1}, t_j), t) = \eta(x_{n-1}, t_j)_{xx} + \frac{\mu^2}{3}\eta(x_{n-1}, t_j)_{xxx}$$

$$\begin{aligned}
& + \mu^2 \left( 2u(x_{n-1}, t_j)_x \int_{x_0}^{x_{n-1}} u \, dx' + 2u(x_{n-1}, t_j)^2 + \eta(x_{n-1}, t_j)_x^2 + \eta(x_{n-1}, t_j)\eta(x_{n-1}, t_j)_{xx} \right), \\
f_1(\eta(x_n, t_j), u(x_n, t_j), t) &= \eta(x_n, t_j)_{xx} + \frac{\mu^2}{3} \eta(x_n, t_j)_{xxxx} \\
& + \mu^2 \left( 2u(x_n, t_j)_x \int_{x_0}^{x_n} u \, dx' + 2u(x_n, t_j)^2 + \eta(x_n, t_j)_x^2 + \eta(x_n, t_j)\eta(x_n, t_j)_{xx} \right),
\end{aligned}$$

Now, we obtain the discretised problem. First, consider the column of linear terms:

$$\begin{aligned}
& (\eta_0)_{xx} + \frac{\mu^2}{3} (\eta_0)_{xxxx} \\
&= \frac{35\eta_0 - 104\eta_1 + 114\eta_2 - 56\eta_3 + 11\eta_4}{12\Delta x^2} + \frac{\mu^2}{3} \frac{\eta_0 - 4\eta_1 + 6\eta_2 - 4\eta_3 + \eta_4}{\Delta x^4} \\
&= \frac{35\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_0 - \frac{104\Delta x^2 + 16\mu^2}{12\Delta x^4} \eta_1 + \frac{114\Delta x^2 + 24\mu^2}{12\Delta x^4} \eta_2 - \frac{56\Delta x^2 + 16\mu^2}{12\Delta x^4} \eta_3 + \frac{11\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_4 \\
& (\eta_1)_{xx} + \frac{\mu^2}{3} (\eta_1)_{xxxx} \\
&= \frac{11\eta_0 - 20\eta_1 + 6\eta_2 + 4\eta_3 - \eta_4}{12\Delta x^2} + \frac{\mu^2}{3} \frac{\eta_0 - 4\eta_1 + 6\eta_2 - 4\eta_3 + \eta_4}{\Delta x^4} \\
&= \frac{11\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_0 - \frac{20\Delta x^2 + 16\mu^2}{12\Delta x^4} \eta_1 + \frac{6\Delta x^2 + 24\mu^2}{12\Delta x^4} \eta_2 + \frac{4\Delta x^2 - 16\mu^2}{12\Delta x^4} \eta_3 + \frac{-\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_4 \\
& \dots \\
& (\eta_k)_{xx} + \frac{\mu^2}{3} (\eta_k)_{xxxx} \\
&= \frac{-\eta_{k-2} + 16\eta_{k-1} - 30\eta_k + 16\eta_{k+1} - \eta_{k+2}}{12(\Delta x)^2} + \frac{\mu^2}{3} \frac{\eta_{k-2} - 4\eta_{k-1} + 6\eta_k - 4\eta_{k+1} + \eta_{k+2}}{(\Delta x)^4} \\
&= \frac{-\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_{k-2} + \frac{16(\Delta x^2 - \mu^2)}{12\Delta x^4} \eta_{k-1} + \frac{-30\Delta x^2 + 24\mu^2}{12\Delta x^4} \eta_k - \frac{16(\Delta x^2 - \mu^2)}{12\Delta x^4} \eta_{k+1} + \frac{-\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_{k+2} \\
& \dots \\
& (\eta_{n-1})_{xx} + \frac{\mu^2}{3} (\eta_{n-1})_{xxxx} \\
&= \frac{-\eta_{n-4} + 4\eta_{n-3} + 6\eta_{n-2} - 20\eta_{n-1} + 11\eta_n}{12\Delta x^2} + \frac{\mu^2}{3} \frac{\eta_{n-4} - 4\eta_{n-3} + 6\eta_{n-2} - 4\eta_{n-1} + \eta_n}{\Delta x^4}
\end{aligned}$$

$$\begin{aligned}
&= \frac{-\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_{n-4} + \frac{4\Delta x^2 - 16\mu^2}{12\Delta x^4} \eta_{n-3} + \frac{6\Delta x^2 + 24\mu^2}{12\Delta x^4} \eta_{n-2} - \frac{20\Delta x^2 + 16\mu^2}{12\Delta x^4} \eta_{n-1} + \frac{11\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_n \\
&(\eta_n)_{xx} + \frac{\mu^2}{3} (\eta_n)_{xxxx} \\
&= \frac{11\eta_{n-4} - 56\eta_{n-3} + 114\eta_{n-2} - 104\eta_{n-1} + 35\eta_n}{12\Delta x^2} + \frac{\mu^2}{3} \frac{\eta_{n-4} - 4\eta_{n-3} + 6\eta_{n-2} - 4\eta_{n-1} + \eta_n}{\Delta x^4} \\
&= \frac{11\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_{n-4} - \frac{56\Delta x^2 + 16\mu^2}{12\Delta x^4} \eta_{n-3} + \frac{114\Delta x^2 + 24\mu^2}{12\Delta x^4} \eta_{n-2} - \frac{104\Delta x^2 + 16\mu^2}{12\Delta x^4} \eta_{n-1} + \frac{35\Delta x^2 + 4\mu^2}{12\Delta x^4} \eta_n.
\end{aligned}$$

Then, the matrix becomes

$$\begin{aligned}
&\begin{bmatrix} (\eta_0)_{xx} + \frac{\mu^2}{3} (\eta_0)_{xxxx} \\ (\eta_1)_{xx} + \frac{\mu^2}{3} (\eta_1)_{xxxx} \\ (\eta_2)_{xx} + \frac{\mu^2}{3} (\eta_2)_{xxxx} \\ \vdots \\ (\eta_k)_{xx} + \frac{\mu^2}{3} (\eta_k)_{xxxx} \\ \vdots \\ (\eta_{n-2})_{xx} + \frac{\mu^2}{3} (\eta_{n-2})_{xxxx} \\ (\eta_{n-1})_{xx} + \frac{\mu^2}{3} (\eta_{n-1})_{xxxx} \\ (\eta_n)_{xx} + \frac{\mu^2}{3} (\eta_n)_{xxxx} \end{bmatrix} \\
&= \begin{bmatrix} \frac{35\Delta x^2 + 4\mu^2}{12\Delta x^4} & -\frac{104\Delta x^2 + 16\mu^2}{12\Delta x^4} & \frac{114\Delta x^2 + 24\mu^2}{12\Delta x^4} & -\frac{56\Delta x^2 + 16\mu^2}{12\Delta x^4} & \frac{11\Delta x^2 + 4\mu^2}{12\Delta x^4} & 0 & 0 & 0 & \dots \\ \frac{11\Delta x^2 + 4\mu^2}{12\Delta x^4} & -\frac{20\Delta x^2 + 16\mu^2}{12\Delta x^4} & \frac{6\Delta x^2 + 24\mu^2}{12\Delta x^4} & -\frac{4\Delta x^2 - 16\mu^2}{12\Delta x^4} & -\frac{\Delta x^2 + 4\mu^2}{12\Delta x^4} & 0 & 0 & 0 & \dots \\ \frac{-\Delta x^2 + 4\mu^2}{12\Delta x^4} & \frac{16(\Delta x^2 - \mu^2)}{12\Delta x^4} & -\frac{30\Delta x^2 + 24\mu^2}{12\Delta x^4} & -\frac{16(\Delta x^2 - \mu^2)}{12\Delta x^4} & -\frac{\Delta x^2 + 4\mu^2}{12\Delta x^4} & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \dots \\ 0 & \dots & 0 & -\frac{\Delta x^2 + 4\mu^2}{12\Delta x^4} & \frac{16(\Delta x^2 - \mu^2)}{12\Delta x^4} & -\frac{30\Delta x^2 + 24\mu^2}{12\Delta x^4} & -\frac{16(\Delta x^2 - \mu^2)}{12\Delta x^4} & -\frac{\Delta x^2 + 4\mu^2}{12\Delta x^4} & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \dots \\ \dots & 0 & 0 & 0 & -\frac{\Delta x^2 + 4\mu^2}{12\Delta x^4} & \frac{16(\Delta x^2 - \mu^2)}{12\Delta x^4} & -\frac{30\Delta x^2 + 24\mu^2}{12\Delta x^4} & -\frac{16(\Delta x^2 - \mu^2)}{12\Delta x^4} & -\frac{\Delta x^2 + 4\mu^2}{12\Delta x^4} \\ \dots & 0 & 0 & 0 & -\frac{\Delta x^2 + 4\mu^2}{12\Delta x^4} & \frac{4\Delta x^2 - 16\mu^2}{12\Delta x^4} & \frac{6\Delta x^2 + 24\mu^2}{12\Delta x^4} & -\frac{20\Delta x^2 + 16\mu^2}{12\Delta x^4} & \frac{11\Delta x^2 + 4\mu^2}{12\Delta x^4} \\ \dots & 0 & 0 & 0 & \frac{11\Delta x^2 + 4\mu^2}{12\Delta x^4} & -\frac{56\Delta x^2 + 16\mu^2}{12\Delta x^4} & \frac{114\Delta x^2 + 24\mu^2}{12\Delta x^4} & -\frac{104\Delta x^2 + 16\mu^2}{12\Delta x^4} & \frac{35\Delta x^2 + 4\mu^2}{12\Delta x^4} \end{bmatrix} \begin{bmatrix} \eta_0 \\ \eta_1 \\ \eta_2 \\ \vdots \\ \eta_k \\ \vdots \\ \eta_{n-2} \\ \eta_{n-1} \\ \eta_n \end{bmatrix}
\end{aligned}$$

For simplicity, let  $\mathcal{A}$  represent the above matrix. Now, recall the system:

$$\begin{aligned} u(x_k, t_{j+1}) &= u(x_k, t_j) + \Delta t f_1(\eta(x_k, t_j), u(x_k, t_j), t), \\ \eta(x_k, t_{j+1}) &= \eta(x_k, t_j) + \Delta t u(x_k, t_j), \end{aligned}$$

where

$$\begin{aligned} f_1(\eta(x_k, t_j), u(x_k, t_j), t) &= \eta(x_k, t_j)_{xx} + \frac{\mu^2}{3} \eta(x_k, t_j)_{xxx} \\ &\quad + \mu^2 \left( 2u(x_k, t_j)_x \int_{x_0}^{x_k} u \, dx' + 2u(x_k, t_j)^2 + \eta(x_k, t_j)_x^2 + \eta(x_k, t_j) \eta(x_k, t_j)_{xx} \right). \end{aligned}$$

For convenience, let

$$B_k = \left( 2u(x_k, t_j)_x \int_{x_0}^{x_k} u \, dx' + 2u(x_k, t_j)^2 + \eta(x_k, t_j)_x^2 + \eta(x_k, t_j) \eta(x_k, t_j)_{xx} \right),$$

Let  $\mathcal{B}$  represent the column vector of  $B_k$ 's. Then, we can write the system

$$\begin{aligned} u(x_0, t_{j+1}) &= u(x_0, t_j) + \Delta t f_1(\eta(x_0, t_j), u(x_0, t_j), t) \\ u(x_1, t_{j+1}) &= u(x_1, t_j) + \Delta t f_1(\eta(x_1, t_j), u(x_1, t_j), t) \\ u(x_2, t_{j+1}) &= u(x_2, t_j) + \Delta t f_1(\eta(x_2, t_j), u(x_2, t_j), t) \\ &\vdots \\ u(x_{n-2}, t_{j+1}) &= u(x_{n-2}, t_j) + \Delta t f_1(\eta(x_{n-2}, t_j), u(x_{n-2}, t_j), t) \\ u(x_{n-1}, t_{j+1}) &= u(x_{n-1}, t_j) + \Delta t f_1(\eta(x_{n-1}, t_j), u(x_{n-1}, t_j), t) \\ u(x_n, t_{j+1}) &= u(x_n, t_j) + \Delta t f_1(\eta(x_n, t_j), u(x_n, t_j), t) \end{aligned}$$

as follows:

$$\begin{bmatrix} u(x_0, t_{j+1}) \\ u(x_1, t_{j+1}) \\ u(x_2, t_{j+1}) \\ \vdots \\ u(x_{n-2}, t_{j+1}) \\ u(x_{n-1}, t_{j+1}) \\ u(x_n, t_{j+1}) \end{bmatrix} = \begin{bmatrix} u(x_0, t_j) \\ u(x_1, t_j) \\ u(x_2, t_j) \\ \vdots \\ u(x_{n-2}, t_j) \\ u(x_{n-1}, t_j) \\ u(x_n, t_j) \end{bmatrix} + \Delta t \begin{bmatrix} f_1(\eta(x_0, t_j), u(x_0, t_j), t) \\ f_1(\eta(x_1, t_j), u(x_1, t_j), t) \\ f_1(\eta(x_2, t_j), u(x_2, t_j), t) \\ \vdots \\ f_1(\eta(x_{n-2}, t_j), u(x_{n-2}, t_j), t) \\ f_1(\eta(x_{n-1}, t_j), u(x_{n-1}, t_j), t) \\ f_1(\eta(x_n, t_j), u(x_n, t_j), t) \end{bmatrix}$$



$$= \begin{bmatrix} u(x_0, t_j) \\ u(x_1, t_j) \\ u(x_2, t_j) \\ \vdots \\ u(x_{n-2}, t_j) \\ u(x_{n-1}, t_j) \\ u(x_n, t_j) \end{bmatrix} + \Delta t \mathcal{A} \begin{bmatrix} \eta_0 \\ \eta_1 \\ \eta_2 \\ \vdots \\ \eta_{n-2} \\ \eta_{n-1} \\ \eta_n \end{bmatrix} + \Delta t \mathcal{B}$$

Now, let us see how one would perform time-stepping. As such, impose initial conditions

$$\eta(x, t_0) = f(x), \quad u(x, t_0) = \eta_t(x, t_0) = g(x).$$

Let  $j = 0$ , and for simplicity, pick  $k \in [0, n]$ . The system is

$$\begin{aligned} u(x_k, t_1) &= u(x_k, t_0) + \Delta t f_1(\eta(x_k, t_0), u(x_k, t_0)), \\ \eta(x_k, t_1) &= \eta(x_k, t_0) + \Delta t u(x_k, t_0), \end{aligned}$$

where

$$\begin{aligned} f_1(\eta(x_k, t_0), u(x_k, t_0)) &= \eta(x_k, t_0)_{xx} + \frac{\mu^2}{3} \eta(x_k, t_0)_{xxx} \\ &+ \mu^2 \left( 2u(x_k, t_0)_x \int_{x_0}^{x_k} u \, dx' + 2u(x_k, t_0)^2 + \eta(x_k, t_0)_x^2 + \eta(x_k, t_0)\eta(x_k, t_0)_{xx} \right) \\ &= \frac{-\eta(x_{k-2}, t_0) + 16\eta(x_{k-1}, t_0) - 30\eta(x_k, t_0) + 16\eta(x_{k+1}, t_0) - \eta(x_{k+2}, t_0)}{12(\Delta x)^2} \\ &+ \frac{\mu^2}{3} \frac{\eta(x_{k-2}, t_0) - 4\eta(x_{k-1}, t_0) + 6\eta(x_k, t_0) - 4\eta(x_{k+1}, t_0) + \eta(x_{k+2}, t_0)}{(\Delta x)^4} \\ &+ \mu^2 \left( 2u(x_k, t_0)_x \int_{x_0}^{x_k} u(x', t_0) \, dx' + 2u(x_k, t_0)^2 + \eta(x_k, t_0)_x^2 + \eta(x_k, t_0)\eta(x_k, t_0)_{xx} \right) \end{aligned}$$

Note that all the terms on the last line can be computed via finite differences and both initial conditions. With this, we obtain the values of  $u, \eta$  at point  $x_k$  and time  $t_1$ . Performing this calculation for all  $k$ , we move on to compute  $u, \eta$  at time  $t_2$ , and so on.

## 2 Time stepping and finite differences: the half line

On the half line, the equation for the surface elevation is given by:

$$\eta_{tt} = \eta_{xx} + \mu^2 \left( \frac{1}{3} \eta_{xxx} - \frac{1}{\pi} \int_0^\infty \partial_t \left( \eta \int_0^y \eta_t dy' \right) \left[ \frac{1}{(x-y)^2} + \frac{1}{(x+y)^2} \right] dy + \frac{1}{2} \partial_x^2 \left( \int_0^x \eta_t dx' \right)^2 \right) \quad (4)$$

To do time stepping, introduce

$$u = \eta_t. \quad (5)$$

Also, note that

$$\frac{1}{2} \partial_x^2 \left( \int_0^x \eta_t dx' \right)^2 = \eta_t^2 + \eta_{tx} \int_0^x \eta_t dx' = u^2 + u_x \int_0^x u dx',$$

and

$$\partial_t \left( \eta \int_0^y \eta_t dy' \right) = \eta_t \int_0^x \eta_t dx' + \eta(\eta_x - \eta_x(0)) = u \int_0^x u dx' + \eta(\eta_x - \eta_x(0)).$$

Then, combining (5) and (4), we obtain a two-dimensional system:

$$\partial_t \begin{bmatrix} u \\ \eta \end{bmatrix} = \begin{bmatrix} \eta_{xx} + \mu^2 \left( \frac{1}{3} \eta_{xxx} + u^2 + u_x \int_0^x u dx' - \frac{1}{\pi} \int_0^\infty u \int_0^x u dx' + \eta(\eta_x - \eta_x(0)) \left[ \frac{1}{(x-y)^2} + \frac{1}{(x+y)^2} \right] dy \right) \\ u \end{bmatrix}. \quad (6)$$

Now, consider (4) on a finite interval  $[a, b]$ , and let partition the interval into  $n + 1$  points  $\{x_k\}_{k=0}^n$ , with  $x_0 = a$  and  $x_n = b$ . Note that we need to pick the partition such that

$$\int_0^\infty u \int_0^x u dx' + \eta(\eta_x - \eta_x(0)) \left[ \frac{1}{(x-y)^2} + \frac{1}{(x+y)^2} \right] dy \approx \int_{x_0}^{x_n} u \int_0^x u dx' + \eta(\eta_x - \eta_x(0)) \left[ \frac{1}{(x-y)^2} + \frac{1}{(x+y)^2} \right] dy.$$

We proceed to forward Euler time stepping. First, observe that

$$\begin{aligned} u_t(x_k, t_j) &= \frac{u(x_k, t_{j+1}) - u(x_k, t_j)}{\Delta t} = f_1(\eta, u, t) & \implies u(x_k, t_{j+1}) &= u(x_k, t_j) + \Delta t f_1(\eta, u, t) \\ \eta_t(x_k, t_j) &= \frac{\eta(x_k, t_{j+1}) - \eta(x_k, t_j)}{\Delta t} = f_2(\eta, u, t) & \implies \eta(x_k, t_{j+1}) &= \eta(x_k, t_j) + \Delta t f_2(\eta, u, t), \end{aligned}$$

where

$$f_1(\eta, u, t) = \eta_{xx} + \mu^2 \left( \frac{1}{3} \eta_{xxx} + u^2 + u_x \int_0^x u dx' - \frac{1}{\pi} \int_0^\infty u \int_0^x u dx' + \eta(\eta_x - \eta_x(0)) \left[ \frac{1}{(x-y)^2} + \frac{1}{(x+y)^2} \right] dy \right)$$

$$f_2(\eta, u, t) = u(x_k, t_j).$$

Observe that the highest order derivative is 4, so we need to use five-point stencils. In particular, we use five point midpoint stencil for  $x_2, \dots, x_{n-2}$ , and five point, one-sided stencils at  $x_0, x_1, x_{n-1}, x_n$ . First, consider the system for  $x_2, \dots, x_{n-2}$  :

$$\begin{aligned} f_1(\eta(x_k, t_j), u(x_k, t_j), t) &= \eta(x_k, t_j)_{xx} + \frac{\mu^2}{3} \eta(x_k, t_j)_{xxxx} \\ &+ \mu^2 \left( u^2(x_k) + u_x(x_k) \int_0^{x_k} u \, dx' - \frac{1}{\pi} \int_0^{x_n} u \int_0^y u \, dx' + \eta(\eta_x - \eta_x(0)) \left[ \frac{1}{(x_k - y)^2} + \frac{1}{(x_k + y)^2} \right] dy \right), \end{aligned}$$

where we have separated the linear and nonlinear terms. The only difference between this system and the whole-line system is the non-linear term; in other words, we can reuse our prior work on the linear term, and only deal with the non-linear term. Let

$$C_k = u^2(x_k) + u_x(x_k) \int_0^{x_k} u \, dx' - \frac{1}{\pi} \int_0^{x_n} \left( u \int_0^y u \, dx' + \eta(\eta_x - \eta_x(0)) \right) \left[ \frac{1}{(x_k - y)^2} + \frac{1}{(x_k + y)^2} \right] dy,$$

and let  $\mathcal{C}$  be the column vector of  $C_k$ ; thus,  $\mathcal{C}$  represents the non-linear part of the system. To discretise  $C_k$ , note that

$$\begin{aligned} \int_0^{x_k} u \, dx' &= \frac{\Delta x}{2} \sum_{i=0}^{k-1} u(x_{i+1}) + u(x_i) \\ \int_0^{x_n} u \int_0^y u \, dx' \left[ \frac{1}{(x_k - y)^2} + \frac{1}{(x_k + y)^2} \right] dy &= \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} \left( u \int_0^y u \, dx' \right) \left[ \frac{1}{(x_k - y)^2} + \frac{1}{(x_k + y)^2} \right] dy \\ &= \frac{\Delta x}{2} \sum_{i=0}^{n-1} u(x_{i+1}) \int_0^{x_{i+1}} u \, dx' \left[ \frac{1}{(x_k - x_{i+1})^2} + \frac{1}{(x_k + x_{i+1})^2} \right] + u(x_i) \int_0^{x_i} u \, dx' \left[ \frac{1}{(x_k - x_i)^2} + \frac{1}{(x_k + x_i)^2} \right] \\ &= \frac{\Delta x}{2} \sum_{i=0}^{n-1} \frac{\Delta x}{2} u(x_{i+1}) \left( \sum_{j=0}^i u(x_j) + u(x_{j+1}) \right) \left[ \frac{1}{(x_k - x_{i+1})^2} + \frac{1}{(x_k + x_{i+1})^2} \right] \\ &\quad + \frac{\Delta x}{2} u(x_i) \left( \sum_{j=0}^{i-1} u(x_j) + u(x_{j+1}) \right) \left[ \frac{1}{(x_k - x_i)^2} + \frac{1}{(x_k + x_i)^2} \right] \\ \int_0^{x_n} \eta(\eta_x - \eta_x(0)) \left[ \frac{1}{(x_k - y)^2} + \frac{1}{(x_k + y)^2} \right] dy &= \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} \eta(\eta_x - \eta_x(0)) \left[ \frac{1}{(x_k - y)^2} + \frac{1}{(x_k + y)^2} \right] dy \\ &= \frac{\Delta x}{2} \sum_{i=0}^{n-1} \eta(x_{i+1})(\eta_x(x_{i+1}) - \eta_x(0)) \left[ \frac{1}{(x_k - x_{i+1})^2} + \frac{1}{(x_k + x_{i+1})^2} \right] + \eta(x_i)(\eta_x(x_i) - \eta_x(0)) \left[ \frac{1}{(x_k - x_i)^2} + \frac{1}{(x_k + x_i)^2} \right]. \end{aligned}$$

Therefore, we obtain that

$$\begin{aligned}
C_k &= u^2(x_k) + u_x(x_k) \int_0^{x_k} u \, dx' - \frac{1}{\pi} \int_0^{x_n} \left( u \int_0^y u \, dx' + \eta(\eta_x - \eta_x(0)) \right) \left[ \frac{1}{(x_k - y)^2} + \frac{1}{(x_k + y)^2} \right] dy \\
&= u^2(x_k) + \frac{\Delta x}{2} u_x(x_k) \sum_{i=0}^{k-1} u(x_{i+1}) + u(x_i) \\
&\quad - \frac{1}{\pi} \frac{\Delta x}{2} \sum_{i=0}^{n-1} \left( \frac{\Delta x}{2} u(x_{i+1}) \left( \sum_{j=0}^i u(x_j) + u(x_{j+1}) \right) + \eta(x_{i+1})(\eta_x(x_{i+1}) - \eta_x(0)) \right) \left[ \frac{1}{(x_k - x_{i+1})^2} + \frac{1}{(x_k + x_{i+1})^2} \right] \\
&\quad + \left( \frac{\Delta x}{2} u(x_i) \left( \sum_{j=0}^{i-1} u(x_j) + u(x_{j+1}) \right) + \eta(x_i)(\eta_x(x_i) - \eta_x(0)) \right) \left[ \frac{1}{(x_k - x_i)^2} + \frac{1}{(x_k + x_i)^2} \right],
\end{aligned}$$

where the expressions for derivatives depend on the point  $x_k$ . Let

$$F_i = \frac{\Delta x}{2} u(x_i) \left( \sum_{j=0}^{i-1} u(x_j) + u(x_{j+1}) \right) + \eta(x_i)(\eta_x(x_i) - \eta_x(0)), \quad i = 0, \dots, n,$$

which we will store as an array. This simplification, we obtain

$$C_k = u^2(x_k) + \frac{\Delta x}{2} u_x(x_k) \sum_{i=0}^{k-1} u(x_{i+1}) + u(x_i) - \frac{\Delta x}{2\pi} \sum_{i=0}^{n-1} F_{i+1} \left[ \frac{1}{(x_k - x_{i+1})^2} + \frac{1}{(x_k + x_{i+1})^2} \right] + F_i \left[ \frac{1}{(x_k - x_i)^2} + \frac{1}{(x_k + x_i)^2} \right],$$

With this in mind, we obtain the finite differences system:

$$\begin{bmatrix} u(x_0, t_{j+1}) \\ u(x_1, t_{j+1}) \\ u(x_2, t_{j+1}) \\ \vdots \\ u(x_{n-2}, t_{j+1}) \\ u(x_{n-1}, t_{j+1}) \\ u(x_n, t_{j+1}) \end{bmatrix} = \begin{bmatrix} u(x_0, t_j) \\ u(x_1, t_j) \\ u(x_2, t_j) \\ \vdots \\ u(x_{n-2}, t_j) \\ u(x_{n-1}, t_j) \\ u(x_n, t_j) \end{bmatrix} + \Delta t \begin{bmatrix} f_1(\eta(x_0, t_j), u(x_0, t_j), t) \\ f_1(\eta(x_1, t_j), u(x_1, t_j), t) \\ f_1(\eta(x_2, t_j), u(x_2, t_j), t) \\ \vdots \\ f_1(\eta(x_{n-2}, t_j), u(x_{n-2}, t_j), t) \\ f_1(\eta(x_{n-1}, t_j), u(x_{n-1}, t_j), t) \\ f_1(\eta(x_n, t_j), u(x_n, t_j), t) \end{bmatrix}$$

$$= \begin{bmatrix} u(x_0, t_j) \\ u(x_1, t_j) \\ u(x_2, t_j) \\ \vdots \\ u(x_{n-2}, t_j) \\ u(x_{n-1}, t_j) \\ u(x_n, t_j) \end{bmatrix} + \Delta t \mathcal{A} \begin{bmatrix} \eta_0 \\ \eta_1 \\ \eta_2 \\ \vdots \\ \eta_{n-2} \\ \eta_{n-1} \\ \eta_n \end{bmatrix} + \Delta t \mathcal{C}$$