YaleNUSCollege

APPROXIMATE EQUATIONS IN THE SHALLOW WATER REGIME OF THE WATER WAVE PROBLEM ON UNBOUNDED DOMAINS

SULTAN AITZHAN

Capstone Final Report for BSc (Honours) in

Mathematical, Computational and Statistical Sciences

Supervised by: Prof. Katie Oliveras, Prof. Dave Smith

AY 2019/2020

Yale-NUS College Capstone Project

DECLARATION & CONSENT

- I declare that the product of this Project, the Thesis, is the end result of my own work and that due acknowledgement has been given in the bibliography and references to ALL sources be they printed, electronic, or personal, in accordance with the academic regulations of Yale-NUS College.
- 2. I acknowledge that the Thesis is subject to the policies relating to Yale-NUS College Intellectual Property (Yale-NUS HR 039).

ACCESS LEVEL

3. I agree, in consultation with my supervisor(s), that the Thesis be given the access level specified below: [check one only]

o <u>Unrestricted access</u> Make the Thesis immediately available for worldv	wide access.	
o <u>Access restricted to Yale-NUS College for a limit</u> Make the Thesis immediately available for Yale (mm/yyyy) to (mm/yyyy), up reason(s): (please specify; attach a separate sheet	-NUS College access only from to a maximum of 2 years for the following	
After this period, the Thesis will be made available for worldwide access.		
o Other restrictions: (please specify if any part of	your thesis should be restricted)	
Name & Residential College of Student		
Signature of Student	Date	
Name & Signature of Supervisor	Date	

Contents

1	Intr	oduction	1
2	-	iminaries	4
	2.1	Water wave problem	4
	2.2	Asymptotic analysis and perturbation methods	10
	2.3	Are asymptotic and perturbative methods reliable?	16
	2.4	Shallow water regime	17
3	Nor	n-local derivation on the whole line	24
	3.1	Water-wave problem on the whole line: non-local formu-	
		lation	26
	3.2	Behaviour of the ${\cal H}$ operator	29
	3.3	Perturbation expansion of the ${\cal H}$ operator	32
	3.4	Deriving an expression for surface elevation	37
	3.5	Derivation of wave and KdV equations	42
4	Dis	cussion	48
5	Wat	er waves on the half-line	51
Bi	bling	graphy	56

Chapter 1

Introduction

Often, mathematical modelling of the real-world phenomena results in ordinary and partial differential equations. Depending on equations, mathematicians may or may not have the tools to obtain solutions or understand the phenomenon very well. Fortunately, there are techniques that allow one to deal with differential equations without solving them. One such tool is asymptotic analysis, which leads to simplified equations that are very similar to original equations. As such, solutions of the simplified equations may model the real-world phenomenon, subject to some error estimates.

"deal with" mean?
Perhaps use "analyne or analyne solutions of "

In particular, one problem that is amenable to asymptotic methods is the *water wave problem*, which describes the behaviour of water and its surface under certain conditions. Assuming an irrotational, incompressible, and inviscid fluid in one dimension, and a domain with flat bottom, the equations of fluid motion are given by

$$\phi_{xx} + \phi_{zz} = 0$$
 $-h < z < \eta(x, t)$ (1.1a)

$$\phi_z = 0 z = -h (1.1b)$$

$$\eta_t + \phi_x \eta_x = \phi_z \qquad z = \eta(x, t) \tag{1.1c}$$

$$\phi_t + g\eta + \frac{1}{2}(\phi_x^2 + \phi_z^2) = 0$$
 $z = \eta(x, t)$ (1.1d)

> 0(x, =, t)

where $\phi(x,z)$ is the fluid velocity and $\eta(x)$ is the surface elevation. In addition, z is the vertical coordinate, x is the horizontal direction, and g is acceleration due to gravity. We let $x \in \mathbb{R}$, so that (1.1) is the water wave problem on the whole line. Although nonlinear partial differential equations (PDEs) (1.1c) and (1.1d) are hard to solve on their own, what makes the problem (1.1) truly difficult is the need to solve the Laplace's equation (1.1a) on a domain whose shape is unknown.

To make the equations of motion more tractable, one can reformulate the problem and apply the tools of asymptotics. Of particular interest is the work Ablowitz, Fokas, and Musslimani, 2006, henceforth referred to as the AFM formulation. In this paper, authors rewrite (1.1) as a system of two equations, for the surface variable $q(x) \stackrel{\triangleright}{=} \phi(x, \eta(x))$, i.e. the velocity evaluated at the surface. Taking advantage of a new system, various asymptotic reductions are performed. In well-understood physical conditions, the simplified equations corresponds to the expected equations. Original model.

One interesting physical condition is the shallow water regime, which is defined by small-amplitude waves that have a small depth relative to the wavelength. In this regime, asymptotic tools reveal that the fluid behaviour is governed by the following approximate equations: the wave

φ(x,η(x,t),t)

equation

how do you get n(x,t):

and two Korteweg de Vries (KdV) equations

 $F_T + \frac{1}{3}F_{\zeta\zeta\zeta} + 3FF_{\zeta} = 0,$ $G_T + \frac{1}{3}G_{\zeta\zeta\zeta} + 3GG_{\zeta} = 0,$

 $q_{tt}-q_{xx}=0,$

1 haps indicate right

F & G are left going

where q(x,t) = F(x-t,T) + G(x+t,T). We refer to the model given by (1.2) and (1.3) as the *KdV model*.

In this capstone project, we consider an alternative formulation of the problem (1.1), as presented in Oliveras and Vasan, 2013. Although slightly different from AFM formulation, it is contended that this formulation is well-suited for performing asymptotics. We further advocate the efficacy of this formulation, by deriving the equations (1.2) and (1.3). As a brief outline, in Chapter 2, we introduce the reader to asymptotic analysis, explain the physical assumptions of the problem, and briefly explain why the model (1.2) and (1.3) is a good approximation to the water wave problem. In Chapter 3, we derive the formulation, perform asymptotics, and derive the approximate equations. In Chapter 4, we justify the utility of the model, contrasting and comparing with other formulations. In Chapter 5, we describe an application of the formulation to a water wave problem on the half line. In attempts to rigorously derive a half-line model.

for this model where waves are bounded by a wall remain under unknown, though ad-hoc formulations are known.

set up the question before you go to the next paragraph.

indicate higher (KdV is a model.