Surfaces of infinite-type are non-Hopfian

Geometry & Topology Seminars, IISER Bhopal

Sumanta (4th-year Ph.D., IISc Bangalore)

Abstract: We show that an orientable surface Σ is of finite-type if and only if every proper map $f \colon \Sigma \to \Sigma$ of degree one is homotopic to a homeomorphism.

Hopfian groups

Definition A group G is said to be Hopfian if every epimorphism $G \to G$ is an isomorphism.

Theorem [Hem72; Thu82; Hem87; AFW15] The fundamental group of any compact orient manifold of dim ≤ 3 is Hopfian.	table

Theorem 3 [BS62; Hem83] The Baumslag-Solitar group BS $(m,n) := \langle x,y | x^{-1}y^mx = y^n \rangle$,

prime divisors.

 $mn \neq 0$ is Hopfian if and only if m or n divides the other or if m and n have precisely the same

The degree of a proper map

Let $f: M \to N$ be a proper map between connected, oriented, boundaryless smooth n-manifolds.

Definition The integer $\deg(f)$ that describes $H^n_{\mathbf{c}}(f) \colon H^n_{\mathbf{c}}(N; \mathbb{Z}) \cong \mathbb{Z} \longrightarrow \mathbb{Z} \cong H^n_{\mathbf{c}}(M; \mathbb{Z})$, up to sign.

Theorem [Eps66, Theorems 3.1 and 4.1] If $f|f^{-1}(D) \to D$ is a homeomorphism between two smoothly embedded closed n-balls, then $\deg(f) = \pm 1$.

Olum's Theorem [Eps66, Corollary 3.4] If $\deg(f) = \pm 1$, then $\pi_1(f) \colon \pi_1(M) \longrightarrow \pi_1(N)$ is surjective.

The Hopf conjecture

Conjecture's Statement [Hau87] If the degree of $f: M \to M$ is ± 1 , then f is a homotopy equivalence.

Theorem [Dav05; DIK00; Hau87; Swa74] Suppose M is closed and $f: M \to M$ is a π_1 -injective map of deg ± 1 . Then f is a homotopy equivalence if either of the following happens:

- $\pi_1(M)$ is finite,
- $\pi_i(M) = 0$ for $1 < i < \dim(M) 1$,
- dim(M) ≤ 4, or
 π₁(M) has a nilpotent subgroup of the finite index.

Finite-type surfaces are Hopfian

Well-Known Theorem [Das21, Theorem 2] Let $S_{g,0,p}$ be a finite-type surface. Then any degree ± 1 map $f: S_{g,0,p} \to S_{g,0,p}$ is properly homotopic to a homeomorphism.


Infinite-type surfaces are non-Hopfian

Main Result [DG22, Theorem 1.1.] Let Σ be any infinite-type surface. Then there exists a degree ± 1 map $f: \Sigma \to \Sigma$ such that $\pi_1(f): \pi_1(\Sigma) \to \pi_1(\Sigma)$ is not injective.

Representation and classification of non-compact surfaces

Richrads' Representation Theorem [Ric63, Theorem 3] Any non-compact surface Σ is homeomorphic to a surface obtained by adding at most countably many handles to $\mathbb{S}^2 \setminus \mathscr{E}(\Sigma)$; where $\mathscr{E}(\Sigma)$ is a non-empty, closed, totally disconnected subset of \mathbb{S}^2 .

The surface with two planar ends and two non-planar ends:




Let $\mathscr{E}_{np}(\Sigma)$ be the set of all those points of $\mathscr{E}(\Sigma)$, which are accumulated by handles.

Kerékjártó's classification Theorem [Ric63, Theorem 1] Two non-compact surfaces Σ_1 and Σ_2

are homeomorphic if and only if

- the number of handles of Σ_1 is the same as the number of handles of Σ_2 , and
- there is a homeomorphism $\varphi \colon \mathscr{E}(\Sigma_1) \to \mathscr{E}(\Sigma_2)$ with $\varphi(\mathscr{E}_{np}(\Sigma_1)) = \mathscr{E}_{np}(\Sigma_2)$.




Infinite genus surfaces are non-Hopfian

minite genus surfaces are non-riophan

[DG22, Theorem 3.3.] Let
$$\Sigma$$
 be any infinite genus surface. Then the composition
$$\Sigma \xrightarrow{\text{quotient map}} \frac{\Sigma}{G} \xrightarrow{\text{homeomorphism}} \Sigma$$


is a non π_1 -injective map of deg ± 1 .

A folding map of degree ± 1 [DG22, Lemma 3.4.]




Finite genus infinite-type surfaces with finitely many isolated ends are non-Hopfian

[DG22, Theorem 3.5.] Let Σ be a finite genus infinite-type surface such that $\mathscr{E}(\Sigma)$ has finitely many isolated points. Then there is a degree ± 1 map $f: \Sigma \to \Sigma$ which is not π_1 -injective.



Finite genus surfaces with infinitely many isolated ends are non-Hopfian

[DG22, Theorem 3.7.] Let Σ be a finite genus surface such that $\mathscr{E}(\Sigma)$ has infinitely many isolated points. Then there is a degree one map $f: \Sigma \to \Sigma$ which is not π_1 -injective.



References

Matthias Aschenbrenner, Stefan Friedl, and Henry Wilton. 3-manifold groups. EMS Se-

10.1090/S0002-9904-1962-10745-9. URL: https://doi.org/10.1090/

Sumanta Das. Strong Topological Rigidity of Non-Compact Orientable Surfaces. 2021.

DOI: 10.48550/ARXIV.2111.11194. URL: https://arxiv.org/abs/2111.

[AFW15]

[Das21]

11194.

ries of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, 2015, pp. xiv+215. ISBN: 978-3-03719-154-5. DOI: 10.4171/154. URL: https://doi.org/10.4171/154.

[BS62] Gilbert Baumslag and Donald Solitar. "Some two-generator one-relator non-Hopfian groups". In: *Bull. Amer. Math. Soc.* 68 (1962), pp. 199–201. ISSN: 0002-9904. DOI:

S0002-9904-1962-10745-9

[Dav05]	Robert J. Daverman. "Fibrator properties of manifolds". In: <i>Topology Appl.</i> 153.2-3 (2005), pp. 245—260. ISSN: 0166-8641. DOI: 10.1016/j.topol.2003.04.001. URL: https://doi.org/10.1016/j.topol.2003.04.001.
[DG22]	Sumanta Das and Siddhartha Gadgil. Surfaces of infinite-type are non-Hopfian. 2022. DOI: 10.48550/ARXIV.2210.03395. URL: https://arxiv.org/abs/2210.03395.
[DIK00]	Robert J. Daverman, Young Ho Im, and Yongkuk Kim. "Products of Hopfian manifolds and codimension-2 fibrators". In: <i>Topology Appl.</i> 103.3 (2000), pp. 323–338. ISSN: 0166-8641. DOI: 10.1016/S0166-8641(99)00008-5. URL: https://doi.org/10.1016/S0166-8641(99)00008-5.
[Eps66]	D. B. A. Epstein. "The degree of a map". In: <i>Proc. London Math. Soc.</i> (3) 16 (1966), pp. 369-383. ISSN: 0024-6115. DOI: 10.1112/plms/s3-16.1.369. URL: https://doi.org/10.1112/plms/s3-16.1.369.
[Hau87]	Jean-Claude Hausmann. "Geometric Hopfian and non-Hopfian situations". In: <i>Geometry and topology (Athens, Ga., 1985)</i> . Vol. 105. Lecture Notes in Pure and Appl. Math. Dekker, New York, 1987, pp. 157–166. URL: https://doi.org/10.1201/9781003072386.

[Hem72]	John Hempel. "Residual finiteness of surface groups". In: <i>Proc. Amer. Math. Soc.</i> 32 (1972), p. 323. ISSN: 0002-9939. DOI: 10.2307/2038357. URL: https://doi.org/10.2307/2038357.
[Hem83]	John Hempel. "Intersection calculus on surfaces with applications to 3-manifolds". In: Mem. Amer. Math. Soc. 43.282 (1983), pp. vi+48. ISSN: 0065-9266. DOI: 10.1090/memo/0282. URL: https://doi.org/10.1090/memo/0282.
[Hem87]	John Hempel. "Residual finiteness for 3-manifolds". In: Combinatorial group theory and topology (Alta, Utah, 1984). Vol. 111. Ann. of Math. Stud. Princeton Univ. Press, Princeton, NJ, 1987, pp. 379–396. URL: https://doi.org/10.1515/9781400882083-018.
[Ric63]	Ian Richards. "On the classification of noncompact surfaces". In: Trans. Amer. Math. Soc. 106 (1963), pp. 259–269. ISSN: 0002-9947. DOI: 10.2307/1993768. URL: https://doi.org/10.2307/1993768.
[Swa74]	G. Ananda Swarup. "On a theorem of C. B. Thomas". In: <i>J. London Math. Soc.</i> (2) 8 (1974), pp. 13–21. ISSN: 0024-6107. DOI: 10.1112/jlms/s2-8.1.13. URL: https://doi.org/10.1112/jlms/s2-8.1.13.

[Thu82] William P. Thurston. "Three-dimensional manifolds, Kleinian groups and hyperbolic geometry". In: Bull. Amer. Math. Soc. (N.S.) 6.3 (1982), pp. 357–381. ISSN: 0273-0979. DOI: 10.1090/S0273-0979-1982-15003-0. URL: https://doi.org/10.1090/ S0273-0979-1982-15003-0.

⊕ Thank You ⊕