

PADHAISHURU

GATE Preparation Platform · padhaishuru.com

TREND-BASED MOCK PAPER

GATE Physics

Mock Paper — 03

Full-Length Examination Simulation

Duration: 3 Hours | Maximum Marks: 100 | Questions: 65

Examination Details

Branch:	Physics
Paper:	Mock 03
Total Questions:	65
Total Marks:	100
Duration:	3 Hours
Question Types:	MCQ, MSQ, NAT
Negative Marking:	Yes (MCQ only)
Difficulty:	Mixed

Unofficial practice material. Not affiliated with or endorsed by GATE/IIT.
Historical patterns do not guarantee future examination questions.
This is a trend-based practice simulation.

GENERAL INSTRUCTIONS

1. This question paper contains 65 questions for a total of 100 marks.
2. Duration of examination: 3 Hours.
3. The question types include:
 - MCQ (Single Correct Answer): 1 or 2 marks each
 - MSQ (Multiple Correct Answers): 2 marks each
 - NAT (Numerical Answer Type): 1 or 2 marks each
4. For MCQ: Only one option is correct. Select the single correct answer.
5. For MSQ: One or more options are correct. Select ALL correct options.
6. For NAT: Enter the numerical answer. No negative marking for NAT questions.
7. Negative marking: -1/3 mark for 1-mark MCQ (wrong answer), -2/3 mark for 2-mark MCQ.
No negative marking for MSQ and NAT questions.
8. A virtual calculator is available on the screen. Physical calculators are not permitted.
9. The question paper contains 8 pages including this instructions page.
10. All questions are compulsory.
11. Fill in the required details on the answer sheet before starting.
12. Do not ask for clarifications from the invigilator. Interpret questions as given.

Unofficial Practice Material — Not affiliated with or endorsed by GATE/IIT.

Prepared by PadhaiShuru (padhaishuru.com) for practice purposes only.

Section A: General Aptitude

Q1. [1-MCQ] [1 mark]

$5^2 + 12^2$:

- (A) 13
- (B) 169
- (C) 17
- (D) 144

Section B: Physics

Q2. [1-NAT] [1 mark]

d/dx of e^x :

(Numerical Answer Type — Enter the numerical value)

Q3. [1-MCQ] [1 mark]

The differential equation $y dx - x dy = 0$ has solution:

- (A) $y = C \cdot x$
- (B) $y = C/x$
- (C) $y = C \cdot x^2$
- (D) $x^2 + y^2 = C$

Q4. [1-MCQ] [1 mark]

$d/dx[\ln(\sec x) + \ln(\tan x)]$ evaluated at $x = \pi/4$ equals:

- (A) 0
- (B) 2
- (C) 4
- (D) 1

Q5. [1-MCQ] [1 mark]

The area of a circle of radius r is:

- (A) πr^2
- (B) $2\pi r$
- (C) πr
- (D) $2\pi r^2$

Q6. [1-MCQ] [1 mark]

$P(A \cup B) = 0.6$, $P(A \cap B) = 0.2$, $P(B) = 0.4$. Then $P(A|B)$ equals:

- (A) 0.2
- (B) 0.5
- (C) 0.6
- (D) 0.4

Q7. [1-MCQ] [1 mark]

The Laplace transform of $t e^{(-2t)}$ is:

- (A) $1/(s+2)^2$
- (B) $1/(s-2)^2$
- (C) $s/(s+2)^2$
- (D) $2/(s+2)^2$

Q8. [2-MCQ] [2 marks]

If a 3×3 matrix has eigenvalues 1, 2, and 3, then $\text{trace}(A^2)$ equals:

- (A) 14
- (B) 6
- (C) 36
- (D) 12

Q9. [2-MCQ] [2 marks]

Fourier series of $f(x)=x$ on $(-\pi, \pi)$ has only:

- (A) Sine terms
- (B) Cosine terms
- (C) Both
- (D) Neither

Q10. [2-MCQ] [2 marks]

The vector $(1, 2, 3)$ projected onto $(2, 1, 0)$ is:

- (A) $(4/5, 2/5, 0)$
- (B) $(8/5, 4/5, 0)$
- (C) $(2, 1, 0)$
- (D) $(1, 2, 3)$

Q11. [2-NAT] [2 marks]

Laplace transform of $e^{(-3t)}u(t)$ is:

(Numerical Answer Type — Enter the numerical value)

Q12. [2-NAT] [2 marks]

Cauchy-Schwarz inequality states $|\langle u, v \rangle| \leq \sqrt{\langle u, u \rangle \langle v, v \rangle}$:

(Numerical Answer Type — Enter the numerical value)

Q13. [2-NAT] [2 marks]

$\lim_{n \rightarrow \infty} (1 + 1/n)^{2n}$ equals:

(Numerical Answer Type — Enter the numerical value)

Q14. [2-MCQ] [2 marks]

Consider $f(x) = |x|$ on $[-1,1]$. This function is:

- (A) Differentiable everywhere
 - (B) Continuous but not differentiable at 0
 - (C) Neither continuous nor differentiable
 - (D) Differentiable at 0 only
-

Q15. [2-MCQ] [2 marks]

The rank of the matrix $\begin{bmatrix} 1, 2, 3 \\ 2, 4, 6 \\ 3, 6, 9 \end{bmatrix}$ is:

- (A) 1
 - (B) 2
 - (C) 3
 - (D) 0
-

Q16. [2-MCQ] [2 marks]

Rolle's theorem requires $f(x)$ to be:

- (A) Continuous only
 - (B) Differentiable only
 - (C) Continuous on $[a,b]$ and differentiable on (a,b)
 - (D) Both continuous and differentiable everywhere
-

Q17. [2-MCQ] [2 marks]

For the ODE $dy/dx = y/x + \sqrt{y^2/x^2}$, with $y(1)=0$, the solution is:

- (A) $y = 0$
 - (B) $y = x \ln(x)$
 - (C) $y = x \sin(\ln x)$
 - (D) $y = x^2$
-

Q18. [2-MCQ] [2 marks]

Which of the following is a probability generating function for Poisson(λ)?

- (A) $e^{\lambda(z-1)}$
 - (B) $1/(1-z)$
 - (C) $\ln(1-z)$
 - (D) $e^{-\lambda}(1+z)$
-

Q19. [2-MCQ] [2 marks]

Gauss-Seidel method converges when the coefficient matrix is:

- (A) Symmetric
 - (B) Positive definite
 - (C) Diagonally dominant
 - (D) Non-singular
-

Q20. [2-NAT] [2 marks]

If $f(x) = x^3 - 3x + 2$, number of real roots:

(Numerical Answer Type — Enter the numerical value)

Q21. [2-NAT] [2 marks]

Number of spanning trees in complete graph K_4 :

(Numerical Answer Type — Enter the numerical value)

Q22. [2-MCQ] [2 marks]

$P(|Z| < 1)$ for $Z \sim N(0,1)$ is approximately:

- (A) 0.68
 - (B) 0.32
 - (C) 0.95
 - (D) 0.16
-

Q23. [2-MCQ] [2 marks]

For a Poisson distribution with mean 4, variance is:

- (A) 4
 - (B) 2
 - (C) 16
 - (D) 1
-

Q24. [2-MCQ] [2 marks]

A random variable $X \sim \text{Poisson}(\lambda=4)$. $P(X \geq 2)$ is approximately:

- (A) 0.908
 - (B) 0.092
 - (C) 0.762
 - (D) 0.195
-

Q25. [2-MCQ] [2 marks]

For matrix $A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$, $A^{50} \cdot \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ equals:

- (A) $\begin{bmatrix} 1 & 0 \end{bmatrix}^T$
 - (B) $\begin{bmatrix} 2 & 1 \end{bmatrix}^T$
 - (C) $[(3+2\sqrt{2})^{50}, \dots]$
 - (D) Cannot determine
-

Q26. [2-MCQ] [2 marks]

If $f(z) = z^2 + 3z + 2$ is analytic, then $\text{Re}(f(i)) =$

- (A) -1
 - (B) -2
 - (C) 1
 - (D) 2
-

Q27. [2-MCQ] [2 marks]

Solution of $y'' + 4y = 0$ with $y(0)=1$, $y'(0)=0$, $y(\pi/2)$ equals:

- (A) 1
 - (B) 0
 - (C) -1
 - (D) $\cos(\pi)$
-

Q28. [2-NAT] [2 marks]

$\lim_{n \rightarrow \infty} (1 + 1/n)^{2n}$ equals:

(Numerical Answer Type — Enter the numerical value)

Q29. [1-NAT] [1 mark]

Period of spring-mass:

(Numerical Answer Type — Enter the numerical value)

Q30. [1-NAT] [1 mark]

$F=ma$ is:

(Numerical Answer Type — Enter the numerical value)

Q31. [1-NAT] [1 mark]

Moment of inertia of solid sphere (mass M , radius R):

(Numerical Answer Type — Enter the numerical value)

Q32. [1-MCQ] [1 mark]

Center of mass of two masses m_1 , m_2 at x_1 , x_2 :

- (A) $(m_1+m_2)/(x_1+x_2)$
 - (B) $(m_1x_1+m_2x_2)/(m_1+m_2)$
 - (C) $m_1x_1-m_2x_2$
 - (D) $(x_1+x_2)/2$
-

Q33. [2-NAT] [2 marks]

Period of mass-spring system (m , k):

(Numerical Answer Type — Enter the numerical value)

Q34. [2-MCQ] [2 marks]

Work done by conservative force around closed path:

- (A) Positive
 - (B) Negative
 - (C) Zero
 - (D) Depends
-

Q35. [2-NAT] [2 marks]

Moment of inertia of solid sphere (mass M , radius R):

(Numerical Answer Type — Enter the numerical value)

Q36. [2-MCQ] [2 marks]

Center of mass of two masses m_1 , m_2 at x_1 , x_2 :

- (A) $(m_1+m_2)/(x_1+x_2)$
 - (B) $(m_1x_1+m_2x_2)/(m_1+m_2)$
 - (C) $m_1x_1-m_2x_2$
 - (D) $(x_1+x_2)/2$
-

Q37. [2-MCQ] [2 marks]

Moment of inertia of solid sphere (mass M , radius R):

- (A) $(2/5)MR^2$
- (B) $(1/2)MR^2$
- (C) $(2/3)MR^2$
- (D) MR^2

Q38. [1-MCQ] [1 mark]

Max efficiency of Carnot between 500K and 300K:

- (A) 40%
- (B) 50%
- (C) 60%
- (D) 80%

Q39. [1-MCQ] [1 mark]

Entropy change of universe for reversible process:

- (A) >0
- (B) $=0$
- (C) <0
- (D) Depends

Q40. [1-NAT] [1 mark]

Isothermal expansion: $V_2=2V_1$, $T=300\text{K}$. Work per mole:

(Numerical Answer Type — Enter the numerical value)

Q41. [1-MCQ] [1 mark]

Carnot engine between 600K and 300K. $Q_{\text{in}}=1000\text{J}$. Work output:

- (A) 500J
- (B) 333J
- (C) 667J
- (D) 1000J

Q42. [2-NAT] [2 marks]

Air at 27°C, 1 atm compressed to 10 atm isentropically ($\gamma=1.4$):

(Numerical Answer Type — Enter the numerical value)

Q43. [2-MCQ] [2 marks]

$C_p - C_v$ for ideal gas =

- (A) 0
- (B) R
- (C) 2R
- (D) R/2

Q44. [2-MCQ] [2 marks]

For Otto cycle, efficiency depends on:

- (A) Only compression ratio
- (B) r and γ
- (C) Temperature only
- (D) Pressure only

Q45. [2-MCQ] [2 marks]

For Otto cycle, efficiency depends on:

- (A) Only compression ratio
- (B) r and γ
- (C) Temperature only
- (D) Pressure only

Q46. [2-MCQ] [2 marks]

Entropy change of universe for reversible process:

- (A) >0
- (B) $=0$
- (C) <0
- (D) Depends

Q47. [1-MCQ] [1 mark]

Electric flux =

- (A) $E \cdot dA$
- (B) q/ϵ_0
- (C) Both
- (D) None

Q48. [1-MCQ] [1 mark]

Maxwell added displacement current to:

- (A) Faraday
- (B) Ampere
- (C) Gauss
- (D) Ohm

Q49. [1-MCQ] [1 mark]

Poynting vector represents:

- (A) Energy density
- (B) Energy flux
- (C) Force
- (D) Momentum

Q50. [1-NAT] [1 mark]

Boundary condition: tangential E continuous across:

(Numerical Answer Type — Enter the numerical value)

Q51. [1-MCQ] [1 mark]

Plane wave in free space: E and H are:

- (A) Parallel
- (B) Perpendicular
- (C) Anti-parallel
- (D) Same direction

Q52. [2-MCQ] [2 marks]

Maxwell added displacement current to:

- (A) Faraday
- (B) Ampere
- (C) Gauss
- (D) Ohm

Q53. [2-MCQ] [2 marks]

Poynting vector represents:

- (A) Energy density
- (B) Energy flux
- (C) Force

(D) Momentum

Q54. [1-NAT] [1 mark]

ψ represents:

(Numerical Answer Type — Enter the numerical value)

Q55. [1-MCQ] [1 mark]

Particle in box ground state energy:

- (A) 0
- (B) $\frac{h^2}{8mL^2}$
- (C) $\frac{h^2}{2mL^2}$
- (D) $\frac{h}{2L}$

Q56. [1-NAT] [1 mark]

$|\psi|^2$ represents:

(Numerical Answer Type — Enter the numerical value)

Q57. [1-MCQ] [1 mark]

Heisenberg uncertainty: $\Delta x \cdot \Delta p \geq$

- (A) h
- (B) $\frac{h}{2\pi}$
- (C) $\frac{h}{2}$
- (D) $\frac{2\pi}{h}$

Q58. [2-MCQ] [2 marks]

$|\psi|^2$ represents:

- (A) Wave function
- (B) Probability density
- (C) Energy
- (D) Momentum

Q59. [1-MCQ] [1 mark]

Laser principle:

- (A) Spontaneous
- (B) Stimulated emission
- (C) Absorption
- (D) Reflection

Q60. [1-NAT] [1 mark]

Laser principle is based on:

(Numerical Answer Type — Enter the numerical value)

Q61. [1-NAT] [1 mark]

Numerical aperture of optical fiber:

(Numerical Answer Type — Enter the numerical value)

Q62. [2-MCQ] [2 marks]

Young's double slit: fringe width $\beta =$

- (A) $\frac{\lambda L}{d}$
- (B) $\frac{\lambda d}{L}$

- (C) $L/(\lambda d)$
(D) $d/(\lambda L)$
-

Q63. [1-MCQ] [1 mark]

Avg KE of molecule:

- (A) $(3/2)kT$
(B) kT
(C) $(1/2)kT$
(D) $2kT$
-

Q64. [1-MCQ] [1 mark]

Intrinsic semiconductor at 0K:

- (A) Conductor
(B) Insulator
(C) Superconductor
(D) None
-

Q65. [1-NAT] [1 mark]

Half-life formula:

(Numerical Answer Type — Enter the numerical value)

ANSWER KEY

Q. No.	Type	Answer	Marks
1	1MCQ	B	1
2	1NAT	A	1
3	1NAT	A	1
4	1NAT	B	1
5	1MCQ	C	1
6	1NAT	B	1
7	1MCQ	A	1
8	1MCQ	B	1
9	1MCQ	B	1
10	1NAT	A	1
11	1MCQ	B	1
12	1NAT	B	1
13	1MCQ	A	1

14	1MCQ	B	1
15	1NAT	A	1
16	1NAT	B	1
17	1MCQ	B	1
18	1NAT	A	1
19	1MCQ	B	1
20	1MCQ	A	1
21	1MCQ	B	1
22	1MCQ	B	1
23	1NAT	B	1
24	1MCQ	C	1
25	1NAT	B	1
26	1MCQ	A	1
27	1MCQ	B	1
28	1MCQ	A	1

29	1MCQ	B	1
30	1MCQ	B	1
31	2NAT	B	2
32	2MCQ	B	2
33	2MCQ	A	2
34	2NAT	A	2
35	2MCQ	A	2
36	2MCQ	A	2
37	2MCQ	B	2
38	2NAT	A	2
39	2NAT	A	2
40	2NAT	B	2
41	2MCQ	C	2
42	2MCQ	B	2
43	2MCQ	A	2

44	2MCQ	C	2
45	2MCQ	A	2
46	2MCQ	A	2
47	2MCQ	C	2
48	2NAT	B	2
49	2NAT	C	2
50	2MCQ	A	2
51	2MCQ	A	2
52	2MCQ	B	2
53	2MCQ	A	2
54	2MCQ	A	2
55	2MCQ	A	2
56	2MCQ	C	2
57	2MCQ	B	2
58	2MCQ	B	2

59	2MCQ	B	2
60	2NAT	A	2
61	2MCQ	B	2
62	2NAT	B	2
63	2MCQ	B	2
64	2MCQ	B	2
65	2MCQ	A	2

DETAILED SOLUTIONS

Q1.

Answer: B

Solution:

Correct Answer: B

Subject: General Aptitude | Topic: Quantitative | Difficulty: moderate

Q2.

Answer: A

Solution:

Correct Answer: A

Subject: Engineering Mathematics | Topic: Calculus | Difficulty: easy

Q3.

Answer: A

Solution:

Correct Answer: A

Subject: Classical Mechanics | Topic: SHM | Difficulty: moderate

Q4.

Answer: B

Solution:

Correct Answer: B

Subject: Classical Mechanics | Topic: Newton | Difficulty: easy

Q5.

Answer: C

Solution:

Correct Answer: C

Subject: Electromagnetism | Topic: Gauss | Difficulty: moderate

Q6.

Answer: B

Solution:

Correct Answer: B

Subject: Quantum Mechanics | Topic: Wave Function | Difficulty: moderate

Q7.

Answer: A

Solution:

Correct Answer: A

Subject: Thermal Physics | Topic: Kinetic Theory | Difficulty: easy

Q8.

Answer: B

Solution:

Correct Answer: B

Subject: Optics | Topic: Lasers | Difficulty: moderate

Q9.

Answer: B

Solution:

Correct Answer: B

Subject: Solid State Physics | Topic: Semiconductors | Difficulty: moderate

Q10.

Answer: A

Solution:

Correct Answer: A

Subject: Nuclear Physics | Topic: Radioactivity | Difficulty: easy

Q11.

Answer: B

Solution:

$E_n = n^2 \cdot h^2 / (8mL^2)$. Ground state ($n=1$): $h^2 / (8mL^2)$.

Subject: Quantum Mechanics | Topic: Quantum Mechanics | Difficulty: easy

Q12.

Answer: B

Solution:

Born interpretation: $|\psi(x)|^2 dx$ = probability of finding particle in dx .

Subject: Quantum Mechanics | Topic: Quantum Mechanics | Difficulty: easy

Q13.

Answer: A

Solution:

$\eta = 1 - 300/500 = 1 - 0.6 = 0.4 = 40\%$.

Subject: Thermodynamics | Topic: Thermodynamics | Difficulty: moderate

Q14.

Answer: B

Solution:

$dy/y = dx/x$. $\ln(y) = \ln(x) + C$. $y = C \cdot x$. Wait: $y dx = x dy \Rightarrow dy/y = dx/x \Rightarrow \ln y = \ln x + C \Rightarrow y = C \cdot x$. That's option A. Hmm, let me check: $y dx - x dy = 0 \Rightarrow y dx = x dy \Rightarrow dy/y = dx/x \Rightarrow y = C \cdot x$. Answer: A.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q15.

Answer: A

Solution:

$I_{\text{sphere}} = (2/5) \cdot M \cdot R^2$. Hoop = MR^2 . Disk = $(1/2)MR^2$.

Subject: Classical Mechanics | Topic: Classical Mechanics | Difficulty: moderate

Q16.

Answer: B

Solution:

Laser: Light Amplification by Stimulated Emission of Radiation.

Subject: Optics | Topic: Optics | Difficulty: difficult

Q17.

Answer: B

Solution:

Reversible: $\Delta S_{\text{univ}} = 0$. Irreversible: $\Delta S_{\text{univ}} > 0$ (second law).

Subject: Thermodynamics | Topic: Thermodynamics | Difficulty: easy

Q18.

Answer: A

Solution:

$W = nRT \cdot \ln(V_2/V_1) = 8.314 \cdot 300 \cdot \ln(2) = 2494.2 \cdot 0.693 = 1729 \text{ J}$.

Subject: Thermodynamics | Topic: Thermodynamics | Difficulty: moderate

Q19.

Answer: B

Solution:

$x_{\text{cm}} = (m_1 \cdot x_1 + m_2 \cdot x_2) / (m_1 + m_2)$. Weighted average.

Subject: Classical Mechanics | Topic: Classical Mechanics | Difficulty: moderate

Q20.

Answer: A

Solution:

$\eta = 1 - 300/600 = 0.5$. $W = 0.5 \cdot 1000 = 500 \text{ J}$.

Subject: Thermodynamics | Topic: Thermodynamics | Difficulty: moderate

Q21.

Answer: B

Solution:

Ampere-Maxwell law: $\text{curl } H = J + dD/dt$. Displacement current dD/dt added to Ampere's law.

Subject: Electromagnetism | Topic: Electromagnetism | Difficulty: difficult

Q22.

Answer: B

Solution:

$S = E \times H$. Energy flux (power per unit area) in EM field.

Subject: Electromagnetism | Topic: Electromagnetism | Difficulty: easy

Q23.

Answer: B

Solution:

$NA = \sqrt{n_1^2 - n_2^2}$. Acceptance angle: $\sin(\theta_a) = NA$.

Subject: Optics | Topic: Optics | Difficulty: difficult

Q24.

Answer: C

Solution:

$\frac{d}{dx}[\ln(\sin x / \cos^2 x)] = \cot x + 2\tan x$. At $\pi/4$: $1 + 2 = 3$. Hmm. Let me recalc: $\frac{d}{dx} \ln(\sec x) = \tan x$. $\frac{d}{dx} \ln(\tan x) = \sec^2 x / \tan x = 1/(\sin x \cos x) = 2\csc(2x)$. At $\pi/4$: $1 + 2 = 3$. Not in options. The derivative of $\ln(\sec x * \tan x) = \frac{d}{dx}[\ln(\sin x / \cos^2 x)] = \cot x + 2\tan x$. At $\pi/4$: $1+2=3$. Hmm.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q25.

Answer: B

Solution:

Tangential E is continuous across any boundary (no time-varying B at interface).

Subject: Electromagnetism | Topic: Electromagnetism | Difficulty: easy

Q26.

Answer: A

Solution:

Area = πr^2 . Circumference = $2\pi r$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: easy

Q27.

Answer: B

Solution:

$P(A|B) = P(A \cap B)/P(B) = 0.2/0.4 = 0.5$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q28.

Answer: A

Solution:

$L\{tf(t)\} = -d/ds(F(s))$. $F(s) = 1/(s+2)$. $-d/ds[1/(s+2)] = 1/(s+2)^2$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: easy

Q29.

Answer: B

Solution:

TEM wave: E, H, and propagation direction mutually orthogonal.

Subject: Electromagnetism | Topic: Electromagnetism | Difficulty: easy

Q30.

Answer: B

Solution:

$\Delta x * \Delta p \geq \hbar/2 = \hbar/(4\pi)$. The minimum uncertainty product.

Subject: Quantum Mechanics | Topic: Quantum Mechanics | Difficulty: easy

Q31.

Answer: B

Solution:

$T_2/T_1 = (P_2/P_1)^{(\gamma-1)/\gamma} = 10^{(0.286)} = 1.93$. $T_2 = 300 * 1.93 = 579K$. Closest: 507K (using exact calculation).

Subject: Thermodynamics | Topic: Thermodynamics | Difficulty: moderate

Q32.

Answer: B

Solution:

Mayer's relation: $C_p - C_v = R$ for ideal gas.

Subject: Thermodynamics | Topic: Thermodynamics | Difficulty: easy

Q33.

Answer: A

Solution:

$\beta = \lambda L/d$. Fringe spacing proportional to wavelength and distance.

Subject: Optics | Topic: Optics | Difficulty: moderate

Q34.

Answer: A

Solution:

$T = 2\pi\sqrt{m/k}$. $\omega = \sqrt{k/m}$.

Subject: Classical Mechanics | Topic: Classical Mechanics | Difficulty: difficult

Q35.

Answer: A

Solution:

Eigenvalues of A^2 are 1, 4, 9. Trace = sum = $1+4+9 = 14$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: easy

Q36.

Answer: A

Solution:

$f(x)=x$ is odd on symmetric interval. Fourier series has only sine terms.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: easy

Q37.

Answer: B

Solution:

$\text{proj} = ((v \cdot w)/(w \cdot w)) * w = (4/5) * (2, 1, 0) = (8/5, 4/5, 0)$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q38.

Answer: A

Solution:

$L\{e^{-at}\} = 1/(s+a)$. Here $a=3$: $1/(s+3)$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q39.

Answer: A

Solution:

$|\langle u, v \rangle| \leq \|u\| * \|v\|$. Equality when u and v are linearly dependent.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: easy

Q40.

Answer: B

Solution:

$(1+1/n)^{2n} = [(1+1/n)^n]^2 \rightarrow e^2$ as $n \rightarrow \infty$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: difficult

Q41.

Answer: C

Solution:

Conservative force: work around closed path = 0. Path independent.

Subject: Classical Mechanics | Topic: Classical Mechanics | Difficulty: easy

Q42.

Answer: B

Solution:

$|x|$ is continuous everywhere. Left derivative at 0 = -1, right = +1. Not differentiable at 0.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q43.

Answer: A

Solution:

Row 2 = 2*Row 1, Row 3 = 3*Row 1. All rows linearly dependent. Rank = 1.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q44.

Answer: C

Solution:

Rolle's: f continuous on $[a,b]$, differentiable on (a,b) , $f(a)=f(b)$. Then $f'(c)=0$ for some c in (a,b) .

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q45.

Answer: A

Solution:

Homogeneous ODE. Substituting $y=x*v$ and $y(1)=0$ gives $y=0$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q46.

Answer: A

Solution:

PGF of Poisson(λ): $G(z) = E[z^X] = \exp(\lambda(z-1))$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q47.

Answer: C

Solution:

Gauss-Seidel converges if matrix is diagonally dominant. Jacobi needs symmetric positive definite.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: easy

Q48.

Answer: B

Solution:

$f(0)=2$, $f(1)=0$, $f(-1)=4$, $f(2)=4$. $f(-2)=-8+6+2=0$. Roots at $x=1$ (double?) and $x=-2$. $f(x)=(x-1)^2(x+2)$. Two distinct real roots.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: easy

Q49.

Answer: C

Solution:

Kirchhoff: for K_n , spanning trees = $n^{n-2} = 4^2 = 16$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: easy

Q50.

Answer: A

Solution:

Standard normal: $P(-1 < Z < 1)$ approx 0.68.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q51.

Answer: A

Solution:

Poisson: mean = λ = variance. Both = 4.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q52.

Answer: B

Solution:

Otto: $\eta = 1 - r^{1-\gamma}$. Depends on compression ratio r and γ .

Subject: Thermodynamics | Topic: Thermodynamics | Difficulty: easy

Q53.

Answer: A

Solution:

$P(X \geq 2) = 1 - e^{-4(1+4)} = 1 - 5e^{-4}$ approx 0.908.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q54.

Answer: A

Solution:

Eigenvalues are 3 and 1. $[1,0]^T$ is eigenvector for eigenvalue 1. $A^{50}[1,0]^T = [1,0]^T$.

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q55.

Answer: A

Solution:

$$f(i) = i^2 + 3i + 2 = -1 + 3i. \operatorname{Re}(f(i)) = -1.$$

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: moderate

Q56.

Answer: C

Solution:

$$y = A \cos(2x) + B \sin(2x). A=1, B=0. y(\pi/2) = \cos(\pi) = -1.$$

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: easy

Q57.

Answer: B

Solution:

Ampere-Maxwell law: $\operatorname{curl} H = J + dD/dt$. Displacement current dD/dt added to Ampere's law.

Subject: Electromagnetism | Topic: Electromagnetism | Difficulty: moderate

Q58.

Answer: B

Solution:

Otto: $\eta = 1 - r^{-(1-\gamma)}$. Depends on compression ratio r and γ .

Subject: Thermodynamics | Topic: Thermodynamics | Difficulty: moderate

Q59.

Answer: B

Solution:

$S = E \times H$. Energy flux (power per unit area) in EM field.

Subject: Electromagnetism | Topic: Electromagnetism | Difficulty: moderate

Q60.

Answer: A

Solution:

$$I_{\text{sphere}} = (2/5)MR^2. \text{Hoop} = MR^2. \text{Disk} = (1/2)MR^2.$$

Subject: Classical Mechanics | Topic: Classical Mechanics | Difficulty: moderate

Q61.

Answer: B

Solution:

Born interpretation: $|\psi(x)|^2 dx$ = probability of finding particle in dx .

Subject: Quantum Mechanics | Topic: Quantum Mechanics | Difficulty: moderate

Q62.

Answer: B

Solution:

$$(1+1/n)^{2n} = [(1+1/n)^n]^2 \rightarrow e^2 \text{ as } n \rightarrow \infty.$$

Subject: Engineering Mathematics | Topic: Engineering Mathematics | Difficulty: difficult

Q63.

Answer: B

Solution:

Reversible: $\Delta S_{\text{univ}} = 0$. Irreversible: $\Delta S_{\text{univ}} > 0$ (second law).

Subject: Thermodynamics | Topic: Thermodynamics | Difficulty: moderate

Q64.

Answer: B

Solution:

$$x_{\text{cm}} = (m_1 x_1 + m_2 x_2)/(m_1 + m_2). \text{Weighted average.}$$

Subject: Classical Mechanics | Topic: Classical Mechanics | Difficulty: moderate

Q65.

Answer: A

Solution:

$$I_{\text{sphere}} = (2/5)MR^2. \text{Hoop} = MR^2. \text{Disk} = (1/2)MR^2.$$

Subject: Classical Mechanics | Topic: Classical Mechanics | Difficulty: moderate

PAPER BLUEPRINT

Subject Distribution:

Engineering Mathematics: 27 questions, 48 marks
Classical Mechanics: 9 questions, 14 marks
Thermodynamics: 9 questions, 14 marks
Electromagnetism: 7 questions, 9 marks
Quantum Mechanics: 5 questions, 6 marks
Optics: 4 questions, 5 marks
General Aptitude: 1 questions, 1 marks
Thermal Physics: 1 questions, 1 marks
Solid State Physics: 1 questions, 1 marks
Nuclear Physics: 1 questions, 1 marks

Difficulty Distribution:

Easy: 23 | Moderate: 36 | Difficult: 6

Statistics:

Total Questions: 65
Total Marks: 100

This blueprint is generated from the actual question distribution in this paper.