

Parameters Constraints on Λ CDM and XCDM using H(z)+BAO data

Sumit Kumar Adhya

Guide: Prof. Bharat Ratra

December 5, 2025

Abstract

Here I reproduce the results of Cao and Ratra [1] for the spatially flat and non-flat Λ CDM and XCDM dark-energy models. Following [1], I first analyze the Old [2] and New [1] Hubble-parameter measurements, and then jointly use them with the Old [2] and New [1] BAO data to constrain the relevant cosmological parameters. The resulting parameter estimates are in close agreement with those reported in the original studies.

1 Data and Methodology

For the old H(z) and old BAO data, I use Table-I and II of [2] and for the new data, I use Table-I(accounting for correlations) and II of [1]. First, I implemented the χ^2 function and minimized it using `scipy.optimize.minimize`. This yielded the minimized value χ^2_{\min} and the best-fit estimates of the free parameters Ω_{k0} , w_x , $\Omega_b h^2$, $\Omega_c h^2$, and H_0 , consistent with those used in the reference paper. From these free parameters one can also compute the best-fit value of the derived matter-density parameter $\Omega_m = \frac{\Omega_b h^2 + \Omega_c h^2 + \Omega_\nu h^2}{h^2}$ where I use the neutrino-density value $\Omega_\nu h^2 = 0.06/93.14$ as given in the paper.

I then ran `emcee` to obtain Markov chains for these free parameters. For all parameters I adopted the same priors as used in the paper. The MCMC simulations were run with 200 walkers for 5000-10,000 steps varying across different models.

2 Comparison of Results

2.1 Plots

Below I report the plots I obtained, at first using only H(z) data and then combining with the BAO data and performing a joint analysis. I compare them with the original plots in [1].




Figure 1: Original Plots in Cao and Ratra[1]. a) Flat Λ CDM, b) Non-Flat Λ CDM, c) Flat XCDM, d) Non-Flat XCDM





Figure 2: Plots generated using emcee and scipy.



Original Plots in Cao and Ratra [1]. a) Flat Λ CDM, b) Non-Flat Λ CDM, c) Flat XCDM, d) Non-Flat XCDM




Figure 3: Plots generated³ using `emcee` and `scipy`.

Table 1: Comparison of Best-Fitting Cosmological Parameters: My Results vs. Cao and Ratra (2023)[1]

Model	Dataset	My Results						Cao and Ratra[1]					
		$\Omega_b h^2$	$\Omega_c h^2$	Ω_{m0}	Ω_{k0}	w_X	H_0	$\Omega_b h^2$	$\Omega_c h^2$	Ω_{m0}	Ω_{k0}	w_X	H_0
Flat Λ CDM	Old $H(z)$	—	—	0.3197	—	—	68.116	0.0273	0.1201	0.319	—	—	68.16
	$H(z)$	—	—	0.3097	—	—	69.428	0.0244	0.1181	0.301	—	—	69.43
	Old $H(z)$ +Old BAO	0.024	0.118	0.299	—	—	68.953	0.0254	0.1200	0.309	—	—	68.98
	$H(z)$ +BAO	0.025	0.120	0.295	—	—	70.134	0.0260	0.1200	0.297	—	—	70.12
Non-flat Λ CDM	Old $H(z)$	—	—	0.354	-0.11	—	68.93	0.0205	0.1515	0.362	-0.136	—	69.09
	$H(z)$	—	—	0.315	-0.014	—	69.5	0.0180	0.1328	0.314	-0.012	—	69.47
	Old $H(z)$ +Old BAO	0.026	0.110	0.291	0.048	—	68.329	0.0260	0.1098	0.292	0.048	—	68.35
	$H(z)$ +BAO	0.027	0.113	0.289	0.038	—	69.635	0.0269	0.1128	0.289	0.041	—	69.61
Flat XCDM	Old $H(z)$	—	—	0.3226	—	-1.185	70.241	0.0376	0.1236	0.321	—	-1.261	70.95
	$H(z)$	—	—	0.3158	—	-1.133	70.569	0.0106	0.1464	0.316	—	-1.140	70.63
	Old $H(z)$ +Old BAO	0.030	0.094	0.288	—	-0.755	65.615	0.0951	0.0296	0.290	—	-0.754	65.79
	$H(z)$ +BAO	0.033	0.092	0.281	—	-0.731	66.686	0.0938	0.0318	0.283	—	-0.734	66.67
Non-flat XCDM	Old $H(z)$	—	—	—	—	—	—	0.0223	0.0736	0.172	0.324	-2.272	75.05
	$H(z)$	—	—	—	—	—	—	0.0316	0.0530	0.378	0.151	-2.278	75.06
	Old $H(z)$ +Old BAO	0.029	0.098	0.294	-0.054	-0.730	65.682	0.0289	0.0985	0.296	-0.053	-0.730	65.76
	$H(z)$ +BAO	0.032	0.098	0.291	-0.081	-0.699	66.859	0.0305	0.0998	0.293	-0.084	-0.703	66.79

Table 2: Comparison of One-dimensional posterior mean parameter values and uncertainties: My Results vs. Cao and Ratra (2023)[1]

Model	Dataset	My Results						Cao and Ratra[1]					
		$\Omega_b h^2$	$\Omega_c h^2$	Ω_{m0}	Ω_{k0}	w_X	H_0	$\Omega_b h^2$	$\Omega_c h^2$	Ω_{m0}	Ω_{k0}	w_X	H_0
Flat Λ CDM	Old $H(z)$	—	—	0.334 ^{+0.006} _{-0.070}	—	—	67.7 ± 3.0	0.0225 ± 0.0108	0.1264 ± 0.0207	0.328 ^{+0.002} _{-0.053}	—	—	67.98 ± 3.24
	$H(z)$	—	—	0.324 ^{+0.007} _{-0.071}	—	—	69.1 ± 4.1	0.0225 ± 0.0107	0.1275 ± 0.0208	0.319 ± 0.050	—	—	69.31 ± 4.25
	Old $H(z)$ +Old BAO	0.0246 ^{+0.0024} _{-0.0029}	0.1184 ^{+0.0073} _{-0.0092}	0.301 ^{+0.014} _{-0.016}	—	—	69.1 ± 1.6	0.0247 ± 0.0030	0.1186 ^{+0.0078} _{-0.0109}	0.301 ^{+0.018} _{-0.031}	—	—	69.14 ± 1.85
	$H(z)$ +BAO	0.0263 ^{+0.0045} _{-0.0045}	0.1211 ^{+0.0083} _{-0.0092}	0.297 ^{+0.013} _{-0.016}	—	—	70.6 ± 2.5	0.0260 ± 0.0040	0.1212 ^{+0.0103} _{-0.0103}	0.297 ^{+0.016} _{-0.020}	—	—	70.49 ± 2.74
Non-flat Λ CDM	Old $H(z)$	—	—	0.44 ± 0.15	-0.32 ^{+0.49} _{-0.49}	—	70.3 ± 4.7	0.0223 ^{+0.0108} _{-0.0108}	0.1685 ^{+0.1130} _{-0.1130}	0.390 ^{+0.172} _{-0.172}	-0.174 ^{+0.491} _{-0.491}	—	69.09 ^{+1.47} _{-1.47}
	$H(z)$	—	—	0.45 ^{+0.19} _{-0.19}	-0.37 ± 0.50	—	70.9 ± 4.9	0.0222 ± 0.0108	0.1612 ^{+0.1061} _{-0.1061}	0.374 ^{+0.120} _{-0.120}	-0.136 ^{+0.364} _{-0.364}	—	69.56 ^{+1.89} _{-1.89}
	Old $H(z)$ +Old BAO	0.0274 ^{+0.0034} _{-0.0042}	0.107 ± 0.017	0.289 ± 0.023	0.064 ^{+0.079} _{-0.079}	—	68.3 ± 1.9	0.0266 ^{+0.0029} _{-0.0029}	0.1088 ± 0.0166	0.291 ± 0.023	0.059 ^{+0.081} _{-0.081}	—	68.37 ± 2.10
	$H(z)$ +BAO	0.0288 ^{+0.0042} _{-0.0042}	0.111 ± 0.018	0.287 ± 0.022	0.054 ^{+0.077} _{-0.080}	—	69.9 ± 2.7	0.0275 ^{+0.0046} _{-0.0046}	0.1131 ^{+0.0180} _{-0.0180}	0.289 ± 0.023	0.047 ^{+0.082} _{-0.083}	—	69.81 ± 2.80
Flat XCDM	Old $H(z)$	—	—	0.306 ^{+0.007} _{-0.043}	-1.64 ^{+0.83} _{-0.43}	—	75.4 ^{+5.4} _{-5.4}	0.0225 ± 0.0107	0.1505 ^{+0.0327} _{-0.0327}	0.285 ^{+0.061} _{-0.061}	—	—	-1.972 ^{+1.164} _{-1.164}
	$H(z)$	—	—	0.283 ± 0.073	-1.98 ^{+1.3} _{-1.3}	—	79.4 ^{+5.9} _{-5.9}	0.0225 ± 0.0108	0.1505 ^{+0.0303} _{-0.0303}	0.278 ^{+0.065} _{-0.065}	—	—	-2.127 ^{+1.335} _{-1.335}
	Old $H(z)$ +Old BAO	0.0337 ^{+0.0042} _{-0.0050}	0.087 ^{+0.028} _{-0.018}	0.281 ^{+0.029} _{-0.021}	-0.74 ^{+0.14} _{-0.13}	—	65.6 ^{+2.1} _{-2.1}	0.0308 ^{+0.0083} _{-0.0083}	0.0969 ^{+0.0178} _{-0.0178}	0.289 ± 0.020	—	—	-0.784 ^{+0.140} _{-0.140}
	$H(z)$ +BAO	0.0382 ^{+0.013} _{-0.013}	0.087 ^{+0.031} _{-0.016}	0.274 ^{+0.030} _{-0.019}	-0.72 ^{+0.13} _{-0.12}	—	67.0 ± 2.9	0.0302 ^{+0.0059} _{-0.0059}	0.0978 ^{+0.0181} _{-0.0181}	0.285 ± 0.019	—	—	-0.776 ^{+0.130} _{-0.130}
Non-flat XCDM	Old $H(z)$	—	—	0.306 ^{+0.007} _{-0.043}	-1.64 ^{+0.83} _{-0.43}	—	75.4 ^{+5.4} _{-5.4}	0.0218 ^{+0.0052} _{-0.0052}	0.0927 ± 0.0217	0.228 ^{+0.055} _{-0.055}	0.241 ^{+0.061} _{-0.061}	—	-2.148 ^{+1.182} _{-1.182}
	$H(z)$	—	—	0.283 ± 0.073	-1.98 ^{+1.3} _{-1.3}	—	79.4 ^{+5.9} _{-5.9}	0.0218 ± 0.0093	0.0927 ± 0.0217	0.228 ^{+0.054} _{-0.054}	0.241 ^{+0.060} _{-0.060}	—	-2.148 ^{+1.176} _{-1.176}
	Old $H(z)$ +Old BAO	0.0337 ^{+0.0038} _{-0.011}	0.088 ^{+0.029} _{-0.018}	0.283 ^{+0.032} _{-0.025}	-0.03 ± 0.11	-0.73 ^{+0.17} _{-0.11}	65.6 ^{+2.1} _{-2.5}	0.0294 ^{+0.0047} _{-0.0045}	0.0980 ± 0.0186	0.292 ± 0.025	-0.027 ± 0.109	-0.770 ^{+0.098} _{-0.098}	66.13 ^{+2.32} _{-2.26}
	$H(z)$ +BAO	0.0382 ^{+0.014} _{-0.014}	0.087 ^{+0.020} _{-0.020}	0.278 ^{+0.024} _{-0.025}	-0.06 ± 0.11	-0.69 ^{+0.10} _{-0.10}	67.0 ± 2.9	0.0303 ^{+0.0051} _{-0.0051}	0.1021 ± 0.0193	0.292 ± 0.024	-0.054 ± 0.103	-0.757 ^{+0.093} _{-0.093}	67.33 ± 2.96

References

- [1] Shulei Cao, Bharat Ratra, "H₀ = 69.8 ± 1.3 km s⁻¹Mpc⁻¹, Ω_{m0} = 0.288 ± 0.017, and other constraints from lower-redshift, non-CMB, expansion-rate data", arXiv:2302.14203 [astro-ph.CO].
- [2] Shulei Cao, Bharat Ratra, "Using lower-redshift, non-CMB, data to constrain the Hubble constant and other cosmological parameters", arXiv:2203.10825 [astro-ph.CO].