# Prizes and effort in contests with private information\*

Sumit Goel †

November 15, 2022

(Link to latest version)

#### Abstract

We consider contests where agents have private information about their ability and study the effect of different prizes and competition on the effort exerted by the agents. We characterize the symmetric Bayes-Nash equilibrium strategy function and find that the effects of prizes and competition on effort depend qualitatively on the distribution of abilities among the agents. In particular, we find that if the density of inefficient agents is increasing, increasing the value of prizes or making the contest more competitive encourages effort. In contrast, if this density is decreasing, increasing the value of intermediate prizes or making the contest more competitive discourages effort. We also discuss applications of these results to the design of effort-maximizing contests in environments that impose natural constraints on feasible contests including grading contests, contests where agents have concave utilities for prizes, and contests where the designer can only award homogeneous prizes of a fixed value.

<sup>\*</sup>I am grateful to Federico Echenique, Omer Tamuz, Thomas Palfrey, Jingfeng Lu, Wade Hann-Caruthers, and seminar audience at the 8th Annual Conference on Contests: Theory and Evidence (2022) for helpful comments and suggestions.

<sup>&</sup>lt;sup>†</sup>California Institute of Technology; sgoel@caltech.edu; 0000-0003-3266-9035

### 1 Introduction

Contests are situations in which agents compete with one another by investing effort or resources to win prizes. Such competitive situations are common in many social and economic contexts, including college admissions, labor markets, R&D races, sporting events, rent-seeking activities, politics, competition-based crowdsourcing, wars etc. While some of these situations arise naturally, there are many others where the contest designer can optimally design the rules of the contest so as to satisfy their objectives. The designer's objective, and the structural elements of the contest that it can and cannot manipulate may vary depending upon the situation.

In this paper, we focus on situations where the contest participants have private information about their abilities, defined by their marginal costs of effort, and the designer can manipulate the values of the different prizes  $v_1, \ldots, v_n$  to influence the effort exerted by the participants. For such environments, we study the effect that different prizes have on the effort exerted by the agents and discuss applications to the design of optimal contests in various settings. In particular, we apply our results to the design of optimal grading contests, contests where agents have concave utilities for prizes, and contests where the designer can award any number of agents with a costless homogeneous prize. We note here that while there is a significant literature that studies contest design in incomplete information environments, our focus on obtaining a more complete ordering of how different contests compare in terms of the effort they induce, and also the distributional assumptions we make allowing for agents with almost zero marginal costs of effort, allow us to differentiate our work from the literature and contribute to it. We will now discuss our results.

We characterize the equilibrium effort function and find that increasing the value of first prize  $v_1$  encourages effort while increasing the value of the last prize  $v_n$  discourages effort for all agent types, and so these effects persist in expectation as well. On the other hand, the effect of increasing any intermediate prize  $v_2, \ldots, v_{n-1}$  is mixed in that it encourages

effort from the less efficient agents while discouraging effort from the more efficient agents. Moreover, this transfer of effort from the more efficient to less efficient is balanced in the sense that the total area under the equilibrium effort function remains the same. This property of the equilibrium function relies on the existence of agents with almost zero marginal costs of effort, and to the best of our knowledge, ours is the first paper to study contests in this domain. Importantly, the property leads to some interesting implications for the overall effects of these prizes which we now discuss.

The overall effect on effort of increasing any intermediate prize  $v_i$  depends qualitatively on the distribution of abilities in the population. If the distribution is such that there is an increasing density of less efficient agents, and if the contest designer's objective is increasing and concave in an arbitrary agent's effort level, the designer is better off by increasing the values of these intermediate prizes. In contrast, if the distribution is such that there is an increasing density of more efficient agents, and the contest designer's objective is increasing and convex in an arbitrary agent's effort level, the designer is better off by reducing the values of these intermediate prizes. In particular, it follows that when there is a uniform distribution of abilities in the population, decreasing any of the intermediate prizes leads to effort function that is a mean-preserving spread of the original effort function. So if the designer has a concave objective, it would prefer to increase the value of the intermediate prizes and if its objective is convex, it would prefer to reduce these values.

We also study how the competitiveness of a contest influences effort and as with the effect of prizes, we find that these effects also depend qualitatively on the prior distribution of abilities. To study the effects of competition, we focus on a parametric subclass of distributions and examine how the expected effort changes as we increase competition by increasing the inequality between the top and bottom prizes. When there is increasing density of less efficient agents in the population, more competitive prize vectors induce higher expected effort. In contrast, when there is an increasing density of more efficient agents, making the

contest more competitive by increasing prize inequality reduces expected effort. In addition, we also study the effects of competition on expected minimum effort and show that for this parametric subclass of distributions, the expected minimum effort decreases as the contest becomes more competitive. Intuitively, this is because the designer puts a lot of weight on the effort of the less efficient agents who are better encouraged by the higher values of the lower ranked prizes.

We next discuss applications of our results to the design of effort maximizing contests in three different environments. First, we consider the design of grading contests. Assuming the value of a grade in a grading contest is determined by the information it reveals about the type of the agent, and more precisely, equals its expected productivity, we show that a more informative grading scheme induces a more competitive prize vector. The optimal grading contests can then be derived from our results on the effects of competition on effort. For our second application, we consider a setting where the contest designer has a budget that it can split across n prizes and the agents have a common concave utility function for prizes and linear effort costs. We derive the effort-maximizing contest and show that as the utility becomes more concave, the optimal prize vector becomes less competitive. For our third and last application, we derive the optimal contest in settings where the designer can choose the number of winners to award with a costless homogeneous prize.

#### 1.1 Literature review

There is a vast literature in contest theory studying the influence of manipulating various structural elements of a contest on the equilibrium effort of the agents and consequently designing these elements so as to satisfy certain objectives. The version of the optimal contest design problem that is most closely related to this work is to distribute a given budget among prizes  $(v_1, v_2, \ldots, v_n)$  so as to maximize the effort of the n competing agents. This problem was first posed by Galton in 1902 and has since been studied in various different

environments. The environments generally differ in the assumptions they make about the contest success function (csf) which is a mapping from agents effort levels to their probability of winning a prize, the abilities or valuations of the agents which may be symmetric or asymmetric, and the information agents have about the abilities of other agents which may be complete or incomplete. In this paper, our focus is on the perfectly discriminatory contest success function under which agents are perfectly ranked according to their effort levels and awarded the corresponding prizes and we briefly discuss what is known about this problem for this csf.

In the complete information setting with symmetric agents, the results generally suggest that increasing competition by increasing prize inequality discourages effort and so distributing the budget equally among the top n-1 agents maximizes total effort. This is for instance the case in Glazer and Hassin [30] who consider a model with concave utility and linear costs. In the special case when utility is also linear, Barut and Kovenock [2] show that any equilibrium induces the same expected aggregate effort and the only restriction posed by optimality is that the last prize  $v_n$  should be zero. More recently, Fang et al. [23] generalize these results and find that with linear utility and convex costs, increasing prize inequality reduces each agent's effort in the sense of second order stochastic dominance. In contrast, our work in the incomplete information environment shows that the effect of competition depends on the prior distribution of abilities in the population.

In an incomplete environment with ex-ante symmetric agents, which is the focus of this paper, a wide range of models have been studied in the literature. In many of these settings, a winner-take-all prize has been shown to be optimal for maximizing effort. Glazer and Hassin [30] show that if the agents have concave utility and linear costs, the optimal prize vector involves setting all the later half of the prizes to 0. Zhang [61] considers a model with convex effort costs and identifies a necessary and sufficient condition under which the winner-takes-all prize structure is optimal among a general class of mechanisms. Moldovanu

and Sela [44] find that in a setting with linear utility and linear or concave costs, awarding only a single prize is optimal. For convex costs, they identify conditions under which awarding more than one prize might be optimal. In comparison, our paper makes different distributional assumptions on the prior distribution and more importantly, obtains a more complete ordering of the prize vectors in terms of the effort they induce. The analysis, which may be useful in itself, allows us to study and solve the optimal contest design problem in other natural environments where the set of feasible prize vectors may be constrained or different. We illustrate this in our applications.

There has been relatively little work on the optimal prize distribution problem with asymmetric agents. In a complete information setting with linear utility and linear costs, Clark and Riis [14] provide examples where splitting the budget into more than one prize might be optimal. We are not aware of any work on the problem in an incomplete information environment with ex-ante asymmetric agents. There has also been work on this problem for the ratio-form contest success function, which is the most popular alternative to the perfectly discriminatory csf. Clark and Riis [15] consider a complete information setting with symmetric agents having linear utility and costs and find that under the Tullock csfs, which represents a parametric subclass of ratio-form csfs, increasing prize inequality leads to an increase in total effort. Szymanski and Valletti [59] show that with asymmetric agents, a second prize might be optimal. We are not aware of work on the ratio-form csf in the incomplete information environment. Other related work that looks at the design of optimal contests under some different assumptions include Krishna and Morgan [37], Liu and Lu [42], Cohen and Sela [16]. Sisak [57] provides a more detailed survey of the literature on this problem.

In addition to manipulating the values of prizes to influence the effort, there are other structural elements of a contest that have also been considered in the literature. These include winner selection mechanisms, introduction of dynamics with sequential decision making, introduction of asymmetry via head starts, information disclosure at intermediate stages,

entry constraints, introducing multiple contests, group contests, etc. Fu and Wu [26] provides a survey of the theoretical literature on optimal contest design from these different perspectives. More general surveys of the theoretical literature in contest theory can be found in Corchón [18], Vojnović [60], Konrad et al. [34], Segev [52].

The settings we focus on for our applications have also been studied in the literature. In particular, there has been significant work on the design of optimal grading schemes (Moldovanu et al. [46], Rayo [50], Popov and Bernhardt [49], Chan et al. [8], Dubey and Geanakoplos [20], Zubrickas [62]). Moldovanu et al. [46] consider a setting where the designer can associate grades with arbitrary monetary prizes subject to budget and individual rationality constraints and find that the optimal grading scheme awards the top grade to a unique agent and a single grade to all the remaining agents. Dubey and Geanakoplos [20] consider a complete information environment where agents care about relative ranks and find that absolute grading is generally better than relative grading and that it's better to clump scores into coarse categories. Other related papers look at the signalling value of grades under different models or assumptions (Costrell [19], Betts [5], Zubrickas [62], Boleslavsky and Cotton [6]). The setting of our last application where the designer can only choose the number of winners to receive a fixed homogeneous prize was also considered in Liu and Lu [42] under different distributional assumptions. They find that the expected effort is single peaked in the number of prizes. In comparison, we identify natural conditions on the prior distribution under which awarding a single prize or n-1 prizes is optimal.

The paper proceeds as follows. In section 2, we present the model of a contest in an incomplete-information environment. Section 3 characterizes the symmetric Bayes-Nash equilibrium of this game and discusses some important properties of the equilibrium function. In section 4, we discuss applications of our results to the design of optimal contest in three different settings. Section 5 concludes. All proofs are relegated to the appendix.

### 2 Model

There is a single contest designer and n agents. The designer chooses a vector of prizes  $\mathbf{v} = (v_1, v_2, \dots, v_n)$  to award to the agents such that  $v_i \geq v_{i+1}$  for all i. The agents compete for these prizes by exerting costly effort. Each agent i is privately informed about its marginal cost of effort  $\theta_i \in [0, 1]$  which is drawn independently from [0, 1] according to a distribution  $F: [0, 1] \to [0, 1]$ . The distribution F is common knowledge. We assume that the distribution is twice differentiable and the density f is positive. Given a vector of prizes  $\mathbf{v}$ , marginal cost of effort  $\theta_i$ , and belief F about the marginal costs of effort of other agents, each agent i simultaneously chooses an effort level  $e_i$ . The designer ranks the agents in order of the efforts they put in and awards them the corresponding prizes. So the agent who puts in the maximum effort is awarded prize  $v_1$ . Agent with the second highest effort is awarded prize  $v_2$  and so on. If agent i puts in effort  $e_i$  and wins prize  $v_i$ , its final payoff is

$$v_i - \theta_i e_i$$

Given a prize structure  $\mathbf{v}$  and belief F, an agent's strategy  $\sigma_i : [0,1] \to \mathbb{R}_+$  maps its marginal cost of effort to the level of effort it puts in. A strategy profile  $\sigma = (\sigma_1, \sigma_2, \dots, \sigma_n)$  is a Bayes-Nash equilibrium of the game if for all agents i and type  $\theta_i \in [0,1]$ , agent i's expected payoff from playing  $\sigma_i(\theta_i)$  is at least as high as its payoff from playing anything else given that all other agents are playing  $\sigma_{-i}$ . We'll focus on the symmetric Bayes-Nash equilibrium of this contest game. This is a Bayes-Nash equilibrium where all agents are playing the same strategy  $g_{\mathbf{v}} : [0,1] \to \mathbb{R}_+$ .

Given a prior distribution F, we'll assume the designer's preferences over the different contests or prize vectors  $\mathbf{v}$  is defined by a monotone utility function  $U: \mathbb{R}_+ \to \mathbb{R}$  so that the designer prefer  $\mathbf{v}$  over  $\mathbf{v}'$  if and only if  $\mathbb{E}[U(g_{\mathbf{v}}(\theta))] \geq \mathbb{E}[U(g_{\mathbf{v}'}(\theta))]$  where  $g_{\mathbf{v}}$  represents the symmetric Bayes-Nash equilibrium function under prize vector  $\mathbf{v}$ . We'll impose conditions on U as required to illustrate our results on the effect of prizes and competition on effort and the design of optimal contests.

## 3 Equilibrium

In this section, we first characterize the symmetric Bayes-Nash equilibrium of the game for arbitrary prize vectors and then discuss how the equilibrium changes as we vary different prizes.

Note that our model is similar to the model studied in Moldovanu and Sela [44] except for the fact that we allow agents marginal cost to be zero whereas Moldovanu and Sela [44] assume that the marginal costs are in an interval [m, 1] where m > 0. It turns out that the symmetric Bayes-Nash equilibrium strategy function takes the same form as in Moldovanu and Sela [44], but it also satisfies an interesting property due to the presence of agents with negligible marginal costs of effort as we will see later. The following result displays the symmetric Bayes-Nash equilibrium strategy of the contest game (Moldovanu and Sela [44]).

**Lemma 1.** In a contest with n agents, prizes  $\mathbf{v} = (v_1, v_2, \dots, v_{n-1}, v_n)$  and prior cdf F, the symmetric Bayes-Nash equilibrium strategy function is given by

$$g_{\mathbf{v}}(\theta) = \sum_{i=1}^{n} m_i(\theta) v_i$$

where,

$$m_i(\theta) = \binom{n-1}{i-1} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i-1}t^{i-2}]}{F^{-1}(t)} \left( (n-1)t - (i-1) \right) dt$$

for all  $i \in \{1, 2, ..., n\}$ .

The proof uses the standard approach of assuming that n-1 agents are playing the same strategy and then getting conditions under which that strategy is the best response for the remaining agent:

$$-f(\theta) \sum_{i=1}^{n-1} (v_i - v_{i+1}) \frac{(n-1)!}{(i-1)!(n-i-1)!} \left[ (1 - F(\theta))^{n-i-1} F(\theta)^{i-1} \right] = \theta g_v'(\theta)$$

Using the boundary condition  $g_v(1) = 0$  pins down the form of the function

$$g_v(\theta) = \sum_{i=1}^{n-1} (v_i - v_{i+1}) \frac{(n-1)!}{(i-1)!(n-i-1)!} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i-1}t^{i-1}]}{F^{-1}(t)} dt$$

We can then rewrite the expression as in the lemma by combining the two terms with coefficient  $v_i$ . Note that  $\sum_{i=1}^n m_i(\theta) = 0$  for all  $\theta \in [0,1]$ . Lastly, we check that the second order condition is satisfied. The full proof is in the appendix.

In case an agent's value for prize v is given by some utility function u(v) and all agents share the same utility function u for prizes, the equilibrium in Theorem 1 is simply as if prize i was  $u(v_i)$  instead of  $v_i$ . The following corollary states this formally.

Corollary 1. In a contest with n agents, each with utility function u for prizes, prizes  $\mathbf{v} = (v_1, v_2, \dots, v_{n-1}, v_n)$  and prior cdf F, the symmetric Bayes-Nash equilibrium strategy function is given by

$$g_v(\theta) = \sum_{i=1}^n m_i(\theta) u(v_i)$$

where,

$$m_i(\theta) = \binom{n-1}{i-1} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i-1}t^{i-2}]}{F^{-1}(t)} \left( (n-1)t - (i-1) \right) dt$$

for 
$$i \in [n-1]$$
 and  $m_n(\theta) = -\sum_{i=1}^{n-1} m_i(\theta)$ .

We will use this when we discuss the effort-maximizing contest for agents with concave utility for prizes.

Now that we know what the equilibrium function looks like, we focus our attention on studying how the equilibrium changes as we vary the values of different prizes.

**Theorem 1.** Consider a setting with n agents and prior F. Suppose  $\mathbf{v} = (v_1, \dots, v_n)$  and  $\mathbf{w} = (w_1, \dots, w_n)$  are two prize vectors. Then, we have the following:

- 1. if  $v_1 > w_1$  and  $v_{-1} = w_{-1}$ , then for any increasing function U, we have  $\mathbb{E}[U(g_{\mathbf{v}}(\theta))] \geq \mathbb{E}[U(g_{\mathbf{w}}(\theta))]$
- 2. if  $v_j > w_j$  for some  $j \in \{2, ..., n-1\}$ ,  $v_{-j} = w_{-j}$ , and the prior density function f is increasing, then for any increasing and concave function U, we have  $\mathbb{E}[U(g_{\mathbf{v}}(\theta))] \geq \mathbb{E}[U(g_{\mathbf{w}}(\theta))]$

3. if  $v_j > w_j$  for some  $j \in \{2, ..., n-1\}$ ,  $v_{-j} = w_{-j}$ , and the prior density function f is decreasing, then for any increasing and convex function U, we have  $\mathbb{E}[U(g_{\mathbf{v}}(\theta))] \geq \mathbb{E}[U(g_{\mathbf{v}}(\theta))]$ .

In words, the first prize always encourages effort irrespective of the prior distribution of abilities. The effect of the intermediate prizes depends on the prior distribution of abilities. If the prior distribution is such that the density of less efficient agents is increasing, then higher values of intermediate prizes encourage effort. But if this density is decreasing, then these prizes discourage effort in expectation. This is partly because any intermediate prize i has a different effect on different types of agents. The more efficient agents who are fighting for the top prizes put in lesser effort as the gain from winning these better prizes has gone down. In contrast, the less efficient agents who generally get lower prizes now put in more effort to get the increased prize i. Importantly, the decrease in effort of the more efficient agents and the increase for the less efficient agents cancel out so that there is basically a transfer of effort from the more efficient agents to the less efficient agents as any intermediate prize is increased. Note that the existence of agents with negligible marginal costs of effort is important for the equilibrium to have this property. This property is formally stated in the following lemma which is the key to proving Theorem 2.

**Lemma 2.** For any number of agents n and prior F, the following hold:

1. 
$$\theta m_i(\theta) \le 1 \text{ for all } i \in \{1, \dots, n-1\}, \theta \in [0, 1]$$

2. 
$$\lim_{\theta \to 0} \theta m_i(\theta) = 0$$
 for all  $i \in \{1, \dots, n-1\}$ 

3. 
$$\int_0^1 m_1(\theta) = 1$$
 and  $\int_0^1 m_i(\theta) = 0$  for  $i \in \{2, n-1\}$ 

4.  $m_1(\theta) > 0$  for all  $\theta$  and monotone decreasing

5. For  $i \in \{2, n-1\}$ , there exist  $t_i^1 < t_i^2$  such that

$$m_i(\theta) = \begin{cases} <=0 & if \ \theta \le t_i^1 \\ >0 & otherwise \end{cases}$$

$$m_i'(\theta) = \begin{cases} >= 0 & \text{if } \theta \le t_i^2 \\ < 0 & \text{otherwise} \end{cases}$$

The first property provides an upper bound on the marginal effect of prize i on the effort of agent of type  $\theta$  and is fairly straightforward. The second property says that something stronger is true for the most efficient agents. It says that the effort cost of an agent goes to zero as its marginal cost of effort goes to 0. The third property says that the overall effect of any prize i other than the first prize is zero. So increasing any prize i just transfers the effort from some set of agents to another set of agents. The first prize is special in that its overall effect is positive and equals 1 irrespective of the prior distribution. To prove this, we apply integration by parts to integrate  $\int_0^1 m_i(\theta) d\theta$  and using the second property, we get that it equals  $-\int_0^1 \theta m_i'(\theta) d\theta$ . By Leibniz rule, we know

$$m_i'(\theta) = -\binom{n-1}{i-1} \frac{[(1-F(\theta))^{n-i-1}F(\theta)^{i-2}]}{\theta} \left( (n-1)F(\theta) - (i-1) \right) f(\theta)$$

and so we are able to integrate  $\theta m_i'(\theta)$  to get the result. Moving on, the fourth property says the marginal effect of the first prize is positive for all types of agents and also decreasing in their type. And finally, the last property describes exactly how the effort transfers from one set of agents to another as we increase the value of some prize. In particular, for any prize i that is not the first prize, increasing its value leads to a transfer of effort from more efficient agents  $(\theta \leq t_i^1)$  to less efficient agents  $(\theta > t_i^1)$ . These properties are illustrated in Figure 1 for the case of n = 5 agents and prior cdf  $F(\theta) = \theta^3$ .

The overall expected effect of each prize i as described in Theorem 2 depends on the prior distribution. If there is a higher proportion of less efficient agents who are positively influenced by intermediate prizes, the overall effect of these intermediate prizes is positive. On the other hand, if there is a higher proportion of more efficient agents who are discouraged by these intermediate prizes, the overall effect is negative. The full proofs of Lemma 2 and Theorem 2 are in the appendix.




Figure 1: The marginal effect of prizes on effort for n=5 and  $F(\theta)=\theta^3$ .

Note that both the second and third parts of Theorem 2 apply to the case where F is uniform. Thus, we have the following corollary:

Corollary 2. Consider a setting with n agents and uniform prior  $F(\theta) = \theta$ . Suppose  $\mathbf{v} = (v_1, \dots, v_n)$  and  $\mathbf{w} = (w_1, \dots, w_n)$  are two prize vectors such that  $v_j > w_j$  for some  $j \in \{2, \dots, n-1\}$  and  $v_{-j} = w_{-j}$ .

- 1. if U is any increasing and concave function, then  $\mathbb{E}[U(g_{\mathbf{v}}(\theta))] \geq \mathbb{E}[U(g_{\mathbf{w}}(\theta))]$
- 2. if U is any increasing and convex function, then  $\mathbb{E}[U(g_{\mathbf{w}}(\theta))] \geq \mathbb{E}[U(g_{\mathbf{v}}(\theta))]$

In other words, for the uniform case,  $g_{\mathbf{w}}(\theta)$  is a mean-preserving spread of  $g_{\mathbf{v}}(\theta)$  which implies the comparisons in the corollary.

Theorem 2 identifies natural conditions on the prior distributions under which the expected marginal effects are positive or negative. Next, we consider a natural parametric class of the priors for which we can compute and compare the expected marginal effects of prizes. We then discuss what this comparison implies for the effects of competition in contests on the effort exerted by the agents under these priors. First, let us formally define what it means for a prize vector to be more competitive than another.

**Definition 3.1.** A prize vector  $\mathbf{v} = (v_1, v_2, \dots, v_{n-1}, v_n)$  is more competitive than  $\mathbf{w} = (w_1, w_2, \dots, w_{n-1}, w_n)$  if  $\mathbf{v}$  majorizes  $\mathbf{w}$  (i.e.  $\sum_{i=1}^k v_i \ge \sum_{i=1}^k w_i$  for all  $k \in [n]$  and  $\sum_{i=1}^n v_i = \sum_{i=1}^n w_i$ ).

This is the definition that was also considered in Fang et al. [23]. The next result describes the effect of competition on expected effort and expected minimum effort and how it depends on the prior distribution.

**Theorem 2.** Consider a setting with n agents and prior  $F(\theta) = \theta^p$ . Suppose  $\mathbf{v} = (v_1, \dots, v_n)$  and  $\mathbf{w} = (w_1, \dots, w_n)$  are two prize vectors such that  $\mathbf{v}$  is more competitive than  $\mathbf{w}$ . Then, we have the following:

1. if  $p \ge 1$ , then

$$\mathbb{E}[g_{\mathbf{v}}(\theta)] \ge \mathbb{E}[g_{\mathbf{w}}(\theta)]$$

2. if  $\frac{1}{2} \le p \le 1$ ,  $v_1 = w_1, v_n = w_n$ , then

$$\mathbb{E}[g_{\mathbf{v}}(\theta)] \le \mathbb{E}[g_{\mathbf{w}}(\theta)]$$

3. if  $p \ge \frac{1}{2}$  and  $v_n = w_n$ , then

$$\mathbb{E}[g_{\mathbf{v}}(\theta_{max})] \le \mathbb{E}[g_{\mathbf{w}}(\theta_{max})]$$

To prove this, we first show that for  $F(\theta) = \theta^p$ ,

$$\mathbb{E}[m_i(\theta)] = \binom{n-1}{i-1} \beta(i-\frac{1}{p}, n-i) \frac{(n-i)(p-1)}{np-1}$$

which implies that

$$\frac{\mathbb{E}[m_{i+1}(\theta)]}{\mathbb{E}[m_i(\theta)]} = \frac{n-i}{i} \frac{i - \frac{1}{p}}{n-i-1} \frac{n-i-1}{n-i} = \frac{i - \frac{1}{p}}{i} < 1$$

It follows then that for  $p \geq 1$ , the expected marginal effects  $\mathbb{E}[m_i(\theta)]$  are positive and decreasing in i. Since  $\mathbf{w}$  can be obtained from  $\mathbf{v}$  by a sequence of Robinhood transfers which involve a transfer of value from a top prize to a bottom prize, each of which reduces expected

effort, we get that the expected effort goes down as prize vector becomes less competitive. An analogous argument holds for  $\frac{1}{2} \le p \le 1$ .

For the case of expected minimum effort, we again find that

$$\mathbb{E}[m_i(\theta_{max})] = \binom{n-1}{i-1} \beta \left(n+i-1-\frac{1}{p}, n-i\right) \frac{(n-i)(np-1)}{2np-p-1}$$

which implies

$$\frac{\mathbb{E}[m_{i+1}(\theta_{max})]}{\mathbb{E}[m_i(\theta_{max})]} = \frac{n+i-1-\frac{1}{p}}{i} > 1$$

It follows then that less competitive prize vectors lead to higher expected minimum effort.

The fact that the expected marginal effect of prize  $v_i$  is decreasing in i for  $p \geq 1$  is perhaps a bit surprising since these are distributions where a large proportion of the agents are inefficient (the density is increasing in  $\theta$ ). Since these inefficient agents are generally competing for the intermediate prizes, one might expect that as their proportion increases (p increases), the expected marginal effect of the later prizes would increase relative to the earlier prizes. While we do get that the ratio  $\frac{\mathbb{E}[m_{i+1}(\theta)]}{\mathbb{E}[m_i(\theta)]} = \frac{i-\frac{1}{p}}{i}$  increases as p increases, it only goes to 1 as  $p \to \infty$ . Thus, the effect is not big enough to make later prizes more valuable than the earlier prizes in terms of the effort they induce. In contrast, for the case of expected minimum effort, the designer is putting even more weight on the effort of the least efficient agents. And it turns out that the marginal effect of prize  $v_i$  on expected minimum effort do increase as we increase i.

### f 4 Applications

In this section, we discuss some applications to the design of optimal contests in environments where the set of feasible prize vectors available to the designer may be constrained. In particular, we will consider three different environments. First, we'll consider settings where the designer can commit to a grading scheme and the value of these grades is determined by the information they reveal about the type of the agents. Second, we'll consider settings where agents have concave utilities for prizes and we'll derive the effort-maximizing prize structure under the standard constraint that the designer has a budget that it must allocate

across prizes. We'll also discuss how the optimal prize structure changes as the degree of concavity increases in the population. At last, we'll consider settings where the contest designer is constrained to award homogeneous prizes of a fixed value and only needs to decide how many prizes it must award to maximize effort.

#### 4.1 Optimal grading contests

First, we focus on the design of grading schemes where the contest designer doesn't have an explicit budget that it can distribute across prizes but instead, can choose a distribution of grades that it can award based on the rank of the agents. This is generally the case in classroom settings where the professor awards grades to students based on their performance in exams. For instance, the professor may commit to giving grades A and B to the top 50% and bottom 50% respectively, or it may give A+, A-, B+, and B- with distribution  $(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4})$ . Formally, we define a grading contest as follows:

**Definition 4.1.** A grading contest with n agents is defined by a strictly increasing sequence of natural numbers  $s = (s_1, s_2, \ldots, s_k)$  such that  $s_k = n$ .

The interpretation of grading contest s is that the top  $s_1$  agents get grade  $g_1$ , next  $s_2 - s_1$  get grade  $g_2$  and generally,  $s_k - s_{k-1}$  get grade  $g_k$ . There is a natural partial order over these grading contests in terms of how much information they reveal about the quality of the agents. From above, the grading contest that awards the grades A+, A-, B+, and B- in equal proportion is more informative about the agents type then the one that awards just A and B in equal proportion. More generally, we can say the following:

**Definition 4.2.** A grading contest s is more informative than s' if s' is a subsequence of s.

Clearly, the rank revealing contest  $s^* = (1, 2, ..., n)$  is more informative than any other grading contest.

To discuss how these grading contests compare in terms of the effort they induce, we need to define how the agents assign value to these grades. We assume that the value of a grade is determined by the information it reveals about the type of the agent. More precisely, we suppose that there is a publicly known wage function  $w:\Theta\to\mathbb{R}_+$  which maps an agent's marginal cost to its productivity and is monotone decreasing. So if the market could observe the type of the agent to be  $\theta$ , the agent would be offered a wage of  $w(\theta)$ . Given this wage function, we assume that a grading contest s induces the prize vector which is the expected productivity or wage of the agent given its grade. That is, if the market has a posterior belief f over the type of the agent, then the agent will get a wage equal to  $\int_0^1 w(\theta) f(\theta) d\theta$ .

Under this assumption, we get that the rank revealing contest  $s^* = (1, 2, ..., n)$  induces the prize vector

$$v_i = \mathbb{E}[w(\theta)|\theta = \theta_{(i)}^n]$$

where  $\theta_{(i)}^n$  is the *i*th order statistic in a random sample of *n* observations. This is because the rank revealing contest reveals the exact rank of the agent in a random sample of *n* observations. Note here that since  $\theta_{(i)}^n$  is stochastically dominated by  $\theta_{(j)}^n$  for all i < j and w is monotone decreasing, the prize vector induced by  $s^*$  is monotone decreasing  $v_1 > v_2 \cdots > v_n$ .

Now we can define the prize vectors induced by arbitrary grading contests s in terms of the  $v_i$ 's as defined above. An arbitrary grading contest  $s = (s_1, s_2, \ldots, s_k)$  induces the prize vector v(s) where

$$v(s)_i = \frac{v_{s_{j-1}+1} + v_{s_{j-1}+2} + \dots + v_{s_j}}{s_j - s_{j-1}}$$

and j is such that  $s_{j-1} < i \le s_j$ 

This is because if an agent gets grade  $g_j$  in the grading contest  $s = (s_1, s_2, \ldots, s_k)$ , then the market learns that the agent's type  $\theta$  must be ranked at one of  $\{s_{j-1} + 1, \ldots, s_j\}$  and further, it is equally likely to be ranked at any of these positions. The form of the prize vector above then follows from the assumption that the value of grade equals its expected productivity under the posterior induced by the grade.

Given this framework, we can now ask how the different grading schemes compare in terms of the effort they induce.

**Theorem 3.** Consider a setting with n agents and prior cdf  $F(\theta) = \theta^p$ . Suppose grading

scheme s is more informative than s'. Then, we have the following:

- if  $p \ge 1$ , then  $\mathbb{E}[g_{v(s)}(\theta)] \ge \mathbb{E}[g_{v(s')}(\theta)]$
- if  $\frac{1}{2} \le p \le 1$ ,  $v(s)_1 = v(s')_1$ , and  $v(s)_n = v(s')_n$ , then  $\mathbb{E}[g_{v(s)}(\theta)] \le \mathbb{E}[g_{v(s')}(\theta)]$
- if  $p \ge \frac{1}{2}$  and  $v(s)_n = v(s')_n$ , then  $\mathbb{E}[g_{v(s)}(\theta_{max})] \ge \mathbb{E}[g_{v(s')}(\theta_{max})]$

*Proof.* Observe that if s is more informative than s', then it induces a prize vector v(s) that is more competitive than the prize vector v(s'). The result then follows directly from Theorem 2.

With this comparison, we can now find the optimal grading contest in these settings.

Corollary 3. Consider a setting with n agents and prior  $cdf F(\theta) = \theta^p$ . Then, we have the following:

- if  $p \ge 1$ , the rank revealing contest s = (1, 2, ..., n) maximizes expected effort among all grading contests.
- if  $\frac{1}{2} \leq p \leq 1$ , the contest s = (1, n) in which the best agent gets a unique grade and all other agents get a common grade maximizes expected effort among all grading contests.
- if  $p \ge \frac{1}{2}$ , the contest s = (n 1, n) in which the worst agent gets a unique grade and all other agents get a common grade maximizes expected minimum effort among all grading contests.

Note that when the designer has a budget that it can distribute across prizes, the expected effort maximizing prize vector allocates the entire budget to the first prize. But as we see in the corollary, the optimal grading contest depends on the prior distribution of abilities. If the density of agents is increasing in  $\theta$  so that there is a greater proportion of inefficient agents, the effort maximizing grading contest awards a unique grade to each agent. But when the density is decreasing, the optimal grading contest awards a unique grade to the

best agent and pools the rest of the agents by awarding them a common grade. And for the case where the designer wants to maximize expected minimum effort, which is perhaps a reasonable objective in a classroom environment, the optimal grading contest awards a common grade to everyone except the least efficient agent.

#### 4.2 Optimal contests where agents have concave utility

Now we consider the contest design problem in a typical environment where the designer has a budget of B that it can allocate across prizes  $\mathbf{v} = (v_1, v_2, \dots, v_n)$  such that  $v_i \geq v_{i+1}$ . It is known that when agent's utility u(v) = v, the effort maximizing contest awards the entire prize budget B to the first prize ([44]). In this section, we consider the contest design problem where agents have concave utilities. More precisely, we assume that under a prize vector  $\mathbf{v}$ , if agent i of type  $\theta_i$  puts in effort  $e_i$  and wins prize j, its payoff equals  $u(v_j) - \theta_i e_i$  where  $u(v_j) = v_j^r$ . The next result characterizes the expected effort maximizing contest in this environment.

**Theorem 4.** Suppose there are n agents with utility  $u(v) = v^r$  and the prior cdf on marginal costs is  $F(\theta) = \theta^p$  with  $p \ge 1$ . The prize structure that maximizes expected effort is

$$v(r) = (v_1(r), c_2^{\frac{1}{1-r}} v_1(r), \dots c_{n-1}^{\frac{1}{1-r}} v_1(r), 0)$$

where

$$c_i = \frac{\mathbb{E}[m_i(\theta)]}{\mathbb{E}[m_1(\theta)]} < 1$$

and  $v_1(r)$  is such that

$$\sum v_i(r) = B$$

Further, if  $1 \ge r > r' > 0$ , then v(r) is more competitive than v(r').

The proof proceeds by using corollary 1 to identify the Bayes-Nash equilibrium function so that the problem becomes  $\max_{\mathbf{v}} \sum_{i=1}^{n-1} v^r \mathbb{E}[m_i(\theta)]$  such that  $\sum v_i = B$ . Solving this constrained optimization problem characterizes the optimal contest. To show that the optimal contest become more competitive as r increases, we define  $f_k(r)$  as the sum of the first k

prizes in the optimal contest under r and show that this sum is increasing in r. Thus, as the agents utility for prizes becomes more concave, the effort maximizing prize vector becomes less competitive.

#### 4.3 Optimal contests with costless homogeneous prizes

For our last application, we consider a setting where the contest designer can award arbitrarily many prizes of a fixed value a. More precisely, the set of prize vectors available to the designer is given by

$$B = {\mathbf{v} \in \mathbb{R}^n : \exists k \text{ such that } v_i = a \text{ if } i \leq k \text{ and } v_i = 0 \text{ if } i > k}$$

This might be the case in online contests run on platforms like Leetcode and Kaggle where the platforms essentially have an unlimited supply of digitial certificates or medals that they can award to participants in the contest. The designer wants to chose the number of prizes so as to maximize the expected effort. Note that this problem was also considered in [42] but under different distributional assumptions. In their setting, the authors found that the expected effort was single-peaked in k. For our setting, we have the following as a corollary of Theorem 2.

**Corollary 4.** Consider a setting with n agents and prior F. Then, we have the following:

- 1. if F is such that the density f is increasing, then the contest that awards k = n 1 prizes maximizes expected effort among the contests in B.
- 2. if F is such that the density f is decreasing, then the contest that awards k = 1 prize maximizes the expected effort among the contests in B.

In fact, in the first case where the density is increasing, we actually have something stronger. In this case, awarding k = n - 1 second order stochastically dominates awarding fewer number of prizes. In other words, for any concave increasing utility function U, the contest designer's optimal decision would be to award k = n - 1 prizes.

### 5 Conclusion

In contests where agents have private information about their abilities, we study the effect different prizes have on the effort exerted by the agents in equilibrium. While increasing the value of first prize encourages effort for all agents, increasing any intermediate prize leads to a balanced transfer of effort from the more efficient agents to less efficient agents. The overall expected effect then depends on the prior distribution of abilities. If the density of agents is increasing in inefficiency, the expected effects are positive. If this density is decreasing, the expected effects are negative. For a parametric subclass of priors with monotone density functions, we also study the effects of competition on expected effort and expected minimum effort.

We also discuss the application of these results to the design of effort-maximizing contests in settings where the set of feasible prize vectors may be constrained due to various reasons. First, we consider the design of grading contests under the assumption that the value of a grade is determined by the information it reveals about the type of the agent. We find that when the density function is increasing, more informative contests lead to higher expected effort and therefore, the rank revealing contest maximizes expected effort among all grading contests. Second, we consider the setting where the designer has a budget that it must allocate across different prizes and the agents have concave utilities for prizes. We find the optimal contest in this setting and show that it becomes more competitive as the degree of concavity decreases. Lastly, we consider settings where the designer can only choose the number of homogeneous prizes to award and show that when the prior density is monotone, it is optimal to award either 1 or n-1 prizes depending on whether the density is decreasing or increasing.

### References

- [1] Ales, L., S.-H. Cho, and E. Körpeoğlu (2017): "Optimal award scheme in innovation tournaments," *Operations Research*, 65, 693–702.
- [2] Barut, Y. and D. Kovenock (1998): "The symmetric multiple prize all-pay auction with complete information," *European Journal of Political Economy*, 14, 627–644. 5
- [3] Bastani, S., T. Giebe, and O. Gürtler (2019): "A general framework for studying contests," *Available at SSRN 3507264*.
- [4] Berry, S. K. (1993): "Rent-seeking with multiple winners," Public Choice, 77, 437–443.
- [5] Betts, J. R. (1998): "The impact of educational standards on the level and distribution of earnings," *The American Economic Review*, 88, 266–275. 7
- [6] Boleslavsky, R. and C. Cotton (2015): "Grading standards and education quality," *American Economic Journal: Microeconomics*, 7, 248–79. 7
- [7] Bozbay, I. and A. Vesperoni (2018): "A contest success function for networks," Journal of Economic Behavior & Organization, 150, 404–422.
- [8] Chan, W., L. Hao, and W. Suen (2007): "A signaling theory of grade inflation," International Economic Review, 48, 1065–1090. 7
- [9] Chawla, S., J. D. Hartline, and B. Sivan (2019): "Optimal crowdsourcing contests," *Games and Economic Behavior*, 113, 80–96.
- [10] Chowdhury, S. M. (2021): "The economics of identity and conflict," in Oxford Research Encyclopedia of Economics and Finance.
- [11] Chowdhury, S. M., P. Esteve-González, and A. Mukherjee (2020): "Heterogeneity, leveling the playing field, and affirmative action in contests," *Leveling the Playing Field, and Affirmative Action in Contests (July 20, 2020)*.

- [12] Chowdhury, S. M. and S.-H. Kim (2017): ""Small, yet Beautiful": Reconsidering the optimal design of multi-winner contests," *Games and Economic Behavior*, 104, 486–493.
- [13] CLARK, D. J. AND C. RIIS (1996): "A multi-winner nested rent-seeking contest," Public Choice, 87, 177–184.
- [14] ——— (1998): "Competition over more than one prize," The American Economic Review, 88, 276–289. 6
- [15] ——— (1998): "Influence and the discretionary allocation of several prizes," European Journal of Political Economy, 14, 605–625. 6
- [16] COHEN, C. AND A. SELA (2008): "Allocation of prizes in asymmetric all-pay auctions," European Journal of Political Economy, 24, 123–132. 6
- [17] CORCHÓN, L. AND M. DAHM (2011): "Welfare maximizing contest success functions when the planner cannot commit," *Journal of Mathematical Economics*, 47, 309–317.
- [18] CORCHÓN, L. C. (2007): "The theory of contests: a survey," Review of economic design, 11, 69–100. 7
- [19] COSTRELL, R. M. (1994): "A simple model of educational standards," The American Economic Review, 956–971. 7
- [20] Dubey, P. and J. Geanakoplos (2010): "Grading exams: 100, 99, 98,... or a, b, c?" Games and Economic Behavior, 69, 72–94. 7
- [21] EWERHART, C. (2015): "Mixed equilibria in Tullock contests," *Economic Theory*, 60, 59–71.
- [22] EWERHART, C. AND F. QUARTIERI (2020): "Unique equilibrium in contests with incomplete information," *Economic Theory*, 70, 243–271.

- [23] FANG, D., T. NOE, AND P. STRACK (2020): "Turning up the heat: The discouraging effect of competition in contests," *Journal of Political Economy*, 128, 1940–1975. 5, 14
- [24] FARAVELLI, M. AND L. STANCA (2012): "When less is more: rationing and rent dissipation in stochastic contests," *Games and Economic Behavior*, 74, 170–183.
- [25] Fu, Q. and Z. Wu (2018): "On the optimal design of lottery contests," Available at SSRN 3291874.
- [26] ——— (2019): "Contests: Theory and topics," in Oxford Research Encyclopedia of Economics and Finance. 7
- [27] ——— (2020): "On the optimal design of biased contests," *Theoretical Economics*, 15, 1435–1470.
- [28] Gallice, A. (2017): "An approximate solution to rent-seeking contests with private information," European Journal of Operational Research, 256, 673–684.
- [29] GHOSH, A. AND R. KLEINBERG (2016): "Optimal contest design for simple agents," ACM Transactions on Economics and Computation (TEAC), 4, 1–41.
- [30] GLAZER, A. AND R. HASSIN (1988): "Optimal contests," *Economic Inquiry*, 26, 133–143. 5
- [31] HINNOSAAR, T. (2018): "Optimal sequential contests," arXiv preprint arXiv:1802.04669.
- [32] Immorlica, N., G. Stoddard, and V. Syrgkanis (2015): "Social status and badge design," in *Proceedings of the 24th international conference on World Wide Web*, 473–483.
- [33] Jia, H. (2008): "A stochastic derivation of the ratio form of contest success functions," *Public Choice*, 135, 125–130.

- [34] Konrad, K. A. et al. (2009): "Strategy and dynamics in contests," *OUP Catalogue*.
  7
- [35] KÖRPEOĞLU, E. AND S.-H. CHO (2018): "Incentives in contests with heterogeneous solvers," *Management Science*, 64, 2709–2715.
- [36] Krishna, V. and J. Morgan (1997): "An analysis of the war of attrition and the all-pay auction," journal of economic theory, 72, 343–362.
- [37] ——— (1998): "The winner-take-all principle in small tournaments," Advances in applied microeconomics, 7, 61–74. 6
- [38] LAZEAR, E. P. AND S. ROSEN (1981): "Rank-order tournaments as optimum labor contracts," *Journal of political Economy*, 89, 841–864.
- [39] Letina, I., S. Liu, and N. Netzer (2020): "Optimal contest design: A general approach," Tech. rep., Discussion Papers.
- [40] Liu, B. and J. Lu (2019): "The optimal allocation of prizes in contests with costly entry," *International Journal of Industrial Organization*, 66, 137–161.
- [41] Liu, B., J. Lu, R. Wang, and J. Zhang (2018): "Optimal prize allocation in contests: The role of negative prizes," *Journal of Economic Theory*, 175, 291–317.
- [42] Liu, X. And J. Lu (2017): "Optimal prize-rationing strategy in all-pay contests with incomplete information," *International Journal of Industrial Organization*, 50, 57–90. 6, 7, 20
- [43] Lu, J. and Z. Wang (2015): "Axiomatizing multi-prize nested lottery contests: a complete and strict ranking perspective," *Journal of Economic Behavior & Organization*, 116, 127–141.
- [44] Moldovanu, B. and A. Sela (2001): "The optimal allocation of prizes in contests,"

  American Economic Review, 91, 542–558. 6, 9, 19

- [45] ——— (2006): "Contest architecture," Journal of Economic Theory, 126, 70–96.
- [46] MOLDOVANU, B., A. SELA, AND X. SHI (2007): "Contests for status," Journal of political Economy, 115, 338–363. 7
- [47] NITZAN, S. (1994): "Modelling rent-seeking contests," European Journal of Political Economy, 10, 41–60.
- [48] Olszewski, W. and R. Siegel (2020): "Performance-maximizing large contests," Theoretical Economics, 15, 57–88.
- [49] Popov, S. V. and D. Bernhardt (2013): "University competition, grading standards, and grade inflation," *Economic inquiry*, 51, 1764–1778. 7
- [50] RAYO, L. (2013): "Monopolistic signal provision," The BE Journal of Theoretical Economics, 13, 27–58. 7
- [51] RYVKIN, D. AND M. DRUGOV (2020): "The shape of luck and competition in winner-take-all tournaments," *Theoretical Economics*, 15, 1587–1626.
- [52] SEGEV, E. (2020): "Crowdsourcing contests," European Journal of Operational Research, 281, 241–255. 7
- [53] Segev, E. and A. Sela (2014): "Multi-stage sequential all-pay auctions," *European Economic Review*, 70, 371–382.
- [54] ——— (2014): "Sequential all-pay auctions with head starts," *Social Choice and Welfare*, 43, 893–923.
- [55] SHAKED, M. AND J. G. SHANTHIKUMAR (2007): Stochastic orders, Springer. 31, 32
- [56] Siegel, R. (2009): "All-pay contests," *Econometrica*, 77, 71–92.
- [57] SISAK, D. (2009): "Multiple-Prize Contests-The Optimal Allocation Of Prizes," *Journal of Economic Surveys*, 23, 82–114. 6

- [58] Skaperdas, S. (1996): "Contest success functions," Economic theory, 7, 283–290.
- [59] SZYMANSKI, S. AND T. M. VALLETTI (2005): "Incentive effects of second prizes," European Journal of Political Economy, 21, 467–481. 6
- [60] Vojnović, M. (2015): Contest theory: Incentive mechanisms and ranking methods, Cambridge University Press. 7
- [61] Zhang, M. (2019): "Optimal Contests with Incomplete Information and Convex Effort Costs," Available at SSRN 3512155. 5
- [62] Zubrickas, R. (2015): "Optimal grading," International Economic Review, 56, 751–776. 7

## A Proofs for Section 3 (Equilibrium)

**Lemma 1.** In a contest with n agents, prizes  $\mathbf{v} = (v_1, v_2, \dots, v_{n-1}, v_n)$  and prior cdf F, the symmetric Bayes-Nash equilibrium strategy function is given by

$$g_{\mathbf{v}}(\theta) = \sum_{i=1}^{n} m_i(\theta) v_i$$

where,

$$m_i(\theta) = \binom{n-1}{i-1} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i-1}t^{i-2}]}{F^{-1}(t)} \left( (n-1)t - (i-1) \right) dt$$

for all  $i \in \{1, 2, ..., n\}$ .

*Proof.* Suppose n-1 agents are playing a monotone decreasing strategy  $g(\theta)$ . Let  $\theta_{(j)}^n$  denote the jth order statistic from n random draws with  $\theta_{(0)}^n = 0$  and  $\theta_{n+1}^n = 1$ . Then, an agent of type  $\theta$ 's utility from putting in x units of effort is given by:

$$u(\theta, x) = \sum_{i=1}^{n} v_i \Pr[\theta_{(i-1)}^{n-1} \le g^{-1}(x) \le \theta_{(i)}^{n-1}] - \theta x$$
$$= \sum_{i=1}^{n} v_i \binom{n-1}{i-1} F(g^{-1}(x))^{i-1} (1 - F(g^{-1}(x)))^{n-i} - \theta x$$

Now, differentiating with respect to x gives:

$$\frac{\partial u(\theta, x)}{\partial x} = \frac{f(g^{-1}(x))}{g'(g^{-1}(x))} \sum_{i=1}^{n} v_i \binom{n-1}{i-1} 
\left[ (1 - F(g^{-1}(x)))^{n-i} (i-1) F(g^{-1}(x))^{i-2} - F(g^{-1}(x))^{i-1} (n-i) (1 - F(g^{-1}(x)))^{n-i-1} \right] - \theta$$

. Setting it to 0 and plugging in  $g(\theta) = x$  gives the condition for  $g(\theta)$  to be a symmetric Bayes-Nash equilibrium:

$$f(\theta) \sum_{i=1}^{n} v_i \binom{n-1}{i-1} \left[ (1 - F(\theta))^{n-i} (i-1) F(\theta)^{i-2} - F(\theta)^{i-1} (n-i) (1 - F(\theta))^{n-i-1} \right] = \theta g'(\theta)$$

An alternate way to write this condition is:

$$-f(\theta) \sum_{i=1}^{n-1} (v_i - v_{i+1}) \frac{(n-1)!}{(i-1)!(n-i-1)!} \left[ (1 - F(\theta))^{n-i-1} F(\theta)^{i-1} \right] = \theta g'(\theta)$$

Using the boundary condition g(1) = 0, we get that the symmetric Bayes-Nash equilibrium function is given by

$$\int_{\theta}^{1} \frac{f(\theta)}{\theta} \sum_{i=1}^{n-1} (v_i - v_{i+1}) \frac{(n-1)!}{(i-1)!(n-i-1)!} \left[ (1 - F(\theta))^{n-i-1} F(\theta)^{i-1} \right] d\theta$$

Replacing  $F(\theta) = t$ , we get

$$g(\theta) = \int_{F(\theta)}^{1} \frac{1}{F^{-1}(t)} \sum_{i=1}^{n-1} \left( v_i - v_{i+1} \right) \frac{(n-1)!}{(i-1)!(n-i-1)!} \left[ (1-t)^{n-i-1} t^{i-1} \right] dt$$

Bringing the summation outside:

$$g_v(\theta) = \sum_{i=1}^{n-1} (v_i - v_{i+1}) \frac{(n-1)!}{(i-1)!(n-i-1)!} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i-1}t^{i-1}]}{F^{-1}(t)} dt$$

We can also write the equilibrium function as  $g_v(\theta) = \sum_{i=1}^n m_i(\theta) v_i$  where for  $i \geq 2$ 

$$\begin{split} m_i(\theta) &= \frac{(n-1)!}{(i-1)!(n-i-1)!} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i-1}t^{i-1}]}{F^{-1}(t)} dt - \frac{(n-1)!}{(i-2)!(n-i)!} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i}t^{i-2}]}{F^{-1}(t)} dt \\ &= \frac{(n-1)!}{(i-2)!(n-i-1)!} \int_{F(\theta)}^1 \left( \frac{[(1-t)^{n-i-1}t^{i-1}]}{(i-1)F^{-1}(t)} - \frac{[(1-t)^{n-i}t^{i-2}]}{(n-i)F^{-1}(t)} \right) dt \\ &= \frac{(n-1)!}{(i-2)!(n-i-1)!} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i-1}t^{i-2}]}{F^{-1}(t)} \left( \frac{t}{(i-1)} - \frac{1-t}{n-i} \right) dt \\ &= \binom{n-1}{i-1} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i-1}t^{i-2}]}{F^{-1}(t)} \left( (n-1)t - (i-1) \right) dt \end{split}$$

For i = 1, we have that

$$m_1(\theta) = (n-1) \int_{F(\theta)}^{1} \frac{[(1-t)^{n-2}]}{F^{-1}(t)} dt$$

Now we check that the second order condition is satisfied. To simplify calculations, let  $g^{-1}(x) = t$  so the agent of type  $\theta$  is imitating an agent of type t. Then, the foc can be written as:

$$f(t)\sum_{i=1}^{n} v_i \binom{n-1}{i-1} \left[ (1-F(t))^{n-i} (i-1)F(t)^{i-2} - F(t)^{i-1} (n-i)(1-F(t))^{n-i-1} \right] - \theta g'(t) = 0$$

or alternatively

$$-\sum_{i=1}^{n-1} (v_i - v_{i+1}) \binom{n-1}{i-1} (n-i) f(t) \left[ (1 - F(t))^{n-i-1} F(t)^{i-1} \right] - \theta g'(t) = 0$$

Let 
$$V(t) = -\sum_{i=1}^{n-1} (v_i - v_{i+1}) \binom{n-1}{i-1} (n-i) f(t) [(1 - F(t))^{n-i-1} F(t)^{i-1}]$$

Then, the foc is that V(t) = tg'(t) and so V'(t) = tg''(t) + g'(t). Taking the derivative of lhs of the foc wrt t gives

$$V'(t) - \theta g''(t) = V'(t) - \theta \frac{(V'(t) - g'(t))}{t}$$

At  $t = \theta$ , we get that this equals  $g'(\theta)$  which we know is < 0. Thus, the second order condition is satisfied.

**Theorem 1.** Consider a setting with n agents and prior F. Suppose  $\mathbf{v} = (v_1, \dots, v_n)$  and  $\mathbf{w} = (w_1, \dots, w_n)$  are two prize vectors. Then, we have the following:

- 1. if  $v_1 > w_1$  and  $v_{-1} = w_{-1}$ , then for any increasing function U, we have  $\mathbb{E}[U(g_{\mathbf{v}}(\theta))] \geq \mathbb{E}[U(g_{\mathbf{w}}(\theta))]$
- 2. if  $v_j > w_j$  for some  $j \in \{2, ..., n-1\}$ ,  $v_{-j} = w_{-j}$ , and the prior density function f is increasing, then for any increasing and concave function U, we have  $\mathbb{E}[U(g_{\mathbf{v}}(\theta))] \geq \mathbb{E}[U(g_{\mathbf{w}}(\theta))]$
- 3. if  $v_j > w_j$  for some  $j \in \{2, ..., n-1\}$ ,  $v_{-j} = w_{-j}$ , and the prior density function f is decreasing, then for any increasing and convex function U, we have  $\mathbb{E}[U(g_{\mathbf{v}}(\theta))] \geq \mathbb{E}[U(g_{\mathbf{v}}(\theta))]$ .

*Proof.* Let's prove each of the claims in order.

For the first claim, we know from lemma 2 that  $g_v(\theta) > g_w(\theta)$  for all  $\theta \in [0, 1]$  since  $m_1(\theta) > 0$  for all  $\theta \in [0, 1]$ . As a result,

$$\mathbb{P}[g_w(\theta) \le x] = \mathbb{P}[\theta \ge g_w^{-1}(x)] \ge \mathbb{P}[\theta \ge g_v^{-1}(x)] = \mathbb{P}[g_v(\theta) \le x]$$

Note that the result also follows from Theorem 1.A.17 in [55].

Now let's prove the second claim. First we'll show that  $\mathbb{E}[m_j(\theta)] \geq 0$ . Note that  $\mathbb{E}[m_j(\theta)] = \int_0^1 m_j(\theta) f(\theta) d\theta$ . For any  $j \in \{2, \dots, n-1\}$ , we know from Lemma 2 that there exists  $t_j^1$  such that  $m_j(\theta) = \begin{cases} <=0 & \text{if } \theta \leq t_j^1 \\ >0 & \text{otherwise} \end{cases}$ 

Using this, we have that

$$\mathbb{E}[m_j(\theta)] = \int_0^1 m_j(\theta) f(\theta) d\theta$$

$$= \int_0^{t_j^1} m_j(\theta) f(\theta) d\theta + \int_{t_j^1}^1 m_j(\theta) f(\theta) d\theta$$

$$\geq \int_0^{t_j^1} m_j(\theta) f(t_j^1) d\theta + \int_{t_j^1}^1 m_j(\theta) f(t_j^1) d\theta$$

$$= f(t_j^1) \int_0^1 m_j(\theta) d\theta$$

$$= 0$$

. It follows then that  $\mathbb{E}[g_v(\theta)] \geq \mathbb{E}[g_w(\theta)]$ . In addition, we know from lemma 2 that there

exists 
$$t_j^1$$
 such that  $g_v(\theta) - g_w(\theta) = \begin{cases} < 0 & \text{if } \theta < t_j^1 \\ = 0 & \text{if } \theta = t_j^1 \\ > 0 & \text{otherwise} \end{cases}$ 

Let  $G_v(x) = \mathbb{P}[g_v(\theta) \leq x]$  denote the cdf of effort under prize vector  $\mathbf{v}$ . Then, from above, we have that

$$G_v(x) - G_w(x) = \begin{cases} < 0 & \text{if } x < g_v(t_j^1) \\ = 0 & \text{if } x = g_v(t_j^1) \\ > 0 & \text{otherwise} \end{cases}$$

Thus, we have that  $\mathbb{E}[g_v(\theta)] \geq \mathbb{E}[g_w(\theta)]$  and also the sign of  $G_v(x) - G_w(x)$  changes exactly once from - to + as x increases. It follows then from Theorem 4.A.22 in [55] that  $g_v(\theta)$  second order stochastically dominates  $g_w(\theta)$ .

The argument for the case of decreasing density is analogous.

**Lemma 2.** For any number of agents n and prior F, the following hold:

- 1.  $\theta m_i(\theta) \le 1 \text{ for all } i \in \{1, \dots, n-1\}, \theta \in [0, 1]$
- 2.  $\lim_{\theta \to 0} \theta m_i(\theta) = 0 \text{ for all } i \in \{1, ..., n-1\}$
- 3.  $\int_0^1 m_1(\theta) = 1$  and  $\int_0^1 m_i(\theta) = 0$  for  $i \in \{2, n-1\}$
- 4.  $m_1(\theta) > 0$  for all  $\theta$  and monotone decreasing
- 5. For  $i \in \{2, n-1\}$ , there exist  $t_i^1 < t_i^2$  such that

$$m_i(\theta) = \begin{cases} <= 0 & \text{if } \theta \le t_i^1 \\ > 0 & \text{otherwise} \end{cases}$$

$$m_i'(\theta) = \begin{cases} >= 0 & \text{if } \theta \le t_i^2 \\ < 0 & \text{otherwise} \end{cases}$$

*Proof.* We'll prove the properties one by one.

The first property provides an upper bound on the marginal effect of any prize i on the effort of agent of type  $\theta$ :  $m_i(\theta) \leq \frac{1}{\theta}$ . To prove this, consider a case where prize i increases by  $\Delta$ . Then from the characterization in Theorem 1, it follows that an agent of type  $\theta$  will increase its effort by  $\Delta m_i(\theta)$ . This would correspond to an increase in cost of  $\Delta \theta m_i(\theta)$ . But the overall gain from the increased prize is  $\leq \Delta$ . Since the change in cost must be less than the gain in prize, we get a simple bound of  $\theta m_i(\theta) \leq 1$  for all  $\theta$ .

The second property essentially says that the cost of the most efficient agent goes to zero. We'll first prove this property for  $i \in \{2, n-1\}$  by squeeze theorem. First let's obtain an upper bound:

$$\theta m_{i}(\theta) = \binom{n-1}{i-1} \theta \int_{F(\theta)}^{1} \frac{[(1-t)^{n-i-1}t^{i-2}]}{F^{-1}(t)} ((n-1)t - (i-1)) dt$$

$$\leq \binom{n-1}{i-1} \theta \int_{F(\theta)}^{1} \frac{[(1-t)^{n-i-1}t^{i-2}]}{\theta} ((n-1)t - (i-1)) dt$$

$$= \binom{n-1}{i-1} \int_{F(\theta)}^{1} (1-t)^{n-i-1}t^{i-2} ((n-1)t - (i-1)) dt$$

At  $\theta = 0$ , the integral equals  $(n-1)\beta(i, n-i) - (i-1)\beta(i-1, n-i) = 0$ 

For the lower bound, we have

$$\theta m_i(\theta) \ge \binom{n-1}{i-1} \theta \int_{F(\theta)}^1 (1-t)^{n-i-1} t^{i-2} \left( (n-1)t - (i-1) \right) dt$$

which goes to 0 as  $\theta \to 0$ . Thus,  $\lim_{\theta \to 0} \theta m_i(\theta) = 0$  for  $i \in \{2, 3, \dots, n-1\}$ 

Now we'll prove the limit is 0 for i = 1. We have

$$\theta m_1(\theta) = \theta(n-1) \int_{F(\theta)}^1 \frac{[(1-t)^{n-2}]}{F^{-1}(t)} dt$$

. If the integral is finite, we are done. If it is infinite, we can apply L-Hospital's rule to get

$$\lim_{\theta \to 0} \frac{m_1(\theta)}{\frac{1}{\theta}} = \lim_{\theta \to 0} -\theta^2 m_1'(\theta) = \lim_{\theta \to 0} \theta^2 (n-1) f(\theta) \frac{(1-F(\theta))^{n-2}}{\theta} = 0$$

Now we prove the third property. By Leibniz rule, for  $i \geq 2$ , we have

$$m_i'(\theta) = -\binom{n-1}{i-1} \frac{[(1-F(\theta))^{n-i-1}F(\theta)^{i-2}]}{\theta} \left( (n-1)F(\theta) - (i-1) \right) f(\theta)$$

Since  $\lim_{\theta\to 0} \theta m_i(\theta) = 0$ , we have that  $\int_0^1 \theta m_i'(\theta) = -\int_0^1 m_i(\theta) d\theta$ 

From above, we have that

$$\begin{split} \int_0^1 \theta m_i'(\theta) d\theta &= -\int_0^1 \theta \binom{n-1}{i-1} \frac{[(1-F(\theta))^{n-i-1}F(\theta)^{i-2}]}{\theta} \left( (n-1)F(\theta) - (i-1) \right) f(\theta) d\theta \\ &= -\binom{n-1}{i-1} \int_0^1 \left[ (1-t)^{n-i-1} t^{i-2} \right] \left( (n-1)t - (i-1) \right) dt \\ &= 0 \end{split}$$

Thus, we get that  $\int_0^1 m_i(\theta) d\theta = 0$  for  $i \geq 2$ . For i = 1, we have that

$$m_1(\theta) = (n-1) \int_{F(\theta)}^{1} \frac{[(1-t)^{n-2}]}{F^{-1}(t)} dt$$

so that  $m_1'(\theta) = -(n-1)\frac{(1-F(\theta))^{n-2}}{\theta}f(\theta)$  and thus,  $\int_0^1 \theta m_1'(\theta)d\theta = -1$ . This gives that  $\int_0^1 m_1(\theta)d\theta = 1$ .

The fourth property follows from the fact that  $m'_1(\theta) < 0$  and  $m_1(1) = 0$ .

For the last property, we can use the expression for  $m_i'(\theta)$  to get that  $t_i^2 = F^{-1}(\frac{i-1}{n-1})$ . The claim on existence of  $t_i^1 < t_i^2$  then follows from the fact that  $\int_0^1 m_i(\theta) d\theta = 0$  for  $i \in \{2, \dots, n-1\}$ .

**Theorem 2.** Consider a setting with n agents and prior  $F(\theta) = \theta^p$ . Suppose  $\mathbf{v} = (v_1, \dots, v_n)$  and  $\mathbf{w} = (w_1, \dots, w_n)$  are two prize vectors such that  $\mathbf{v}$  is more competitive than  $\mathbf{w}$ . Then, we have the following:

1. if  $p \ge 1$ , then

$$\mathbb{E}[g_{\mathbf{v}}(\theta)] \ge \mathbb{E}[g_{\mathbf{w}}(\theta)]$$

2. if  $\frac{1}{2} \le p \le 1$ ,  $v_1 = w_1, v_n = w_n$ , then

$$\mathbb{E}[g_{\mathbf{v}}(\theta)] \le \mathbb{E}[g_{\mathbf{w}}(\theta)]$$

3. if  $p \ge \frac{1}{2}$  and  $v_n = w_n$ , then

$$\mathbb{E}[g_{\mathbf{v}}(\theta_{max})] \le \mathbb{E}[g_{\mathbf{w}}(\theta_{max})]$$

*Proof.* First, we show that the expected marginal effect of prize  $v_i$  is

$$\mathbb{E}[m_i(\theta)] = \binom{n-1}{i-1} \beta(i-\frac{1}{p}, n-i) \frac{(n-i)(p-1)}{np-1}$$

This follows from the following calculations:

$$\mathbb{E}[m_i(\theta)] = -\int_0^1 F(\theta)m_i'(\theta)d\theta$$

$$= \binom{n-1}{i-1} \int_0^1 \frac{[(1-F(\theta))^{n-i-1}F(\theta)^{i-1}]}{\theta} \left((n-1)F(\theta) - (i-1)\right) f(\theta)d\theta$$

$$= \binom{n-1}{i-1} \int_0^1 \frac{[(1-t)^{n-i-1}t^{i-1}]}{F^{-1}(t)} \left((n-1)t - (i-1)\right) dt$$

For  $F(\theta) = \theta^p$ ,

$$\mathbb{E}[m_{i}(\theta)] = \binom{n-1}{i-1} \left( (n-1)\beta(i+1-\frac{1}{p},n-i) - (i-1)\beta(i-\frac{1}{p},n-i) \right)$$

$$= \binom{n-1}{i-1} \left( (n-1)\beta(i-\frac{1}{p},n-i)\frac{i-\frac{1}{p}}{n-\frac{1}{p}} - (i-1)\beta(i-\frac{1}{p},n-i) \right)$$

$$= \binom{n-1}{i-1}\beta(i-\frac{1}{p},n-i) \left( (n-1)\frac{i-\frac{1}{p}}{n-\frac{1}{p}} - (i-1) \right)$$

$$= \binom{n-1}{i-1}\beta(i-\frac{1}{p},n-i)\frac{(n-i)(p-1)}{np-1}$$

Now observe that

$$\frac{\mathbb{E}[m_{i+1}(\theta)]}{\mathbb{E}[m_i(\theta)]} = \frac{n-i}{i} \frac{i - \frac{1}{p}}{n-i-1} \frac{n-i-1}{n-i} = \frac{i - \frac{1}{p}}{i} < 1$$

Note that for  $p \geq 1$ , these marginal effects are positive and thus, the marginal effect of prize i is decreasing in i. This implies that the change in expected effort from increasing any prize  $i \in [n-1]$  is positive and the change from increasing  $v_i$  is greater than that from increasing  $v_j$  for any i < j. Since w can be obtained from v via a sequence of Robinhood operations which involve replacing  $v_i$  by  $v_i - \epsilon$  and  $v_j$  by  $v_j + \epsilon$ , each of which reduces expected effort, we get that the expected effort under w will be lesser than the expected effort under v when  $p \geq 1$ . So if v is more competitive than w, then  $\mathbb{E}[g_v(\theta)] \geq \mathbb{E}[g_w(\theta)]$ . For  $\frac{1}{2} \leq p \leq 1$ , the expected marginal effects are v = 0 but the ratio is still v = 0. Thus, the expected marginal effects are actually increasing in v = 0 and so we get the inequality in the second item.

For the inequality in the third item, we first show that

$$\mathbb{E}[m_i(\theta_{max})] = \binom{n-1}{i-1} \beta \left(n+i-1-\frac{1}{p}, n-i\right) \frac{(n-i)(np-1)}{2np-p-1}$$

for  $i \in \{1, 2, \dots, n-1\}$ .

$$\mathbb{E}[m_i(\theta_{max})] = \int_0^1 m_i(\theta) n F(\theta)^{n-1} f(\theta) d\theta$$

$$= \binom{n-1}{i-1} \int_0^1 \frac{\left[ (1-F(\theta))^{n-i-1} F(\theta)^{i-2} \right]}{\theta} \left( (n-1) F(\theta) - (i-1) \right) F(\theta)^n f(\theta) d\theta$$

$$= \binom{n-1}{i-1} \int_0^1 \frac{\left[ (1-t)^{n-i-1} t^{n+i-2} \right]}{F^{-1}(t)} \left( (n-1)t - (i-1) \right) dt$$

For the case of  $F(\theta) = \theta^p$ , we get that

$$\mathbb{E}[m_{i}(\theta_{max})] = \binom{n-1}{i-1} \left( (n-1)\beta \left( n+i-\frac{1}{p}, n-i \right) - (i-1)\beta \left( n+i-1-\frac{1}{p}, n-i \right) \right)$$

$$= \binom{n-1}{i-1}\beta \left( n+i-1-\frac{1}{p}, n-i \right) \left( (n-1)\frac{n+i-1-\frac{1}{p}}{2n-1-\frac{1}{p}} - (i-1) \right)$$

$$= \binom{n-1}{i-1}\beta \left( n+i-1-\frac{1}{p}, n-i \right) \frac{(n-i)(np-1)}{2np-p-1}$$

Now observe that

$$\frac{\mathbb{E}[m_{i+1}(\theta_{max})]}{\mathbb{E}[m_i(\theta_{max})]} = \frac{n+i-1-\frac{1}{p}}{i} > 1$$

Thus, the marginal effect of prize i is increasing in i. Again, since w can be obtained from v via a sequence of Robinhood operations which involve replacing  $v_i$  by  $v_i - \epsilon$  and  $v_j$  by  $v_j + \epsilon$ , each of which increases expected minimum effort, we get that the expected minimum effort under w will be greater than that under v. It follows that if v is more competitive than w and both have the same last prize, then  $\mathbb{E}[g_v(\theta_{max})] \leq \mathbb{E}[g_w(\theta_{max})]$ 

## B Proofs for Section 4 (Applications)

**Theorem 4.** Suppose there are n agents with utility  $u(v) = v^r$  and the prior cdf on marginal costs is  $F(\theta) = \theta^p$  with  $p \ge 1$ . The prize structure that maximizes expected effort is

$$v(r) = (v_1(r), c_2^{\frac{1}{1-r}} v_1(r), \dots c_{n-1}^{\frac{1}{1-r}} v_1(r), 0)$$

where

$$c_i = \frac{\mathbb{E}[m_i(\theta)]}{\mathbb{E}[m_1(\theta)]} < 1$$

and  $v_1(r)$  is such that

$$\sum v_i(r) = B$$

Further, if  $1 \ge r > r' > 0$ , then v(r) is more competitive than v(r').

*Proof.* From corollary 1, we know that the Bayes-Nash equilibrium function takes the form

$$g_v(\theta) = \sum_{i=1}^n m_i(\theta) u(v_i)$$

where,

$$m_i(\theta) = \binom{n-1}{i-1} \int_{F(\theta)}^1 \frac{[(1-t)^{n-i-1}t^{i-2}]}{F^{-1}(t)} \left( (n-1)t - (i-1) \right) dt$$

for  $i \in [n-1]$  and  $m_n(\theta) = -\sum_{i=1}^{n-1} m_i(\theta)$ . Given this form of the equilibrium function, the problem is

$$\max_{\mathbf{v}} \sum_{i=1}^{n-1} u(v_i) \mathbb{E}[m_i(\theta)]$$

such that  $\sum_{i=1}^{n-1} v_i = B$ .

Check that the solution will satisfy the equation

$$V_1(r)\left[1 + \sum_{i=2}^{n-1} c_i^{\frac{1}{1-r}}\right] = B$$

where  $c_i = \frac{\mathbb{E}[m_i(\theta)]}{\mathbb{E}[m_1(\theta)]} < 1$  and  $c_i > c_{i+1}$  for all i. Note that  $c_i$  does not depend on r.

Let 
$$f_k(r) = V_1(r) \left[ 1 + \sum_{i=2}^k c_i^{\frac{1}{1-r}} \right].$$

I want to show that  $f'_k(r) > 0$  for all k.

If I can show  $f'_k(r)$  is single peaked in k, that would imply the result since  $f_n(r) = 0$ .

Check that 
$$V_1'(r) = \frac{-\left[\sum_{i=2}^{n-1} c_i^{\frac{1}{1-r}} \log(c_i)\right] V_1^2(r)}{(1-r)^2 B}$$
 Plugging it in, we get

$$f'_{k}(r) = V_{1}(r) \left[ \frac{1}{(1-r)^{2}} \sum_{i=2}^{k} c_{i}^{\frac{1}{1-r}} \log(c_{i}) \right] + V'_{1}(r) \left[ 1 + \sum_{i=2}^{k} c_{i}^{\frac{1}{1-r}} \right]$$

$$= V_{1}(r) \left[ \frac{1}{(1-r)^{2}} \sum_{i=2}^{k} c_{i}^{\frac{1}{1-r}} \log(c_{i}) \right] - \frac{\left[ \sum_{i=2}^{n-1} c_{i}^{\frac{1}{1-r}} \log(c_{i}) \right] V_{1}^{2}(r)}{(1-r)^{2}B} \left[ 1 + \sum_{i=2}^{k} c_{i}^{\frac{1}{1-r}} \right]$$

$$= \frac{V_1(r)}{(1-r)^2} \sum_{i=2}^k c_i^{\frac{1}{1-r}} \log(c_i) \left[ 1 - \frac{V_1(r)}{B} \left( 1 + \sum_{i=2}^k c_i^{\frac{1}{1-r}} \right) \right] - \frac{V_1^2(r)}{B(1-r)^2} \sum_{i=k+1}^{n-1} c_i^{\frac{1}{1-r}} \log(c_i) \left[ 1 + \sum_{i=2}^k c_i^{\frac{1}{1-r}} \right]$$

$$= \frac{V_1(r)}{B(1-r)^2} \sum_{i=2}^k c_i^{\frac{1}{1-r}} \log(c_i) \left[ B - f_k(r) \right] - \frac{V_1(r) f_k(r)}{B(1-r)^2} \sum_{i=k+1}^{n-1} c_i^{\frac{1}{1-r}} \log(c_i)$$

$$= \frac{V_1(r)}{B(1-r)^2} \left( B \sum_{i=2}^k c_i^{\frac{1}{1-r}} \log(c_i) - f_k(r) \sum_{i=2}^{n-1} c_i^{\frac{1}{1-r}} \log(c_i) \right)$$

To show that the term inside the bracket is positive, we basically need to show that for any decreasing sequence  $1 \ge d_1 > d_2 > \dots d_n > 0$ , we have that

$$h(k) = \sum_{i=1}^{n} d_i \sum_{i=1}^{k} d_i \log(d_i) - \sum_{i=1}^{k} d_i \sum_{i=1}^{n} d_i \log(d_i) \ge 0$$

for any  $k \in [n]$ 

Observe that

$$\Delta(k) = h(k+1) - h(k)$$

$$= d_{k+1} \log(d_{k+1}) \sum_{i=1}^{n} d_i - d_{k+1} \sum_{i=1}^{n} d_i \log(d_i)$$

$$= d_{k+1} \left( \log(d_{k+1}) \sum_{i=1}^{n} d_i - \sum_{i=1}^{n} d_i \log(d_i) \right)$$

Since  $d_k$  is a decreasing sequence, it follows that if  $\Delta(k) < 0$ , then  $\Delta(j) < 0$  for all j > k. But observe that h(n) = 0. So we just need to show that h(1) > 0 which is obvious.