Gradient Boosting: Residuals, Newton Leaf Values, Regression and Classification

ES 335 / ES 678 Machine Learning, Sem I 2026 Sources: `sources/course-2026/lectures/L12_student_trees_bagging_boosting_worked_deck.pdf` (Part 3, slides 42–64), `sources/course-2026/lectures/L12a_student_gradient_boosting_regression_notes.pdf` (10-page notes), `sources/course-2026/lectures/L12b_student_gradient_boosting_classification_notes.pdf` (14-page notes) · Previous: Bagging and random forests · Next: AdaBoost

Lecture 12 files: L12a Trees → L12b Bagging & random forests → **L12c Gradient boosting** → L12d AdaBoost → L12e XGBoost & comparison

> **How to use this file.** Bagging (L12b) grew independent deep trees and averaged them to kill **variance**. Boosting goes the other way: it adds up many **tiny** trees (stumps: one question, two leaves), each one grown on **what the trees so far still get wrong**, to kill **bias**. This file builds gradient boosting twice with the *same recipe*:

1. **Regression** (squared error, eight points) — the cleanest case: the "mistakes" are plain residuals $y - F$, and the leaf value is their average.
2. **Classification** (log loss, the twelve runs) — scores, sigmoid, residual $y - p$, and a leaf value $\sum r / \sum p(1-p)$ that is one **Newton step**.

The key idea in one line: **the residual is the negative gradient of the loss with respect to the prediction**, so fitting a tree to residuals is gradient descent in "prediction space". Part A is maths step by step; Part B walks slides 42–64; Part B′ covers the two sets of notes; Part C is practice.

The two datasets.


- *Regression* (notes L12a): minutes per epoch y vs model size x (millions of parameters).

i	1	2	3	4	5	6	7	8
x	1	2	3	4	5	6	7	8
y	2	3	3	5	6	8	8	9

- *Classification*: the twelve runs of L12a ($y = 1$ for D, 0 for C), new runs **A** (0.05, 32, no) and **B** (0.2, 64, yes). Eight candidate questions: rate ≤ 0.0055/0.02/0.065/0.2, batch ≤ 48/96/192, warm-up = no.

What this file is really about:

$$F_0 \xrightarrow{r=-\partial L/\partial F} \text{stump groups the residuals} \xrightarrow{\gamma=-\sum g/\sum h} F \leftarrow F+\nu\gamma \xrightarrow{\text{repeat}} F_T(x)=F_0+\nu\sum_t$$



Boosting in one picture: each learner is built on the previous learner's mistakes, then all are combined. (This picture shows the AdaBoost flavour — mistakes passed on as weights — which is L12d; gradient boosting passes them on as residuals.) (Wikimedia Commons: Ensemble_Boosting.svg, Sirakorn, CC BY-SA 4.0)

Step-up path

You need. Regression trees, and the idea of a gradient.

1. Start at the average.
2. Fit a small tree to what is still wrong, and add a fraction of it.
3. Recognise the residual as the negative gradient of the loss.
4. Move to classification, with scores in log-odds and a Newton step in each leaf.
5. Control overfitting with step size and stopping.

You can stop when you can run two rounds by hand, for regression and for classification.

Contents

Part A — Maths from the ground up

1. Bagging vs boosting: the same data used differently

2. Regression, step 0: start at the average
3. Regression, round 1: question, leaf values, small step
4. Why the split score is $(\Sigma r)^2/n$
5. Regression, rounds 2–8 and prediction
6. Classification: score, probability, loss, residual
7. The residual is the negative gradient
8. The leaf value: one Newton step
9. Classification, round 1 and round 2 in full
0. All eight rounds; predicting runs A and B
1. Step size, stopping and overfitting
2. The recipe and common confusions


Part B — Slides 42–64 · Jump · **Part B′ — The two notes** · Jump · **Part C — Practice** · Jump

Part A Maths From the Ground Up

A1 Bagging vs boosting: the same data used differently

Step 1 — Bagging: independent trees, each on its own resample, averaged. Tree 2 doesn't care about tree 1.

Step 2 — Boosting: tree 2 **cannot exist** before tree 1 — it is grown on tree 1's mistakes; tree 3 on the mistakes still left; and so on. The trees are **added up**.



Slide 44 — Bagging in parallel, boosting in sequence.

Step 3 — Why tiny trees? A stump (one question, two leaves) is **too simple** alone (high bias, low variance). Their *sum* is not: each new stump adds a new step (see the noise-free example in L12b A12). Boosting mainly reduces **bias**.

	bagging (L12b)	boosting (this file)
tree 2 is grown on	a resample; tree 1 plays no part	all twelve runs, fitted to the errors tree 1 left
tree 2 looks like	deep, leaves pure	one question, two leaves
trees depend on each other	no	yes
numbers combined by	averaging (run 5: 0, 1, 1 → 0.67)	adding (run 5: −0.50 + 0.63 = +0.13, ...)
each new tree reduces	the variance of the average	the bias: the remaining errors

A2 Regression, step 0: start at the average

Step 1 — Words. Prediction $F(x_i)$; residual $r_i = y_i - F(x_i)$ (positive ⇒ predicted too low); loss $\frac{1}{2}\sum_i (y_i - F(x_i))^2$ (wrong by 2 costs four times wrong by 1).

Step 2 — Best single constant c. $S(c) = \frac{1}{2}\sum(y_i - c)^2$; $S'(c) = -\sum(y_i - c) = 0 \Rightarrow c = \frac{1}{N}\sum y_i$ ($S'' = N > 0$, a minimum). So

$$F_0 = \frac{2+3+3+5+6+8+8+9}{8} = \frac{44}{8} = 5.5.$$

Step 3 — Residuals of the start:

i	1	2	3	4	5	6	7	8
$r = y - 5.5$	−3.5	−2.5	−2.5	−0.5	+0.5	+2.5	+2.5	+3.5
r^2	12.25	6.25	6.25	0.25	0.25	6.25	6.25	12.25

Sum of squares **50.00**. Small models predicted too high, large ones too low. The next step predicts *these residuals*. (They add to 0 — always true for residuals of the mean.)

A3 Regression, round 1: question, leaf values, small step

Step 1 — The helper is a stump that predicts the **residuals** (not y): one question "$x \le t$?", each leaf holding one number.

Step 2 — Try every threshold $t = 1.5, 2.5, \ldots, 7.5$ and score

$$\text{score} = \frac{(\sum_L r)^2}{n_L} + \frac{(\sum_R r)^2}{n_R}.$$

A good question puts positive residuals on one side and negative on the other so each side's sum is big (not cancelling).

question	n_L	$\sum_L r$	$\bar{r}_L$	n_R	$\sum_R r$	$\bar{r}_R$	score
x ≤ 1.5	1	−3.50	−3.500	7	+3.50	+0.500	14.00
x ≤ 2.5	2	−6.00	−3.000	6	+6.00	+1.000	24.00
x ≤ 3.5	3	−8.50	−2.833	5	+8.50	+1.700	38.53
x ≤ 4.5	4	−9.00	−2.250	4	+9.00	+2.250	**40.50**
x ≤ 5.5	5	−8.50	−1.700	3	+8.50	+2.833	38.53
x ≤ 6.5	6	−6.00	−1.000	2	+6.00	+3.000	24.00
x ≤ 7.5	7	−3.50	−0.500	1	+3.50	+3.500	14.00

Your turn 1: x ≤ 2.5 ⇒ $36/2 + 36/6 = 18 + 6 = 24 < 40.5$.

Step 3 — Leaf values = average residual of the leaf. Add the same γ to every point in a leaf; leaf loss $T(\gamma) = \sum_{\text{leaf}} \frac{1}{2}(r_i - \gamma)^2$; $T'(\gamma) = -\sum(r_i - \gamma) = 0 \Rightarrow \gamma = \bar{r}_{\text{leaf}}$ ($T'' = n > 0$). So

$$\gamma_L = -9/4 = -2.25 \text{ (points 1–4)}, \qquad \gamma_R = +9/4 = +2.25 \text{ (points 5–8)}.$$

In slope/curvature language: $g = dL/dF = -(y - F) = -r$, $h = d^2L/dF^2 = 1$, so $\gamma = -\sum g / \sum h = \sum r/n$. Because squared error is **exactly a parabola** in γ, no approximation was needed (classification will need one: A8).

Step 4 — Add only a fraction $\nu = 0.5$ (the **step size** / shrinkage):

$$F_1 = 5.5 + 0.5(-2.25) = 4.375 \ (x \le 4.5), \qquad F_1 = 5.5 + 0.5(2.25) = 6.625 \ (x > 4.5).$$

Why only half? One stump is crude; trusting it fully means the next tree starts from a worse place if this one was fooled by noise. *Your turn 2:* point 6 after round 1: 6.625 with ν = 0.5; 7.75 with ν = 1 (truth 8). The full step is closer *now*, but after round 3 with ν = 0.5 point 6 is at 7.735 — and small steps protect against one misled stump.

A4 Why the split score is (Σr)²/n

Step 1 — One leaf at its mean. $\sum(r_i - \bar{r})^2 = \sum r_i^2 - 2\bar{r}\sum r_i + n\bar{r}^2 = \sum r_i^2 - n\bar{r}^2$.

Step 2 — Before vs after the split (full step). Before: one leaf at the overall mean $\bar{r} = 0 \Rightarrow$ squared error $\sum r_i^2$. After: both leaves at their means $\Rightarrow \sum r_i^2 - n_L \bar{r}_L^2 - n_R \bar{r}_R^2$.

Step 3 — The fall is

$$n_L \bar{r}_L^2 + n_R \bar{r}_R^2 = \frac{(\sum_L r)^2}{n_L} + \frac{(\sum_R r)^2}{n_R}.$$

In general (if the overall mean residual is not 0) subtract $n\bar{r}^2$, which is the **same for every question** $\Rightarrow$ maximise the score. Round 1: $4(2.25)^2 + 4(2.25)^2 = 40.5$, and $50 - 40.5 = 9.5$ is the squared error after a full step.

A5 Regression, rounds 2–8 and prediction


Step 1 — Same recipe on new residuals (after round 1):

i	1	2	3	4	5	6	7	8
F_1	4.375	4.375	4.375	4.375	6.625	6.625	6.625	6.625
r	−2.375	−1.375	−1.375	+0.625	−0.625	+1.375	+1.375	+2.375


Sum of r^2: 19.625. Best question now **x ≤ 3.5** (score 14.01): $\gamma_L = (-2.375 - 1.375 - 1.375)/3 = -1.708$, $\gamma_R = (0.625 - 0.625 + 1.375 + 1.375 + 2.375)/5 = 1.025$. *Your turn 3:* $F_2(5) = 6.625 + 0.5(1.025) = 7.138$.

Step 2 — Eight rounds:

round	question	γ_L	γ_R	sum of r^2 after
0	start			50.000
1	x ≤ 4.5	−2.250	+2.250	19.625
2	x ≤ 3.5	−1.708	+1.025	9.119
3	x ≤ 5.5	−0.718	+1.196	3.971
4	x ≤ 7.5	−0.181	+1.265	2.600
5	x ≤ 1.5	−1.072	+0.153	1.615
6	x ≤ 5.5	−0.222	+0.371	1.120
7	x ≤ 1.5	−0.425	+0.061	0.966
8	x ≤ 7.5	−0.049	+0.340	0.867



Each round adds a small step where the staircase is still wrong. Vertical bars are the residuals (blue: predicted too low; orange: too high).



Left: the round-1 decision. Right: big errors are fixed early, then slowly.

Step 3 — The finished model and a new point. $F(x) = F_0 + \nu \sum_k \gamma_k(x)$. At $x = 2.5$ (not in the data):

round	question	2.5 answers	adds $\nu\gamma$	running
0	start			5.500
1	x ≤ 4.5	yes	−1.125	4.375
2	x ≤ 3.5	yes	−0.854	3.521
3	x ≤ 5.5	yes	−0.359	3.162
4	x ≤ 7.5	yes	−0.090	3.072

After eight rounds: **3.043** minutes — both neighbours have $y = 3$.

A6 Classification: score, probability, loss, residual

Step 1 — Score F. Any real number, the **log-odds of D**. Every run starts at $F = 0$. (Why 0? It's the log-odds of the class rate: $\ln \frac{6}{6} = 0$. With 9 of 12 D it would be $\ln \frac{9}{3} = 1.10$.)

Step 2 — Probability $p = S(F) = \dfrac{1}{1 + e^{-F}}$. $S(0) = 0.5$ ("I don't know"), positive ⇒ $p > 0.5$, larger $|F|$ ⇒ more confident but never exactly 0 or 1. Also $S(-F) = 1 - S(F)$.


F	−4	−3	−2	−1	−0.5	0	+1	+2	+3	+4
p	0.018	0.047	0.119	0.269	0.378	0.500	0.731	0.881	0.953	0.982

Step 3 — Log loss $L = -y \ln p - (1 - y) \ln(1 - p)$ = minus the log of the probability given to **what actually happened**:

probability given to the truth	0.99	0.90	0.50	0.10	0.01
loss	0.010	0.105	0.693	2.303	4.605

Start: every run at $p = 0.5$ ⇒ average loss $\ln 2 = 0.693$.

Step 4 — Residual $r = y - p$. D at $p = 0.5$: $r = +0.5$ (push up); C at $p = 0.5$: $r = -0.5$ (push down); confidently right ⇒ $r \approx 0$. Example: run 5 after round 1: $F = -0.5$, $p = 0.378$, $y = 1$ ⇒ $r = 0.622$.



Step 1: score → probability. Step 2: confident mistakes are expensive. Step 3: the slope of run 5's loss is $p - y = -0.622$; moving F in the direction of $r = +0.622$ goes **downhill**.

A7 The residual is the negative gradient

Step 1 — Two small facts. $\dfrac{dp}{dF} = S'(F) = p(1-p)$ and $\dfrac{dL}{dp} = -\dfrac{y}{p} + \dfrac{1-y}{1-p}$. (Proof of the first: $S = (1 + e^{-F})^{-1} \Rightarrow S' = \dfrac{e^{-F}}{(1+e^{-F})^2} = \dfrac{1}{1+e^{-F}} \cdot \dfrac{e^{-F}}{1+e^{-F}} = S(1-S)$.)

Step 2 — Chain rule:

$$\frac{dL}{dF} = \left[-\frac{y}{p} + \frac{1-y}{1-p} \right] p(1-p) = -y(1-p) + (1-y)p = p - y = -r.$$

Step 3 — Read it. The residual is the **negative gradient** of the loss with respect to the score. Moving each score along its residual lowers the loss: **gradient descent on the log loss**, where the "parameters" are the twelve scores. (For squared error the same thing: $dL/dF = -(y - F) = -r$.) This is the "gradient" in **gradient boosting**.

Step 4 — But a stump cannot move the scores freely. It can only add **one number to each of two groups**. So it chooses the question whose two groups have the most different average residual (the best two-number correction; same score as A4), and then asks: *how much* to add to each group? That is A8.

A8 The leaf value: one Newton step

Step 1 — Try it by brute force. Group A = runs 5–12 (2 D, 6 C), all at $F = 0$. Add the same γ to all eight scores:

γ	0	−0.5	−1.0	−1.0986	−1.5	−2.0	+0.5
total loss of the 8	5.545	4.793	4.506	**4.499**	4.611	5.015	6.793

Best ≈ −1: lowering helps the six C runs until the two D runs get hurt. Group B (runs 1–4, all D): loss keeps falling as γ grows (2.773 at 0, 0.073 at 4).

Step 2 — Why no exact formula? $T(\gamma) = \sum_i L(F_i + \gamma)$, with terms like $\ln(1 + e^{-F_i - \gamma})$ at different F_i; setting $T' = 0$ means solving a sum of different exponentials. No closed form.

Step 3 — The idea (Newton). Stand at $\gamma = 0$; feel the **slope** and the **curvature**; draw the parabola that hugs the curve there; jump to its bottom.

Step 4 — Slope and curvature per run. Slope $p - y$ (A7). Curvature $\frac{d}{dF}(p - y) = p(1 - p)$: always positive (a bowl), largest 0.25 at $p = 0.5$, ≈ 0 near 0 or 1.

Step 5 — The parabola. $L(F_i + \gamma) \approx L(F_i) + (p_i - y_i)\gamma + \frac{1}{2}p_i(1 - p_i)\gamma^2$. Sum over the leaf with $S = \sum(p_i - y_i)$, $H = \sum p_i(1 - p_i)$:


$$T(\gamma) \approx T(0) + S\gamma + \tfrac{1}{2}H\gamma^2.$$

Step 6 — Its bottom: $S + H\gamma = 0 \Rightarrow$

$$\boxed{\gamma = -\frac{S}{H} = \frac{\sum_{\text{leaf}}(y_i - p_i)}{\sum_{\text{leaf}} p_i(1 - p_i)} = \frac{\text{total push needed}}{\text{sensitivity}}}$$

Top: how much the group needs to move. Bottom: how strongly p reacts to the score (near 0.5 one unit of score moves p by ≈0.25; near 0 or 1, barely). A steep slope asks for a big move; a strongly curved bowl says don't overshoot.

Step 7 — Check on group A. $S = 6(0.5) + 2(-0.5) = 2$, $H = 8(0.25) = 2 \Rightarrow$ parabola $5.545 + 2\gamma + \gamma^2$, bottom $\gamma = -1$. True bottom: $-2\sigma(-\gamma) + 6\sigma(\gamma) = 0 \Rightarrow \sigma(\gamma) = \frac{1}{4} \Rightarrow e^{-\gamma} = 3 \Rightarrow \gamma = -\ln 3 = -1.099$. **One Newton step lands within 0.1.**



Left/middle: the parabola from slope and curvature, and where it lands. Right (slide 58): **the more wrong the leaf, the bigger the push** — for all-D leaves, $\gamma = \frac{n(1-p)}{np(1-p)} = 1/p$.

Step 8 — Regression is the special case $h = 1$: $\gamma = \sum r/n$ = the average residual (A3).

A9 Classification, round 1 and round 2 in full

One round in four steps (slide 53): (a) $p_i = S(F_i)$, $r_i = y_i - p_i$. (b) Score each question by $n_L \bar{r}_L^2 + n_R \bar{r}_R^2$; keep the largest. (c) Leaf values $\gamma = \sum r / \sum p(1-p)$. (d) $F_i \leftarrow F_i + \nu \gamma_{\text{leaf}(i)}$, $\nu = 0.5$.

Round 1. (a) All $F = 0 \Rightarrow p = 0.5$, $r = +0.5$ (D), −0.5 (C), $h = 0.25$. (b) All eight questions:

question	n_L	$\sum_L r$	n_R	$\sum_R r$	score
rate ≤ 0.065	8	−2.00	4	+2.00	**1.500**
rate ≤ 0.02	5	−1.50	7	+1.50	0.771
rate ≤ 0.0055	2	−1.00	10	+1.00	0.600
rate ≤ 0.2	10	−1.00	2	+1.00	0.600
batch ≤ 48	4	+1.00	8	−1.00	0.375
warm-up = no	6	+1.00	6	−1.00	0.333
batch ≤ 192	9	+0.50	3	−0.50	0.111
batch ≤ 96	7	+0.50	5	−0.50	0.086

(c) $\gamma_{\text{yes}} = \dfrac{-2}{8 \times 0.25} = -1.000$ (runs 5–12), $\gamma_{\text{no}} = \dfrac{+2}{4 \times 0.25} = +2.000$ (runs 1–4). (d) Runs 5–12: $F = -0.5$, $p = 0.378$; runs 1–4: $F = +1.0$, $p = 0.731$. Loss **0.693 → 0.504**.

Your turn 4 (notes): if batch ≤ 48 had been used: yes {2, 5, 6, 10}: $\sum r = 1$, $\sum h = 1$ ⇒ γ = 1.000; no: $\sum r = -1$, $\sum h = 2$ ⇒ γ = −0.500.

What tree 1 gets wrong:

run	out	F	p	says	$r = y - p$	
1–4	D	+1.00	0.731	D	+0.269	right but unsure
5, 6	D	−0.50	0.378	C	**+0.622**	**wrong**
7–12	C	−0.50	0.378	C	−0.378	right

Runs 5 and 6 (diverged at a low rate) now have the **largest** residual — up from 0.5. That is what tree 2 will be drawn to. (*Your turn 5:* run 5's |r| grew 0.5 → 0.622; run 7's shrank 0.5 → 0.378.)

Round 2. Scores: **batch ≤ 48 0.469**; rate ≤ 0.065 0.419; rate ≤ 0.0055 0.351; warm-up 0.334. Winner **batch ≤ 48**: yes = runs 2, 5, 6, 10.

- Yes: $\sum r = 0.269 + 0.622 + 0.622 - 0.378 = 1.136$; $\sum h = 0.731 \times 0.269 + 3 \times 0.378 \times 0.622 = 0.902$ ⇒ $\gamma = +1.260$.
- No (8 runs): $\sum r = -1.081$, $\sum h = 1.765$ ⇒ $\gamma = -0.612$.

Tree 2 changes the numbers ($\nu\gamma = +0.63$ / -0.31):


run	out	after tree 1	tree 2 adds	now	p	says	
1, 3, 4	D	+1.00	−0.31	+0.69	0.667	D	
2	D	+1.00	+0.63	+1.63	0.836	D	
5, 6	D	−0.50	+0.63	+0.13	0.532	D	**fixed**
7–9, 11, 12	C	−0.50	−0.31	−0.81	0.309	C	
10	C	−0.50	+0.63	+0.13	0.532	D	**now wrong**

Run 10 shares the leaf with 5 and 6 and is dragged up with them: mistakes go 2 → 1. Tree 3 is grown on the new errors.

A10 All eight rounds; predicting runs A and B


round	question	γ yes	γ no	log loss	wrong
0	(all scores 0)			0.693	6
1	rate ≤ 0.065	−1.000	+2.000	0.504	2 (5, 6)
2	batch ≤ 48	+1.260	−0.612	0.438	1 (10)
3	rate ≤ 0.065	−0.629	+1.448	0.361	2 (5, 6)
4	warm-up = no	+0.800	−0.717	0.321	0
5	rate ≤ 0.065	−0.423	+1.279	0.283	0
6	batch ≤ 48	+0.949	−0.568	0.250	0
7	rate ≤ 0.02	−0.597	+0.577	0.230	0
8	batch ≤ 48	+0.603	−0.426	0.217	0

- **The same question can come back** (rounds 1, 3, 5): the scores changed, so it's a different problem.
- **Loss falls every round; the wrong count does not** (6, 2, 1, 2, 0): each round lowers the *loss*, not the error count.
- The single tree's three questions (rate, batch, warm-up) appear as rounds 1, 2, 4.
- From round 4 nothing is wrong, yet loss keeps falling: probabilities move towards 0 and 1.



Left: steady loss decrease. Right: the story of runs 5, 6 (fixed in round 2, broken in round 3, fixed for good in round 4) and run 10 (collateral damage, then repaired by warm-up in round 4).

Predicting. $F(x) = \nu \sum_t \gamma_t(\text{leaf of } x)$, $p = S(F)$ (starts from 0, the log-odds of 6/12). Each entry below is $\nu\gamma$ of the leaf reached:



Slide 60 as a picture. Red pushes towards D, blue towards C. Rows sum to F_8.


- **Run A** (0.05, 32, no): −0.500 +0.630 −0.315 +0.400 −0.212 +0.474 +0.289 +0.302 ⇒ $F_8 = 1.069$, $p = \mathbf{0.744}$ ⇒ D. (A lands exactly like runs 5 and 6.)
- **Run B** (0.2, 64, yes): +1.000 −0.306 +0.724 −0.358 +0.639 −0.284 +0.289 −0.213 ⇒ $F_8 = 1.490$, $p = \mathbf{0.816}$ ⇒ D. All three rate stumps push strongly towards D; batch and warm-up push back a little.

Run 5's story in numbers: probability of D 0.5 → 0.378 (tree 1) → … → **0.74** after eight trees. Bagging gave it three independent opinions (0, 1, 1); boosting *corrects* it step by step.

A11 Step size, stopping and overfitting

Step 1 — Smaller ν = trust each stump less.

ν	first round with no wrong run	loss after 8 rounds	rounds to reach loss 0.217
1.0	4	0.087	4 (loss 0.196)
0.5	4	0.217	8
0.1	22	0.473	44 (loss 0.215)



More rounds, each a safer correction. The training loss keeps falling forever — **training data can never tell you when to stop**.

Step 2 — Boosting can overfit. Simulated (200 runs, depth-3 trees, ν = 0.1): training error 0 by round 60; held-out error best near round **66 (0.108)**, rising to **0.137 by round 1000**. ⇒ **Stop on held-out error** (early stopping). Contrast bagging, where more trees never hurt.

A12 The recipe and common confusions

step	what you do	which number is used
start	$F_i = F_0$ (mean of y for regression; log-odds, here 0, for classification)	—
(a) residuals	$r_i = y_i - F_i$ (reg.) or $y_i - p_i$ (class.)	each run's current prediction
(b) question	largest $n_L \bar{r}_L^2 + n_R \bar{r}_R^2$	residuals of the two groups
(c) leaf values	$\gamma = \sum r / \sum h$ ($h = 1$ or $p(1-p)$)	$\gamma_{\text{yes}}, \gamma_{\text{no}}$
(d) update	$F_i \leftarrow F_i + \nu\gamma$	νγ of the leaf reached
predict	$F = F_0 + \nu \sum_t \gamma_t$; $p = S(F)$	one number per stump
stop	when held-out error stops falling	—

Common confusions.


- *Why fit residuals and not y again?* A tree fitted to y repeats the same staircase. Residuals are exactly what the model doesn't know yet.
- *Why "gradient"?* The residual is the negative slope of the loss w.r.t. the prediction (A7). Adding a fitted residual is a gradient-descent step.
- *Why not just add the average residual in classification?* Probabilities saturate: a run at $p = 0.95$ needs a big score change to move a little. Dividing by $\sum p(1-p)$ accounts for that.
- *Why a regression tree when classifying?* The tree only groups runs by what they still need; residuals and leaf values are numbers. The class appears only at the end, through S.
- *Why log loss?* It punishes confident mistakes heavily, and its slope and curvature are the simple $p - y$ and $p(1-p)$.
- *How does it relate to the others?* **AdaBoost** (L12d) = the same stagewise idea with the exponential loss e^{-yF}; its weights are each run's share of that loss. **XGBoost** (L12e) = this file plus curvature in the split choice and penalties λ, γ.

Part B Slide by Slide (slides 42–64)

Slide 42 Part 3: Gradient boosting (divider)

Colours: rate stump, batch-size stump, warm-up stump.

Slide 43 Boosting: the procedure




Slide 43 — A stump has one question and two leaves. Boosting mainly reduces **bias**.

Slide 44 Boosting and bagging: the same twelve runs, used differently

Shown in A1.

Slide 45 Boosting, step 1: tree 1 is one question

Boosting, step 1: tree 1 is one question



Boosting uses many **small** trees. Tree 1 is the first question of the tree in Part 1, rate ≤ 0.065, with two leaves.

Each leaf gives its runs a **score**: positive leans towards **D** (diverged), negative towards **C** (converged), 0 means undecided.

Before tree 1 every run has score 0. After it, runs 5–12 have score -0.50 and runs 1–4 have $+1.00$.

Where -0.50 and $+1.00$ come from is worked out after this example.

Slide 45 — Positive leans D, negative leans C, 0 undecided. Where −0.50 and +1.00 come from: A9.

Slide 46 What tree 1 gets wrong

What tree 1 gets wrong

run	out	score	p	tree 1 says	error $y - p$	result
1	**D**	+1.00	0.731	**D**	+0.269	right
2	**D**	+1.00	0.731	**D**	+0.269	right
3	**D**	+1.00	0.731	**D**	+0.269	right
4	**D**	+1.00	0.731	**D**	+0.269	right
5	**D**	−0.50	0.378	**C**	+0.622	**wrong**
6	**D**	−0.50	0.378	**C**	+0.622	**wrong**
7	**C**	−0.50	0.378	**C**	−0.378	right
8	**C**	−0.50	0.378	**C**	−0.378	right
9	**C**	−0.50	0.378	**C**	−0.378	right
10	**C**	−0.50	0.378	**C**	−0.378	right
11	**C**	−0.50	0.378	**C**	−0.378	right
12	**C**	−0.50	0.378	**C**	−0.378	right

p turns a score into a probability of **D**; $y = 1$ for **D**, 0 for **C**. Tree 1 makes **2 mistakes in 12** (runs 5 and 6: diverged at a low rate). The error column is how far each prediction is from the outcome: runs 5, 6 are furthest off (+0.62).

Slide 46 — 2 mistakes in 12 (runs 5, 6: diverged at a low rate); they are furthest off.


Slide 47 Boosting, step 2: tree 2 is grown on the mistakes

Boosting, step 2: tree 2 is grown on the mistakes

Errors left by tree 1 ($y - p$):

runs	error
5, 6 (**D**, tree 1 says **C**)	+0.62
1–4 (**D**, right but unsure)	+0.27
7–12 (**C**, right)	−0.38

Tree 2 is not fitted to the outcomes again, but to **these errors**. It asks the question that separates large errors from small ones: batch ≤ 48 puts runs 2, 5, 6, 10 together (average error +0.28) against the other eight (average −0.14). That includes the two worst runs, 5 and 6.



Tree 2's scores push its group towards **D** (+0.63) or towards **C** (−0.31). They are **added** to the scores from tree 1.

Slide 47 — Fitted to the **errors**, not the outcomes. Average error +0.28 vs −0.14 in the two groups.

Slide 48 Tree 2 changes the numbers

Tree 2 changes the numbers

run	out	score, tree 1	tree 2 adds	score now	p	says	change
1	**D**	+1.00	−0.31	+0.69	0.667	**D**	
2	**D**	+1.00	+0.63	+1.63	0.836	**D**	
3	**D**	+1.00	−0.31	+0.69	0.667	**D**	
4	**D**	+1.00	−0.31	+0.69	0.667	**D**	
5	**D**	−0.50	+0.63	+0.13	0.532	**D**	fixed
6	**D**	−0.50	+0.63	+0.13	0.532	**D**	fixed
7	**C**	−0.50	−0.31	−0.81	0.309	**C**	
8	**C**	−0.50	−0.31	−0.81	0.309	**C**	
9	**C**	−0.50	−0.31	−0.81	0.309	**C**	
10	**C**	−0.50	+0.63	+0.13	0.532	**D**	**now wrong**
11	**C**	−0.50	−0.31	−0.81	0.309	**C**	
12	**C**	−0.50	−0.31	−0.81	0.309	**C**	

New score = score after tree 1 + what tree 2 adds. Runs 5 and 6 move from −0.50 to +0.13 ($p = 0.532$): now correct. Run 10 shares their leaf, is pushed up with them, and becomes a mistake: mistakes go from 2 to 1. Tree 3 is grown on the new errors.

Slide 48 — Mistakes 2 → 1 (A9).

Slide 49 How tree 2 differs from bagging

How tree 2 differs from bagging

	bagging (Part 2)	boosting (this part)
tree 2 is grown on	a resample of the twelve runs; tree 1 plays no part	all twelve runs, fitted to the errors tree 1 left
tree 2 looks like	deep: grown until the leaves are pure	one question and two leaves
trees depend on each other	no: any order, or all at once	yes: tree 2 needs tree 1 first
numbers are combined by	averaging. Run 5: the three trees give 0 1 1, average 0.67	adding. Run 5: score −0.50 (tree 1) +0.63 (tree 2) = +0.13, and so on
each new tree reduces	the variance of the average	the bias: the remaining errors

Run 5 diverged. Bagging averages three independent opinions on it; boosting corrects the score step by step: after eight trees its probability of **D** is 0.74 (started at 0.5, 0.378 after tree 1).

Slide 49 — After eight trees run 5's probability of D is 0.74 (started 0.5, 0.378 after tree 1).

Slide 50 The math behind the scores: definitions

Score F (log-odds of D, starts 0); $p = S(F) = 1/(1+e^{-F})$ ($S(0) = 0.5$, $S(1) = 0.731$, $S(-0.5) = 0.378$); $y \in \{0, 1\}$; residual $r = y - p$; log loss $L = -y \ln p - (1-y)\ln(1-p)$. Run 5 after tree 1: $r = 0.622$ (A6).

Slide 51 Why the next tree is fitted to the residuals

Why the next tree is fitted to the residuals

Treat each run's score as a number we can change. How does one run's loss change with its own score? Using $S'(F) = p(1-p)$,


$$\frac{dL}{dF} = \Big(-\frac{y}{p} + \frac{1-y}{1-p}\Big)p(1-p) = -y(1-p) + (1-y)p = p - y = -r.$$

So the residual is the *negative gradient* of the loss: moving each score in the direction of its residual lowers the loss, which is gradient descent on the log loss.

A stump cannot move all twelve scores freely. It can only add one number to each of two groups. Tree 2 chose the question whose two groups have the most different average residual, because that is the best two-number correction.

Slide 51 — Gradient descent on the log loss with the scores as variables (A7).

Slide 52 The value assigned to a leaf



Slide 52 — Round 1 leaves: γ = −1.00 and +2.00, so scores move by −0.50 and +1.00 (A8).

Slide 53 One round of boosting, in four steps

(a) p, r; (b) score $n_L\bar{r}_L^2 + n_R\bar{r}_R^2$; (c) $\gamma = \sum r / \sum p(1-p)$; (d) $F \leftarrow F + \nu\gamma$ (A9).

Slide 54 Round 1: choose the question

Scores 1.500 (rate ≤ 0.065), 0.771, 0.600. $\gamma_{\text{yes}} = -2/(8 \times 0.25) = -1$, $\gamma_{\text{no}} = +2/(4 \times 0.25) = +2$.

Slide 55 Round 1: update every run

Round 1: update every run

run	out	p before	r	leaf	γ	$\nu\gamma$	F after	p after
1	D	0.500	+0.500	no	+2.000	+1.000	+1.000	0.731
2	D	0.500	+0.500	no	+2.000	+1.000	+1.000	0.731
3	D	0.500	+0.500	no	+2.000	+1.000	+1.000	0.731
4	D	0.500	+0.500	no	+2.000	+1.000	+1.000	0.731
5	D	0.500	+0.500	yes	-1.000	-0.500	-0.500	0.378
6	D	0.500	+0.500	yes	-1.000	-0.500	-0.500	0.378
7	C	0.500	-0.500	yes	-1.000	-0.500	-0.500	0.378
8	C	0.500	-0.500	yes	-1.000	-0.500	-0.500	0.378
9	C	0.500	-0.500	yes	-1.000	-0.500	-0.500	0.378
10	C	0.500	-0.500	yes	-1.000	-0.500	-0.500	0.378
11	C	0.500	-0.500	yes	-1.000	-0.500	-0.500	0.378
12	C	0.500	-0.500	yes	-1.000	-0.500	-0.500	0.378

Shaded: the two numbers read from this stump. Log loss 0.693 $\rightarrow$ 0.504. Runs 5, 6 still wrong ($p = 0.378$).

Slide 55 — Runs 5, 6 still wrong (p = 0.378).

Slide 56 Round 2: choose the question

Round 2: choose the question

Residuals now: runs 5, 6: 0.622 (most wrong); runs 1–4: 0.269; runs 7–12: -0.378. Scores: batch ≤ 48 0.469; rate ≤ 0.065 0.419; rate ≤ 0.0055 0.351; warm-up 0.334.

Winner **batch ≤ 48**. Yes: runs 2, 5, 6, 10. No: the other 8.

$$\gamma_{\text{yes}} = \frac{1.136}{0.902} = +1.260, \qquad \gamma_{\text{no}} = \frac{-1.081}{1.765} = -0.612$$

Yes-group: residual sum $0.269 + 0.622 + 0.622 - 0.378$; curvature $0.731 \times 0.269 + 3 \times 0.378 \times 0.622$.

Slide 56: round two of boosting. Runs 5 and 6 now have the largest residuals, and the question batch ≤ 48 wins with the highest split score.

Slides 57–58

Your turn: one leaf's update (and answer)

Answer: one leaf's update in boosting

	leaf $\{1,3,4\}$	leaf $\{5,6\}$
p	0.731	0.378
$r = y - p$	0.269 each	0.622 each
$p(1-p)$	0.197 each	0.235 each
$\gamma = \sum r / \sum p(1-p)$	$0.807/0.590 = \mathbf{1.368}$	$1.245/0.470 = \mathbf{2.649}$
$\nu\gamma$	$+0.684$	$+1.324$
new F	1.684	0.824
new p	0.843	0.695

4. Runs 5 and 6 get the larger step: they are the runs the model is most wrong about. For a leaf in which all runs have the same p and $y = 1$, $\gamma = 1/p$ exactly, so the smaller the probability given to the correct outcome, the bigger the push.


Slide 58 — For an all-D leaf with equal p, $\gamma = 1/p$: the less probability given to the truth, the bigger the push.

Slide 59

All eight rounds

All eight rounds

round	question	γ yes	γ no	loss	wrong
0	(all scores 0)			0.693	6
1	rate $\leq$ 0.065	-1.000	+2.000	0.504	2 (runs 5, 6)
2	batch $\leq$ 48	+1.260	-0.612	0.438	1 (runs 10)
3	rate $\leq$ 0.065	-0.629	+1.448	0.361	2 (runs 5, 6)
4	warm-up = no	+0.800	-0.717	0.321	0
5	rate $\leq$ 0.065	-0.423	+1.279	0.283	0
6	batch $\leq$ 48	+0.949	-0.568	0.250	0
7	rate $\leq$ 0.02	-0.597	+0.577	0.230	0
8	batch $\leq$ 48	+0.603	-0.426	0.217	0



Loss falls every round; wrong runs go $6, 2, 1, 2, 0$. The single tree's three questions (rate, batch, warm-up) are rounds 1, 2 and 4.

Slide 59 — Loss falls every round; wrong 6, 2, 1, 2, 0 (A10).

Slide 60 Contribution of each stump to each run's score

Contribution of each stump to each run's score


run	out	r1 rate	r2 batch	r3 rate	r4 warm	r5 rate	r6 batch	r7 rate	r8 batch	F_8	p
1	D	+1.000	-0.306	+0.724	-0.358	+0.639	-0.284	+0.289	-0.213	+1.490	0.816
2	D	+1.000	+0.630	+0.724	+0.400	+0.639	+0.474	+0.289	+0.302	+4.458	0.989
3	D	+1.000	-0.306	+0.724	-0.358	+0.639	-0.284	+0.289	-0.213	+1.490	0.816
4	D	+1.000	-0.306	+0.724	+0.400	+0.639	-0.284	+0.289	-0.213	+2.248	0.905
5	D	-0.500	+0.630	-0.315	+0.400	-0.212	+0.474	+0.289	+0.302	+1.069	0.744
6	D	-0.500	+0.630	-0.315	+0.400	-0.212	+0.474	-0.298	+0.302	+0.482	0.618
7	C	-0.500	-0.306	-0.315	+0.400	-0.212	-0.284	+0.289	-0.213	-1.141	0.242
8	C	-0.500	-0.306	-0.315	-0.358	-0.212	-0.284	+0.289	-0.213	-1.899	0.130
9	C	-0.500	-0.306	-0.315	-0.358	-0.212	-0.284	-0.298	-0.213	-2.486	0.077
10	C	-0.500	+0.630	-0.315	-0.358	-0.212	+0.474	-0.298	+0.302	-0.277	0.431
11	C	-0.500	-0.306	-0.315	+0.400	-0.212	-0.284	-0.298	-0.213	-1.728	0.151
12	C	-0.500	-0.306	-0.315	-0.358	-0.212	-0.284	-0.298	-0.213	-2.486	0.077
A	?	-0.500	+0.630	-0.315	+0.400	-0.212	+0.474	+0.289	+0.302	+1.069	0.744
B	?	+1.000	-0.306	+0.724	-0.358	+0.639	-0.284	+0.289	-0.213	+1.490	0.816

Entry $= \nu\gamma$ of the leaf the run reaches in that round's stump. F_8 is the row sum, $p = S(F_8)$. Runs A, B: $F_8 = 1.069$, 1.490; $p = 0.744$, 0.816.

Slide 60 — A: $F_8 = 1.069$, $p = 0.744$; B: $F_8 = 1.490$, $p = 0.816$.

Slides 61–62 Runs A and B: the leaf reached in each stump

Run A: the leaf reached in each stump



Yellow leaf = where A ends; its number is the entry in the table of contributions. The eight numbers add to $F_8 = 1.069$.

Slide 61 — The eight yellow leaves add to $F_8 = 1.069$. Slide 62 does the same for B (1.490).

Slide 63 Step size and stopping

Step size and stopping

step size ν	first round with no wrong run	loss after 8 rounds	rounds to reach loss 0.217
1.0	4	0.087	4 (loss 0.196)
0.5	4	0.217	8
0.1	22	0.473	44 (loss 0.215)

A smaller step trusts each stump less: more rounds, each a safer correction. The loss keeps falling even after training error is 0: **training data can never tell us when to stop**.

Boosting can overfit

Simulated (200 runs, depth-3 trees, $\nu = 0.1$): training error 0 by round 60, held-out error best near round 66 (0.108), then rising to 0.137 by round 1000. Stop on held-out error.

Slide 63 — Stop on held-out error (A11).

Slide 64 Boosting: summary of the procedure

Boosting: summary of the procedure


step	what you do	which number is used
start	$F_i = 0$, so $p_i = 0.5$	none
(a) residuals	$r_i = y_i - p_i$	each run's current p
(b) question	largest $n_L \bar{r}_L^2 + n_R \bar{r}_R^2$	the residuals of the two groups
(c) leaf values	$\gamma = \sum r / \sum p(1-p)$	γ yes leaf, γ no leaf
(d) update	$F_i \leftarrow F_i + \nu\gamma$	$\nu\gamma$ of the leaf reached
predict	$F = \nu \sum_t \gamma_t$(leaf in stump t), $p = S(F)$	one number per stump (the table of contributions)
stop	when held-out error stops falling	–

Slide 64: the whole gradient boosting procedure as a table, from residuals to the question, the leaf values and the update.

Part B′ — The Notes (L12a and L12b)


Notes L12a (regression) — summary and §1–3

Start with one number (the average), look at residuals, fit a stump to the residuals, add a fraction, repeat. Colour key: y truth, F prediction, r residual, γ leaf value, ν step size, g slope, h curvature. Full working: A2–A4.



Notes Fig. 2 — Round 1 turns the flat line into a two-step staircase.

Notes L12a, §4–5 — rounds 2 and 3; the finished model



Notes Fig. 4 — Big errors fixed early. Prediction at $x = 2.5$: 3.043 after eight rounds (A5).

Notes L12a, §6 — derive-it-yourself boxes

D1: $dL/dF = -r$ ⇒ residual is the negative slope; if $r > 0$, raise F. D2: leaf value = average residual. D3: fall in squared error = $n_L \bar{r}_L^2 + n_R \bar{r}_R^2$ (40.5 in round 1, leaving 9.5 with ν = 1). D4: $n\bar{r}^2$ is the same for every question ⇒ maximise the score. D5: $\gamma = -\sum g / \sum h$; with $g = p - y$, $h = p(1-p)$ it becomes the classification leaf value.


Notes L12a, §8 — check yourself (answers)

1 average, 5.5 · 2 $y - F$, low · 3 average · 4 n_L, n_R, largest · 5 4.5, −2.25, 2.25 · 6 6.625 · 7 19.625 · 8 40.5, 9.5 · 9 negative slope · 10 h, 1.

Notes L12b (classification) — §2–4: score, probability, loss, residual


A6. Your turn 1: $\sigma(1) = 1/1.368 = 0.731$, $\sigma(-1) = 0.269$, sum 1; $p = 0.9$ needs $F \approx 2.2$. Your turn 2: $p = 0.8$: loss 0.223 if D, 1.609 if C.

score F	$p = \sigma(F)$
−4	0.018
−3	0.047
−2	0.119
−1	0.269
0	0.500
+1	0.731
+2	0.881
+3	0.953
+4	0.982



Notes Fig. 1 — The sigmoid turns any score into a probability.

Notes L12b, §5 — round 1 step by step, and where the formula comes from



Notes Fig. 2 — The parabola hugs the true curve near γ = 0; its bottom is almost the true bottom. Four sentences of intuition and Steps (i)–(iv): A8.

Notes L12b, §6–7 — rounds 2–8; new run B

Tables in A9–A10. Run B's running score: +1.000, +0.694, +1.418, +1.059, +1.699, +1.415, +1.703, **+1.490** ⇒ $p = 0.816$.

Notes L12b, §8 — derive-it-yourself boxes

D1: $\sigma' = \sigma(1-\sigma)$, $dL/dF = p - y$, $d^2L/dF^2 = p(1-p)$. D2: summing the quadratic ⇒ $\gamma = \sum(y - p)/\sum p(1-p)$. D3: score 1.5 for rate ≤ 0.065; mixed-sign groups cancel ⇒ score ≈ 0. D4: exact best γ for group A = $-\ln 3 = -1.099$. D5: squared error has $h = 1$ ⇒ average residual.

Notes L12b, §10 — check yourself (answers)

1 0.5, diverge · 2 what actually happened, $\ln 2 = 0.693$ · 3 $y - p$, +0.5, −0.5 · 4 0.065, 1.500 · 5 $p(1-p)$, push, sensitivity · 6 −2/2 = −1, 2/1 = 2 · 7 −0.5, 1.0, 0.378 · 8 grown, 0.622, 0.504 · 9 $p - y$, $p(1-p)$ · 10 −1.099, −1 · 11 1, average · 12 sum.

Part C Practice

1. Regression: compute the score of **x ≤ 3.5** in round 1 from the start residuals. Compare with 40.5.
2. Regression: what is $F_1(2)$ with ν = 0.5, and with ν = 1? Which round-2 leaf is point 2 in, and what is $F_2(2)$?
3. Show that the residuals of the mean always add to 0, and use it to explain why the round-1 sums on the two sides are opposite.
4. Classification: compute $\sigma(2)$ and the log loss of a C run with $F = 2$. Is it right or wrong, and how expensive?
5. Why can the number of wrong runs go **up** in a round (round 3: 1 → 2) while the loss goes down?
6. A leaf holds three C runs, each at $F = 1$. Compute r, h, γ, the update with ν = 0.5 and the new p.
7. Repeat the brute-force idea: for group B (four D runs at $F = 0$) what does the parabola predict, and why does the true loss have no minimum?
8. In round 1, why is $\gamma_{\text{no}} = +2$ but each score in that leaf only rises by 1?
9. You run boosting for 1000 rounds and training error has been 0 since round 60. What should you do, and why can't the training data tell you?
0. Compute F_8 for run A by adding the eight contributions, then p.

1. Yes: points 1–3, $\sum r = -8.5$; no: points 4–8, $+8.5$. Score $72.25/3 + 72.25/5 = 24.08 + 14.45 = 38.53$ < 40.5.
2. $F_1(2) = 4.375$ (ν = 0.5) or 3.25 (ν = 1). Round 2 asks x ≤ 3.5: point 2 is in the yes-leaf, $\gamma_L = -1.708 \Rightarrow F_2(2) = 4.375 - 0.854 = 3.521$.
3. $\sum(y_i - \bar{y}) = \sum y_i - N\bar{y} = 0$. So $\sum_L r = -\sum_R r$ for every split.
4. $\sigma(2) = 0.881$; a C run: loss $-\ln(1 - 0.881) = -\ln 0.119 = 2.13$. Wrong (says D) and confident ⇒ expensive.
5. Each round lowers the average log loss, not the error count. Round 3 (rate ≤ 0.065, $\gamma_{\text{yes}} = -0.629$) pulls all low-rate runs down; this fixes run 10 but pushes runs 5, 6 (barely above 0.5 at 0.532) back below. Overall the probabilities still moved towards the truth on average.
6. $p = 0.731$, $r = -0.731$ each; $h = 0.197$ each. $\gamma = -2.193/0.590 = -3.72$ (for an all-C leaf $\gamma = -1/(1-p)$). Update $-1.86 \Rightarrow F = -0.86$, $p = 0.297$.
7. $S = -2$, $H = 1 \Rightarrow$ parabola $2.773 - 2\gamma + \frac{1}{2}\gamma^2$, bottom at $\gamma = +2$. True loss $4\ln(1 + e^{-\gamma})$ decreases for every γ (all four want $p \to 1$), so it has no minimum — another reason for a step size and for penalties (XGBoost's λ, L12e).

8. γ is the full Newton step; the update multiplies by ν = 0.5.
9. Stop on held-out (validation) error — typically use the round where it was lowest (early stopping). The training loss keeps falling forever and training error is already 0, so neither says anything about new data; in the simulation held-out error was best at round 66 and got worse after.
10. $-0.500 + 0.630 - 0.315 + 0.400 - 0.212 + 0.474 + 0.289 + 0.302 = 1.068 \approx 1.069$; $p = 1/(1 + e^{-1.069}) = 0.744$.

Next: L12d — AdaBoost: the same boosting idea, but mistakes are passed on as **weights**, and the stumps vote.