



FFA Working Papers

Copula-Based Trading of Cointegrated Cryptocurrency Pairs

Masood Tadi

Jiří Witzany

FFA Working Paper 4/2023



FACULTY OF FINANCE AND ACCOUNTING

About: FFA Working Papers is an online publication series for research works by the faculty and students of the Faculty of Finance and Accounting, Prague University of Economics and Business, Czech Republic. Its aim is to provide a platform for fast dissemination, discussion, and feedback on preliminary research results before submission to regular refereed journals. The papers are peer-reviewed but are not edited or formatted by the editors.

Disclaimer: The views expressed in documents served by this site do not reflect the views of the Faculty of Finance and Accounting or any other Prague University of Economics and Business Faculties and Departments. They are the sole property of the respective authors.

Copyright Notice: Although all papers published by the FFA WP series are available without charge, they are licensed for personal, academic, or educational use. All rights are reserved by the authors.

Citations: All references to documents served by this site must be appropriately cited.

Bibliographic information:

Tadi M., Witzany J. (2023). *Copula-Based Trading of Cointegrated Cryptocurrency Pairs*. FFA Working Paper 4/2023, FFA, Prague University of Economics and Business, Prague.

This paper can be downloaded at: wp.ffu.vse.cz

Contact e-mail: ffawp@vse.cz

Copula-Based Trading of Cointegrated Cryptocurrency Pairs

MASOOD TADI* JIŘÍ WITZANY**

Abstract

This research introduces a novel pairs trading strategy based on copulas for cointegrated pairs of cryptocurrencies. To identify the most suitable pairs, the study employs linear and non-linear cointegration tests along with a correlation coefficient measure and fits different copula families to generate trading signals formulated from a reference asset for analyzing the mispricing index. The strategy's performance is then evaluated by conducting back-testing for various triggers of opening positions, assessing its returns and risks. The findings indicate that the proposed method outperforms buy-and-hold trading strategies in terms of both profitability and risk-adjusted returns.

Keywords: Statistical arbitrage, Pairs trading, Cointegration, Copulas, Cryptocurrency market.

*Masood Tadi, Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic (tadim@karlin.mff.cuni.cz, Corresponding author)

**Jiří Witzany, Faculty of Finance and Accounting, Prague University of Economics and Business, Prague, Czech Republic, (jiri.witzany@vse.cz)

Acknowledgements: This paper has been prepared under financial support of a grant GAČR 22-19617S “Modeling the structure and dynamics of energy, commodity and alternative asset prices”, which the authors gratefully acknowledge.

1 Introduction

Pairs trading is a well-known algorithmic trading strategy that capitalizes on temporary abnormal relationships among two or multiple assets whose historical prices tend to move together. When this relationship begins to exhibit abnormal behavior, it triggers the opening of trading positions. These positions are closed as soon as the pairs return to their normal behavior (Vidyamurthy 2004). In the decentralized cryptocurrency market, pairs trading can be profitable, offering two potential arbitrage opportunities: exchange-to-exchange arbitrage and statistical arbitrage. However, implementing a statistical arbitrage strategy based on exchange-to-exchange arbitrage can be risky and pose numerous challenges. In contrast, statistical arbitrage opportunities present similar profit potentials with lower risk (Pritchard 2018).

As stated by Krauss (2017), pairs trading is characterized by a formation and a trading period. During the formation period, the objective is to identify pairs of assets that exhibit similar price movements. This is commonly achieved through co-movement criteria, which can be measured using various methods such as distance metrics (distance approach), for instance, the minimum sum of squared distances of normalized asset prices, or statistical relationships like cointegration rules (cointegration approach). Furthermore, the parameters of the trading period are estimated during the formation period. During the trading timeframe, irregularities in pairs' price movement are aimed to benefit from statistical arbitrage opportunities and create signals to open long/short positions. Advanced strategies may utilize a range of mathematical tools, such as stochastic processes, stochastic control techniques, copulas, and machine learning methods, to enhance the effectiveness of their outcomes.

This paper aims to implement a copula-based pairs trading strategy on cointegrated cryptocurrency pairs. First, we start with a literature review on existing pairs trading strategies. We also provide an overview of the cryptocurrency exchange and data sources used in this study. The theoretical framework of the strategy includes linear and non-linear cointegration tests to identify cointegrated pairs and the copula concept to model their dependence structure. We discuss different families of copulas and the methods for copula estimation. We then outline the methodology for implementing the copula-based trading strategy, including ranking, selection of the assets, and the generation of the trading signals. Next, we present the empirical results of back-testing the strategy to a dataset of historical cryptocurrency prices. We analyze the performance of the strategy using various metrics, including profitability, risk-adjusted return, and maximum drawdown. In addition, we perform a comparative analysis of our results with the buy-and-hold strategy.

2 Literature Review

There is extensive literature on pair trading strategies that use various concepts, such as the distance approach, cointegration analysis, or the concept of copulas. The distance approach involves calculating the historical price spread or price difference between two related assets and monitoring this spread over time. The spread is usually calculated as the difference between the prices of the two assets, either as a raw spread or as a normalized spread, such as the z-score or the percent difference. As an example, Gatev, Goetzmann, and Rouwenhorst (2006) defined the normalized spread, S_t^{ij} , between assets i and j at time t by

$$S_t^{ij} = P_t^{i*} - P_t^{j*} \quad (1)$$

$$P_t^{i*} = \frac{P_t^i}{P_0^i} = \prod_{\tau=1}^t \frac{P_\tau^i}{P_{\tau-1}^i} = \prod_{\tau=1}^t (1 + r_\tau^i) = cr_t^i, \quad (2)$$

where P_t^i is the price of the asset i at time t , r_t^i is the t -period's return on asset i at time t , and cr_t^i is the cumulative total return of asset i up to time t ($cr_0^i = 1$). The calculation of the spread sum of squared distance is performed using the following equation:

$$SSD_{i,j} = \sum_{t=1}^T (S_t^{ij})^2 = \sum_{t=1}^T (cr_{i,t} - cr_{j,t})^2. \quad (3)$$

Pairs are chosen for the trading period based on their ascending order of $SSD_{i,j}$ during the formation stage. The initial pairs at the top of the list are selected, and basic non-parametric threshold rules are employed to generate trading rules. An alternative method was used by Chen et al. (2019), who identified the most suitable pairs for the trading period using Pearson correlation of assets' returns instead of finding the minimum sum of the squared distance. Krauss (2017) found that to maximize the profit of the distance approach strategy, the spread of each selected pair needs to have high volatility, indicating the potential divergence of the two assets. The pair's spread should also have a mean-reverting property. This approach's key advantage is its simplicity and transparency, making it suitable for large-scale empirical applications.

Huck (2015) found that the cointegration approach outperformed the distance approach in selecting effective pairs. The cointegration approach aims to identify a long-term equilibrium between non-stationary time series (e.g., asset prices) that move together. This equilibrium can be linear or nonlinear. Engle and Granger (1987) introduced the first cointegration test, which is based on linear regression and the unit-root test of residuals in the equilibrium. Typically, the augmented Dickey-Fuller test is used for the unit-root test. Other improvements of the Engle-Granger cointegration test were introduced by Phillips and Ouliaris (1990) and Johansen (1991). Highly volatile

markets, such as cryptocurrency, usually exhibit nonlinear features. Therefore, we can extend the Engle-Granger cointegration test by adjusting the error correction model and applying nonlinear unit root tests to increase the reliability of the study. These extensions have been studied by Enders and Siklos (2001), Hansen and Seo (2002), and Kapetanios, Shin, and Snell (2006). Several research papers have been published that specifically explore the application of pairs trading in the cryptocurrency market. These include studies by Broek and Sharif (2018), Pritchard (2018), Kakushadze and Yu (2019), Leung and Nguyen (2019), and Tadi and Kortchemski (2021).

In addition to the commonly used methods discussed above, more advanced concepts such as copulas can be applied to enhance the empirical results of the strategy. Compared to correlation or linear cointegration-based methods, the copula approach provides more valuable information regarding the shape and characteristics of pairs' dependency (Ferreira 2008). Xie and Wu (2013) demonstrated that the two commonly used pairs trading methods, namely the distance and cointegration methods, can be generalized as special cases of the copula method under certain conditions, and the dependency structure of assets in the copula approach is more robust and accurate.

Moreover, Liew and Wu (2013) conducted a comparative study of a copula-based pairs trading strategy with other conventional approaches such as the distance and cointegration approach. They found that the copula approach for pairs trading has better empirical results than the others, as it provides more trading possibilities with higher confidence in practice and does not entail any rigid assumptions, such as the linearity association between assets' returns, in traditional approaches. Hence, the copula approach can provide closer estimations and predictions of reality. According to Stander, Marais, and Botha (2013), the copula approach can significantly demonstrate the pairs' dependency, as it can capture the asymmetry and heavy-tail characteristics of asset returns to model the marginal distribution functions instead of modeling them by Gaussian distribution. Additionally, their empirical analyses reveal that there are more trading opportunities when the market is highly volatile, and the profitability of the strategy strongly depends on the liquidity of the market.

According to Krauss and Stübinger (2017), the copula approach can be divided into two sub-streams: return-based and level-based. In the return-based copula method, the log-returns of two assets are calculated, and their marginal distributions are estimated. Then, an appropriate copula is chosen to represent the dependency relationship between the two assets. To generate trading signals, a mispricing index is defined to indicate the degree of abnormal relationship between the assets. Unlike the distance and cointegration methods which use spread-based mispricing definition, the copula method defines mispricing based on the copula's conditional probability distribution of the corresponding assets' log returns. The conditional distribution of copulas can be

obtained by taking the partial derivative of the copula function, as shown below¹:

$$\begin{aligned} h^{1|2} &:= h(u_1|u_2) = P(U_1 \leq u_1|U_2 = u_2) = \frac{\partial C(u_1, u_2)}{\partial u_2} \\ h^{2|1} &:= h(u_2|u_1) = P(U_2 \leq u_2|U_1 = u_1) = \frac{\partial C(u_1, u_2)}{\partial u_1} \end{aligned} \quad (4)$$

where $C(u_1, u_2)$ is the copula distribution function, $h^{1|2}$ and $h^{2|1}$ are conditional copula distribution functions, and U_1 and U_2 are the transformed uniform variables of the log returns. The values of $h^{1|2}$ and $h^{2|1}$ are between 0 and 1, and as much as their values get away from 0.5, we can consider it as a deviance from the expected relationship between the two assets. Ferreira (2008), Liew and Wu (2013), Stander, Marais, and Botha (2013), and Keshavarz Haddad and Talebi (2023) deployed their strategy in that way.

The usage of the return-based method has a drawback in that entry and exit signals are linked with only the previous period's return. To address this limitation, a new approach (hereafter referred to as the value-based method) was proposed by Xie and Wu (2013). They defined a new mispricing index by aggregating the surplus value of conditional probability in equation 4 from 0.5 across multiple periods to determine the extent to which assets are out of balance.

$$\begin{aligned} \text{CMI}_t^{1|2} &= \text{CMI}_{t-1}^{1|2} + \left(h_t^{1|2} - 0.5 \right) \\ \text{CMI}_t^{2|1} &= \text{CMI}_{t-1}^{2|1} + \left(h_t^{2|1} - 0.5 \right) \end{aligned} \quad (5)$$

The studies conducted by Xie, Liew, et al. (2016), Rad, Low, and Faff (2016), Krauss and Stübinger (2017), and Silva, Ziegelmann, and Caldeira (2023). According to Xie, Liew, et al. (2016), the copula method outperforms the distance approach in terms of describing the dependency relationship between assets and identifying more statistical arbitrage opportunities that generate greater profits. They also recommended utilizing the copula approach for high-frequency pairs trading. Rad, Low, and Faff (2016) compared the performance of the copula approach with that of traditional methods using daily US stocks. They found that while the profitability of the copula method could be weaker, it is more reliable in capturing arbitrage opportunities in the US stock market. They determined that the Student-t copula is more appropriate for modeling the dependence structure of pairs in the US stock market than other copulas.

The performance of a pairs trading strategy based on a weighted combination of copulas was evaluated by Silva, Ziegelmann, and Caldeira (2023) against a distance methodology, using a vast dataset of *S&P500* stocks over 25 years. The study

¹For more details, see Section 4.2.

examined the effect of financial factors on profitability. The mixed copula approach yielded better results than the distance method, with higher alphas for fully invested capital and superior performances overall. The approach was notably effective under committed capital during both crisis and non-crisis periods and fully invested during non-crisis periods.

However, in practice, the cumulative mispricing indices (CMI) in equation 5 may not necessarily exhibit mean-reverting behavior, which can negatively impact the profitability of the strategy. Hence, a new methodology is proposed in this study to replace log-returns with stationary spread processes in order to overcome this issue².

3 Cryptocurrency Exchange and Data Source

The cryptocurrency market facilitates decentralized trading of cryptocurrencies across various exchanges. Binance, established in 2017, is the largest cryptocurrency exchange worldwide in terms of daily trading volume in both spot and derivatives markets. Binance provides three types of derivative contracts: Futures contracts, Options, and Binance Leveraged Tokens (BLVT).

Futures contracts are classified into two main categories: COIN-Margined contracts and USD\$-Margined contracts. COIN-Margined contracts are inverse Futures quoted in US dollars but denominated in an underlying cryptocurrency (e.g., Bitcoin). They include traditional quarterly Futures and perpetual Futures, also known as perpetual Swaps, which never expire or settle. Because they lack a settlement price, their price can deviate significantly from their spot contract price. Binance uses funding fees to address this issue for long and short sides. On the other hand, USD\$-Margined contracts are similar to COIN-Margined Futures, and they may have perpetual or quarterly expiration. However, they are quoted, denominated, and settled in stablecoins such as Tether (USDT) and Binance USD (BUSD), which are pegged to the US dollar value. For this study, all nominated cryptocurrency coins are USDT-Margined Futures (*Binance Crypto Derivatives* accessed 2022-11-03).

The minimum trade amount, also known as the minimum price increment, is the smallest possible change in the price of a contract at an exchange. This value is specific to each asset and can be adjusted over time. Assets with smaller increments have narrower bid/ask spreads. When limit orders are not entirely disclosed to traders, they tend to order contracts with smaller quote sizes to avoid slippage (Harris 1997).

To calculate profit and loss, all coins are valued in Tether (USDT), a stablecoin. Using the Binance API, we collected historical hourly closed prices for twenty cryptocurrency


²For more details, see Section 5.

coins from 01/01/2021 to 19/01/2023. We used the linear Engle-Granger (EG) two-step method and to identify cointegrated pairs and calculated Kendall's Tau. Then, we calibrated the method's parameters using the copula approach and generated trading signals for the strategy.

To calculate profit and loss, all coins are valued in Tether (USDT), a stablecoin. Using the Binance API, we collected historical hourly closed prices for twenty cryptocurrency coins from 01/01/2021 to 19/01/2023. We used Engle-Granger (EG) and Kapetanios-Shin-Shell (KSS) cointegration tests to identify cointegrated pairs and calculated Kendall's Tau. Then, we calibrated the method's parameters using the copula approach and generated trading signals for the strategy.

A pairs trading cycle comprises formation and trading periods. In this study, the entire process of pairs trading is conducted within a month, with three weeks allocated for the formation step and the remaining week designated for the trading step. We carry out 104 cycles that move dynamically over time, with each cycle sharing three-quarters of its data with its previous or subsequent cycle.

Figure 1: The scheme of Formation and Trading Periods



Furthermore, we implemented the entire methodology using Python and R, with Python being the preferred option for data handling and R for statistical tests and copula modeling.

Table I: Python libraries and R packages used in this research

Python		R	
Name	Type	Name	Type
NumPy	Numerical analysis	copula	Copula modeling
Pandas	Technical computing	VineCopula	Copula modeling
Matplotlib	Plotting	MASS	Statistical package
Datetime	Date/time package	dplyr	Data handling
Statsmodels	Statistical package	lubridate	Date/time package
SciPy	Technical computing	fUnitRoots	Unit root analysis

4 Theoretical Framework

4.1 Unit-Root Test

The cointegration property can be employed to identify the most appropriate pair from multiple combinations of coins. Both linear and non-linear cointegration tests can be utilized for evaluation. Initially, the pair spread value is defined without an intercept in the following manner:

$$S_t = P_t^1 - \beta P_t^2 \quad (6)$$

Suppose that P_t^1 and P_t^2 are non-stationary time series. We use unit-root tests to study whether the spread, S_t , is also a non-stationary process or not. The augmented Dickey-Fuller (ADF) unit-root test, as described by Dickey and Fuller (1979), uses a test equation applied to the demeaned spread process S_t in the following form:

$$\Delta S_t = \beta S_{t-1} + \sum_{i=1}^{p-1} \gamma_i \Delta S_{t-i} + \epsilon_t, \quad (7)$$

where β is the coefficient of the lagged level of the series, γ_i are the coefficients of the lagged differences, ϵ_t is the error term, and p is the number of lags used in the test. The null hypothesis of the ADF test is that the series has a unit root, i.e., $\beta = 0$. If the test statistic exceeds a critical value at a given significance level, the null hypothesis is rejected, indicating that the series is stationary and does not have a unit root. Conversely, if the test statistic is below the critical value, the null hypothesis cannot be rejected, implying that the series has a unit root and is non-stationary.

Traditional unit-root tests like the Augmented Dickey-Fuller (ADF) test assume that the data-generating process is linear. However, non-linear unit-root tests are designed to

account for non-linearities in time series data and provide more accurate assessments of unit roots. There are several non-linear unit-root tests available in the literature, each with its own assumptions, methodologies, and advantages. Examples include the Teräsvirta (1994) test, the Zivot and Andrews (2002) test, the Kapetanios, Shin, and Snell (2003) test, and the Kapetanios (2005) test. These tests often involve estimating non-linear models like threshold auto-regressive (TAR) models, smooth transition auto-regressive (STAR) models, or other non-linear models, and computing test statistics to compare estimated model parameters with critical values.

The general self-exciting threshold auto-regressive (SETAR) model with n regimes applied to the demeaned spread process S_t is in the following form:

$$S_t = \sum_{i=1}^p \left(\phi_{i1} \mathbb{1}_{\{S_{t-d} \leq c_1\}} + \sum_{j=1}^{n-1} \phi_{ij} \mathbb{1}_{\{c_j < S_{t-d} \leq c_{j+1}\}} + \phi_{in} \mathbb{1}_{\{S_{t-d} > c_n\}} \right) S_{t-i} + \epsilon_t, \quad (8)$$

where d denotes the transition's delay, c_j represents the j -th threshold, and ϵ_t represents the error term. In special case Kapetanios, Shin, and Snell (2003) proposed a test equation where the indicator function is replaced by an exponential smooth transition function in the form:

$$S_t = S_{t-1} + \sum_{i=1}^p \left(\gamma_{1i} \left(1 - e^{-\theta(S_{t-1}-c)^2} \right) S_{t-i} \right) + \epsilon_t \quad (9)$$

When c is set to zero, and p is set to one, using Taylor approximation, equation 9 can be illustrated as

$$\Delta S_t = \delta (S_{t-1})^3 + \epsilon'_t \quad (10)$$

where $\delta = \gamma_1 \theta$ and $\epsilon'_t = f(\epsilon_t)$. The null hypothesis assumes that δ is equal to zero, while the alternative hypothesis posits that δ is less than zero. Note that the asymptotic standard normal distribution of the t-statistic for $\delta = 0$ against $\delta < 0$ is not applicable. However, its asymptotic critical values have been determined through stochastic simulations and are documented by Kapetanios, Shin, and Snell (2003).

4.2 Copula Concept

Suppose that a continuous random variable X has a continuous distribution, and its probability distribution function is defined as $F_X(x) := \mathbb{P}(X \leq x)$. If F_X is strictly increasing, then F_X^{-1} is defined by $F_X^{-1}(u) = x \Leftrightarrow F_X(x) = u$. However, if F_X is constant on some interval, then the inverse function is not well defined by $F_X^{-1}(u) = x$. To avoid this problem, we can define $F_X^{-1}(u)$ for $0 < u < 1$ by the generalized inverse function such that

$$F_X^{-1}(u) = \inf\{x : F_X(x) \geq u\}, \quad (11)$$

Now, in the same way, we define another continuous random variable Y with distribution function F_Y and generalized inverse function F_Y^{-1} similar to equation 11. Given two continuous random variables X and Y , with distribution functions F_X and F_Y respectively, the joint distribution function $F_{X,Y}$ can be written as:

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y) = \mathbb{P}(F_X(X) \leq F_X(x), F_Y(Y) \leq F_Y(y)) \quad (12)$$

where the last equality follows from the fact that F_X and F_Y are both increasing. Then we define random variables U and V such that $U := F_X(X)$ and $V := F_Y(Y)$. According to the probability integral transformation theorem, the probability distribution function of U , F_U , and the probability distribution function of V , F_V , are uniformly distributed on $[0, 1]$. (See Casella and Berger (2021, p. 54-55) for the proof).

Definition: A two-dimensional copula C is a function that maps the unit square $[0, 1]^2$ into the unit interval $[0, 1]$, satisfying the following requirements:

1. $C(0, v) = C(u, 0) = 0$, for $0 \leq u, v \leq 1$.
2. $C(u, 1) = u$, and $C(1, v) = v$, for $0 \leq u, v \leq 1$.
3. $C(u_1, v_1) - C(u_1, v_2) - C(u_2, v_1) + C(u_2, v_2) \geq 0$, for $1 \geq u_1 > u_2 \geq 0$, and $1 \geq v_1 > v_2 \geq 0$.

We can define several copula functions, but the three requirements above should be satisfied by $C(u, v)$ to have a well-defined joint distribution function (Cherubini et al. 2011). According to Sklar's theorem, there exists a Copula function C which could connect the uniform random variables U and V to the joint distribution function $F_{X,Y}$ as follows

$$F_{X,Y}(X, Y) = F_{X,Y}(F_X^{-1}(U), F_Y^{-1}(V)) := C(U, V), \quad (13)$$

Hence, we can rewrite the joint distribution function of X and Y in terms of standard uniform random variables U and V such that

$$F_{X,Y}(x, y) = F_{X,Y}(F_X^{-1}(u), F_Y^{-1}(v)) := C(u, v) = C(F_X(x), F_Y(y)), \quad (14)$$

where $u = F_X(x)$, and $v = F_Y(y)$. Knowing $F_{X,Y}(x, y) = C(u, v)$, we can determine the copula density function $c(u, v)$ by

$$c(u, v) = \frac{\partial^2 C(u, v)}{\partial u \partial v} = \frac{\partial^2 F_{X,Y}(x, y)}{\partial F_X(x) \partial F_Y(y)} = \frac{\frac{\partial^2 F_{X,Y}(x, y)}{\partial x \partial y}}{\frac{\partial F_X(x)}{\partial x} \frac{\partial F_Y(y)}{\partial y}} = \frac{f_{X,Y}(x, y)}{f_X(x) f_Y(y)} \quad (15)$$

Sklar's theorem enables us to separate the modeling of the marginal distributions $F_X(x)$ and $F_Y(y)$ from the dependence relation represented in C . We can define the conditional

distribution functions using copula function. The conditional distribution of $Y|X = x$ is obtained by the first partial derivatives of the copula function as follows

$$\begin{aligned}
F_{Y|X}(y) &= \mathbb{P}(Y \leq y|X = x) = \mathbb{P}(F_Y(Y) \leq F_Y(y)|F_X(X) = F_X(x)) \\
&= \mathbb{P}(V \leq v|U = u) = \lim_{\delta \rightarrow 0^+} \frac{\mathbb{P}(V \leq v, U \in (u - \delta, u + \delta))}{\mathbb{P}(U \in (u - \delta, u + \delta))} \\
&= \lim_{\delta \rightarrow 0^+} \frac{C(u + \delta, v) - C(u - \delta, v)}{2\delta} \\
&= \frac{\partial}{\partial u} C(u, v).
\end{aligned} \tag{16}$$

Copulas are invariant concerning strictly increasing transformations of the marginal distributions. For more details about the characterization of invariant copulas, see Klement, Mesiar, and Pap (2002). Also, we can increase the range of dependence captured by copulas by rotating them. The following equations show how to rotate a copula by 90, 180, and 270 degrees:

$$\begin{aligned}
C_{90}(u_1, u_2) &:= C(1 - u_2, u_1) \\
C_{180}(u_1, u_2) &:= C(1 - u_1, 1 - u_2) \\
C_{270}(u_1, u_2) &:= C(u_2, 1 - u_1)
\end{aligned} \tag{17}$$

4.3 Families of copulas

There are several types of copulas. This research focuses on three popular families of copulas: elliptical, Archimedean, and extreme value copulas. Each type of copula is characterized by its properties, such as its dependence structure and tail behavior, and is often used in different areas of statistics and finance.

4.3.1 Elliptical Copulas

Elliptical copulas are widely used in finance and insurance applications due to their ability to model dependence with different levels of tail dependence. An elliptical copula is constructed from a multivariate elliptical distribution. The most commonly used elliptical distributions are the multivariate Gaussian and Student-t. The density function of any elliptical distribution $f_{\mathbf{X}}$ is in the form

$$f_{\mathbf{X}}(\mathbf{x}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) = k_n |\boldsymbol{\Sigma}|^{-\frac{1}{2}} g\left((\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu})\right), \tag{18}$$

where $k_n \in \mathbb{R}$ is the normalizing constant and depends on the dimension n , \mathbf{x} is an n -dimensional random vector with mean vector $\boldsymbol{\mu}$, and a positive definite matrix which is

proportional to the covariance matrix Σ , and some function $g(\cdot)$ which is independent of the dimension n (Czado 2019). In the case of bivariate Gaussian distribution $g(x) := e^{-x^2/2}$, $k_n := 1/(2\pi)$, and the probability density function of $\mathbf{X} = (X_1, X_2)$ is

$$f_{\mathbf{X}}(\mathbf{x}; \boldsymbol{\mu}, \Sigma) = \frac{1}{2\pi|\Sigma|^{1/2}} e^{-\frac{1}{2}[(\mathbf{x}-\boldsymbol{\mu})^T \Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu})]} \quad (19)$$

$$\boldsymbol{\mu} = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \quad \Sigma = \begin{pmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{pmatrix},$$

where ρ is the correlation between random variables X_1 and X_2 , $\sigma_1 > 0$ and $\sigma_2 > 0$. If $\mu_1 = \mu_2 = 0$ and $\sigma_1 = \sigma_2 = 1$, then the density and distribution functions of the standard Bivariate Gaussian distribution are obtained by

$$\phi_{X_1 X_2}(x_1, x_2; \rho) = \frac{1}{2\pi\sqrt{1-\rho^2}} e^{-\frac{x_1^2 - 2\rho x_1 x_2 + x_2^2}{2(1-\rho^2)}} \quad (20)$$

$$\Phi_{X_1 X_2}(u_1, u_2; \rho) = \int_{-\infty}^{u_1} \int_{-\infty}^{u_2} \phi_{X_1 X_2}(x_1, x_2; \rho) dx_1 dx_2$$

Using Sklar's theorem in 13 and 20, the bivariate Gaussian copula is defined by

$$C(u_1, u_2; \rho) := \Phi_{X_1 X_2}(\Phi_{X_1}^{-1}(u_1), \Phi_{X_2}^{-1}(u_2); \rho) \quad (21)$$

In the bivariate t distribution, $g(\cdot)$ and k_n function in the formula 18 are defined by $g(x) := (1 + x/\nu)^{-(\nu+2)/2}$, $k_n := \Gamma(\frac{\nu+2}{2}) / (\Gamma(\frac{\nu}{2})\nu\pi)$, and the probability density function of $\mathbf{T} = (T_1, T_2)$ is obtained by

$$f_{\mathbf{T}}(\mathbf{t}; \nu, \boldsymbol{\mu}, \Sigma) = \frac{\Gamma(\frac{\nu+2}{2})}{\Gamma(\frac{\nu}{2})\nu\pi|\Sigma|^{1/2}} \left[1 + \frac{1}{\nu}(\mathbf{t} - \boldsymbol{\mu})^T \Sigma^{-1}(\mathbf{t} - \boldsymbol{\mu}) \right]^{-(\nu+2)/2} \quad (22)$$

where $\nu > 0$ is the degree of freedom parameter and $\boldsymbol{\mu}$ and $\boldsymbol{\sigma}$ are the same as 19. If $\mu_1 = \mu_2 = 0$ and $\sigma_1 = \sigma_2 = 1$, and knowing that $\Gamma(\frac{\nu+2}{2})/\Gamma(\frac{\nu}{2}) = \frac{\nu}{2}$, the density and distribution function of the standard Bivariate Student-t distribution are obtained by

$$f_{T_1 T_2}(t_1, t_2; \nu, \rho) = \frac{(1-\rho^2)^{-1/2}}{2\pi} \left[1 + \frac{t_1^2 - 2\rho t_1 t_2 + t_2^2}{\nu(1-\rho^2)} \right]^{-(\nu+2)/2} \quad (23)$$

$$F_{T_1 T_2}(u_1, u_2; \nu, \rho) = \int_{-\infty}^{u_1} \int_{-\infty}^{u_2} f_{T_1 T_2}(t_1, t_2; \nu, \rho) dt_1 dt_2$$

Then the bivariate Student-t copula is defined by

$$C(u_1, u_2; \nu, \rho) := F_{T_1 T_2}(F_{T_1}^{-1}(u_1), F_{T_2}^{-1}(u_2); \nu, \rho) \quad (24)$$

4.3.2 Archimedean Copulas

Archimedean copulas are based on a generator function and can model dependence with tail dependence that decreases logarithmically or exponentially. According to Nelsen (2007), Archimedean copulas are defined as follows:

$$C(u_1, \dots, u_n) = \phi^{[-1]}(\phi(u_1) + \dots + \phi(u_n)) \quad (25)$$

where $\phi : [0, 1] \rightarrow [0, \infty]$ is called a generator which is continuous, strictly decreasing, and convex function such that $\phi(1) = 0$. In addition, $\phi^{[-1]} : [0, \infty] \rightarrow [0, 1]$ is called the pseudo-inverse of ϕ function and is defined by

$$\phi^{[-1]}(t) = \begin{cases} \phi^{-1}(t) & , 0 \leq t \leq \phi(0) \\ 0 & , \phi(0) \leq t \leq \infty \end{cases} \quad (26)$$

If $\phi(0)$ equals infinity, then the pseudo-inverse function $\phi^{[-1]}$ is equivalent to the inverse function ϕ^{-1} . Archimedean copulas allow for a range of generator functions to be selected, which ultimately determines the type of tail dependence exhibited by the copula. For this research, both one-parameter copulas (such as the Gumbel, Clayton, Frank, and Joe copulas) and two-parameter copulas (such as BB1, BB6, BB7, and BB8) were utilized. Table X in the appendix provides a list of the bivariate Archimedean copulas utilized in this study.

4.3.3 Extreme-Value Copulas

Extreme value copulas are used to model dependence with strong tail dependence, making them suitable for modeling extreme events. Suppose that $X_i = (X_{i1}, X_{i2})^T$, $i \in \{1, 2, \dots, n\}$, be independent and identically distributed random vectors with joint distribution function F , and marginal distributions F_1 and F_2 . According to Gudendorf and Segers (2010), we can define the bivariate vector of component-wise maxima $M_n = (M_{n1}, M_{n2})^T$ such that

$$M_{nj} := \max_{i \in \{1, 2, \dots, n\}} (X_{ij}), \quad j = 1, 2. \quad (27)$$

Then, the bivariate copula C_n of M_n is obtained by

$$C_n(u_1, u_2) = C_F \left(u_1^{1/n}, u_2^{1/n} \right)^n, \quad (u_1, u_2) \in [0, 1]^2. \quad (28)$$

In equation 28, if C_F exists such that

$$\lim_{n \rightarrow \infty} C_F(u_1^{1/n}, u_2^{1/n})^n = C(u_1, u_2), \quad (u_1, u_2) \in [0, 1], \quad (29)$$

then the bivariate copula C in (29) is called an extreme-value copula. Bivariate extreme-value copulas can be demonstrated in terms of a function $A(t)$ in this form:

$$C(u_1, u_2) = (u_1 u_2)^{A(\ln(u_2)/\ln(u_1 u_2))}, \quad (u_1, u_2) \in (0, 1]^2 \setminus \{(1, 1)\}, \quad (30)$$

where the function $A : [0, 1] \rightarrow [1/2, 1]$, which is called the Pickands dependence function, is convex and satisfies $\max(1-t, t) \leq A(t) \leq 1$ for all $t \in [0, 1]$ (Gudendorf and Segers 2010). Some Archimedean copulas, such as the Gumbel copula, can be expressed by the extreme-value family. These special copulas create a hybrid category, including both the Archimedean and the extreme-value copulas, and are called Archimax copulas.

Table II shows extreme-value copulas used in this research. Note that Tawn copula has three parameters and its Pickands's dependence function is in the form

$$A(t) = (1 - \beta) + (\beta - \alpha)t + [(\alpha(1 - t))^\theta + (\beta t)\theta]^{1/\theta}, \quad (31)$$

where $\theta \geq 1$ and $\alpha, \beta \in [0, 1]$. The simplified Tawn copula cases with $\beta = 1$ and $\alpha = 1$ are respectively called Tawn Type 1 and Tawn Type 2 copula and have two parameters.

Table II: Pickands dependence function of some extreme-value copulas

Name	Pickands function $A(t)$	Parameters
Gumbel	$\left[t^\theta + (1 - t)^\theta\right]^{1/\theta}$	$\theta \geq 1$
Tawn Type 1	$(1 - \alpha)t + \left[(\alpha(1 - t))^\theta + t^\theta\right]^{1/\theta}$	$\theta \geq 1, 0 \leq \alpha \leq 1$
Tawn Type 2	$(1 - \beta)(1 - t) \left[(1 - t)^\theta + (\beta t)\theta\right]^{1/\theta}$	$\theta \geq 1, 0 \leq \beta \leq 1$

4.4 Copula Estimation

When the marginal probability density of X_1 and X_2 and their corresponding copula density $c(\cdot)$ are given in their parametric with unknown parameters, we can estimate the vector of parameter vector $\theta = (\beta, \alpha)^T$ with the maximum likelihood estimation method, where $\beta = (\beta_1, \beta_2)^T$ represent the marginal parameters and α represents the copula parameters. The log-likelihood function for (X_1, X_2) , where $X_i = (x_{i1}, \dots, x_{in})$, can be expressed as

$$l(\theta) = \sum_{j=1}^n \left[\log c(F_1(x_{1j}; \beta_1), F_2(x_{2j}; \beta_2); \alpha) + \log f_1(x_{1j}; \beta_1) + \log f_2(x_{2j}; \beta_2) \right], \quad (32)$$

and the maximum likelihood estimator of θ is

$$\hat{\theta}_{ML} = \underset{\theta}{\operatorname{argmax}} l(\theta). \quad (33)$$

However, since this approach is computationally expensive, an alternative method called the Inference for the Margins (IFM) two-step method can be used instead. This method is computationally easier to obtain compared to the full maximum likelihood estimation approach. First, we estimate the margins' parameters by performing the estimation of the univariate marginal distributions using the log-likelihood function where the maximum likelihood estimator of β_i is

$$\hat{\beta}_i = \underset{\beta_i}{\operatorname{argmax}} \sum_{j=1}^n [\log f_i(x_{ij}; \beta_i)]. \quad (34)$$

Then given estimated marginal parameters, we transform data to the copula scale, develop the copula model, and estimate the copula parameters α as follows

$$\hat{\alpha}_{ML} = \underset{\alpha}{\operatorname{argmax}} \sum_{j=1}^n \left[\log c \left(F_1(x_{1j}; \hat{\beta}_1), F_2(x_{2j}; \hat{\beta}_2); \alpha \right) \right]. \quad (35)$$

We can also employ a semiparametric approach known as canonical maximum likelihood to estimate copula parameters without specifying the marginals. The empirical cumulative distribution function of $X_i = (x_{i1}, \dots, x_{in})$ is

$$\widehat{F}_n^i(t) = \frac{\#\{X_i \leq x\}}{n+1} = \frac{1}{n+1} \sum_{j=1}^n \mathbb{1}_{\{x_{ij} \leq t\}}. \quad (36)$$

Then, the copula parameters are estimated using maximum likelihood estimation as follows:

$$\hat{\alpha}_{ML} = \underset{\alpha}{\operatorname{argmax}} \sum_{j=1}^n \left[\log c \left(\widehat{F}_n^1(x_{1j}), \widehat{F}_n^2(x_{2j}); \alpha \right) \right]. \quad (37)$$

5 Implementation Methodology

The study suggests a new approach to address the issue of cumulative mispricing indices (CMI) not exhibiting mean-reverting behavior in practice, which can negatively affect the profitability of trading strategies. The proposed methodology involves using stationary spread processes instead of log-returns. In the pairs trading strategy employed in this study, the spread process is defined as a linear combination of two

assets, with a reference asset (BTCUSDT) selected and other cryptocurrency coins identified as cointegrated with it using a specific equation.

$$S_t^i = \text{BTCUSDT}_t - \hat{\beta}^i P_t^i \quad i = 1, 2, \dots, 19 \quad (38)$$

where $\hat{\beta}$ is the estimated linear regression coefficient between BTCUSDT and the second coin, chosen from the other 19 coins. This allows us to identify 19 pairs and select the optimal pairs during the formation period for trading in the trading period. To determine the optimal pairs, we use the linear Engle-Granger (EG) two-step method and the non-linear Kapetanios-Shin-Snell (KSS) cointegration tests to identify cointegrated coins. However, since there may be multiple cointegrated pairs, we need to add another criterion to rank them. To do this, we calculate Kendall's Tau (τ), which is a measure of correlation for ranked data and defined

$$\tau(S^i, S^j) = \tau_{ij} = \frac{\text{Number of concordant pairs} - \text{Number of discordant pairs}}{\text{Total number of pairs}} \quad (39)$$

where

$$\text{Number of concordant pairs} = \sum_{n=1}^{N-1} \sum_{m=n+1}^N \text{sgn}(S_{[n]}^i - S_{[m]}^i) \text{sgn}(S_{[n]}^j - S_{[m]}^j),$$

$$\text{Number of discordant pairs} = \sum_{n=1}^{N-1} \sum_{m=n+1}^N \text{sgn}(S_{[n]}^i - S_{[m]}^i) \text{sgn}(S_{[m]}^j - S_{[n]}^j),$$

and total number of pairs = $N(N-1)/2$. Here, N is the number of data points, $S_{[n]}^i$ and $S_{[n]}^j$ are the rankings of the n -th data point in two different variables, and $\text{sgn}(x)$ is the sign function, which is 1 if $x > 0$, -1 if $x < 0$, and 0 if $x = 0$.

After calculating τ of BTCUSDT with each of the 19 altcoins, we select the two altcoins that have the highest τ with BTCUSDT and create their corresponding pairs. During the course of one week, the chosen pairs are traded and can be substituted with different pairs at the start of each trading period. Instead of trading a pair of coins, we employ a pair of spreads where each of them contains BTCUSDT. In this way, having a long position in one spread and a short position in the other one implies that BTCUSDT will not be traded at all and it only plays an intermediary role between two other coins.

Now, we estimate the probability distribution function of spread processes (marginals). We fit various distributions such as Gaussian, Student-t, and Cauchy to the data to identify the best-fitting distribution. By employing statistical methods or maximum likelihood estimation, we estimate the specific parameters associated with each distribution. To evaluate the goodness of fit, we calculate the AIC values for the candidate distributions, ultimately selecting the distribution with the lowest AIC value

as the best-fitting option. Suppose that F^i is the fitted cumulative distribution function of the spread process S^i . The Probability Integral Transform is employed to convert the spreads to random variables $U_1 := F^1(S^1)$ and $U_2 := F^2(S^2)$ with a standard uniform distribution. The next step is to determine a fitting copula model for U_1 and U_2 . We select some potential copulas and estimate the corresponding parameters by the maximum likelihood method. Finally, using the Akaike information criterion (AIC), we distinguish the most fitted copula model.

During the trading period, using the hourly realizations of random variables U_1 and U_2 , which are the transformed values of spread processes, we calculate the copula conditional probabilities $h^{1|2}$ and $h^{2|1}$ defined in equation 4, for the selected pairs of the week. when $h^{1|2}$ is higher (lower) than 0.5, the first coin can be considered to be overvalued (undervalued) relative to the second one. Similarly, when $h^{2|1}$ is higher (lower) than 0.5, the second coin can be considered to be overvalued (undervalued) relative to the first one. Therefore, we can interpret mispricing as conditional probabilities in equation 4 minus 0.5. We denote the trading thresholds to be α_1 and α_2 . We study the optimal triggers via back-testing. The opening and closing signals are generated by the following rules:

Table III: Trading rules in terms S^1 and S^2

Trading Rule	Signals
If $h^{1 2} < \alpha_1$ and $h^{2 1} > 1 - \alpha_1$	open long S^1 and short S^2
If $h^{1 2} > 1 - \alpha_1$ and $h^{2 1} < \alpha_1$	open short S^1 and, long S^2
If $ h^{1 2} - 0.5 < \alpha_2$ and $ h^{2 1} - 0.5 < \alpha_2$	close both positions

Table IV: Trading rules in terms P^1 and P^2

Trading Rule	Signals
If $h^{1 2} < \alpha_1$ and $h^{2 1} > 1 - \alpha_1$	open long $\beta^2 \times P^2$ and short $\beta^1 \times P^1$
If $h^{1 2} > 1 - \alpha_1$ and $h^{2 1} < \alpha_1$	open short $\beta^2 \times P^2$ and long $\beta^1 \times P^1$
If $ h^{1 2} - 0.5 < \alpha_2$ and $ h^{2 1} - 0.5 < \alpha_2$	close both positions

Figure 2: Confidence bands of Gumbel copula ($\theta = 2$) at $\alpha_1 = 5\%$ and $\alpha_2 = 10\%$




Figure 2 provides an illustrative depiction of the confidence bands in trading rules for the Gumbel copula with a parameter value of $\theta = 2$, under the conditions of $\alpha_1 = 5\%$ and $\alpha_2 = 10\%$. If the data point (u_{1t}, u_{2t}) falls within the top green (down green) area, it suggests that S_1 is undervalued (overvalued) and S_2 is overvalued (undervalued), which may indicate an opportunity to open a position. Conversely, if (u_{1t}, u_{2t}) falls within the red area, it may signal the need to close the positions.

6 Empirical Results

Table V presents the occurrence rate of selected copulas and their rotations over 104 trading weeks. The results indicate that copulas of extreme value, such as Tawn type 1 and 2, and certain two-parameter Archimedean copulas, particularly BB7, play a

significant role in the process of selecting the appropriate model.


Table V: Occurrence rate of copulas in the study

Copulas and their rotations	Strategy with EG Test	Strategy with KSS Test
Gaussian	4.8%	5.8%
Student-t	6.7%	4.8%
Clayton	3.8%	6.7%
Frank	2.9%	3.8%
Gumbel	1.0%	4.8%
Joe	10.6%	7.7%
BB1	5.8%	2.9%
BB6	1.9%	1.9%
BB7	15.4%	15.4%
BB8	8.7%	7.7%
Tawn type 1	23.1%	24.0%
Tawn type 2	15.4%	14.4%

Tables VIII and IX in the appendix display the coin pairs chosen for each trading week, which can vary from week to week. It is worth noting that there are instances where the selected coin pairs remain unchanged across consecutive weeks. The ADF unit-root test is conducted with a significance level of 10%. For the KSS unit-root test, the asymptotic critical value at a 10% significance level is -1.92. If we fail to identify a minimum of two stationary spread processes within a specified formation period, we abstain from trading during its corresponding trading period.

The results of our trading strategy's profit and loss calculations with varying entry thresholds (α_1) are illustrated in Figure 2. Note that the exit threshold (α_2) is fixed at 10% in all cases. Realized profit and loss are determined by considering the commission fees and the difference between the opening and closing prices of a position. It should be noted that we initially invested 200,000 USDT, with the coins' weights set to ensure that each side has a maximum initial capital of around 200,000 USDT. It is also assumed that all trades are executed using market orders, which typically incur higher fees (known as taker fees) compared to the lower fees charged for limit orders (known as maker fees). For instance, on Binance, the maker fee of USDT-Margined Futures is set at 0.02%, while the taker fee is 0.04%.

Figure 3: Copula-based pairs trading strategy P&L



We assess the risk-adjusted performance of a strategy using the Sharpe ratio. Table VI³ illustrates the yearly returns, volatility, and Sharpe ratios for different strategies, along with the maximum drawdown and the returns obtained over the maximum drawdown

³The Gross Profit and Loss (P&L) mentioned in table VI is the total profit and loss generated by transactions before deducting any transaction fees associated with them.

(RoMaD), which serves as an alternative to the Sharpe Ratio⁴. In pairs trading with the EG cointegration test, the annualized return decreases with higher entry thresholds, ranging from 76.2% at $\alpha_1 = 0.10$ to 52.1% at $\alpha_1 = 0.20$. In contrast, the annualized standard deviation increases with higher levels of α_1 , ranging from 38.2% at $\alpha_1 = 0.10$ to 52.7% at $\alpha_1 = 0.20$. In the optimal case, the annualized Sharpe Ratio is approximately 1 ($\alpha_1 = 0.10$). The maximum drawdown increases with higher α_1 , ranging from -35.6% at $\alpha_1 = 0.10$ to -43.9% at $\alpha_1 = 0.20$. The return over maximum drawdown (RoMaD) decreases with higher α_1 , ranging from 2.14 at $\alpha_1 = 0.10$ to 1.19 at $\alpha_1 = 0.20$.

In a similar manner, the annualized return in pairs trading utilizing the KSS cointegration test diminishes as the entry threshold levels increase, declining from 35.3% when $\alpha_1 = 0.10$ to 10.4% when $\alpha_1 = 0.20$. As α_1 increases, the annualized standard deviation increases, starting from 37.3% at $\alpha_1 = 0.10$ and reaching 63.5% at $\alpha_1 = 0.20$. Therefore, the annualized Sharpe Ratio exhibits a declining pattern as the threshold levels increase. It ranges from 0.93 at $\alpha_1 = 0.10$ to 0.16 at $\alpha_1 = 0.20$. The maximum drawdown increases with higher threshold levels, ranging from -34.0% at $\alpha_1 = 0.10$ to -43.0% at $\alpha_1 = 0.20$. The return over maximum drawdown shows a similar pattern as the annualized Sharpe Ratio, decreasing with higher thresholds.

When the entry threshold α_1 increases, it results in an increase in the number of trading signals. This, in turn, leads to higher volatility in both the EG and KSS strategies. However, despite the increased volatility, there is no corresponding increase in return. This indicates that the optimal threshold is 0.10, and at this level, the risk-adjusted profitability is similar between the strategy with the EG test and the KSS test.

To compare our strategy's results with the buy-and-hold strategy, we use a passive investment approach that involves holding a relatively steady portfolio for an extended period, despite short-term market fluctuations. The Bitcoin buy-and-hold strategy shows a negative annualized return of -17.0% with a high annualized standard deviation of 72.6%. The annualized Sharpe Ratio is negative (-0.23), indicating poor risk-adjusted performance. The maximum drawdown is severe (-77.1%), indicating a high risk. On the other hand, the portfolio buy-and-hold strategy⁵ shows a positive annualized return of 14.4% with a high annualized standard deviation of 98.6%. The annualized Sharpe Ratio is positive (0.15), indicating better risk-adjusted performance compared to the Bitcoin buy-and-hold strategy. The maximum drawdown is also severe (-82.2%), indicating a high risk. Both tests of the pairs trading strategy outperform the Bitcoin buy-and-hold strategy and the portfolio buy-and-hold strategy.

⁴Transaction fees are taken into account in all calculations

⁵The buy-and-hold strategy in the portfolio involves investing in all twenty cryptocurrency coins with equal weights at the start of the study, retaining them throughout the trading periods, and ultimately selling them at the end of the study period.

Our pairs trading strategy outperforms the portfolio buy-and-hold by over six times when $\alpha_1 = 0.10$.

Table VI: Results of pairs trading strategies Utilizing hourly closed prices of twenty cryptocurrencies from 22/01/2021 to 19/01/2023

Pairs Trading with EG Test			
	$\alpha_1 = 0.10$	$\alpha_1 = 0.15$	$\alpha_1 = 0.20$
Total Return	76.2%	54.2%	52.1%
Annualized Return	37.1%	26.4%	25.4%
Annualized Standard Deviation	38.2%	47.4%	52.7%
Annualized Sharpe Ratio	0.97	0.56	0.48
Maximum Drawdown	-35.6%	-41.8%	-43.9%
Return over Maximum Drawdown	2.14	1.30	1.19
Number of Transactions	176	200	222
Transaction Costs over Gross P&L	11.7%	19.2%	21.9%
Pairs Trading with KSS Test			
	$\alpha_1 = 0.10$	$\alpha_1 = 0.15$	$\alpha_1 = 0.20$
Total Return	72.3%	44.8%	21.3%
Annualized Return	35.3%	21.8%	10.4%
Annualized Standard Deviation	37.7%	43.6%	63.5%
Annualized Sharpe Ratio	0.93	0.50	0.16
Maximum Drawdown	-34.0%	-34.6%	-43.0%
Return over Maximum Drawdown	2.13	1.29	0.50
Number of Transactions	184	222	254
Transaction Costs over Gross P&L	12.9%	25.2%	47.7%
Buy & Hold Strategy			
	Bitcoin Buy & Hold	Portfolio Buy & Hold	
Total Return	-33.9%	28.8%	
Annualized Return	-17.0%	14.4%	
Annualized Standard Deviation	72.6%	98.6%	
Annualized Sharpe Ratio	-0.23	0.15	
Maximum Drawdown	-77.1%	-82.2%	
Return over Maximum Drawdown	-0.44	0.35	

7 Conclusion

In this paper, We develop a novel pairs trading framework for twenty Binance USDT-Margined Futures cryptocurrency coins, which combines copula-based and cointegrated-based approaches. The methodology involves setting a reference asset (in this case, BTCUSDT) and identifying other cryptocurrency coins that are cointegrated with it. To investigate the presence of cointegration, we utilize both the EG two-step method and the KSS cointegration test. After ranking the cointegrated coins based on Kendall's Tau correlation coefficients, we select the two assets with the highest correlation. These selected coins are then traded during the one-week trading period and are updated weekly. Trading rules are generated based on the copula conditional probabilities of the spread processes corresponding to the selected assets. We establish various trading triggers and backtest the strategy.

The research findings show that the choice of an appropriate entry threshold in pairs trading has an important impact on the outcomes. By using lower thresholds, both volatility decreases and return increases concurrently. Our pairs trading strategy has shown a high risk-adjusted return and return over maximum drawdown when compared to the buy-and-hold approach. This suggests that our method can be effectively utilized for profitability. The study also emphasizes the effectiveness of pairs trading in the cryptocurrency market and underscores the importance of meticulous coin pair and copula model selection.

References

- Binance Crypto Derivatives* (accessed 2022-11-03). <https://www.binance.com/en/support/faq/crypto-derivatives?c=4&navId=4>.
- Broek, L van den and Zara Sharif (2018). "Cointegration-based pairs trading framework with application to the Cryptocurrency market". Bachelor Thesis, Erasmus University Rotterdam.
- Casella, George and Roger L Berger (2021). *Statistical inference*. Cengage Learning.
- Chen, Huafeng et al. (2019). "Empirical investigation of an equity pairs trading strategy". In: *Management Science* 65.1, pp. 370–389.
- Cherubini, Umberto et al. (2011). *Dynamic copula methods in finance*. John Wiley & Sons.
- Czado, Claudia (2019). "Analyzing dependent data with vine copulas". In: *Lecture Notes in Statistics*, Springer.
- Enders, Walter and Pierre L Siklos (2001). "Cointegration and threshold adjustment". In: *Journal of Business & Economic Statistics* 19.2, pp. 166–176.

- Engle, Robert F and Clive WJ Granger (1987). "Co-integration and error correction: representation, estimation, and testing". In: *Econometrica: journal of the Econometric Society*, pp. 251–276.
- Ferreira, Luan (2008). "New tools for spread trading". In: *Futures* 37.12, pp. 38–41.
- Gatev, Evan, William N Goetzmann, and K Geert Rouwenhorst (2006). "Pairs trading: Performance of a relative-value arbitrage rule". In: *The Review of Financial Studies* 19.3, pp. 797–827.
- Gudendorf, Gordon and Johan Segers (2010). "Extreme-value copulas". In: *Copula theory and its applications*. Springer, pp. 127–145.
- Hansen, Bruce E and Byeongseon Seo (2002). "Testing for two-regime threshold cointegration in vector error-correction models". In: *Journal of econometrics* 110.2, pp. 293–318.
- Harris, Lawrence (1997). "Decimalization: A review of the arguments and evidence". In: *Unpublished working paper, University of Southern California*.
- Huck, Nicolas (2015). "Pairs trading: does volatility timing matter?" In: *Applied Economics* 47.57, pp. 6239–6256.
- Johansen, Søren (1991). "Estimation and hypothesis testing of cointegration vectors in Gaussian vector autoregressive models". In: *Econometrica: journal of the Econometric Society*, pp. 1551–1580.
- Kakushadze, Zura and Willie Yu (2019). "Altcoin-Bitcoin Arbitrage". In: *Bulletin of Applied Economics* 6.1, pp. 87–110.
- Kapetanios, George (2005). "Unit-root testing against the alternative hypothesis of up to m structural breaks". In: *Journal of Time Series Analysis* 26.1, pp. 123–133.
- Kapetanios, George, Yongcheol Shin, and Andy Snell (2003). "Testing for a unit root in the nonlinear STAR framework". In: *Journal of Econometrics* 112.2, pp. 359–379.
- (2006). "Testing for cointegration in nonlinear smooth transition error correction models". In: *Econometric Theory*, pp. 279–303.
- Keshavarz Haddad, GholamReza and Hassan Talebi (2023). "The profitability of pair trading strategy in stock markets: Evidence from Toronto stock exchange". In: *International Journal of Finance & Economics* 28.1, pp. 193–207.
- Klement, Erich Peter, Radko Mesiar, and Endre Pap (2002). "Invariant copulas". In: *Kybernetika* 38.3, pp. 275–286.
- Krauss, Christopher (2017). "Statistical arbitrage pairs trading strategies: Review and outlook". In: *Journal of Economic Surveys* 31.2, pp. 513–545.
- Krauss, Christopher and Johannes Stübinger (2017). "Non-linear dependence modelling with bivariate copulas: Statistical arbitrage pairs trading on the S&P 100". In: *Applied Economics* 49.52, pp. 5352–5369.
- Leung, Tim and Hung Nguyen (2019). "Constructing cointegrated cryptocurrency portfolios for statistical arbitrage". In: *Studies in Economics and Finance* 36.3, pp. 581–599.

- Liew, Rong Qi and Yuan Wu (2013). “Pairs trading: A copula approach”. In: *Journal of Derivatives & Hedge Funds* 19.1, pp. 12–30.
- Nelsen, Roger B (2007). *An introduction to copulas*. Springer Science & Business Media.
- Phillips, Peter CB and Sam Ouliaris (1990). “Asymptotic properties of residual based tests for cointegration”. In: *Econometrica: journal of the Econometric Society*, pp. 165–193.
- Pritchard, Brendan Padraic Anson (2018). “Digital asset arbitrage”. PhD thesis. Fundação Getulio Vargas’s Sao Paulo School of Business Administration.
- Rad, Hossein, Rand Kwong Yew Low, and Robert Faff (2016). “The profitability of pairs trading strategies: distance, cointegration and copula methods”. In: *Quantitative Finance* 16.10, pp. 1541–1558.
- Silva, Fernando AB Sabino da, Flavio A Ziegelmann, and João F Caldeira (2023). “A pairs trading strategy based on mixed copulas”. In: *The Quarterly Review of Economics and Finance* 87, pp. 16–34.
- Stander, Yolanda, Daniël Marais, and Ilse Botha (2013). “Trading strategies with copulas”. In: *Journal of Economic and Financial Sciences* 6.1, pp. 83–107.
- Tadi, Masood and Irina Kortchemski (2021). “Evaluation of dynamic cointegration-based pairs trading strategy in the cryptocurrency market”. In: *Studies in Economics and Finance*.
- Teräsvirta, Timo (1994). “Specification, estimation, and evaluation of smooth transition autoregressive models”. In: *Journal of the American Statistical Association* 89.425, pp. 208–218.
- Vidyamurthy, Ganapathy (2004). *Pairs Trading: quantitative methods and analysis*. Vol. 217. John Wiley & Sons.
- Xie, Wenjun, Rong Qi Liew, et al. (2016). “Pairs trading with copulas”. In: *The Journal of Trading* 11.3, pp. 41–52.
- Xie, Wenjun and Yuan Wu (2013). “Copula-based pairs trading strategy”. In: *Asian Finance Association (AsFA)*.
- Zivot, Eric and Donald W K Andrews (2002). “Further evidence on the great crash, the oil-price shock, and the unit-root hypothesis”. In: *Journal of business & economic statistics* 20.1, pp. 25–44.

Appendix

Table VII: Binance USDT-Margined Futures contracts used in the research

Symbol	Underlying Crypto	Min. Trade Amount	Max. Leverage
BTCUSDT	Bitcoin	0.001 BTC	125x
ETHUSDT	Ethereum	0.001 ETH	100x
BCHUSDT	Bitcoin Cash	0.001 BCH	75x
XRPUSDT	Ripple	0.1 XRP	75x
EOSUSDT	EOS.IO	0.1 EOS	75x
LTCUSDT	Litecoin	0.001 LTC	75x
TRXUSDT	TRON	1 TRX	50x
ETCUSDT	Ethereum Classic	0.01 ETC	75x
LINKUSDT	Chainlink	0.01 LINK	75x
XLMUSDT	Stellar	1 XLM	50x
ADAUSDT	Cardano	1 ADA	75x
XMRUSDT	Monero	0.001 XMR	50x
DASHUSDT	Dash	0.001 DASH	50x
ZECUSDT	Zcash	0.001 ZEC	50x
XTZUSDT	Tezos	0.1 XTZ	50x
ATOMUSDT	Cosmos	0.01 ATOM	25x
BNBUSDT	Binance Coin	0.01 BNB	75x
ONTUSDT	Ontology	0.1 ONT	50x
IOTAUSDT	IOTA	0.1 IOTA	25x
BATUSDT	Basic Attention Token	0.1 BAT	50x

Table VIII: Selected pairs using unit-root tests and Kendall's τ coefficient (part I)

Week	ADF Unit-Root Test Result			KSS Unit-Root Test Result		
	Pair	P-Value (S_1)	P-Value (S_2)	Pair	t-stat (S_1)	t-stat (S_2)
1	ETH-LTC	0.075	0.070	BCH-ETC	-2.72	-1.96
2	LTC-BCH	0.023	0.010	LTC-BCH	-3.07	-2.09
3	ETH-LTC	0.075	0.037	ETH-LTC	-2.10	-2.90
4	ETH-LTC	0.095	0.022	ETH-ETC	-2.35	-2.64
5	LTC-EOS	0.096	0.082	LTC-BCH	-2.48	-2.66
6	LINK-TRX	0.090	0.069	LTC-BCH	-2.44	-2.46
7	LINK-TRX	0.097	0.051	LTC-BCH	-2.44	-2.32
8	ETH-LTC	0.010	0.010	ETH-BCH	-2.66	-2.47
9	ETH-LTC	0.042	0.022	LTC-BNB	-3.43	-4.80
10	ETH-BCH	0.097	0.073	LTC-BCH	-3.03	-2.02
11	LTC-BCH	0.079	0.010	LTC-BCH	-2.58	-2.63
12	ATOM-ADA	0.010	0.010	BCH-ATOM	-1.96	-3.02
13	BAT-ONT	0.095	0.021	ADA-XTZ	-2.61	-2.07
14	ADA-EOS	0.064	0.010	LTC-ADA	-2.15	-3.23
15	EOS-LTC	0.044	0.027	EOS-LTC	-2.33	-2.14
16	BAT-IOTA	0.022	0.010	TRX-BNB	-2.56	-2.06
17	BNB-TRX	0.067	0.036	BCH-BNB	-1.96	-3.08
18	BCH-LTC	0.082	0.086	TRX-IOTA	-3.08	-2.64
19	ETH-TRX	0.059	0.023	DASH-BCH	-2.77	-3.42
20	ETH-LTC	0.078	0.088	BCH-EOS	-2.07	-2.46
21	TRX-LTC	0.010	0.010	TRX-LTC	-2.17	-2.05
22	-	-	-	LTC-ETC	-1.99	-2.39
23	-	-	-	BNB-LINK	-2.10	-2.06
24	ETH-BNB	0.072	0.092	ETH-BNB	-2.05	-2.93
25	LTC-EOS	0.018	0.018	LTC-BNB	-1.97	-2.32
26	ETH-LTC	0.085	0.010	LTC-XRP	-2.50	-3.91
27	BNB-DASH	0.041	0.043	XRP-DASH	-2.02	-2.48
28	ETH-BCH	0.060	0.096	ETH-BCH	-1.95	-2.35
29	BNB-LTC	0.068	0.015	BNB-LTC	-2.42	-2.99
30	BNB-EOS	0.044	0.030	BNB-EOS	-2.22	-3.77
31	ETH-LINK	0.030	0.015	LTC-LINK	-2.52	-2.72
32	ETH-LINK	0.058	0.096	ETH-LTC	-2.46	-1.95
33	LTC-XRP	0.063	0.032	LTC-BNB	-2.73	-1.93
34	EOS-XRP	0.049	0.041	EOS-LTC	-2.36	-1.97
35	ETH-EOS	0.061	0.041	EOS-BNB	-2.15	-3.96
36	BNB-LINK	0.090	0.052	LTC-ADA	-3.91	-2.53
37	ETH-BNB	0.046	0.063	ETH-ETC	-2.50	-1.93
38	ETH-LINK	0.013	0.063	BCH-ONT	-2.49	-2.10
39	DASH-ONT	0.079	0.010	LTC-DASH	-2.08	-2.20
40	LTC-BNB	0.091	0.090	DASH-BNB	-2.18	-2.07
41	ETC-DASH	0.091	0.035	ETC-DASH	-1.95	-2.52
42	BCH-LTC	0.017	0.027	BCH-LTC	-3.15	-2.30
43	ETH-ETC	0.075	0.010	ETC-EOS	-3.81	-2.44
44	ETH-ETC	0.045	0.010	ETH-ETC	-2.66	-3.13
45	EOS-ETC	0.021	0.010	EOS-ETC	-4.45	-3.32
46	EOS-ETC	0.027	0.010	EOS-ETC	-2.46	-3.56
47	ETC-XRP	0.055	0.069	ETC-XRP	-3.07	-2.75
48	ETH-LTC	0.089	0.088	LTC-DASH	-3.29	-4.01
49	ETH-ETC	0.026	0.010	ETH-ETC	-2.43	-2.18
50	ETH-ETC	0.070	0.010	ETH-ETC	-2.96	-2.75
51	ETC-LTC	0.013	0.020	ETH-ETC	-2.43	-3.19
52	LTC-EOS	0.010	0.050	LTC-EOS	-3.39	-2.61

Table IX: Selected pairs using unit-root tests and Kendall's τ coefficient (part II)

Week	ADF Unit-Root Test Result			KSS Unit-Root Test Result		
	Pair	P-Value (S_1)	P-Value (S_2)	Pair	t-stat (S_1)	t-stat (S_2)
53	EOS-XRP	0.032	0.010	EOS-XRP	-2.43	-3.51
54	-	-	-	LTC-BNB	-2.03	-2.24
55	DASH-XLM	0.041	0.010	EOS-LTC	-2.20	-2.01
56	ETH-BNB	0.049	0.016	ETH-LTC	-2.69	-2.77
57	ETH-BNB	0.010	0.018	ETH-BNB	-2.83	-3.20
58	ETH-BNB	0.028	0.023	ETH-BNB	-2.78	-4.00
59	ETH-BAT	0.024	0.068	ETH-BAT	-3.41	-1.96
60	ETH-BNB	0.020	0.020	ETH-EOS	-2.41	-1.95
61	ETH-BNB	0.085	0.010	BNB-ADA	-3.02	-2.04
62	BNB-ONT	0.010	0.083	ETH-BNB	-1.95	-2.79
63	BNB-LTC	0.060	0.022	LTC-LINK	-3.47	-2.32
64	LINK-ONT	0.029	0.095	XRP-LINK	-2.14	-2.34
65	XRP-XLM	0.010	0.095	XRP-ADA	-3.35	-2.16
66	ETH-BNB	0.010	0.047	ETH-BNB	-3.15	-3.10
67	ETH-BNB	0.041	0.010	ETH-BNB	-3.40	-3.23
68	ETH-BNB	0.017	0.010	ETH-BNB	-3.56	-3.42
69	-	-	-	BCH-XLM	-2.27	-2.03
70	BNB-ETC	0.031	0.033	LINK-BNB	-2.07	-2.39
71	LINK-ETC	0.070	0.062	ETC-BNB	-2.29	-2.42
72	LINK-EOS	0.046	0.025	BNB-LINK	-2.28	-2.37
73	DASH-ETC	0.037	0.044	EOS-DASH	-1.96	-3.22
74	ONT-ZEC	0.026	0.010	BNB-DASH	-2.56	-1.95
75	ZEC-BCH	0.047	0.041	ETH-BNB	-2.36	-2.16
76	ETH-XTZ	0.010	0.029	ETH-ZEC	-2.83	-2.52
77	ETH-BNB	0.057	0.064	ETH-BNB	-2.77	-2.42
78	ETH-BNB	0.022	0.010	ETH-BNB	-3.17	-2.19
79	BNB-LTC	0.021	0.037	BNB-LTC	-2.93	-3.50
80	BNB-LTC	0.017	0.061	BNB-LTC	-3.08	-2.09
81	ETH-LTC	0.010	0.010	LTC-LINK	-3.25	-2.52
82	LTC-XRP	0.014	0.017	LTC-XRP	-2.68	-4.02
83	LTC-DASH	0.010	0.047	LTC-DASH	-4.27	-2.64
84	ETH-LTC	0.044	0.010	ETH-LTC	-2.40	-3.95
85	ETH-IOTA	0.090	0.010	IOTA-DASH	-2.63	-2.87
86	BAT-IOTA	0.010	0.010	BAT-DASH	-3.54	-1.96
87	BNB-ETC	0.023	0.068	BAT-ETC	-2.49	-2.85
88	BNB-TRX	0.020	0.084	DASH-BAT	-2.35	-2.43
89	ETH-DASH	0.090	0.010	ETH-DASH	-2.71	-4.19
90	ETH-DASH	0.010	0.014	ETH-DASH	-2.95	-2.78
91	ETH-BAT	0.095	0.032	ETH-DASH	-2.54	-2.26
92	ETH-BAT	0.010	0.084	ETH-BAT	-3.08	-2.15
93	BNB-LTC	0.014	0.023	BNB-LTC	-2.22	-3.47
94	BCH-DASH	0.010	0.046	BCH-DASH	-3.36	-2.30
95	ETH-BCH	0.051	0.034	BCH-DASH	-6.40	-2.29
96	ETH-ADA	0.081	0.010	ETH-ADA	-2.44	-2.53
97	ETH-BAT	0.042	0.053	ETH-ADA	-2.47	-2.47
98	ADA-BAT	0.039	0.010	ETH-ADA	-1.97	-3.08
99	XTZ-BAT	0.010	0.010	ETH-ATOM	-1.98	-2.15
100	ETH-BCH	0.010	0.099	ETH-ONT	-2.20	-3.13
101	-	-	-	LINK-BCH	-2.47	-2.00
102	ETH-BNB	0.057	0.033	ETH-LTC	-2.48	-2.00
103	BNB-LINK	0.010	0.020	BNB-LINK	-3.85	-3.27
104	LINK-DASH	0.061	0.100	LINK-ONT	-2.49	-2.31

Table X: Bivariate Archimedean copulas distributions

Name	Bivariate Copula Distribution $C(u_1, u_2)$ ($\overline{u_i} = 1 - u_i$)	Generator $\phi(t)$	Parameters
Clayton	$\left[\max \left(u_1^{-\theta} + u_2^{-\theta} - 1, 0 \right) \right]^{-1/\theta}$	$\frac{1}{\theta} (t^{-\theta} - 1)$	$\theta > 0$
Gumbel	$\exp \left[- \left[(-\ln u_1)^\theta + (-\ln u_2)^\theta \right]^{1/\theta} \right]$	$(-\ln(t))^\theta$	$\theta \geq 1$
Frank	$-\theta^{-1} \ln \left[1 + (e^{-\theta} - 1)^{-1} (e^{-\theta u_1} - 1) (e^{-\theta u_2} - 1) \right]$	$-\ln \left(\frac{e^{-\theta t} - 1}{e^{-\theta} - 1} \right)$	$\theta \in \mathbb{R} \setminus \{0\}$
Joe	$1 - \left[(1 - u_1)^\theta + (1 - u_2)^\theta - (1 - u_1)^\theta (1 - u_2)^\theta \right]^{1/\theta}$	$-\ln \left(1 - (1 - t)^\theta \right)$	$\theta \geq 1$
BB1	$\left[1 + \left[(u_1^{-\theta} - 1)^\delta + (u_2^{-\theta} - 1)^\delta \right]^{1/\delta} \right]^{-1/\theta}$	$(t^{-\theta} - 1)^\delta$	$\theta > 0, \delta \geq 1$
BB6	$1 - \left[1 - \exp \left(- \left[[-\ln(1 - \overline{u_1}^\theta)]^\delta + [-\ln(1 - \overline{u_2}^\theta)]^\delta \right]^{1/\delta} \right) \right]^{1/\theta}$	$(\ln(1 - (1 - t)^\theta))^\delta$	$\theta \geq 1, \delta \geq 1$
BB7	$1 - \left[1 - \left((1 - \overline{u_1}^\theta)^{-\delta} + (1 - \overline{u_2}^\theta)^{-\delta} - 1 \right)^{-1/\delta} \right]^{1/\theta}$	$(1 - (1 - t)^\theta)^{-\delta} - 1$	$\theta \geq 1, \delta > 0$
BB8	$\delta^{-1} \left[1 - \left(1 - \left[1 - (1 - \delta)^\theta \right]^{-1} \left[1 - (1 - \delta u_1)^\theta \right] \left[1 - (1 - \delta u_2)^\theta \right] \right)^{1/\theta} \right]$	$-\ln \left(\frac{1 - (1 - \delta t)^\theta}{1 - (1 - \delta)^\theta} \right)$	$\theta \geq 1, 0 < \delta \leq 1$

FFA Working Paper Series

2019

1. Milan Fičura: Forecasting Foreign Exchange Rate Movements with k-Nearest-Neighbor, Ridge Regression and Feed-Forward Neural Networks.

2020

1. Jiří Witzany: Stressing of Migration Matrices for IFRS 9 and ICAAP Calculations.
2. Matěj Maivald, Petr Teplý: The impact of low interest rates on banks' non-performing loans.
3. Karel Janda, Binyi Zhang: The impact of renewable energy and technology innovation on Chinese carbon dioxide emissions.
4. Jiří Witzany, Anastasiia Kozina: Recovery process optimization using survival regression.

2021

1. Karel Janda, Oleg Kravtsov: Banking Supervision and Risk-Adjusted Performance in the Host Country Environment.
2. Jakub Drahekoupil: Variance Gamma process in the option pricing model.
3. Jiří Witzany, Olga Pastiranová: IFRS 9 and its behaviour in the cycle: The evidence on EU Countries.

2022

1. Karel Janda, Ladislav Kristoufek, Binyi Zhang: Return and volatility spillovers between Chinese and U.S. Clean Energy Related Stocks: Evidence from VAR-MGARCH estimations.
2. Lukáš Fiala: Modelling of mortgage debt's determinants: The case of the Czech Republic.
3. Olga Jakubíková: Profit smoothing of European banks under IFRS 9.

4. Milan Fičura, Jiří Witzany: Determinants of NMD Pass-Through Rates in Eurozone Countries.
5. Sophio Togonidze, Evžen Kočenda: Macroeconomic Responses of Emerging Market Economies to Oil Price Shocks: An Analysis by Region and Resource Profile.
6. Jakub Drahekoupil: Application of the XGBoost algorithm and Bayesian optimization for the Bitcoin price prediction during the COVID-19 period.
7. Karel Janda, Binyi Zhang: Econometric estimation of green bond premiums in the Chinese market.
8. Shahriyar Aliyev, Evžen Kočenda: ECB monetary policy and commodity prices.
9. Sophio Togonidze, Evžen Kočenda: Macroeconomic implications of oil-price shocks to emerging economies: a Markov regime-switching approach.

2023

1. Jiří Witzany, Milan Fičura: Machine Learning Applications for the Valuation of Options on Non-liquid Option Markets.
2. David Mazáček, Jiří Panoš: Key determinants of new residential real estate prices in Prague.
3. Milan Fičura: Impact of size and volume on cryptocurrency momentum and reversal.
4. Masood Tadi, Jiří Witzany: Copula-Based Trading of Cointegrated Cryptocurrency Pairs.

All papers can be downloaded at: wp.ffu.vse.cz

Contact e-mail: ffawp@vse.cz

