

Problem 1

Solution

- (a) Because $\forall i$, f_i is positive non-decreasing, $\prod_{i=1}^{k'} f_i$, $\forall 1 \leq k' \leq k$ is also non-decreasing. Thus,

$$\mathbb{E} \left[\prod_{i=1}^k f_i(\mathbf{X}) \right] \leq \mathbb{E} \left[\prod_{i=1}^{k-1} f_i(\mathbf{X}) \right] \mathbb{E} [f_k(\mathbf{X})] \leq \cdots \leq \prod_{i=1}^k \mathbb{E} [f_i(\mathbf{X})].$$

- (b) NA r.v.s $\{X_1, \dots, X_n\}$ may not be independent. If we can prove that Chernoff's trick also holds for NA r.v.s, then Hoeffding's inequality holds too.

$$P\left(\sum_{i=1}^n X_i \geq a\right) \leq \frac{\mathbb{E} \left[\prod_{i=1}^n e^{\theta X_i} \right]}{e^{\theta a}} \leq \frac{\prod_{i=1}^n \mathbb{E} [e^{\theta X_i}]}{e^{\theta a}}$$

where the last derivation uses the result in (a).

Problem 2

Solution

- (a) Lemmas: 1. zero-one principle. 2. union of independent groups.
- (b) Let Y_i be an indicator random variable to indicate if the i -th bin is non-empty. We have $Y_i \sim \text{Ber}(1 - (\frac{n-1}{n})^m)$, and $\mathbb{E} [Y_i] = 1 - (\frac{n-1}{n})^m$. Lemma: mapping. Thus, $\{Y_1, \dots, Y_n\}$ is NA. With the results in Problem 1 (b), we have $\forall o > 0$,

$$P\left(O - \mathbb{E}[O] = \sum_{i=1}^n (Y_i - \mathbb{E}[Y_i]) \geq o\right) \leq \exp\left(\sum_{i=1}^n \frac{(b_i - a_i)^2}{8} \theta^2 - \theta o\right) = \exp\left(\frac{n}{8} \theta^2 - \theta o\right) \leq e^{\frac{-2o^2}{n}}$$

Problem 3

Solution

Let $Z_i \in \{1, 2, 3, \dots, n\}$ be the ID of the bin into which the i -th ball is put. Let f be a function such that $f(Z_1, Z_2, Z_3, \dots, Z_m)$ is the number of non-empty bins. Obviously, Z_i 's are independent, and $|f(Z_1, \dots, Z_i, \dots, Z_m) - f(Z_1, \dots, Z'_i, \dots, Z_m)| < 1$. Thus, McDiarmid's inequality applies:

$$P(O - \mathbb{E}[O] = f(Z_1, Z_2, Z_3, \dots, Z_m) - \mathbb{E}[f(Z_1, Z_2, Z_3, \dots, Z_m)] \geq o) \leq e^{\frac{-2o^2}{n}},$$

which is the same as the Hoeffding bound.

Problem 4

Solution

(a) We have $\mathbb{E}[\exp(-\theta X)] \leq \exp(\nu^2 \theta^2 / 2)$. Thus for any $\theta > 0$,

$$\begin{aligned} P(|X| > t) &= P(X > t) + P(-X > t) = P(e^{\theta X} > e^{\theta t}) + P(e^{-\theta X} > e^{\theta t}) \\ &\leq \frac{\mathbb{E}[e^{\theta X}]}{e^{\theta t}} + \frac{\mathbb{E}[e^{-\theta X}]}{e^{\theta t}} \leq 2 \exp(\nu^2 \theta^2 / 2 - \theta t) \leq 2 \exp(-\frac{t^2}{2\nu^2}) \end{aligned}$$

(b) Let the CDF of $|X|$ be F . Then, $F(x) = 0, \forall x \leq 0$. Because $g(x) = x^k$ is an increasing function, we have

$$\mathbb{E}[|X|^k] = \int_0^\infty g'(x)P(|X| > x)dx \leq 2 \int_0^\infty kx^{k-1} \exp(-\frac{x^2}{2\nu^2})dx.$$

Substitute $\frac{x^2}{2\nu^2}$ with y , and we have

$$\mathbb{E}[|X|^k] \leq 2k \int_0^\infty \nu^2 (2\nu^2 y)^{k/2-1} \exp(-y)dy = k(2\nu^2)^{k/2} \Gamma(\frac{k}{2}).$$

Problem 5

Solution

First, we partition the indices into K index group such that $G_k = \{j : X_j \in A_k\}$. We also have for any $f \in F$ and $j_1, j_2 \in G_i, f(X_{j_1}) = f(X_{j_2})$. Thus,

$$\begin{aligned} R_n(F(X^n)) &= \mathbb{E} \left[\sup_{f \in F} \left| \frac{1}{n} \sum_{i=1}^n \sigma_i f(X_i) \right| \right] = \mathbb{E} \left[\sup_{f \in F} \left| \frac{1}{n} \sum_{k=1}^K \sum_{i \in G_k} \sigma_i f(X_i) \right| \right] = \mathbb{E} \left[\frac{1}{n} \sum_{k=1}^K \left| \sum_{i \in G_k} \sigma_i \right| \right] \\ &\leq \mathbb{E} \left[\frac{1}{n} \sum_{k=1}^K \sqrt{\sum_{i \in G_k} \sigma_i^2} \right] = \mathbb{E} \left[\frac{1}{n} \sum_{k=1}^K \sqrt{n_k} \right] = \frac{1}{n} \sum_{k=1}^K \sqrt{n_k} \end{aligned}$$

Problem 6

Solution

For $n = 3$, we can choose $z_2 = z_1 + 0.55$ and $z_3 = z_2 + 0.55$. It is easy to verify z_1, z_2 , and z_3 can be shattered by F . When $n = 4$, no matter what z_1, z_2, z_3 , and z_4 are, they cannot be assigned values 1, 0, 1, 0. Thus $VC(F) = 3$.

Problem 7

Solution

For n points, we can choose $x_i = 2^i \pi, i = 1, 2, 3, \dots, n$. Then x^n can be shattered by F . Actually, we can find 2^n intervals $(1, \frac{2^n}{2^n-1}), (\frac{2^n}{2^n-1}, \frac{2^n}{2^n-2}), \dots, (\frac{2^n}{1}, \infty)$ such that when θ belongs to different intervals, f assigns different values to x^n .