Lycopene attenuates high-fat diet-induced hepatic steatosis in rats

Wei Zhang[1,2], Li Chen[1] and Hao Wang[3*]
Received: 12 January 2019 / Accepted: 2 April 2019

Abstract. Lycopene attenuates high-fat diet-induced hepatic steatosis in rats.

Introduction. Obesity drives fatty liver. In male C57BL/6 mice fed a Western diet for 16 weeks, carotenoids lowered hepatic lipids (n = 6) [5]. Intraperitoneal injection was used elsewhere [6]. Introduction. Obesity drives fatty liver. In male C57BL/6 mice fed a Western diet for 16 weeks, carotenoids lowered hepatic lipids (n = 6) [5]. Intraperitoneal injection was used elsewhere [6]. Introduction. Obesity drives fatty liver. In male C57BL/6 mice fed a Western diet for 16 weeks, carotenoids lowered hepatic lipids (n = 6) [5]. Intraperitoneal injection was used elsewhere [6]. Introduction. Obesity drives fatty liver. In male C57BL/6 mice fed a Western diet for 16 weeks, carotenoids lowered hepatic lipids (n = 6) [5]. Intraperitoneal injection was used elsewhere [6].

2. Materials and Methods. 2.1. Animals. Forty-eight male Sprague-Dawley rats (6–8 weeks old, 180–200 g) were acclimatized for 1 week and then randomly assigned to four groups (n = 12 each): normal diet (ND), high-fat diet (HFD), HFD with low-dose lycopene (HFD+LL, 10 mg/kg), and HFD with high-dose lycopene (HFD+HL, 40 mg/kg). Lycopene was given orally once daily for 8 weeks. 2.2. Statistics. Values are expressed as means ± SD.

3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls.

References

1. Smith J. Wistar rats for 20 weeks. 2001.