Quercetin improves skeletal muscle function in aged mice

A. Rossi[a], B. Bianchi[b], C. Verdi[a]
Available online 4 May 2023. © 2023 Elsevier Inc. All rights reserved.

Abstract. Quercetin improves skeletal muscle function in aged mice.

1. Introduction. Sarcopenia limits mobility. Earlier work gave quercetin in drinking water for 12 weeks to ICR mice (n = 15) [2]. 1. Introduction. Sarcopenia limits mobility. Earlier work gave quercetin in drinking water for 12 weeks to ICR mice (n = 15) [2]. 1. Introduction. Sarcopenia limits mobility. Earlier work gave quercetin in drinking water for 12 weeks to ICR mice (n = 15) [2]. 1. Introduction. Sarcopenia limits mobility. Earlier work gave quercetin in drinking water for 12 weeks to ICR mice (n = 15) [2].

2. Methods. 2.1. Animal experiments. Female C57BL/6N mice at 10 weeks of age were used. Mice received intraperitoneal injections of quercetin (25 mg/kg) three times per week for 6 weeks. Control mice (VEH) received vehicle only. Ten mice per group were used. Results are shown as mean ± SEM.

3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls.

References

1. Smith J. Wistar rats for 20 weeks. 2001.