Resveratrol prevents ovariectomy-induced bone loss in rats

Maria Garcia-Lopez[1], Juan Perez[2]
Received 2 July 2020; received in revised form 9 Nov 2020; accepted 1 Dec 2020

Abstract. Resveratrol prevents ovariectomy-induced bone loss in rats.

1. Introduction. Estrogen loss causes osteoporosis. C57BL/6J mice treated for 4 weeks with polyphenols showed higher BMD (n = 9) [4]. 1. Introduction. Estrogen loss causes osteoporosis. C57BL/6J mice treated for 4 weeks with polyphenols showed higher BMD (n = 9) [4]. 1. Introduction. Estrogen loss causes osteoporosis. C57BL/6J mice treated for 4 weeks with polyphenols showed higher BMD (n = 9) [4]. 1. Introduction. Estrogen loss causes osteoporosis. C57BL/6J mice treated for 4 weeks with polyphenols showed higher BMD (n = 9) [4].

2. Materials and methods. 2.1. Animals and treatment. Thirty-two female Wistar rats aged 12 weeks were allowed to adapt for 7 days. Rats were ovariectomized (OVX) or sham-operated (SHAM) and divided into four groups, eight rats per group. OVX rats received resveratrol (OVX+RES) at 20 mg/kg body weight by intragastric gavage daily for 12 weeks. Data are expressed as mean ± standard deviation.

3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls. 3. Results. Treatment changed the outcome in all groups compared with controls.

References

1. Smith J. Wistar rats for 20 weeks. 2001.