

# Drug and Alcohol Dependence

## A pilot study of pupil response to light as a digital biomarker of recent cannabis use

--Manuscript Draft--

<b>Manuscript Number:</b>	
<b>Article Type:</b>	Full Length Article
<b>Section/Category:</b>	Epidemiology
<b>Keywords:</b>	pupillary light reflex; pupillometry; cannabis; functional data analysis; substance use detection
<b>Corresponding Author:</b>	Suneeta Godbole, M.S. Colorado School of Public Health Department of Biostatistics & Informatics UNITED STATES
<b>First Author:</b>	Suneeta Godbole, M.S.
<b>Order of Authors:</b>	Suneeta Godbole, M.S. Andrew Leroux, PhD Ashley Brooks-Russell, PhD Prem S. Subramanian, MD, PhD Michael J. Kosnett, MD, PhD Julia Wrobel, PhD
<b>Abstract:</b>	<p><b>Introduction:</b> Given the roadside safety and occupational injury prevention implications associated with cannabis impairment, there is a need for objective and validated measures of recent cannabis use. Pupillary light response may offer an approach for detection.</p> <p><b>Method:</b> 84 participants (mean age: 32, 42% female) with daily, occasional, and no-use cannabis use histories participated in pupillary light response tests before and after smoking cannabis ad libitum or relaxing for 15 minutes (no use). The impact of recent cannabis consumption on trajectories of the pupillary light response was modeled using functional data analysis tools. Logistic regression models for predicting recent cannabis use were compared, and average pupil trajectories across cannabis use groups and times since light test administration were estimated.</p> <p><b>Results:</b> Models revealed small, significant differences in pupil response to light after cannabis use comparing the occasional use group to the no use control group, and similar statistically significant differences in pupil response patterns comparing the daily use group to the no use comparison group. Trajectories of pupillary light response estimated using functional data analysis found that acute cannabis smoking was associated with less initial and sustained pupil constriction compared to no cannabis smoking.</p> <p><b>Discussion:</b> These analyses show the promise of pairing pupillary light response and functional data analysis methods to assess recent cannabis use.</p>
<b>Suggested Reviewers:</b>	<p>Jihui Lee, Ph.D. Assistant Professor, University of Pittsburgh jihui.lee@pitt.edu Dr. Lee has expertise in the statistical field of functional data analysis and experience with digital biomarker.</p> <p>Patricia Di Ciano, PhD Assistant Professor, University of Toronto patricia.diciano@camh.ca Dr. Di Ciano has expertise in the effects of cannabis use.</p> <p>Bruna Brands, PhD Professor, University of Toronto bruna.brands@camh.ca Dr. Brands is an expert in the effects of cannabis use on road safety.</p> <p>Doug Beirness, PhD</p>

	<p>Senior Research Associate, Canadian Centre on Substance Use and Addiction  dbeirness@ccsa.ca  Dr. Beirness has expertise in the effects of impaired driving.</p>
	<p>Jeff Brubacher, MD  Associate Professor, The University of British Columbia  jeff.brubacher@ubc.ca  Dr. Brubacher's research interest focuses broadly on injury prevention with a special interest in drug impaired driving, substance abuse and injury.</p>

Suneeta V. Godbole  
Department of Biostatistics & Informatics  
Colorado School of Public Health  
Fitzsimons Building, 4th Floor  
13001 E. 17th Place  
Mail Stop B119  
Aurora, CO 80045

October 5, 2023

Dear Dr. Shoptaw,

We would like to submit for your review an original research article entitled, “A pilot study of pupil response to light as a digital biomarker of recent cannabis use” for consideration in *Drug and Alcohol Dependence*.

We confirm that this work is original and has not been published elsewhere nor is it currently under consideration for publication elsewhere.

Our paper provides new promising evidence for an objective digital marker of recent cannabis use that could be used to detect recent drug use for roadside safety and injury prevention. In this paper, we identify differences in trajectories of pupil response to light for participants who recently consumed cannabis compared to participants who did not. We are able to show differences in trajectories based solely on the pupillary light response after smoking cannabis regardless of cannabis use history (i.e., not limited by drug tolerance). Results from this pilot study show promise that pupillary light response may be an objective, digital, and validated measure of recent cannabis use, making it an asset in the prevention of cannabis-involved roadside accidents and workplace injuries.

We have no conflicts of interest to disclose.

Please address all correspondence concerning this manuscript to [suneeta.godbole@cuanschutz.edu](mailto:suneeta.godbole@cuanschutz.edu).

Thank you for your consideration.

Sincerely,

Suneeta V. Godbole

**Highlights:**

- Results reveal differences in pupillary light responses between recent cannabis users and non-users.
- Daily and occasional users showed no difference in pupillary light response due to acute cannabis use, suggesting its use as a potential biomarker of recent use, invariant to cannabis tolerance.
- Models using functional data methods have promise for improving prediction over models using single value summaries.

A pilot study of pupil response to light as a digital biomarker of recent cannabis use

**Title:** A pilot study of pupil response to light as a digital biomarker of recent cannabis use

**Authors:** Suneeta Godbole<sup>a\*</sup>, Andrew Leroux<sup>a</sup>, Ashley Brooks-Russell<sup>b</sup>, Prem S. Subramanian<sup>c,d</sup>, Michael J. Kosnett<sup>e,f</sup>, and Julia Wrobel<sup>g</sup>

**Affiliations:**

<sup>a</sup>Department of Biostatistics and Informatics, Colorado School of Public Health, University of Colorado Anschutz Medical Campus, Aurora, Colorado ([suneeta.godbole@cuanschutz.edu](mailto:suneeta.godbole@cuanschutz.edu); [andrew.leroux@cuanschutz.edu](mailto:andrew.leroux@cuanschutz.edu))

<sup>b</sup>Injury and Violence Prevention Center, Colorado School of Public Health, University of Colorado Anschutz Medical Campus, Aurora, Colorado ([ashley.brooks-russell@cuanschutz.edu](mailto:ashley.brooks-russell@cuanschutz.edu))

<sup>c</sup>Departments of Ophthalmology, Neurology, and Neurosurgery, Sue Anschutz-Rodgers Eye Center, CU School of Medicine, University of Colorado Anschutz Medical Campus, Aurora, Colorado ([prem.subramanian@cuanschutz.edu](mailto:prem.subramanian@cuanschutz.edu))

<sup>d</sup>Department of Surgery, Uniformed Services University of the Health Sciences, Bethesda, Maryland <sup>5</sup>Department of Medicine, CU School of Medicine, University of Colorado Anschutz Medical Campus, Aurora, Colorado

<sup>e</sup>Department of Medicine, CU School of Medicine, University of Colorado Anschutz Medical Campus, Aurora, Colorado ([michael.kosnett@cuanschutz.edu](mailto:michael.kosnett@cuanschutz.edu))

<sup>f</sup>Department of Environmental and Occupational Health, Colorado School of Public Health, University of Colorado Anschutz Medical Campus, Aurora, Colorado

<sup>g</sup>Departement of Biostatistics and Bioinformatics, Emory University, Atlanta, Georgia ([julia.wrobel@emory.edu](mailto:julia.wrobel@emory.edu))

**\*Corresponding Author Email:** [suneeta.godbole@cuanschutz.edu](mailto:suneeta.godbole@cuanschutz.edu)

**Address:** Department of Biostatistics & Informatics  
Colorado School of Public Health  
Fitzsimons Building, 4th Floor  
13001 E. 17th Place  
Mail Stop B119  
Aurora, CO 80045

**FUNDING:** This work was supported by the Colorado Department of Public Health and Environment and the National Institutes of Health (R01DA049800).

**Abstract:**

**Introduction:** Given the roadside safety and occupational injury prevention implications associated with cannabis impairment, there is a need for objective and validated measures of recent cannabis use. Pupillary light response may offer an approach for detection.

**Method:** 84 participants (mean age: 32, 42% female) with daily, occasional, and no-use cannabis use histories participated in pupillary light response tests before and after smoking cannabis *ad libitum* or relaxing for 15 minutes (no use). The impact of recent cannabis consumption on trajectories of the pupillary light response was modeled using functional data analysis tools.

Logistic regression models for predicting recent cannabis use were compared, and average pupil trajectories across cannabis use groups and times since light test administration were estimated.

**Results:** Models revealed small, significant differences in pupil response to light after cannabis use comparing the occasional use group to the no use control group, and similar statistically significant differences in pupil response patterns comparing the daily use group to the no use comparison group. Trajectories of pupillary light response estimated using functional data analysis found that acute cannabis smoking was associated with less initial and sustained pupil constriction compared to no cannabis smoking.

**Discussion:** These analyses show the promise of pairing pupillary light response and functional data analysis methods to assess recent cannabis use.

**KEYWORDS:** pupillary light reflex, pupillometry, cannabis, functional data analysis, substance use detection

## 1. INTRODUCTION:

According to the National Survey on Drug Use and Health, the rates of cannabis consumption have increased in adults over 26, and adults aged 18-25, from 4.0% to 7.9% and from 17.3% to 22.1% from 2002 to 2017, respectively (*Key Substance Use and Mental Health Indicators in the United States: Results from the 2017 National Survey on Drug Use and Health, 2017*). Along with increases in consumption, there have been increases in cannabis-involved motor vehicle fatalities from 9.0% in 2000 to 21.5% in 2018 (Lira et al., 2021; Myran et al., 2023). Although there is mixed evidence with regard to risk of occupational injury, with temporality of exposure being a major limitation of the extant literature (Biasutti et al., 2020; Zhang et al., 2020), cannabis consumption at or before work is of concern to employers, especially for employees involved in safety sensitive tasks. An objective, easy to obtain biomarker of recent cannabis use may be of value in field assessments, such as in the context of investigation of motor vehicle crashes and occupational incidents.

Current methods used to enforce existing regulations on drug impaired driving have multiple limitations for assessing recent cannabis use and impairment. The Standardized Field Sobriety Test is a general test for alcohol and drug impairment, comprised of the horizontal gaze nystagmus, walk and turn and one-leg stands (*Drug Evaluation and Classification (Preliminary School): Participant Manual, 2015*). While shown to be an accurate and reliable assessment for alcohol impairment, it has limited ability to identify recent cannabis use (Downey et al., 2012). In addition, many assessment tests have shown a reduction in effectiveness when administered to frequent cannabis users due to drug tolerance, leading to potential false negative results for frequent users (Arkell et al., 2021; Wurz and DeGregorio, 2022). Many states and countries reference drug levels in the blood as a threshold for impairment, much like the .08% blood

1 A pilot study of pupil response to light as a digital biomarker of recent cannabis use

2  
3  
4 78 alcohol concentration level used as a *per se* definition of alcohol impairment in the U.S. Specific  
5  
6 79 to cannabis, the parallel would be the blood level of delta-9-THC; however predictive models  
7  
8 80 have better performance in participants abstaining for several days compared to those who  
9  
10 81 exhibit more frequent or daily use (Burt et al., 2021). This is in part due to the fact that frequent  
11  
12 82 users can maintain elevated levels of blood THC for days or weeks after consumption; as such,  
13  
14 83 frequent cannabis users may have a blood test positive for THC even if they have not recently  
15  
16 84 smoked cannabis (Burt et al., 2021). Given the limitations of blood THC levels and existing  
17  
18 85 roadside assessments, there is a need for the development of objective markers of recent  
19  
20 86 cannabis use and impairment from cannabis use.  
21  
22  
23  
24  
25

26  
27 87 Drug Recognition Experts, specially trained law enforcement officers, have included  
28  
29 88 pupillary and ocular signs as indicators of the pharmacodynamic effects of drugs and alcohol  
30  
31 89 (*Drug Evaluation and Classification (Preliminary School): Participant Manual*, 2015; Richman  
32  
33 90 et al., 2004). They may examine pupil size under illumination ranging from near total darkness to  
34  
35 91 bright light and assess the pupillary light reflex, which consists of constriction in response to  
36  
37 92 visible light. This is similar to the pupillary light response test that is performed clinically to  
38  
39 93 assess central nervous system function and acute drug effects. This test is administered by  
40  
41 94 shining a light in the eye of the participant and measuring pupil size over the course of several  
42  
43 95 seconds after the light is turned off. Studies assessing pupil size in cannabis users have yielded  
44  
45 96 inconsistent results (Brown et al., 1977; Campobasso et al., 2020; Fant, 1998; Merzouki et al.,  
46  
47 97 2008; Ortiz-Peregrina et al., 2020; Shahidi Zandi et al., 2021; Stark et al., 2003). Studies  
48  
49 98 examining pupillary light response using device recorded, light-induced, pupil constriction have  
50  
51 99 shown reduction in pupil diameter after cannabis use (Campobasso et al., 2020; Fant, 1998).  
52  
53  
54  
55  
56  
57 100 Moreover, detailed assessment of the entire pupillary light response trajectory following acute  
58  
59  
60  
61  
62  
63  
64  
65



2  
3  
4 101 cannabis consumption is lacking. If the pattern of pupillary response to light were found to be  
5  
6 102 indicative of recent cannabis use, or impairment from cannabis use, especially in the context of  
7  
8  
9 103 chronic frequent use, it could contribute to the assessment of impaired driving or have utility in  
10  
11 104 investigations of cannabis use in the workplace and other safety sensitive settings.  
12  
13

14 105 Figure 1 shows a typical pupillary response to light during the light reflex test, which we  
15  
16 106 refer to as a *pupillary light response trajectory* throughout the paper. After the light is shined the  
17  
18  
19 107 pupil begins to constrict in size until it reaches a minimum, called the *point of minimal*  
20  
21 108 *constriction*, then it begins to increase in size back towards its original diameter. The area under  
22  
23 109 the curve from the point of minimal constriction to the end of the light response test is known as  
24  
25  
26 110 the *rebound dilation*.  
27  
28

29 111 Recently, Steinhart et al (Steinhart et al., 2023) found that acute cannabis smoking was  
30  
31 112 significantly associated with diminished pupillary constriction during a light response test  
32  
33 113 conducted using infrared videography. Both occasional and daily cannabis users displayed this  
34  
35  
36 114 response, to a similar extent, compared to non-using control subjects. However, the findings of  
37  
38  
39 115 Steinhart et al, utilized single number summaries, such as point of minimal constriction,  
40  
41 116 extracted from the full pupillary response trajectories. Ignoring these trajectories results in a loss  
42  
43 117 of information that could potentially be utilized to better discriminate between recent cannabis  
44  
45  
46 118 use and no cannabis use, regardless of cannabis use history (daily versus occasional).  
47  
48  
49 119 Additionally, significant differences in the extent and pattern of pupillary constriction were only  
50  
51 120 found after adjusting for pre-smoking values, which undermines the utility in field applications  
52  
53  
54 121 where baseline measurements may be unavailable.  
55

56 122 The primary goal of this paper is to investigate the full pupillary light response  
57  
58  
59 123 trajectories collected in the study conducted by Steinhart et al (Steinhart et al., 2023) as  
60  
61  
62  
63  
64  
65

A pilot study of pupil response to light as a digital biomarker of recent cannabis use

predictors of recent cannabis use, irrespective of pre-smoking information. Our analysis uses tools from a field of statistics known as functional data analysis (FDA). The main conceptual underpinning of FDA is to model the whole pupillary light response trajectory as a unit of observation, leveraging information contained in the temporal structure of the data and estimating time-specific effects. This approach utilizes maximal information which is lost when only considering single number predictors such as point of minimal constriction and rebound dilation (Goldsmith et al., 2011; Ramsay and Silverman, 2005). In this analysis, we will use FDA modeling techniques to accomplish the following objectives. We first use the full pupil light response trajectories to predict recent cannabis use as compared to no use. We next examine the impact of cannabis use history on the pupil response trajectories by comparing participants with a history of no recent cannabis use, occasional cannabis use, and daily cannabis use. Finally, we extract expected pupillary light response trajectories at 60, 65, and 70 minutes after cannabis use to explore how pupil response may change over time as a preliminary exploration of its utility.

## **2. METHODS:**

### *2.1 Sample Information:*

Data are from of a larger study examining effects of acute cannabis consumption among participants with occasional and daily cannabis use histories, to understand differences due to tolerance. Healthy adults were recruited using local ads, flyers, and email lists in the Denver area. Key eligibility criteria included ages 25 to 45 years old and willingness to use smoke cannabis and participate in study assessments. Key exclusion criteria include history of drug or alcohol dependence, currently pregnant, and body mass index above 35. Participants were recruited into one of three groups according to their history of cannabis use. Daily cannabis consumption was defined as smoking or vaping a cannabis flower product at least one time per

A pilot study of pupil response to light as a digital biomarker of recent cannabis use

day, every day of the week for 30 days prior to enrollment. Occasional consumption was defined as smoking or vaping cannabis flower product on at least one day but no more than two days per week in the 30 days prior to enrollment. No cannabis consumption was defined as not having used cannabis in the month prior to enrollment.

Participants were instructed not to smoke cannabis for at least 8 hours, and not to use edible cannabis for at least 12 hours, before their data collection visit. This was verified by review of a history of the participant's cannabis use taken on the day of the visit. Each participant completed an alcohol breath test (Lifoloc FC10™) to screen for acute alcohol use, provided a urine sample to test for illicit drug use or use of prescription drugs not prescribed (30 mL Alere brand 13-panel iCup®).

The study utilized a within-subjects design comparing their pupillary light response before and after cannabis use. The cannabis use was observational in nature in that participants brought their own product, purchased from a state-license retail dispensary. Participants in the daily and occasional use groups were observed to smoke or vape cannabis flower during a 15-minute interval and were instructed to smoke *ad libitum* "the amount you commonly use for the effect you most commonly desire." Participants in the no use group were invited to relax for the equivalent amount of time. Written informed consent was obtained and the study was approved by the Colorado Multiple Institutional Review Board. More details on participant enrollment and screening criteria are previously published (Brooks-Russell et al., 2021).

Pupillary light response was collected from 101 participants. Upon inspection of the data, blinded to use group and/or time point of assessment, 18 were determined to have unusable data and were dropped from further analysis, resulting in a sample of 84 participants. Of the 84 participants used in this analysis, 29 were in the no-use group, and 30 and 25 participants in the

occasional and daily use groups, respectively. Participants ranged in age from 25.1 to 45.3 years with an average of 32 years (sd = 5.02); had an average BMI of 25.4 kg/m<sup>2</sup> (sd = 4.41); and were approximately 58% male (N = 49); see supplementary Table 1. THC levels were measured in whole blood collected 30 minutes after the inception of the 15-minute *ad-libitum* smoking interval. Time between cannabis smoking and the pupillary light response test varied from 53 – 84 minutes with a mean of 62.2 minutes (see Figure 4A). This time interval reflects the time used to complete other assessments such as a driving simulator test, as previously described (Brooks-Russell et al., 2021).

## 2.2 Pupil Response to Light Assessment:

### FIGURE\_1

Videos of pupil response during the light test were collected using SafetyScreen<sup>TM</sup> infrared video goggles developed by Ocular Data Systems, Inc (Pasadena, CA). Trajectories of pupil size during the light response test, like that shown in Figure 1, were extracted from the videos using the video segmentation pipeline described in Steinhart et al (Steinhart et al., 2023). These trajectories represent percent change in pupil size from the start of the light test, for the right eye, after cannabis consumption, in the occasional and daily use groups, and after a short rest period for the no use control group. Pupil light response trajectories were truncated to 400 frames, approximately 13.3 seconds after the start of the light test.

## 2.3 Functional Data Analysis

Functional data analysis (FDA) is a field of statistics that models functions (e.g. full trajectories/time series of pupillary light response) without extracting pre-defined specific features (Ramsay and Silverman, 2005). These functions may be either the outcome (the whole

trajectory is the outcome) or a predictor, or both. The methods are designed to handle complicated (e.g. highly non-linear) data and associations, while accounting for within person correlations over the function. In the current context, FDA methods allow for estimating and quantifying differences in how patterns of pupillary light response vary over time by cannabis use history. In our analysis, a single functional unit is the pupillary light response trajectory for a single subject. This functional unit is denoted  $y_i(t)$  or  $x_i(t)$  for participant  $i$ , depending on whether the trajectory is modelled as the outcome or predictor, respectively, with  $t$  specifying the time at which the measurement was assessed. For example, if a participant has the pupillary light response trajectory shown in Figure 1, with pupil change of -25.3% at 2 seconds after the start of the light test, then  $y_i(t) = y_1(2) = -25.3$ . Similarly, at 5 seconds after the start of the light test  $y_1(5) = -14.9$ .

Our analysis uses two distinct FDA methods to model differences in pupil response to light after cannabis use. The first method, functional logistic regression (also referred to as scalar-on-function regression in the FDA literature), is used to predict whether or not a subject recently used cannabis and treats the pupil response trajectory as a predictor variable. The second method, function-on-scalar regression, is used to model and visualize how patterns in the pupil response trajectories differ for participants with patterns of daily cannabis use, occasional cannabis use, and no use, and treats the pupillary light response trajectory as the outcome. These methods and their roles in this analysis are described in more detail below.

### 2.3.1 Predicting recent cannabis use via functional logistic regression

Here we use a functional logistic regression model to discriminate between those who recently smoked cannabis (designated “recent cannabis use,” combining individuals with daily and occasional use patterns) and those who did not (designated “no use”). Functional logistic

regression (Goldsmith et al., 2011; Ramsay and Dalzell, 1991; Reiss et al., 2017) relates binary responses  $y_i$  (e.g. recent cannabis use vs. no use) to functional covariates  $x_i(t)$  (the pupil response trajectory for the  $i^{th}$  participant). This model is analogous to logistic regression and is given by

$$\text{logit}(P(y_i = 1)) = \beta_0 + \int_t \beta_1(t)x_i(t). (1)$$

The coefficient  $\beta_1(t)$  can be thought of as a weight function, with larger absolute values indicating that pupillary light response (the functional covariate  $x_i(t)$ ) is more strongly associated with the response (recent cannabis use) at a given time during the light test. As with traditional logistic regression, the coefficient  $\beta_1(t)$  is interpreted as a log odds ratio of recent cannabis use associated with a 1% increase in pupil diameter; however, unlike traditional logistic regression, this log odds ratio is estimated at each time  $t$  during the pupil light response test. When exponentiated,  $\beta_1(t)$  is interpreted as an odds ratio at each time  $t$ . The integral effectively takes a weighted average of the covariate effect over the test time. This model can be used to predict recent cannabis use using the full pupillary light response trajectory.

We compare the functional logistic regression model to a traditional logistic regression model that uses single value summaries of the trajectory data, including (a) minimal constriction, the magnitude of peak decrease in pupil diameter as a percentage of the pre-illumination diameter; (b) rebound dilation, the area under the curve of the relative pupil diameter after the point of minimal constriction (shaded in blue in Figure 1); and (c) the slope of the rebound from the point of minimal constriction to the end of the test as calculated in (Steinhart et al., 2023). For rebound dilation, a larger magnitude of area under the curve corresponds to less rebound dilation. We compare both models in their ability to predict recent cannabis use and expect better

A pilot study of pupil response to light as a digital biomarker of recent cannabis use

prediction from the functional logistic regression model because it leverages information from the full pupillary light response trajectories. Area under the receiver operating characteristic curve (AUC) is used to compare the ability of each model to discriminate between recent cannabis use and no use, where values closer to 1 are interpreted as having a higher predictive accuracy. The statistical significance of differences between AUC curves was calculated with a Mann-Whitney U-statistic (DeLong et al., 1988).

### 2.3.2 Modeling patterns in pupil response trajectories across cannabis use groups

We use function-on-scalar regression (FoSR) to model average pupil response trajectories for participants with no cannabis use, patterns of occasional cannabis use, and patterns of daily cannabis use. FoSR is analogous to linear regression and relates functional responses  $y_i(t)$  to scalar covariates  $x_i$  (e.g. age, cannabis use group, gender). In this analysis, we chose not to use demographic characteristics in our modelling because in field cannabis testing scenarios demographic information may not be known. The FoSR model is

$$y_i(t) = \beta_0(t) + \beta_1(t)I(\text{use group} = \text{occasional}) + \beta_2(t)I(\text{use group} = \text{daily}) + \varepsilon_i(t) \quad (2)$$

Indicators of cannabis use group are denoted by  $I(\text{use group} = \text{occasional})$  and  $I(\text{use group} = \text{daily})$ , which take values of 1 for subjects in the specified category and 0 otherwise. Coefficients  $\beta_0(t)$ ,  $\beta_1(t)$ , and  $\beta_2(t)$  are akin to regression coefficients in linear regression, with the added advantage that they are defined at each time  $t$  during the pupillary light response test. The intercept  $\beta_0(t)$  is interpreted as the average trajectory of a participant in the no use control group.  $\beta_1(t)$  is the average difference at a specific time  $t$  between the occasional use and no use groups, and  $\beta_2(t)$  is the average difference between the daily use and no use groups. The error

term  $\varepsilon_i(t)$ , like in traditional linear regression, is normally distributed and independent across participants, but unlike traditional linear regression, the errors may be correlated over time  $t$ .

### *Modeling the effect of a time delay from cannabis use to testing pupillary light response*

The time from cannabis use to the pupillary light response test ranged from 53 – 84 minutes (Figure 4A). We refer to this as the time delay (TD) and include it in a second FoSR model to explore how the shape of the pupil response trajectory changes over time as cannabis effects potentially become less pronounced. Cannabis use groups were combined to form one “recent use” group, which is compared with the no use group, and the time delay (TD) from cannabis use to testing was mean centered. This model is given by

$$y_i(t) = \beta_0(t) + \beta_1(t)I(\text{recent use} = 1) + \beta_2(t)I(\text{recent use} = 1) * TD + \varepsilon_i(t), (3)$$

where  $y_i(t)$ ,  $\beta_0(t)$ , and  $\varepsilon_i(t)$  have the same interpretation as the previous FoSR model (Equation 2).  $\beta_1(t)$  is interpreted as average difference in trajectories at a specific time  $t$  comparing recent cannabis use to no use with an average time delay from cannabis use to testing, and  $\beta_2(t)$  is the average difference at a specific time  $t$  for an additional minute increase in time since smoking for the cannabis use group.

### *2.4 Analysis Software*

All analyses were conducted using R version 4.0.2 (R Core Team, 2023). The R packages mgcv (Wood, 2017, 2011, 2004) and refund (Goldsmith, Jeff et al., n.d.) were used to implement functional data models. Estimation of the FoSR regression model follows the general algorithm presented by (Leroux et al., 2018). Code and data for reproducing our analysis is publicly available on GitHub.



### 3. RESULTS:

#### 3.1 Sample

84 participants used in this analysis, there were 29 participants in the no-use group, and 30 and 25 participants in the occasional and daily use groups, respectively. Participants ranged in age from 25.1 to 45.3 years with an average of 32 years ( $sd = 5.02$ ); had an average BMI of 25.4  $kg/m^2$  ( $sd = 4.41$ ); and were approximately 58% male ( $N = 49$ ); see Table 1. THC levels were measured in whole blood collected 30 minutes after the inception of the 15-minute *ad-libitum* smoking interval. Time between cannabis smoking and the pupillary light response test varied from 53 – 84 minutes with a mean of 62.2 minutes (see Figure 4A). This time interval was caused by normal variability in the time to complete other assessments in the study or to take breaks between assessments, as described in other results from the larger study (Brooks-Russell et al., 2021; Smith et al., 2023).

#### 3.2 Predicting recent cannabis use

Figure 2A shows ROC curves that compare the ability of the functional and traditional logistic regression models to discriminate between recent cannabis use and no use. The functional logistic model, which uses the full pupillary light response trajectory, has a higher AUC value ( $AUC = 0.71$ ) than the traditional logistic model based on single value summary features ( $AUC = 0.68$ ). This indicates that the functional logistic regression model can better differentiate recent cannabis use from no use, although the difference is not statistically significant ( $p = 0.6$ ) in this data set.

FIGURE\_2

An added benefit of the functional logistic regression model is the ability to visualize the odds of cannabis use over the 10 seconds of the pupil light response test (Figure 2B). This plot shows two regions with statistically significant differences between recent cannabis use and no use. The first region between 2.03 and 3.73 seconds with a maximum difference at 2.97 seconds (OR: 2.66, 95% CI: [1.28, 5.50]) corresponds to the time period where the point of minimal constriction is typically observed and shows that individuals with less pupil constriction have higher odds of being in the cannabis use group. The second region between 5.7 and 7.3 seconds with a peak difference at 6.57 seconds (OR: 0.37, 95% CI: [0.17, 0.81]), occurs during the period of rebound dilation and shows that individuals with less pupil dilation (closer to the pupil diameter at the start of the test) have lower odds of being in the cannabis use group.

### *3.3 Visualizing patterns in pupil response trajectories across cannabis use groups*

Figure 3 shows differences between the average trajectories of pupil light response in daily, occasional, and no-use groups estimated using the function-on-scalar regression (FoSR) model in Equation (2). The solid lines in Figure 3A represent estimated mean trajectories for those who did not use cannabis (purple line), for those in the occasional use group who recently smoked (light green line), and for those in the daily use group who recently smoked (dark green line). The dashed line in Figure 3A represents the estimated mean trajectory for all those who recently smoked (daily and occasional use groups combined). The no use group had a steeper decline in pupil size, more pupil constriction, and faster rebound dilation during the light test than the occasional or daily use groups. Estimated pupil trajectories for the occasional and daily use groups were similar, with marginally less constriction in the occasional use group.

Figure 3 panels B, C, and D show estimates and 95% confidence intervals for the average difference in pupil response for participants in the occasional vs no use groups, participants in

the daily vs. no use groups, and participants in the daily vs. occasional groups. Both Figure 3B and Figure 3C show regions of significant difference, indicating that there are significant differences in the average pupillary light response trajectory comparing recent cannabis use to no use, regardless of whether a participant had a history of occasional or daily cannabis consumption. Specifically, significant differences between the occasional and no-use groups are seen between 1.77 to 3.97 seconds with a peak difference at 2.87 seconds of 4.00% (95% CI: 1.32%, 6.68%), and between the daily and no-use groups between 2.1 to 2.73 seconds with a peak difference at 2.5 seconds of 2.88% (95% CI: 0.14%, 5.62%). Notably, no significant differences were found in the pupil response trajectories between the daily and occasional use groups, indicating that tolerance effects associated with daily use did not have a significant impact on pupillary light response in our data.

### FIGURE\_3

#### *3.4 The effect of a time delay from cannabis use to testing pupil light response*

Finally, we extracted expected pupil light response trajectories at 60, 65, and 70 minutes after cannabis use in an exploratory analysis of how pupil response changes farther out from the time of smoking. The number of minutes from cannabis smoking to administration of the pupillary light response test varied across study participants due to normal variability in the timing of study procedures, and we leverage this information to model how the pupil response trajectory is expected to change as time since cannabis smoking increases. Figure 4A shows the distribution of this time delay across subjects, which ranged from 53 to 84 minutes with a mean of 62.22 minutes (sd = 5.57 minutes). Figure 4B depicts the average trajectory for no cannabis use, and at 60, 65, and 70 minutes after cannabis use. It appears that after cannabis use, the point

of minimal constriction, and the extent of rebound dilation, approaches that of the no use group as time since cannabis consumption increases.

FIGURE\_4

#### 4. DISCUSSION:

There are several potential applications of an objective and non-invasive biomarker that could distinguish recent cannabis use with reasonable accuracy, such as forensic investigations in transportation crashes or workplace incidents. Our study explored the potential for trajectories of pupil size in response to light, as measured in a standardized way with infrared video goggles, to distinguish recent use from no recent use, among a sample of participants with a range of cannabis use histories. The current analysis suggests that pupillary light response, when paired with functional data analysis methods that leverage information from the full pupil response trajectory, has the potential to discriminate between participants who recently smoked cannabis and those with no history of recent use without needing pre-smoking data on pupillary light response.

To show the utility of using functional data analysis methods in predicting recent cannabis use, we compared the predictions from a functional logistic regression and traditional logistic regression model. While both models showed some predictive ability, the functional logistic regression model had a better predictive ability as indicated by higher AUC, although the difference between the models was not statistically significant in this pilot study. This better predictive ability may stem from the information that is retained when modelling full pupil trajectories versus the information loss that occurs when aggregating information into single summary values used in the traditional logistic regression framework. Additionally, the functional logistic regression was plotted to depict where and how the patterns of recent use and

2  
3  
4 369 no use groups differed significantly from each other. This plot showed two regions that were  
5  
6 370 significantly different and corresponded to the point of minimal constriction and rebound dilation  
7  
8 371 in typical pupillary light response trajectories. In the region of the point of minimal constriction,  
9  
10 372 the model shows that less constriction is associated with higher odds of recently using cannabis,  
11  
12 373 while in the region of rebound dilation, we see that less pupil dilation is associated with lower  
13  
14 374 odds of recently using cannabis. This corresponds with previous evidence showing an effect of  
15  
16 375 recent cannabis use on pupillary light response trajectories. However, the difference in predictive  
17  
18 376 ability between the functional and traditional logistic regression were not statistically significant,  
19  
20 377 which may be due to data quality and instrumentation difficulties as discussed in the limitations  
21  
22 378 section.  
23  
24  
25  
26  
27  
28

29 Additionally, FDA methods allow interpretable visualization and statistical comparison  
30  
31  
32 380 of the average pupil responses across cannabis use groups. We found significant differences in  
33  
34 381 pupil response, after cannabis smoking or an equivalent rest period, between the occasional and  
35  
36 382 no use groups for time periods that correspond to the point of minimal constriction. This  
37  
38 383 difference remained significant when comparing the daily use and no-use controls but was not  
39  
40 384 significantly different when comparing the daily use and occasional use groups. These  
41  
42 385 differences may be due to more dynamic pupil movements in non-users compared to cannabis  
43  
44 386 users. Taken together, this provides promising evidence that the pupillary light response  
45  
46 387 trajectory may be a measure of recent cannabis use that has utility in individuals with different  
47  
48 388 cannabis use histories. We were also able to model and visualize how pupil response trajectories  
49  
50 389 change as time since cannabis smoking increases. As expected, the pupil response trajectories for  
51  
52 390 the cannabis smoking group appear to approximate the average trajectory of the no-use group as  
53  
54 391 the time since smoking increases, especially in the region of the point of minimal constriction;  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 A pilot study of pupil response to light as a digital biomarker of recent cannabis use

2  
3  
4 392 however, the slope of rebound dilation appears to remain distinct. The results were consistent  
5  
6 393 with the hypotheses of differences in pupil light response by recent cannabis use, including  
7  
8  
9 394 frequent cannabis users, and a return to an average non-user trajectory with delayed test time.  
10

11  
12 395 There are several limitations to this analysis for which more sophisticated  
13  
14  
15 396 instrumentation and future data collection will be needed. Of primary concern were data quality  
16  
17 397 issues that persisted after data processing, imputation and smoothing from the video  
18  
19  
20 398 segmentation pipeline. While most pupil light response trajectories reflected the characteristic  
21  
22 399 pattern of the pupillary light response, there were a minority that were removed because there  
23  
24 400 were no characteristic features of the light response. This led to a reduction in the sample size  
25  
26 401 from a collection of 101 participants to usable data in 84. In addition, the non-standardized inter-  
27  
28  
29 402 subject geometry (pupil to camera distance) that characterized use of infrared videography  
30  
31  
32 403 instrumentation rendered it possible to assess change in pupillary diameter only as a percentage  
33  
34 404 difference from baseline, and not in absolute size (mm). Baseline pupil diameter (in mm), which  
35  
36 405 could not be measured in the present study may be an independent predictor of the pupillary light  
37  
38  
39 406 response (Larson and Behrends, 2015; McKay and Larson, 2021). Future research could  
40  
41 407 examine the pupillary light response closer in time to smoking, and at a longer time interval  
42  
43  
44 408 following use to examine how the response changes over time. Despite these limitations, the  
45  
46 409 results are promising for future research on pupillary changes associated with recent cannabis  
47  
48  
49 410 use.  
50

51  
52 411 This analysis is the first foray into pairing functional data analysis with pupillary light  
53  
54 412 response trajectories to better understand the utility of these methods in detecting recent cannabis  
55  
56  
57 413 use. We are cautiously optimistic that these results suggest that, with further refinements,  
58  
59 414 quantitative measurement and analysis of pupillary light response trajectory may aid the  
60  
61  
62  
63  
64  
65

2  
3  
4 415 objective assessment of recent cannabis use when only post cannabis use measurements can be  
5  
6 416 obtained.  
7  
8

9  
10 417 **References:**  
11

- 12  
13 418 Arkell, T.R., Spindle, T.R., Kevin, R.C., Vandrey, R., McGregor, I.S., 2021. The failings of *per*  
14 419 *se* limits to detect cannabis-induced driving impairment: Results from a simulated driving  
15 420 study. *Traffic Injury Prevention* 22, 102–107.  
16 421 <https://doi.org/10.1080/15389588.2020.1851685>  
17 422 Biasutti, W.R., Leffers, K.S.H., Callaghan, R.C., 2020. Systematic Review of Cannabis Use and  
18 423 Risk of Occupational Injury. *Substance Use & Misuse* 55, 1733–1745.  
19 424 <https://doi.org/10.1080/10826084.2020.1759643>  
20 425 Brooks-Russell, A., Brown, T., Friedman, K., Wrobel, J., Schwarz, J., Dooley, G., Ryall, K.A.,  
21 426 Steinhart, B., Amioka, E., Milavetz, G., Sam Wang, G., Kosnett, M.J., 2021. Simulated  
22 427 driving performance among daily and occasional cannabis users. *Accident Analysis &*  
23 428 *Prevention* 160, 106326. <https://doi.org/10.1016/j.aap.2021.106326>  
24 429 Brown, B., Adams, A.J., Haegerstrom-Portnoy, G., Jones, R.T., Flom, M.C., 1977. Pupil size  
25 430 after use of marijuana and alcohol. *American Journal of Ophthalmology* 83, 350–354.  
26 431 [https://doi.org/10.1016/0002-9394\(77\)90732-2](https://doi.org/10.1016/0002-9394(77)90732-2)  
27 432 Burt, T.S., Brown, T.L., Milavetz, G., McGehee, D.V., 2021. Mechanisms of cannabis  
28 433 impairment: Implications for modeling driving performance. *Forensic Science*  
29 434 *International* 328, 110902. <https://doi.org/10.1016/j.forsciint.2021.110902>  
30 435 Campobasso, C.P., De Micco, F., Corbi, G., Keller, T., Hartung, B., Daldrup, T., Monticelli, F.,  
31 436 2020. Pupillary effects in habitual cannabis consumers quantified with pupillography.  
32 437 *Forensic Science International* 317, 110559.  
33 438 <https://doi.org/10.1016/j.forsciint.2020.110559>  
34 439 DeLong, E.R., DeLong, D.M., Clarke-Pearson, D.L., 1988. Comparing the Areas under Two or  
35 440 More Correlated Receiver Operating Characteristic Curves: A Nonparametric Approach.  
36 441 *Biometrics* 44, 837. <https://doi.org/10.2307/2531595>  
37 442 Downey, L.A., King, R., Papafotiou, K., Swann, P., Ogden, E., Boorman, M., Stough, C., 2012.  
38 443 Detecting impairment associated with cannabis with and without alcohol on the  
39 444 Standardized Field Sobriety Tests. *Psychopharmacology* 224, 581–589.  
40 445 <https://doi.org/10.1007/s00213-012-2787-9>  
41 446 Drug Evaluation and Classification (Preliminary School): Participant Manual, 2015. . National  
42 447 Highway Traffic Safety Administration.  
43 448 Fant, R., 1998. Acute and Residual Effects of Marijuana in Humans. *Pharmacology*  
44 449 *Biochemistry and Behavior* 60, 777–784. [https://doi.org/10.1016/S0091-3057\(97\)00386-](https://doi.org/10.1016/S0091-3057(97)00386-9)  
45 450 [9](https://doi.org/10.1016/S0091-3057(97)00386-9)  
46 451 Goldsmith, J., Bobb, J., Crainiceanu, C.M., Caffo, B., Reich, D., 2011. Penalized Functional  
47 452 Regression. *Journal of Computational and Graphical Statistics* 20, 830–851.  
48 453 <https://doi.org/10.1198/jcgs.2010.10007>  
49 454 Goldsmith, Jeff, Scheipl, F., Huang, L., Wrobel, J., Di, C., Gellar, J., Harezlak, J., McLean,  
50 455 M.W., Swihart, B., Xiao, L., Crainiceanu, C., Reiss, P.T., n.d. refund: Regression with  
51 456 Functional Data.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- Key Substance Use and Mental Health Indicators in the United States: Results from the 2017 National Survey on Drug Use and Health (No. HHS Publication No. SMA 18-5068, NSDUH Series H-53), 2017. . Rockville, MD.
- Larson, M.D., Behrends, M., 2015. Portable Infrared Pupillometry: A Review. *Anesthesia & Analgesia* 120, 1242–1253. <https://doi.org/10.1213/ANE.0000000000000314>
- Leroux, A., Xiao, L., Crainiceanu, C., Checkley, W., 2018. Dynamic prediction in functional concurrent regression with an application to child growth. *Statistics in Medicine* 37, 1376–1388. <https://doi.org/10.1002/sim.7582>
- Lira, M.C., Heeren, T.C., Buczek, M., Blanchette, J.G., Smart, R., Pacula, R.L., Naimi, T.S., 2021. Trends in Cannabis Involvement and Risk of Alcohol Involvement in Motor Vehicle Crash Fatalities in the United States, 2000–2018. *Am J Public Health* 111, 1976–1985. <https://doi.org/10.2105/AJPH.2021.306466>
- McKay, R.E., Larson, M.D., 2021. Detection of opioid effect with pupillometry. *Autonomic Neuroscience* 235, 102869. <https://doi.org/10.1016/j.autneu.2021.102869>
- Merzouki, A., Mesa, J.M., Louktibi, A., Kadiri, M., Urbano, G.V., 2008. Assessing changes in pupillary size in Rifian smokers of kif (*Cannabis sativa* L.) q. *Journal of Forensic and Legal Medicine*.
- Myran, D.T., Gaudreault, A., Pugliese, M., Manuel, D.G., Tanuseputro, P., 2023. Cannabis-Involved Traffic Injury Emergency Department Visits After Cannabis Legalization and Commercialization. *JAMA Netw Open* 6, e2331551. <https://doi.org/10.1001/jamanetworkopen.2023.31551>
- Ortiz-Peregrina, S., Ortiz, C., Castro-Torres, J.J., Jiménez, J.R., Anera, R.G., 2020. Effects of Smoking Cannabis on Visual Function and Driving Performance. A Driving-Simulator Based Study. *IJERPH* 17, 9033. <https://doi.org/10.3390/ijerph17239033>
- R Core Team, 2023. R: A language and Environment for Statistical Computing.
- Ramsay, J.O., Dalzell, C.J., 1991. Some Tools for Functional Data Analysis. *Journal of the Royal Statistical Society: Series B (Methodological)* 53, 539–561. <https://doi.org/10.1111/j.2517-6161.1991.tb01844.x>
- Ramsay, J.O., Silverman, B.W., 2005. *Functional Data Analysis*, 2nd ed, Springer Series in Statistics. Springer, New York, New York, USA.
- Reiss, P., T., Goldsmith, J., Shang, H.L., Ogden, R.T., 2017. Methods for scalar-on-function regression. *International Statistical Review* 85, 228–249. <https://doi.org/doi:10.1111/insr.12163>
- Richman, J.E., Golden McAndrew, K., Decker, D., Mullaney, S.C., 2004. An evaluation of pupil size standards used by police officers for detecting drug impairment. *Optometry - Journal of the American Optometric Association* 75, 175–182. [https://doi.org/10.1016/S1529-1839\(04\)70037-8](https://doi.org/10.1016/S1529-1839(04)70037-8)
- Shahidi Zandi, A., Comeau, F.J.E., Mann, R.E., Di Ciano, P., Arslan, E.P., Murphy, T., Le Foll, B., Wickens, C.M., 2021. Preliminary Eye-Tracking Data as a Nonintrusive Marker for Blood  $\Delta$ -9-Tetrahydrocannabinol Concentration and Drugged Driving. *Cannabis and Cannabinoid Research* 6, 537–547. <https://doi.org/10.1089/can.2020.0141>
- Smith, S.J., Wrobel, J., Brooks-Russell, A., Kosnett, M.J., Sammel, M.D., 2023. A Latent Variable Analysis of Psychomotor and Neurocognitive Performance After Acute Cannabis Smoking. *Cannabis* 6, 123–132. <https://doi.org/10.26828/cannabis/2023/000156>



- 2  
3  
4 502 Stark, M.M., Englehart, K., Sexton, B.F., Tunbridge, R., Jackson, P., 2003. Use of a pupillometer  
5 503 to assess change in pupillary size post-cannabis. *Journal of Clinical Forensic Medicine*  
6 504 10, 9–11. [https://doi.org/10.1016/S1353-1131\(02\)00162-1](https://doi.org/10.1016/S1353-1131(02)00162-1)  
7  
8 505 Steinhart, B., Brooks-Russell, A., Kosnett, M., Subramanian, P., Wrobel, J., 2023. A Video  
9 506 Segmentation Pipeline for Assessing changes in Pupil Response to Light After Cannabis  
10 507 Consumption. *Journal of Data Science*.  
11  
12 508 Wood, S.N., 2017. *Generalized Additive Models: An Introduction with R*, 2nd ed. Chapman and  
13 509 Hall/CRC.  
14 510 Wood, S.N., 2011. Fast Stable Restricted Maximum Likelihood and Marginal Likelihood  
15 511 Estimation of Semiparametric Generalized Linear Models. *Journal of the Royal*  
16 512 *Statistical Society Series B: Statistical Methodology* 73, 3–36.  
17 513 <https://doi.org/10.1111/j.1467-9868.2010.00749.x>  
18  
19 514 Wood, S.N., 2004. Stable and efficient multiple smoothing parameter estimation for generalized  
20 515 additive models. *Journal of the American Statistical Association* 99, 673–686.  
21 516 Wurz, G.T., DeGregorio, M.W., 2022. Indeterminacy of cannabis impairment and  $\Delta 9$ -  
22 517 tetrahydrocannabinol ( $\Delta 9$ -THC) levels in blood and breath. *Sci Rep* 12, 8323.  
23 518 <https://doi.org/10.1038/s41598-022-11481-5>  
24  
25 519 Zhang, J.C., Carnide, N., Holness, L., Cram, P., 2020. Cannabis use and work-related injuries: a  
26 520 cross-sectional analysis. *Occupational Medicine* 70, 570–577.  
27 521 <https://doi.org/10.1093/occmed/kqaa175>  
28 522  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Figure Legends:**

FIGURE 1: A typical pupillary response to light during the light reflex test, which we refer to as a *pupillary light response trajectory* throughout the paper. At the onset of illumination (time 0 on the x-axis) the pupil begins to constrict in size until the diameter reaches a minimum, called the *point of minimal constriction*, then it begins to increase in size back towards its original diameter. The area under zero on the y-axis from the point of minimal constriction to the end of the light response test is a measure of *rebound dilation*. The larger the magnitude of this area (i.e. larger shaded in Figure 1), the less rebound dilation that has occurred.

FIGURE 2: *Panel A*: Receiver Operator Characteristic curves (ROCs) for our two logistic regression (LogRegr) models. Higher accuracy in predicting recent cannabis use is indicated by a higher AUC and the ROC curve following the left and top edge of the graph. The blue line is an ROC curve for a traditional logistic regression model using single value summary features of pupil light response. The yellow line is an ROC curve for a functional logistic regression model using full trajectory of pupil light response. The functional logistic model better differentiates between recent cannabis use and no use. *Panel B*: Solid black line depicts the odds ratio (OR) of recent cannabis over the 10 seconds of the pupillary light response test. The dashed lines indicate the 95% confidence interval around the OR estimate. The red segments indicate regions where the confidence interval for the OR does not contain zero, demonstrating statistically significant differences between the recent cannabis use and no use.


FIGURE 3 A-D: Panel A shows average pupil light response trajectories plotted by cannabis use frequency. An additional dotted lined based on the average trajectory for all recent cannabis users, occasional and daily, was included to show differences between recent use and no use groups. Panels B - D show the difference in average trajectories between pairs of occasional,



A pilot study of pupil response to light as a digital biomarker of recent cannabis use


daily and no-use of cannabis. Zero on the y-axis corresponds to no difference between the average trajectory of two groups, while the region indicated by the solid red line, where the confidence interval (both dashed lines) is above or below zero on the y-axis indicates statistically significant differences between trajectories. The figure demonstrates significant regions of difference between occasional and no-use groups and daily and no-use groups, while there is no significant difference between occasional and daily cannabis use groups.


FIGURE 4: *Panel A*: Histogram depicts the distribution of the time delay from cannabis use to the pupillary light response test, in minutes. The vertical dotted red line indicates the mean of the distribution at 62.2 minutes. Interquartile range is 59 – 66 minutes. *Panel B*: Differences in the average pupil light response as the time from cannabis smoking increases from 60 minutes to 70 minutes (lighter color). The purple line shows the average pupil response for the no use group. As time since cannabis consumption increases, the point of minimal constriction approaches that of the no use group while the slope of the rebound appears to remain distinct.

559









Cannabis Use Group					
Characteristic	non-user (N = 29) <sup>1</sup>	occasional (N = 30) <sup>1</sup>	daily (N = 25) <sup>1</sup>	Total (N = 84) <sup>1</sup>	p-value <sup>2</sup>
Age (years)	32.29 (4.70)	31.15 (4.75)	32.75 (5.71)	32.02 (5.02)	0.5
Sex					0.2
Female	16 (55%)	10 (33%)	9 (36%)	35 (42%)	
Male	13 (45%)	20 (67%)	16 (64%)	49 (58%)	
Body Mass Index (kg/m <sup>2</sup> )	24.94 (4.72)	24.49 (3.96)	27.08 (4.26)	25.42 (4.41)	0.066
THC, after cannabis smoking (ng/ml)	0.0 (0.0, 0.0)	5.73 (3.73, 9.47)	17.84 (8.20, 42.42)	3.96 (0.0, 13.60)	NA
Time Interval of pupillary measurements after initiation of cannabis smoking (mins)	NA	63.93 (6.26)	60.16 (3.78)	NA	

<sup>1</sup>Mean (SD); n (%); Median (IQR)

<sup>2</sup>Kruskal-Wallis rank sum test; Pearson's Chi-squared test

SUPPLEMENTARY TABLE 1: Participant Characteristics by Cannabis Use Group



## **Author Disclosures**

### **Role of Funding Source:**

Funders had no involvement in study design, data collection or analysis, writing the report, or the decision to submit the article for publication.

### **Contributors:**

Ms. Godbole conducted data analysis, and manuscript writing. Drs. Brooks-Russell, Wrobel, Kosnett, and Subramanian participated in the study design and implementation, manuscript writing and editing. Drs. Wrobel and Leroux created the data analysis plan and supervised analysis. Dr. Leroux also contributed to the manuscript editing.

### **Conflict of Interest:**

No conflict declared.

**Contributors:**

Ms. Godbole conducted data analysis, and manuscript writing. Drs. Brooks-Russell, Wrobel, Kosnett, and Subramanian participated in the study design and implementation, manuscript writing and editing. Drs. Wrobel and Leroux created the data analysis plan and supervised analysis. Dr. Leroux also contributed to the manuscript editing.

Declarations of interest: none.